



Clinical Characteristics and Risk Factors for Mortality of COVID-19 Patients With Diabetes in Wuhan, China: A Two-Center, Retrospective Study

Qiao Shi,^{1,2} Xiaoyi Zhang,³ Fang Jiang,⁴ Xuanzhe Zhang,² Ning Hu,² Chibu Bimu,² Jiarui Feng,⁵ Su Yan,² Yongjun Guan,² Dongxue Xu,³ Guangzhen He,⁶ Chen Chen,² Xingcheng Xiong,¹ Lei Liu,¹ Hanjun Li,¹ Jing Tao,¹ Zhiyong Peng,³ and Weixing Wang²

Diabetes Care 2020;43:1382–1391 | <https://doi.org/10.2337/dc20-0598>

OBJECTIVE

Diabetes is common in COVID-19 patients and associated with unfavorable outcomes. We aimed to describe the characteristics and outcomes and to analyze the risk factors for in-hospital mortality of COVID-19 patients with diabetes.

RESEARCH DESIGN AND METHODS

This two-center retrospective study was performed at two tertiary hospitals in Wuhan, China. Confirmed COVID-19 patients with diabetes ($N = 153$) who were discharged or died from 1 January 2020 to 8 March 2020 were identified. One sex- and age-matched COVID-19 patient without diabetes was randomly selected for each patient with diabetes. Demographic, clinical, and laboratory data were abstracted. Cox proportional hazards regression analyses were performed to identify the risk factors associated with the mortality in these patients.

RESULTS

Of 1,561 COVID-19 patients, 153 (9.8%) had diabetes, with a median age of 64.0 (interquartile range 56.0–72.0) years. A higher proportion of intensive care unit admission (17.6% vs. 7.8%, $P = 0.01$) and more fatal cases (20.3% vs. 10.5%, $P = 0.017$) were identified in COVID-19 patients with diabetes than in the matched patients. Multivariable Cox regression analyses of these 306 patients showed that hypertension (hazard ratio [HR] 2.50, 95% CI 1.30–4.78), cardiovascular disease (HR 2.24, 95% CI 1.19–4.23), and chronic pulmonary disease (HR 2.51, 95% CI 1.07–5.90) were independently associated with in-hospital death. Diabetes (HR 1.58, 95% CI 0.84–2.99) was not statistically significantly associated with in-hospital death after adjustment. Among patients with diabetes, nonsurvivors were older (76.0 vs. 63.0 years), most were male (71.0% vs. 29.0%), and they were more likely to have underlying hypertension (83.9% vs. 50.0%) and cardiovascular disease (45.2% vs. 14.8%) (all P values <0.05). Age ≥ 70 years (HR 2.39, 95% CI 1.03–5.56) and hypertension (HR 3.10, 95% CI 1.14–8.44) were independent risk factors for in-hospital death of patients with diabetes.

CONCLUSIONS

COVID-19 patients with diabetes had worse outcomes compared with the sex- and age-matched patients without diabetes. Older age and comorbid hypertension independently contributed to in-hospital death of patients with diabetes.

¹Department of Pancreatic Surgery, Renmin Hospital of Wuhan University, Wuhan, China

²Department of General Surgery, Renmin Hospital of Wuhan University, Wuhan, China

³Department of Critical Care Medicine, Zhongnan Hospital of Wuhan University, Wuhan, China

⁴Department of Anesthesiology, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, China

⁵Department of Medical Management, Renmin Hospital of Wuhan University, Wuhan, China

⁶Department of Endocrinology, Zhongnan Hospital of Wuhan University, Wuhan, China

Corresponding authors: Zhiyong Peng, pengzy5@hotmail.com, and Weixing Wang, sate.lite@163.com

Received 22 March 2020 and accepted 28 April 2020

This article contains supplementary material online at <https://doi.org/10.2337/figshare.12210008>.

This article is part of a special article collection available at <https://care.diabetesjournals.org/collection/diabetes-and-COVID19>.

Q.S., Xi.Z., and F.J. contributed equally.

© 2020 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <https://www.diabetesjournals.org/content/license>.

In December 2019, pneumonia of an unknown cause was detected in Wuhan, China, which was later named coronavirus disease 2019 (COVID-19) by the World Health Organization. The virus that caused this epidemic was named severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). The outbreak of COVID-19 swept across China and other countries, arousing global concern. As of 24 April 2020, a total of 2,626,321 COVID-19 cases were confirmed worldwide, and 181,938 patients had died (1). Compared with severe acute respiratory syndrome (SARS) and Middle East respiratory syndrome (MERS), COVID-19 has a lower mortality among confirmed cases. However, elderly patients with underlying comorbidities, including diabetes, hypertension, and coronary heart disease, are at greater risk of poor outcomes (2,3).

Diabetes is one of the leading causes of morbidity, and it causes enormous health and financial burdens worldwide (4). Connections between diabetes and increased susceptibility to infections, including respiratory tract, urinary tract, and soft-tissue infections, have long been accepted (5). The available evidence demonstrates that diabetes predisposes people to developing infectious diseases, and patients with diabetes are at greater risk of infection-related mortality (6,7). Furthermore, diabetes has been associated with a poor prognosis and increased pneumonia-associated mortality (8,9). Previous studies demonstrated that diabetes is one of the major comorbidities in COVID-19 patients. Wang et al. (10) and Guan et al. (11) reported that patients with diabetes accounted for 10.1% and 7.4% of COVID-19 patients, respectively. Recent publications showed that 20–30% of nonsurviving COVID-19 patients had underlying diabetes (3,12). This evidence indicates that COVID-19 patients with diabetes might be at a higher risk of death. Thus, the clinical characteristics and risk factors for in-hospital mortality of COVID-19 patients with diabetes need to be explored.

In this study, we aimed to describe the demographic features, clinical data, treatments, and outcomes of COVID-19 patients with diabetes. We also compared the characteristics and risk factors for in-hospital death of the patients who had diabetes with those of age- and sex-matched patients without diabetes.

RESEARCH DESIGN AND METHODS

Study Design

This two-center, retrospective study was conducted at Renmin Hospital of Wuhan University and Zhongnan Hospital of Wuhan University, which are two major tertiary hospitals in Wuhan that serve as government-designated hospitals for the treatment of COVID-19. All the confirmed COVID-19 patients with diabetes who were discharged or died from 1 January 2020 to 8 March 2020 were identified. The patients with diabetes included in our study had a clear diagnosis of diabetes by their physicians on the electronic medical records. The age of those with diabetes (64.0 [interquartile range [IQR], 56.0–72.0] years) in our study was significantly different from the overall COVID-19 population (47.0 [IQR, 35.0–58.0] years) in China (11). Older age and male sex have also been demonstrated to be associated with in-hospital death of COVID-19 patients (3,12). Thus, to adjust age and sex, an age- (± 2 years) and sex-matched COVID-19 patient without diabetes was randomly selected for each patient with diabetes according to previously reported similar methods (13,14). Whenever more than one patient was available for each patient with diabetes, the match was randomly selected from those available. This study was approved by the Institutional Ethics Boards of Renmin Hospital of Wuhan University (No. WDRY2020-K060) and Zhongnan Hospital of Wuhan University (No. 2020020), Wuhan, China. The ethic committees in these two hospitals waived informed consent. Oral consent was obtained when we contacted patients or their families for information about patients' diabetes history.

Data Abstraction

Data abstracted included age, sex, exposure history, history of diabetes, other underlying comorbidities (hypertension, cardiovascular disease, cerebrovascular disease, chronic pulmonary disease, chronic kidney disease, chronic liver disease, and malignancy), onset of symptoms (fever, cough, dyspnea, myalgia, etc.), vital signs at admission (heart rate, respiratory rate, and mean artery pressure), laboratory parameters on admission (blood glucose level, white blood cell count, lymphocyte count, procalcitonin, triglyceride, etc.), random blood glucose (RBG), chest computed tomographic (CT) scans, complications

(acute respiratory distress syndrome [ARDS], acute cardiac injury, acute kidney injury [AKI], shock, and secondary infections), medications for treatment (antiviral, antibacterial agents, corticosteroids, Ig), treatment strategies (supplemental oxygen, noninvasive mechanical ventilation, invasive mechanical ventilation, continuous renal replacement therapy [CRRT], and extracorporeal membrane oxygenation [ECMO]), and date of discharge or death. Duration from the onset of symptoms to admission, diagnosis of COVID-19, and total hospital length of stay were also recorded. Data were abstracted using the electronic medical record systems in Renmin Hospital of Wuhan University and Zhongnan Hospital of Wuhan University. All data were reviewed by a team of experienced physicians. Any uncertain or missing records were addressed through communication with the involved health care providers, the patients, or their families. Also included in our study were 14 patients with diabetes and 9 patients without diabetes in a previous publication (10).

Definitions

The diagnosis of COVID-19 was performed according to the World Health Organization interim guidance (15). The method of detection of SARS-CoV-2 using throat swabs and RT-PCR was reported previously (10). Diabetes was defined according to the guidelines of American Diabetes Association (16). Pregnant women with gestational diabetes or patients with glucocorticoid-induced hyperglycemia were identified and excluded from this study. ARDS was defined according to the Berlin definition (17). Cardiac injury was reported if serum levels of myocardial injury biomarkers (e.g., ultrasensitive troponin I) were >99 th percentile of the upper reference. AKI was diagnosed according to the Kidney Disease: Improving Global Outcomes definition (18). Shock was defined according to the 2016 Third International Consensus Definition for Sepsis and Septic Shock (19). Secondary infections were diagnosed when positive cultures of pathogens were obtained from lower respiratory tract specimens or blood samples after admission (3). The calculation of average RBG was based on all the available RBG test results of the patients during this hospitalization. Average RBG = $(RBG_1 + RBG_2 + RBG_3 \dots + RBG_n)/n$.

Table 1—Characteristics, laboratory findings, complications, treatments, and outcomes of COVID-19 patients with diabetes and sex- and age-matched patients without diabetes

	Patients without diabetes (n = 153)	Patients with diabetes (n = 153)	P value	
Age, years	65.0 (56.0–72.0)	64.0 (56.0–72.0)	0.872	
Sex			1.000	
Female	78 (51.0)	78 (51.0)	–	
Male	75 (49.0)	75 (49.0)	–	
Exposure history	31 (20.3)	18 (11.8)	0.043	
Smoking	9 (5.9)	7 (4.6)	0.608	
Drinking	7 (4.6)	6 (3.9)	0.777	
Comorbidities				
Hypertension	44 (28.8)	87 (56.9)	<0.001	
Cardiovascular disease	17 (11.1)	32 (20.9)	0.019	
Cerebrovascular disease	2 (1.3)	12 (7.8)	0.006	
Chronic pulmonary disease	13 (8.5)	8 (5.2)	0.258	
Chronic kidney disease	6 (3.9)	6 (3.9)	1.000	
Chronic liver disease	4 (2.6)	5 (3.3)	1.000	
Malignancy	6 (3.9)	8 (5.2)	0.584	
Signs and symptoms				
Fever	118 (77.1)	120 (78.4)	0.783	
Cough	78 (51.0)	95 (62.1)	0.050	
Dyspnea	60 (39.2)	52 (34.0)	0.342	
Myalgia	16 (10.5)	22 (14.4)	0.298	
Headache	6 (3.9)	3 (2.0)	0.501	
Diarrhea	23 (15.0)	18 (11.8)	0.401	
Nausea or vomiting	5 (3.3)	8 (5.2)	0.395	
Anorexia	82 (53.6)	81 (52.9)	0.909	
Fatigue	79 (51.6)	95 (62.1)	0.065	
From onset symptom to, days				
Hospital admission	10.0 (7.0–15.0)	11.0 (7.0–18.0)	0.693	
Confirmation of SARS-CoV-2	8.0 (5.0–12.0)	8.0 (4.0–14.0)	0.963	
Hospital length of stay, days	15.0 (9.0–22.0)	15.0 (8.0–22.0)	0.507	
ICU admission	12 (7.8)	27 (17.6)	0.010	
Respiratory rate, rpm	20.0 (18.0–21.0)	20.0 (18.0–21.0)	0.719	
Heart rate, bpm	84.0 (76.0–92.0)	83.0 (78.0–92.0)	0.665	
Mean arterial pressure, mmHg	95.0 (87.0–103.0)	94.0 (87.0–102.0)	0.445	
CT manifestations, area of lung injury				
<25%	81/118 (68.6)	74/121 (61.2)	0.225	
25–50%	19/118 (16.1)	21/121 (17.4)	0.795	
50–75%	10/118 (8.5)	19/121 (15.7)	0.087	
>75%	6/118 (5.1)	8/121 (6.6)	0.615	
Laboratory findings on admission	<u>Normal range</u>			
White blood cell count, $\times 10^9/L$	3.5–9.5	5.9 (4.1–7.5)	5.6 (4.5–8.0)	0.514
Neutrophil count, $\times 10^9/L$	1.8–6.3	3.6 (2.5–5.4)	3.8 (2.8–6.3)	0.193
Lymphocyte count, $\times 10^9/L$	1.1–3.2	1.1 (0.8–1.5)	1.0 (0.7–1.5)	0.189
Platelet count, $\times 10^9/L$	125–350	217.0 (164.0–278.0)	193.0 (141.0–267.0)	0.052
C-reactive protein, mg/L	0–10	16.8 (5.0–62.8)	23.3 (5.0–85.2)	0.178
Prothrombin time, s	9–13	12.0 (11.4–12.7)	12.0 (11.5–13.1)	0.328
D-dimer, ng/mL	0–550	570.0 (270.0–1,540.0)	683.5 (270.0–2,344.0)	0.551
ALT, units/L	9–50	22.5 (16.0–38.3)	25.0 (16.0–40.0)	0.888
Creatinine, $\mu\text{mol/L}$	57–111	62.1 (51.8–77.5)	65.2 (52.1–84.0)	0.276
eGFR, mL/min	>90	94.6 (82.8–106.6)	93.6 (74.5–104.2)	0.163
Total cholesterol, mmol/L	<5.2	4.1 (3.4–4.9)	3.8 (3.2–4.3)	0.006
Triglyceride, mmol/L	<1.70	1.25 (1.01–1.67)	1.37 (1.06–1.73)	0.223
pH	7.35–7.45	7.41 (7.37–7.43)	7.41 (7.36–7.45)	0.549
Pao ₂ , mmHg	80–100	88.0 (69.3–113.0)	68.5 (48.5–90.8)	0.011
Spo ₂ , %	95–100	97.0 (94.0–99.0)	96.0 (90.0–98.0)	0.093
Glucose, mmol/L	3.9–6.1	5.7 (4.8–7.3)	9.4 (6.9–13.3)	<0.001
Lactate, mmol/L	0.5–1.5	2.3 (1.6–2.8)	2.0 (1.5–2.9)	0.626
Procalcitonin, ng/mL	<0.1	0.05 (0.04–0.10)	0.06 (0.05–0.24)	0.010
Ultrasensitive troponin I, ng/mL	0–0.04	0.006 (0.006–0.019)	0.007 (0.006–0.034)	0.104
CD3 ⁺ cell count, μL	723–2,737	670.0 (448.0–929.0)	581.0 (307.5–1,013.5)	0.143

Continued on p. 1385

Table 1—Continued

		Patients without diabetes (<i>n</i> = 153)	Patients with diabetes (<i>n</i> = 153)	<i>P</i> value
CD4 ⁺ cell count, /μL	404–1,612	380.0 (264.0–570.5)	365.0 (202.5–633.5)	0.517
CD8 ⁺ cell count, /μL	220–1,129	242.0 (130.0–361.5)	164.0 (98.0–303.0)	0.026
CD19 ⁺ cell count, /μL	80–616	131.0 (85.5–210.0)	143.0 (84.5–214.5)	0.610
CD16 ⁺ 56 ⁺ cell count, /μL	84–724	128.5 (69.8–189.3)	119.0 (72.0–201.0)	0.492
Complications				
ARDS		17 (11.1)	38 (24.8)	0.002
AKI		5 (3.3)	19 (12.4)	0.003
Acute cardiac injury		26 (17.0)	47 (30.7)	0.005
Shock		16 (10.5)	32 (20.9)	0.012
Secondary infection		17 (11.1)	37 (24.2)	0.003
Treatments				
Antiviral therapy		146 (95.4)	148 (96.7)	0.556
Antibiotic therapy		112 (73.2)	122 (79.7)	0.178
Glucocorticoid therapy		53 (34.6)	54 (35.3)	0.905
Ig therapy		35 (22.9)	32 (20.9)	0.678
Supplemental oxygen		92 (60.1)	91 (59.5)	0.907
Noninvasive mechanical ventilation		8 (5.2)	21 (13.7)	0.011
Invasive mechanical ventilation		4 (2.6)	11 (7.2)	0.064
CRRT		1 (0.7)	3 (1.9)	0.623
ECMO		0	1 (0.7)	1.000
Prognosis				
Death		16 (10.5)	31 (20.3)	—
Survival		137 (89.5)	122 (79.7)	—

Data are expressed as median (IQR), *n* (%), or *n/N* (%), where *N* is available total cases. Boldface *P* values are statistically significant (*P* < 0.05). ALT, alanine aminotransferase; eGFR, estimated glomerular filtration rate.

The primary outcome was the in-hospital mortality of COVID-19 patients with diabetes and risk factors for the death of patients with diabetes. The secondary outcomes were the clinical characteristics, laboratory findings, incidence of complications, and the differences of risk factors for in-hospital death in COVID-19 patients with and without diabetes.

Statistical Analysis

We made no assumptions regarding missing data. Categorical variables are described as frequencies and percentages based on the available data. Continuous variables are described as the medians and IQRs. We used the Pearson χ^2 test, the Mann-Whitney test, and the Fisher exact test for comparisons between patients with and without diabetes and between survivors and nonsurvivors, as appropriate.

To explore the risk factors associated with in-hospital death for the 306 COVID-19 patients and to assess whether diabetes was an independent risk factor for death, a multivariable Cox proportional hazards regression model was performed. Four variables, including diabetes, hypertension, cardiovascular disease, and chronic pulmonary disease, were included in the model. Age and sex were not included because the patients with and without

diabetes were matched on age and sex. We also conducted Cox regression analyses to identify risk factors for in-hospital death of patients with diabetes and matched patients without diabetes. Considering the death toll was not large in our study, and to avoid overfitting in the model, five variables, including age, sex, underlying hypertension, cardiovascular disease, and chronic pulmonary disease, were chosen for the final regression models. All the variables included in the final models were based on clinical and scientific understanding, previous findings, and the results of univariable analyses. Variables were excluded from the Cox regression models if the number of events was too small (i.e., chronic kidney disease, cerebrovascular disease).

Survival curves for patients with diabetes and matched patients without diabetes were developed using the Kaplan-Meier method with the log-rank test. The statistical analyses were conducted with SPSS (version 25.0), GraphPad Prism (version 5.0), and R (version 3.6.1) software. A two-sided *P* value < 0.05 was considered statistically significant.

RESULTS

Of 1,561 COVID-19 patients, 153 patients (42 from Zhongnan Hospital and 111

from Renmin Hospital) with diabetes were included, and the prevalence of diabetes was 9.8%. The median age of the patients with diabetes was 64.0 (IQR, 56.0–72.0) years. The patients with and without diabetes were well matched for age and sex, with men and women represented approximately equally (Table 1). More patients with diabetes reported no exposure history but had a higher prevalence of hypertension (56.9% vs. 28.8%), cardiovascular disease (20.9% vs. 11.1%), and cerebrovascular disease (7.8% vs. 1.3%) (all *P* values < 0.05). Fever and cough were the most common onset of symptoms in both groups. Patients with diabetes were more likely to require intensive care unit (ICU) admission (17.6% vs. 7.8%, *P* < 0.05). The characteristic ground-glass opacity on CT images is shown in Supplementary Fig. 1. Among laboratory findings, patients with diabetes had lower levels of cholesterol (3.8 vs. 4.1 mmol/L), Pao₂ (68.5 vs. 88.0 mmHg) and CD8⁺ cell count (164.0 vs. 242.0/μL), and higher levels of blood glucose (9.4 vs. 5.7 mmol/L) and procalcitonin (0.06 vs. 0.05 ng/mL) (all *P* values < 0.05). Patients with diabetes were more likely to have ARDS (24.8% vs. 11.1%), acute cardiac injury (30.7% vs. 17.0%), secondary

Table 2—Characteristics, laboratory findings, complications, treatments, and outcomes of survivors and nonsurvivors among COVID-19 patients with diabetes and matched patients without diabetes

	Patients without diabetes		Patients with diabetes	
	Survivors (n = 137)	Nonsurvivors (n = 16)	Survivors (n = 122)	Nonsurvivors (n = 31)
Age, years	63.0 (56.0–70.0)	72.0 (68.0–81.0)*	63.0 (56.0–69.0)	76.0 (65.0–82.0)†
Sex				
Female	68 (49.6)	10 (62.5)	69 (56.6)	9 (29.0)†
Male	69 (50.4)	6 (37.5)	53 (43.4)	22 (71.0)†
Exposure history	28 (20.4)	3 (18.8)	14 (11.5)	4 (12.9)
Smoking	9 (6.6)	0	4 (3.3)	3 (9.7)
Drinking	7 (5.1)	0	5 (4.1)	1 (3.2)
Comorbidities				
Hypertension	38 (27.7)	6 (37.5)	61 (50.0)	26 (83.9)†
Cardiovascular disease	14 (10.2)	3 (18.8)	18 (14.8)	14 (45.2)†
Cerebrovascular disease	0	2 (12.5)*	7 (5.7)	5 (16.1)
Chronic pulmonary disease	10 (7.3)	3 (18.8)	4 (3.3)	4 (12.9)
Chronic kidney disease	3 (2.2)	3 (18.8)*	4 (3.3)	2 (6.5)
Chronic liver disease	4 (2.9)	0	5 (4.1)	0
Malignancy	4 (2.9)	2 (12.5)	6 (4.9)	2 (6.5)
Signs and symptoms				
Fever	106 (77.4)	12 (75.0)	95 (77.9)	25 (80.6)
Cough	69 (50.4)	9 (56.3)	77 (63.1)	18 (58.1)
Dyspnea	49 (35.8)	11 (68.8)*	35 (28.7)	17 (54.8)†
Myalgia	15 (10.9)	1 (6.3)	18 (14.8)	4 (12.9)
Headache	6 (4.4)	0	3 (2.5)	0
Diarrhea	20 (14.6)	3 (18.8)	17 (13.9)	1 (3.2)
Nausea or vomiting	5 (3.6)	0	6 (4.9)	2 (6.5)
Anorexia	71 (51.8)	11 (68.8)	63 (51.6)	18 (58.1)
Fatigue	70 (51.1)	9 (56.3)	73 (59.8)	22 (71.0)
From onset symptom to, days				
Hospital admission	10.0 (7.0–15.0)	10.0 (7.0–14.0)	11.0 (7.0–20.0)	10.0 (5.0–15.0)
Confirmation of SARS-CoV-2	8.0 (4.0–12.0)	9.0 (6.0–12.0)	8.0 (4.0–14.0)	9.0 (3.0–14.0)
Hospital length of stay, days	16.0 (10.0–22.0)	4.0 (3.0–8.0)*	17.0 (11.0–23.0)	6.0 (2.0–11.0)†
ICU admission	2 (1.5)	10 (62.5)*	7 (5.7)	20 (64.5)†
Respiratory rate, rpm	20.0 (18.0–20.0)	21.0 (17.0–34.0)	20.0 (18.0–20.0)	21.0 (18.0–29.0)†
Heart rate, bpm	85.0 (76.0–92.0)	79.0 (70.0–97.0)	82.0 (78.0–91.0)	88.0 (82.0–108.0)†
Mean arterial pressure, mmHg	95.0 (87.0–102.0)	98.0 (87.0–115.0)	94.0 (87.0–100.0)	95.0 (86.0–103.0)
CT manifestations, area of lung injury				
<25%	81/115 (70.4)	0*	68/105 (64.8)	6/16 (37.5)†
25–50%	19/115 (16.5)	0	16/105 (15.2)	5/16 (31.3)
50–75%	10/115 (8.7)	0	18/105 (17.3)	1/16 (6.3)
>75%	3/115 (2.6)	3/3 (100.0)*	4/105 (3.8)	4/16 (25.0)†
Laboratory findings	<u>Normal range</u>			
Average RBG, mmol/L	<11.1	NA	NA	7.6 (6.2–10.4)
HbA _{1c} , %	3.6–6.0	NA	NA	7.9 (6.6–9.1)
HbA _{1c} , mmol/mol	16.0–42.0	NA	NA	63.0 (49.0–76.0)
White blood cell count, ×10 ⁹ /L	3.5–9.5	5.2 (4.0–7.1)	9.6 (6.9–13.4)*	5.4 (4.4–7.1)
Neutrophil count, ×10 ⁹ /L	1.8–6.3	3.3 (2.5–4.9)	8.0 (5.1–11.7)*	3.5 (2.6–4.9)
Lymphocyte count, ×10 ⁹ /L	1.1–3.2	1.1 (0.9–1.6)	0.5 (0.4–1.2)*	1.2 (0.8–1.7)
Platelet count, ×10 ⁹ /L	125–350	222.0 (165.0–289.0)	205.0 (136.0–226.0)	202.0 (147.0–279.5)
C-reactive protein, mg/L	0–10	10.8 (5.0–47.4)	115.2 (34.3–192.2)*	11.6 (5.0–62.5)
Prothrombin time, s	9–13	12.0 (11.3–12.5)	12.7 (12.0–14.4)*	11.9 (11.4–12.7)
D-dimer, mg/L	0–550	520.0 (250.0–1,102.3)	4,330.0 (1,510.0–17,140.0)*	495.0 (240.5–1,306.3)
ALT, units/L	9–50	22.0 (15.8–38.0)	29.0 (20.3–71.5)	25.0 (16.0–35.5)
Creatinine, μmol/L	57–111	61.0 (51.0–76.3)	73.5 (59.9–97.8)*	61.8 (49.0–73.3)
eGFR, mL/min	>90	97.4 (85.2–107.7)	73.0 (56.3–90.1)*	97.3 (84.7–106.4)
Total cholesterol, mmol/L	<5.2	4.1 (3.4–5.0)	3.9 (3.1–4.2)	3.8 (3.3–4.3)
Triglyceride, mmol/L	<1.70	1.3 (1.0–1.7)	1.4 (1.0–1.6)	1.3 (1.0–1.7)
pH	7.35–7.45	7.41 (7.39–7.43)	7.37 (7.28–7.48)	7.40 (7.37–7.46)
Pao ₂ , mmHg	80–100	91.5 (74.3–115.5)	55.5 (39.5–74.8)*	89.0 (60.0–122.0)

Continued on p. 1387

Downloaded from <http://diabetesjournals.org/care/article-pdf/43/7/1382/630127/doi200598.pdf> by guest on 24 April 2024

Table 2—Continued

		Patients without diabetes		Patients with diabetes	
		Survivors (n = 137)	Nonsurvivors (n = 16)	Survivors (n = 122)	Nonsurvivors (n = 31)
SpO ₂ , %	95–100	97.0 (95.0–99.0)	87.0 (78.3–89.8)*	97.0 (94.0–98.0)	89.0 (79.0–93.0)†
Glucose, mmol/L	<11.1	5.7 (4.8–7.0)	7.1 (5.1–12.2)	8.5 (6.6–12.1)	12.7 (8.8–18.6)†
Lactate, mmol/L	0.5–1.5	2.1 (1.3–2.7)	2.4 (2.1–6.0)*	1.8 (1.5–2.5)	2.3 (1.9–3.3)†
Procalcitonin, ng/mL	<0.1	0.05 (0.04–0.07)	0.14 (0.10–0.48)*	0.05 (0.04–0.10)	0.47 (0.15–1.40)†
Ultrasensitive troponin I, ng/mL	0–0.04	0.006 (0.006–0.010)	0.06 (0.014–0.506)*	0.006 (0.006–0.012)	0.070 (0.018–0.189)†
CD3 ⁺ cell count, /μL	723–2,737	706.0 (491.5–1,004.5)	266.5 (173.8–579.8)*	657.5 (431.0–1,035.3)	297.0 (139.0–433.0)†
CD4 ⁺ cell count, /μL	404–1,612	396.0 (293.0–599.0)	130.5 (92.0–369.8)*	442.5 (264.5–676.0)	130.0 (103.0–277.0)†
CD8 ⁺ cell count, /μL	220–1,129	268.0 (157.0–396.0)	106.5 (42.0–212.5)*	221.0 (128.3–312.0)	68.0 (52.0–156.0)†
CD19 ⁺ cell count, /μL	80–616	139.0 (91.5–221.5)	88.5 (54.8–175.0)	149.5 (111.3–237.0)	75.0 (45.0–163.0)†
CD16 ⁺ 56 ⁺ cell count, /μL	84–724	132.5 (71.8–196.3)	51.0 (24.3–124.0)*	137.5 (81.3–224.8)	100.0 (40.0–157.0)†
Complications					
ARDS		1 (0.7)	16 (100.0)*	7 (5.7)	31 (100.0)†
AKI		1 (0.7)	4 (25.0)*	3 (2.5)	16 (51.6)†
Acute cardiac injury		15 (10.9)	11 (68.8)*	17 (13.9)	30 (96.8)†
Shock		0	16 (100.0)*	2 (1.6)	30 (96.8)†
Secondary infection		19 (13.9)	9 (56.3)*	17 (13.9)	20 (64.5)†
Treatments					
Antiviral therapy		130 (94.9)	16 (100.0)	117 (95.9)	31 (100.0)
Antibiotic therapy		96 (70.1)	16 (100.0)*	91 (74.6)	31 (100.0)†
Glucocorticoid therapy		40 (29.2)	13 (81.3)*	37 (30.3)	17 (54.8)†
Ig therapy		28 (20.4)	7 (43.8)	22 (18.0)	10 (32.3)
Supplemental oxygen		76 (55.5)	16 (100.0)*	65 (53.3)	26 (83.9)†
Noninvasive mechanical ventilation		1 (0.7)	7 (43.8)*	5 (4.1)	16 (51.6)†
Invasive mechanical ventilation		0	4 (25.0)*	2 (1.6)	9 (29.0)†
CRRT		0	1 (6.3)	1 (0.8)	2 (6.5)
ECMO		0	0	1 (0.8)	0

Data are expressed as median (IQR), n (%), or n/N (%), where N is available total cases. ALT, alanine aminotransferase; eGFR, estimated glomerular filtration rate; NA, not available. * $P < 0.05$ (statistically significant) between nonsurvivors and survivors among patients without diabetes. † $P < 0.05$ (statistically significant) between nonsurvivors and survivors among patients with diabetes.

infections (24.2% vs. 11.1%), shock (20.9% vs. 10.5%), and AKI (12.4% vs. 3.3%) (all P values < 0.05). Antiviral therapy, antibacterial therapy, and supplemental oxygen were the most common treatments in both groups. Among all of the treatment strategies, only noninvasive mechanical ventilation (13.7% vs. 5.2%, $P < 0.05$) was applied more commonly in patients with diabetes. Death was more common in patients with diabetes (20.3% vs. 10.5%, $P < 0.05$).

Diabetes history regarding duration, previous glycemic condition, previous glucose control methods, and complications are reported in Supplementary Table 1. Among patients with diabetes who died compared with survivors (Table 2), they were older (76.0 vs. 63.0 years), most were male (71.0% vs. 29.0%), and were more likely to have hypertension (83.9% vs. 50.0%), cardiovascular disease (45.2% vs. 14.8%), and present with dyspnea (54.8% vs. 28.7%) (all P values < 0.05). CT analysis of the lungs in nonsurvivors revealed a higher proportion of patients with $> 75\%$ involvement and fewer with

$< 25\%$ involvement compared with survivors. Compared with survivors, nonsurvivors had higher respiratory rate (21 vs. 20 rpm) and heart rate (88 vs. 82 bpm) (all P values < 0.05). Nonsurvivors were more likely to be admitted to the ICU (64.5% vs. 5.7%) but had a significantly shorter hospital length of stay (6.0 vs. 17.0 days) (all P values < 0.05). Average RBG (13.6 vs. 7.6 mmol/L, $P < 0.05$) and numerous laboratory parameters, including lymphocyte count, D-dimer concentration, triglyceride level, and oxygen saturation (SpO₂) among others, distinguished nonsurvivors from survivors (Table 2). The daily average RBG during the first consecutive 5 days of hospitalization tracked in 13 nonsurvivors and 32 survivors showed that nonsurvivors had a significantly higher average RBG (except day 2) (Supplementary Fig. 2A). Compared with survivors, nonsurvivors were more likely to have ARDS (100.0% vs. 5.7%), acute cardiac injury (96.8% vs. 13.9%), shock (96.8% vs. 1.6%), secondary infections (64.5% vs. 13.9%), and AKI (51.6% vs. 2.5%) (all P

values < 0.05). With regard to treatments, nonsurvivors were more often treated with antibiotics (100.0% vs. 74.6%), glucocorticoids (54.8% vs. 30.3%), supplemental oxygen (83.9% vs. 53.3%), noninvasive mechanical ventilation (51.6% vs. 4.1%), and invasive mechanical ventilation (29.0% vs. 1.6%) (all P values < 0.05).

Among patients with diabetes, 30 patients with the highest average RBG and 30 patients with the lowest average RBG during hospitalization were analyzed (Supplementary Fig. 2B and C). The median average RBG levels were 16.4 (IQR, 14.9–17.6) mmol/L and 5.9 (IQR, 5.3–6.1) mmol/L, respectively. Half of the patients with the highest RBG had sustained detectable SARS-CoV-2 virus until death, while the RT-PCR tests turned negative in most patients with the lowest average RBG. In addition, we found that 16 of 30 patients with the highest average RBG died, whereas only 1 of 30 patients with the lowest average RBG died.

Among matched patients without diabetes who died compared with survivors

Table 3—Cox regression analyses of risk factors for in-hospital death of COVID-19 patients with diabetes and matched patients without diabetes

Variables	Patients without diabetes (N = 153)		Patients with diabetes (N = 153)	
	Univariable HR (95% CI)	Multivariable HR (95% CI)	Univariable HR (95% CI)	Multivariable HR (95% CI)
Demographics and clinical characteristics				
Age ≥70 years	5.28 (1.83–15.21)*	5.87 (1.88–18.33)*	4.87 (2.29–10.34)*	2.39 (1.03–5.56)*
Male sex	0.65 (0.24–1.79)	0.46 (0.16–1.32)	2.56 (1.18–5.56)*	2.10 (0.95–4.65)
Hypertension	1.58 (0.57–4.37)	1.06 (0.35–3.20)	4.48 (1.72–11.69)*	3.10 (1.14–8.44)*
Cardiovascular disease	1.83 (0.52–6.42)	1.08 (0.25–4.70)	3.79 (1.86–7.70)*	1.87 (0.88–4.00)
Chronic pulmonary disease	2.40 (0.68–8.51)	1.21 (0.28–5.15)	3.76 (1.30–10.89)*	2.77 (0.90–8.54)
Laboratory findings				
Average RBG, mmol/L	NA	—	1.23 (1.15–1.32)*	—
White blood cell count, ×10 ⁹ /L	1.05 (1.01–1.09)*	—	1.17 (1.09–1.26)*	—
Neutrophil count, ×10 ⁹ /L	1.31 (1.19–1.44)*	—	1.24 (1.15–1.34)*	—
Lymphocyte count, ×10 ⁹ /L	0.61 (0.25–1.52)	—	0.12 (0.04–0.31)*	—
Platelet count, ×10 ⁹ /L	0.993 (0.99–0.999)*	—	1.00 (0.99–1.00)	—
C-reactive protein, mg/L	1.01 (1.008–1.02)*	—	1.01 (1.00–1.02)*	—
Prothrombin time, s	1.78 (1.36–2.33)*	—	1.12 (1.03–1.22)*	—
Creatinine, μmol/L	1.02 (1.01–1.03)*	—	1.01 (1.00–1.01)*	—
eGFR, mL/min	0.96 (0.94–0.98)*	—	0.97 (0.96–0.98)*	—
Total cholesterol, mmol/L	0.57 (0.32–0.10)*	—	0.94 (0.59–1.48)	—
Triglyceride, mmol/L	0.85 (0.42–1.70)	—	1.73 (1.22–2.45)*	—
Pao ₂ , mmHg	0.97 (0.95–0.99)*	—	0.96 (0.94–0.99)*	—
SpO ₂ , %	0.95 (0.93–0.98)*	—	0.94 (0.92–0.97)*	—
Glucose, mmol/L	1.13 (1.02–1.25)*	—	1.12 (1.05–1.19)*	—
Lactate, mmol/L	1.32 (1.09–1.61)*	—	1.39 (1.13–1.70)*	—
Procalcitonin, ng/mL	2.13 (1.43–3.17)*	—	1.18 (1.04–1.35)*	—
CD3 ⁺ cell count, /μL	0.995 (0.992–0.998)*	—	0.997 (0.995–0.999)*	—
CD4 ⁺ cell count, /μL	0.994 (0.990–0.998)*	—	0.994 (0.991–0.998)*	—
CD8 ⁺ cell count, /μL	0.992 (0.986–0.998)*	—	0.994 (0.989–0.999)*	—
CD19 ⁺ cell count, /μL	1.000 (1.000–1.001)	—	0.991 (0.984–0.998)*	—
CD16 ⁺ 56 ⁺ count, /μL	0.987 (0.977–0.998)*	—	0.998 (0.994–1.002)	—

eGFR, estimated glomerular filtration rate; NA, not available. **P* < 0.05 (statistically significant).

(Table 2), they were older (72.0 vs. 63.0 years), more likely to have cerebrovascular disease (12.5% vs. 0), chronic kidney disease (18.8% vs. 2.2%), and present with dyspnea (68.8% vs. 35.8%) (all *P* values < 0.05). Nonsurvivors were more likely to be admitted to the ICU (62.5% vs. 1.5%, *P* < 0.05). Laboratory findings, treatments, and complications are reported in Table 2.

Among the included 306 patients, multivariable analyses (Supplementary Table 2) showed that hypertension (HR 2.50, 95% CI 1.30–4.78), cardiovascular disease (HR 2.24, 95% CI 1.19–4.23), and chronic pulmonary disease (HR 2.51, 95% CI 1.07–5.90) were independent risk factors for in-hospital death. Diabetes (HR 1.58, 95% CI 0.84–2.99) was not independently associated with death after adjusting for covariables. In univariable analyses for patients with diabetes (Table 3), 23 variables, including age, male sex, hypertension, chronic pulmonary disease, cardiovascular disease, higher average RBG, and decreased

lymphocyte count, among others, were related to death. In multivariable analyses, age ≥70 years (HR 2.39, 95% CI 1.03–5.56) and hypertension (HR 3.10, 95% CI 1.14–8.44) were independent risk factors for in-hospital death of patients with diabetes. For matched patients without diabetes (Table 3), Cox regression analyses indicated that age ≥70 years (HR 5.87, 95% CI 1.88–18.33) was independently associated with death. The survival curves of COVID-19 patients with diabetes and matched patients without diabetes are shown in Fig. 1.

CONCLUSIONS

This study analyzed the characteristics of COVID-19 patients with diabetes and sex- and age-matched patients without diabetes and identified the risk factors associated with in-hospital death of these patients. Among patients hospitalized with COVID-19, the prevalence of diabetes was 9.8%. Patients with diabetes had more underlying comorbidities, were more likely to suffer complications, had a

higher proportion of ICU admissions, and more deaths compared with sex- and age-matched patients without diabetes. However, underlying hypertension, cardiovascular disease, and chronic pulmonary disease, rather than diabetes, were independently associated with in-hospital death of COVID-19 patients. Among patients with diabetes, age ≥70 years and underlying hypertension were independent risk factors for in-hospital death.

The most common symptoms of SARS-CoV-2 infection in patients with diabetes included fever, cough, and dyspnea, which were consistent with recent publications (20,21). A previous study showed that diabetes tripled the risk of hospitalization after pandemic influenza A H1N1 and significantly increased the risks of ICU admission and mortality (14,22). In our study, patients with diabetes had a significantly higher mortality and more severe disease, as verified by the higher proportion of ICU cases and the higher incidences of ARDS and multiple organ dysfunction syndrome as well as




Figure 1—Survival curves of COVID-19 patients with diabetes and matched patients without diabetes. The gray and pink areas represent 95% CIs.

secondary infections even after sex and age adjustment.

The worse outcomes of COVID-19 patients with diabetes could be associated with underlying comorbidities. A nationwide analysis in China showed that the most prevalent comorbidity in COVID-19 patients was hypertension (16.9%), followed by diabetes (8.2%), and that the underlying diseases were associated with adverse outcomes of COVID-19 patients (23). In our study, hypertension and cardiovascular disease were more common in patients with diabetes. Although diabetes itself was not independently associated with death of COVID-19 patients in our multivariable analyses, diabetes and other comorbidities, which included cardiovascular disease and hypertension, were often closely related, and the effect of these factors could not be considered separately. Diabetes and hypertension often coexist and may act synergistically to promote adverse clinical events (24,25). The persistent hyperglycemic condition and metabolism changes in diabetes together with coexisting hypertension lead

to microvascular and macrovascular changes and form a vicious cycle that further contributes to cardiovascular events (26). Recent study also suggested that hypertension was associated with increased risk of severe and fatal COVID-19, and ACE inhibitors reduced the mortality in COVID-19 patients with hypertension (27,28). This evidence further supported our finding that hypertension was independently associated with death among patients with diabetes. Thus, patients with diabetes with underlying comorbidities, especially hypertension, should attract more attention. In sex- and age-matched patients without diabetes, Cox regression analyses did not reveal underlying comorbidities as risk factors associated with death. This might be explained by the absence of diabetes and the relatively lower prevalence of other comorbidities in the matched population.

A recent publication demonstrated that old age was an independent predictor of mortality in COVID-19 patients (3). The age-dependent decreases in cellular and humoral immune function

in elderly patients have been reported before, especially with regard to adaptive immune function (29). In our study, nonsurvivors among patients with diabetes were older compared with survivors and had obvious lymphopenia as demonstrated by significantly lower numbers of T and B cells. Furthermore, the high risks of elderly patients with diabetes could be due to their poor overall health condition and greater number of comorbidities.

Besides the above-mentioned independent risk factors, other potential risk factors related to death were also identified by our univariable analyses. Recent studies revealed lymphopenia as an important characteristic of SARS-CoV-2 infection, especially in critically ill and deceased patients (2,12). Besides, infection and destruction of lymphocytes by the SARS-CoV-2 virus have been proven (30). Thus, the destruction of lymphocytes by the SARS-CoV-2 virus might be speculated but needs to be further investigated. Lymphopenia was also noticed in our study, especially in patients with diabetes. This might be explained by the previous findings that diabetes and hyperglycemia could impair the innate and adaptive immunity (31,32).

During hospitalization, underlying diabetes, illness severity, and medical treatments could contribute to the hyperglycemic condition. Hyperglycemia has been proven to be associated with increased risks of in-hospital complications and in-hospital death (33,34). In our study, we also noticed higher blood glucose levels during hospitalization in patients with diabetes who did not survive than in survivors. Thus, frequent monitoring of blood glucose and the use of oral glucose-lowering medication or insulin would be important routine procedures for patients with diabetes.

Besides the findings in our study, some other factors might also contribute to the mortality of COVID-19 patients, among which obesity would be a potential candidate. A recent study in the New York area showed that 41.7% of the patients were obese, and an increasing number of reports have linked obesity to more severe COVID-19 illness (35–37).

Despite the importance of the aforementioned findings, the current study, however, has some limitations. Firstly, data collection relied on electronic

medical records. CT images of some patients transferred from other hospitals were not available in the electronic medical record systems of these two hospitals. Some important indicators were not tested in all patients. Thus, the missing data might lead to bias. Secondly, these two hospitals are designated for treatment of patients with relatively severe infection, which to some extent might lead to a higher mortality. Thirdly, the sample size and relatively small number of deaths might influence the interpretation of our findings. Fourthly, the possibility of obesity as a contributor to death in COVID-19 patients was not investigated due to the lack of information on BMI. Fifthly, patients in this study were selected on the basis of diabetes status and death or discharge. Some patients who remained alive in these two hospitals at the time of the analysis were not included in this study. Finally, because a matched design was used in our study, the roles of sex and age in the death of patients with diabetes versus those without could not be examined and should be addressed further.

In summary, the findings of our study suggested that COVID-19 patients with diabetes had worse outcomes compared with the sex- and age-matched patients without diabetes. Diabetes was not independently associated with in-hospital death, while hypertension, cardiovascular disease, and chronic pulmonary disease played more important roles in contributing to the mortality of COVID-19 patients. In-hospital death among COVID-19 patients with diabetes was associated with hypertension and advanced age, whereas only older age was independently associated with death among matched patients without diabetes. The need for early monitoring and supportive care should be addressed in these patients at high risks.

Acknowledgments. The authors thank all patients and health providers involved in this study and thank Chuanhua Yu and Lu Ma (Wuhan University School of Health Sciences) for help in statistical analysis.

Funding. This study was funded by the National Natural Science Foundation of China (No. 81800574, No. 81870442).

The funding sources for this study had no role in the study design, data collection, analyses, or interpretation, or writing of the manuscript.

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

Author Contributions. Q.S., Xi.Z., and F.J. drafted the manuscript. Q.S., Xi.Z., Xu.Z., and N.H.

researched the literature. Xi.Z., F.J., and G.H. contributed to the statistical analysis. Xu.Z., N.H., C.B., J.F., S.Y., Y.G., and D.X. collected the epidemiological and clinical data. C.C., X.X., L.L., H.L., and J.T. contributed to editing the manuscript, radiological analysis, and figure construction. Z.P. and W.W. conceived and supervised the study. All authors reviewed and approved the final version of the manuscript. Z.P. and W.W. are the guarantors of this work and, as such, had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

References

- World Health Organization. Coronavirus disease 2019 (COVID-19): situation report, 95 [Internet], 2020. Available from https://www.who.int/docs/default-source/coronavirus/situation-reports/20200424-sitrep-95-covid-19.pdf?sfvrsn=e8065831_4. Accessed 24 April 2020
- Yang X, Yu Y, Xu J, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study [published correction appears in *Lancet Respir Med* 2020;8:e26]. *Lancet Respir Med*. 2020;8:475–481
- Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395:1054–1062
- Pearson-Stuttard J, Blundell S, Harris T, Cook DG, Critchley J. Diabetes and infection: assessing the association with glycaemic control in population-based studies. *Lancet Diabetes Endocrinol* 2016;4:148–158
- Joshi N, Caputo GM, Weitekamp MR, Karchmer AW. Infections in patients with diabetes mellitus. *N Engl J Med* 1999;341:1906–1912
- Shah BR, Hux JE. Quantifying the risk of infectious diseases for people with diabetes. *Diabetes Care* 2003;26:510–513
- Bertoni AG, Saydah S, Brancati FL. Diabetes and the risk of infection-related mortality in the U.S. *Diabetes Care* 2001;24:1044–1049
- Kornum JB, Thomsen RW, Riis A, Lervang HH, Schønheyder HC, Sørensen HT. Type 2 diabetes and pneumonia outcomes: a population-based cohort study. *Diabetes Care* 2007;30:2251–2257
- Falguera M, Pifarre R, Martin A, Sheikh A, Moreno A. Etiology and outcome of community-acquired pneumonia in patients with diabetes mellitus. *Chest* 2005;128:3233–3239
- Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA*. 2020;323:1061–1069
- Guan WJ, Ni ZY, Hu Y, et al. Clinical characteristics of 2019 novel coronavirus infection in China. *N Engl J Med*. 2020;382:1708–1720
- Chen T, Wu D, Chen H, et al. Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. *BMJ* 2020;368:m1091
- Lovshin JA, Bjornstad P, Lovblom LE, et al. Atherosclerosis and microvascular complications: results from the Canadian study of longevity in type 1 diabetes. *Diabetes Care* 2018;41:2570–2578
- Cortes Garcia M, Sierra Moros MJ, Santa-Olalla Peralta P, Hernandez-Barrera V, Jimenez-Garcia R, Pachon I. Clinical characteristics and outcomes of diabetic patients who were hospitalized with 2009 pandemic influenza A H1N1 infection. *J Infect* 2012;64:218–224
- World Health Organization. Clinical management of severe acute respiratory infection when novel coronavirus (2019-nCoV) infection is suspected: interim guidance [Internet], 2020. Available from <https://apps.who.int/iris/bitstream/handle/10665/330893/WHO-nCoV-Clinical-2020.3-eng.pdf?sequence=1&isAllowed=y>. Accessed 28 January 2020
- American Diabetes Association. 2. Classification and diagnosis of diabetes: *Standards of Medical Care in Diabetes—2019*. *Diabetes Care* 2019;42(Suppl. 1):S13–S28
- Ranieri VM, Rubenfeld GD, Thompson BT, et al.; ARDS Definition Task Force. Acute respiratory distress syndrome: the Berlin Definition. *JAMA* 2012;307:2526–2533
- Kellum JA, Lameire N; KDIGO AKI Guideline Work Group. Diagnosis, evaluation, and management of acute kidney injury: a KDIGO summary (part 1). *Crit Care* 2013;17:204
- Rhodes A, Evans LE, Alhazzani W, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock: 2016. *Crit Care Med* 2017;45:486–552
- Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395:497–506
- Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020;395:507–513
- Allard R, Leclerc P, Tremblay C, Tannenbaum TN. Diabetes and the severity of pandemic influenza A (H1N1) infection. *Diabetes Care* 2010;33:1491–1493
- Guan W, Liang W, Zhao Y, et al. Comorbidity and its impact on 1590 patients with Covid-19 in China: a nationwide analysis. *Eur Respir J*. 26 March 2020 [Epub ahead of print]. DOI: 10.1183/13993003.00547-2020
- Adler AI, Stratton IM, Neil HA, et al. Association of systolic blood pressure with macrovascular and microvascular complications of type 2 diabetes (UKPDS 36): prospective observational study. *BMJ* 2000;321:412–419
- Chen G, McAlister FA, Walker RL, Hemmelgarn BR, Campbell NRC. Cardiovascular outcomes in Framingham participants with diabetes: the importance of blood pressure. *Hypertension* 2011;57:891–897
- Climie RE, van Sloten TT, Bruno RM, et al. Macrovasculature and microvasculature at the crossroads between type 2 diabetes mellitus and hypertension. *Hypertension* 2019;73:1138–1149
- Lippi G, Wong J, Henry BM. Hypertension and its severity or mortality in Coronavirus Disease 2019 (COVID-19): a pooled analysis. *Pol Arch Intern Med* 2020;130:304–309
- Meng J, Xiao G, Zhang J, et al. Renin-angiotensin system inhibitors improve the clinical outcomes of COVID-19 patients with hypertension. *Emerg Microbes Infect* 2020;9:757–760

29. Opal SM, Girard TD, Ely EW. The immunopathogenesis of sepsis in elderly patients. *Clin Infect Dis* 2005;41(Suppl. 7):S504–S512
30. Gu J, Gong E, Zhang B, et al. Multiple organ infection and the pathogenesis of SARS. *J Exp Med* 2005;202:415–424
31. Muller LM, Gorter KJ, Hak E, et al. Increased risk of common infections in patients with type 1 and type 2 diabetes mellitus. *Clin Infect Dis* 2005;41:281–288
32. Pozzilli P, Leslie RD. Infections and diabetes: mechanisms and prospects for prevention. *Diabet Med* 1994;11:935–941
33. McAlister FA, Majumdar SR, Blitz S, Rowe BH, Romney J, Marrie TJ. The relation between hyperglycemia and outcomes in 2,471 patients admitted to the hospital with community-acquired pneumonia. *Diabetes Care* 2005;28:810–815
34. Umpierrez GE, Isaacs SD, Bazargan N, You X, Thaler LM, Kitabchi AE. Hyperglycemia: an independent marker of in-hospital mortality in patients with undiagnosed diabetes. *J Clin Endocrinol Metab* 2002;87:978–982
35. Richardson S, Hirsch JS, Narasimhan M, et al. Presenting characteristics, comorbidities, and outcomes among 5700 patients hospitalized with COVID-19 in the New York city area. *JAMA*. 22 April 2020 [Epub ahead of print]. DOI: 10.1001/jama.2020.6775
36. Lighter J, Phillips M, Hochman S, et al. Obesity in patients younger than 60 years is a risk factor for Covid-19 hospital admission. *Clin Infect Dis*. 9 April 2020 [Epub ahead of print]. DOI: 10.1093/cid/ciaa415
37. Zheng KI, Gao F, Wang XB, et al. Obesity as a risk factor for greater severity of COVID-19 in patients with metabolic associated fatty liver disease. *Metabolism*. 19 April 2020 [Epub ahead of print]. DOI: 10.1016/j.metabol.2020.154244