



# High-Normal Protein Intake Is Not Associated With Faster Renal Function Deterioration in Patients With Type 2 Diabetes: A Prospective Analysis in the DIALECT Cohort

Diabetes Care 2022;45:35–41 | <https://doi.org/10.2337/dc21-1211>

Milou M. Oosterwijk,<sup>1</sup> Dion Groothof,<sup>2</sup> Gerjan Navis,<sup>2</sup> Stephan J.L. Bakker,<sup>2</sup> and Gozewijn D. Laverman<sup>1,3</sup>

## OBJECTIVE

To study the prospective association between dietary protein intake and renal function deterioration in patients with type 2 diabetes (T2D).

## RESEARCH DESIGN AND METHODS

Prospective analyses were performed in data of 382 patients of the Diabetes and Lifestyle Cohort Twente (DIALECT) study. Dietary protein intake was determined by the Maroni equation from 24-h urinary urea excretion. Renal function deterioration was defined as need for renal replacement therapy or a persistent increase of  $\geq 50\%$  in serum creatinine. Cox proportional hazards models were used to calculate hazard ratios (HRs) for the association between dietary protein intake and renal function deterioration. Threshold levels represent the dietary protein intake at which there was a significantly increased and reduced hazard of renal function deterioration.

## RESULTS

Renal function deterioration occurred in 53 patients (14%), with a median follow-up duration of 6 (interquartile range 5–9) years. Mean dietary protein intake was  $91 \pm 27$  g/day ( $1.22 \pm 0.33$  g/kg ideal body weight/day). Dietary protein intake was inversely associated with renal function deterioration (HR 0.62 [95% CI 0.44–0.90]). Patients with an intake  $< 92$  g/day had an increased hazard for renal function deterioration (HR 1.44 [95% CI 1.00–2.06]), while patients with an intake  $> 163$  g/day had a decreased hazard for renal function deterioration (HR 0.42 [95% CI 0.18–1.00]). Regarding dietary protein intake per kilogram body weight, patients with an intake  $< 1.08$  g/kg/day had an increased hazard for renal function deterioration (HR 1.63 [95% CI 1.00–2.65]).

## CONCLUSIONS

In patients with T2D, unrestricted dietary protein intake was not associated with an increased hazard of renal function deterioration. Therefore, substituting carbohydrates with dietary protein is not contraindicated as a part of T2D management, although it may have a positive effect on body weight while minimizing loss of muscle mass.

<sup>1</sup>Department of Internal Medicine/Nephrology, Ziekenhuis Groep Twente, Almelo, the Netherlands

<sup>2</sup>Department of Internal Medicine, Division of Nephrology, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands

<sup>3</sup>Biomedical Signals and Systems, University of Twente, Enschede, the Netherlands

Corresponding author: Milou M. Oosterwijk, [mi.oosterwijk@zgt.nl](mailto:mi.oosterwijk@zgt.nl)

Received 9 June 2021 and accepted 27 September 2021

© 2021 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <https://www.diabetesjournals.org/journals/pages/license>.

Because a high dietary protein intake has traditionally been considered as a factor that could enhance progression of chronic kidney disease (CKD) (1), dietary guidelines advise to limit dietary protein intake or even to maintain a dietary protein restriction in advanced CKD (2). In type 2 diabetes (T2D), renal function impairment is a prominent complication that occurs in 20–40% of patients (3,4).

Regarding diet in T2D, there is a strong tendency to focus on carbohydrate content (5). A low-carbohydrate diet is recommended in diabetes guidelines to manage glycemic control and to prevent complications (6,7). However, reducing carbohydrates will usually be accompanied by a shift toward increased dietary protein intake, which raises concerns about the effects of such a diet on kidney health (8). The current Kidney Disease: Improving Global Outcomes (KDIGO) guidelines for patients with T2D and CKD (estimated glomerular filtration rate [eGFR] <60 mL/min/1.73 m<sup>2</sup>) not treated with dialysis suggest to maintain a protein intake that does not exceed 0.8 g/kg/day (9). It remains unsettled, however, whether a higher protein intake would jeopardize renal function in the long-term in T2D with or without CKD. In this study, we investigated the prospective association between dietary protein intake and renal function deterioration in patients with T2D treated in secondary care.

## RESEARCH DESIGN AND METHODS

### Study Design

We performed prospective analyses using data of the Diabetes and Lifestyle Cohort Twente (DIALECT) study (10). DIALECT is an observational prospective cohort study performed in Ziekenhuis Groep Twente Hospital (Almelo and Hengelo, the Netherlands) that investigates the effect of lifestyle, including dietary habits, and medication use on outcomes in patients with T2D. Patients in DIALECT were included between September 2009 and January 2016. The study was performed according to the guidelines of good clinical practice and the Declaration of Helsinki. Written informed consent was obtained from all subjects before participation. The study has been approved by the local institutional review boards (METC registration

numbers NL57219.044.16 and 1009.68020) and is registered in The Netherlands Trial Register (NTR trial code 5855).

### Patients

The study population consists of patients with T2D aged  $\geq 18$  years treated in the outpatient clinic of the Ziekenhuis Groep Twente Hospital. Patients requiring renal replacement therapy or patients with insufficient knowledge of the Dutch language were excluded from participation. From the initial 433 patients with T2D with or without CKD included, we excluded those with missing follow-up data on renal function deterioration ( $n = 2$ ) and with missing dietary protein intake data ( $n = 49$ ), leaving 382 patients for analysis.

### Dietary Assessment

Dietary protein intake was determined at baseline from urea excretion in 24-h urine by the Maroni equation:  $6.25 * ([0.18 * \text{urinary urea excretion (mmol/24 h)}] + [0.031 * \text{body weight (kg)}]) + \text{urinary protein excretion (g/24 h)}$  (11,12). Ideal body weight was used to estimate dietary protein intake in g/kg/day on the basis of a BMI of 25 kg/m<sup>2</sup> (13). DIALECT is a strictly observational study, and participants did not receive dietary counseling other than their usual care.

### Outcome Measurement

Renal function deterioration was defined as need for renal replacement therapy (i.e., kidney transplantation, initiation of peritoneal dialysis or hemodialysis) or a persistent increase of  $\geq 50\%$  in serum creatinine from the baseline visit for at least 3 months (14). Serum creatinine was routinely measured every 3 months. In case of nephrectomy, the surgery date was used as the censor date. Until time to surgery, patients were not at risk for a chronic renal function deterioration.

### Covariates

During the study visit at which baseline assessments were performed, we collected information relevant to medical condition and pharmacological treatment. Weight (kg), height (cm), and waist and hip circumference (cm) were measured and BMI calculated (kg/m<sup>2</sup>). Body surface

area (BSA) was determined with the DuBois equation:  $\text{weight (kg)}^{0.425} * \text{height (m)}^{0.725} * 0.007184$  (15). Information about lifestyle exposures (e.g., smoking, physical activity) was collected by a self-administered questionnaire. Physical activity was assessed by the previously validated Short Questionnaire to Assess Health Enhancing Physical Activity questionnaire (16). An activity score was calculated on the basis of minutes of activity per day multiplied by an intensity factor. From these data, we scored which patients met the Dutch healthy exercise norm of 30-min moderate intensity activity a day for at least 5 days a week (17).

Patients were asked to collect 24-h urine to obtain the urinary excretion of urea and sodium as objective measure of protein and sodium intake, respectively, by multiplying these concentrations with the volume of the 24-hour urine collection. Prevalent proteinuria was defined as  $\geq 150$  mg/24 h. For proper collection of the 24-h urine sample, patients were instructed to dispose of the first morning void and thereafter collect all urine in the provided canister, including the first morning void of the next day. Patients were instructed to store the canister in a dark, cool place, preferably in a refrigerator. Nonfasting venous blood was used for routine laboratory tests. Renal function at baseline was estimated with the serum creatinine-based Chronic Kidney Disease Epidemiology Collaboration equation from 2009 (18). Blood pressure was measured in a supine position by an automated device (Dinamap; GE Medical Systems, Milwaukee, WI) for 15 minutes with a 1-minute interval. Mean systolic and diastolic blood pressure of the last three measurements was used for further analysis.

### Procedures

At baseline, laboratory tests and 24-h urine collection were used to identify the exposure variable and potential confounders. Urinary urea excretion was used to determine dietary protein intake, and urinary protein excretion was used to assess prevalent proteinuria ( $\geq 150$  mg/24 h). Serum creatinine was assessed to estimate renal function because low eGFR at baseline is an independent risk factor for renal function deterioration among patients with T2D (19). During the follow-up period,

routine laboratory tests were used to identify a persistent increase of  $\geq 50\%$  in serum creatinine from baseline visit for at least 3 months. The need for renal replacement therapy was derived from the electronic patient file.

### Statistical Analysis

All statistical analyses were performed using R 4.0.3 software (R Foundation for Statistical Computing, Vienna, Austria). Normally distributed data are presented as mean (SD), skewed variables are presented as median (interquartile range [IQR]), and categorical variables are presented as number (percent). A two-tailed  $P < 0.05$  was considered to indicate statistical significance.

Univariable and multivariable Cox proportional hazards models were used to calculate hazard ratios (HRs) for the association between dietary protein intake and renal function deterioration, expressed per incremental SD. Survival time was calculated as the difference in days between the baseline visit and date of renal function deterioration or censoring date, defined as the date of the last available routine laboratory test. Competing risk analysis was performed in which patients were censored in case of all-cause mortality. The proportional hazards assumption was verified visually with plots of the scaled Schoenfeld residuals and was not violated in any of the models. Potential nonlinear effects of dietary protein intake on the baseline hazard of renal function deterioration were modeled using natural cubic splines with 2 df, where boundary knots were set to the 5th and 95th percentiles of dietary protein intake. Potential modification of the effect of dietary protein intake on the baseline hazard for renal function deterioration by sex was explored by introduction of a product term into the model. Multivariable Wald tests were used to select an appropriate structure of the linear predictor. Potential confounders were selected on the basis of relevant differences in characteristics in the baseline table, biological plausibility, and previous literature.

The multivariable model was adjusted for age, sex, baseline eGFR, and prevalent proteinuria at baseline ( $\geq 150$  mg/24 h). Results were graphically depicted to facilitate their interpretation. HRs for different levels of dietary protein intake

were established by comparing the hazard of renal function deterioration for a hypothetical male patient of median age (65 years) with median eGFR (84 mL/min/1.73 m<sup>2</sup>) and prevalent proteinuria ( $\geq 150$  mg/24 h) with the hazard of renal function deterioration for a hypothetical patient who had the mean value for every covariate (including sex and prevalent proteinuria). Threshold levels represent the dietary protein intakes at which there was a significantly increased and reduced hazard of renal function deterioration on the basis of the 95% CI. In addition, we computed the expected cumulative incidence of renal function deterioration for two hypothetical patients, with dietary protein intakes below the value associated with an increased hazard of renal function deterioration (75 g/day) and above the value associated with a reduced hazard of renal function deterioration (180 g/day). Sensitivity analyses were performed in which BSA was added to the multivariable model, with blood pressure and use of antihypertensive agents added to the multivariable model, with use of urinary urea excretion as exposure instead of dietary protein intake, and with cystatin C–based eGFR instead of serum creatinine–based eGFR.

## RESULTS

### Baseline Characteristics

Mean  $\pm$  SD age of the 382 participants (59% male) was 63  $\pm$  9 years. Mean  $\pm$  BMI, eGFR, and dietary protein intake were 32.8  $\pm$  5.8 kg/m<sup>2</sup>, 78  $\pm$  24 mL/min/1.73 m<sup>2</sup>, and 91  $\pm$  27 g/day (i.e., 1.22  $\pm$  0.33 g/kg/day), respectively (Table 1). Median (IQR) frequency rate of serum creatinine assessment was 3 (2–5) measurements per year. After a median (IQR) follow-up duration of 6 (5–9) years, 53 patients experienced renal function deterioration. Age (HR 1.39 [95% CI 1.02–1.89]), prevalent macrovascular complications (HR 2.07 [95% CI 1.20–3.55]), proteinuria (HR 1.98 [95% CI 1.09–3.59]), and use of any antihypertensive treatment (HR 2.41 [95% CI 1.09–5.34]) were associated with an increased hazard of renal function deterioration. In contrast, diastolic blood pressure (HR 0.72 [95% CI 0.55–0.97]); eGFR (HR 0.70 [95% CI 0.54–0.92]); and dietary protein intake, when expressed in g/day (HR 0.68 [95% CI 0.51–0.92]) and in g/kg/day (HR 0.71

[95% CI 0.53–0.96]), were associated with a reduced hazard of renal function deterioration.

### Dietary Protein Intake and Renal Function Deterioration


There was no indication for a nonlinear association between dietary protein intake and renal function deterioration ( $P_{\text{nonlinear}} = 0.26$ ). Adjusted for multiple potential confounders, dietary protein intake remained inversely associated with renal function deterioration (HR 0.62 [95% CI 0.44–0.90]) (Fig. 1A). The effect of dietary protein intake on renal function deterioration did not differ between males and females ( $P_{\text{interaction}} = 0.27$ ). Patients with an intake  $< 92$  g/day had an increased hazard for renal function deterioration (HR 1.44 [95% CI 1.00–2.06]), while patients with an intake  $> 163$  g/day had a decreased hazard of renal function deterioration (HR 0.42 [95% CI 0.18–1.00]). To facilitate clinical interpretation of the magnitude of the difference in hazards associated with low-versus high-protein intake, the expected cumulative incidence of renal function deterioration was graphically depicted for two hypothetical patients, i.e., one with a protein intake of 75 g/day and one with a protein intake of 180 g/day (Fig. 1B). After 4 years of follow-up, a protein intake of 75 g/day was associated with a higher expected cumulative incidence of renal function deterioration than 180 g/day. In a subgroup analysis of patients with CKD ( $n = 85$ , 22%), the point estimate for the association between protein intake and renal function deterioration was of similar magnitude, albeit not statistically significant (HR 0.59 [95% CI 0.32–1.08]).

### Protein Intake per kg Body Weight and Renal Function Deterioration

Dietary protein intake in g/kg/day was inversely associated with renal function deterioration after accounting for relevant confounders. Patients with an intake  $< 1.08$  g/kg/day ( $n = 121$ , 32%) had an increased hazard of renal function deterioration (Fig. 2). Prevalent proteinuria ( $\geq 150$  mg/24 h,  $n = 221$ ) was associated with an increased hazard for renal function deterioration (HR 2.16 [95% CI 1.16–4.03]) (Table 2). Age, sex, and baseline eGFR were not associated with renal function deterioration after accounting for relevant confounders.







**Figure 2**—Association between dietary protein intake (g/kg/day) and the hazard of renal function deterioration. HRs for different levels of dietary protein intake were established by comparing the hazard of renal function deterioration for a hypothetical male patient of median age (65 years) with median eGFR (84 mL/min/1.73 m<sup>2</sup>) and prevalent proteinuria ( $\geq 150$  mg/24 h) with the hazard of renal function deterioration for a hypothetical patient who had the mean value for every covariate (including sex and prevalent proteinuria). The HR is shown by the solid line and the associated pointwise 95% CIs are represented by the shaded area. The null effect (HR 1) is shown by the orange dashed line. The red dashed lines represent the dietary protein intakes at which there was a significantly increased hazard of renal function deterioration.

focus on a lower blood pressure target in patients with proteinuria. In line with this notion, the percentage of patients using antihypertensive treatments is high in our population (73%).

The strength of the DIALECT cohort is that it reflects the real world of patients with T2D, with minimal loss to follow-up, and it was specifically designed to evaluate lifestyle characteristics such as dietary habits. Our study population appears to be a representative T2D population since renal function impairment was present in 22% of the patients compared with a prevalence rate of 20–40% in other T2D populations (3,4). In addition, we were able to assess dietary protein intake objectively, which is more reliable than

subjective dietary protein assessment. Another strength is that we were able to adjust for baseline kidney function on the basis of cystatin C–based eGFR, which is independent of muscle mass, although follow-up data on cystatin C–based renal function were not available. Obviously, our observational design limits us in determining causation, and we were not able to check whether our results are mainly the result of dietary protein intake. Also, 24-h urine specimens might have been collected with error. Finally, protein intake was assessed at baseline only, and we did not monitor whether patients had changed their dietary protein intake during the follow-up period.

**Table 2—Multivariable HRs for the association between dietary protein intake and renal function deterioration in patients with T2D included in the DIALECT population**

	HR	95% CI	P
Dietary protein intake (g/kg/day)	0.67	0.48–0.94	0.017
Age (years)	1.16	0.82–1.64	0.40
Sex (female)	0.98	0.56–1.72	0.95
eGFR (mL/min/1.73 m <sup>2</sup> )	0.83	0.60–1.14	0.24
Prevalent proteinuria (yes)	2.16	1.16–4.03	0.015

The multivariable model was adjusted for age, sex, baseline eGFR, and prevalent proteinuria at baseline ( $>150$  mg/24 h). Continuous variables were expressed per incremental SD.

With regard to the potential impact on population health, substituting carbohydrates with dietary protein seems a rational approach in T2D management, which may have a positive effect on body weight while minimizing loss of muscle mass (31). A specific note is that dietary recommendations should take into account individual nutritional needs. Patients with T2D and CKD often have multiple comorbid diseases, which requires more complex dietary recommendations. In addition, to conform a sustained effect of dietary protein intake over the long term, assessment of dietary compliance will be a challenging aspect.

**Acknowledgments.** The authors thank Christina Gant, Meander Medical Center, the Netherlands, and Heleen Binnenmars, University Medical Center Groningen, the Netherlands, for general contributions to DIALECT, including patient inclusion.

**Duality of Interest.** G.D.L. reports grants and personal fees from Sanofi, AstraZeneca, Astellas, and Novo Nordisk and personal fees from Janssen outside the submitted work. No other potential conflicts of interest relevant to this article were reported.

**Author Contributions.** M.M.O. performed statistical analyses and wrote the manuscript. D.G. supervised the statistical analyses, interpreted the outcome measures, contributed to the discussion, and reviewed/edited the manuscript. G.N. and S.J.L.B. contributed to the discussion and reviewed/edited the manuscript. G.D.L. was principal investigator of this study, coordinated the study, contributed to the discussion, and reviewed/edited the manuscript. M.M.O. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

**References**

1. Fouque D, Laville M. Low protein diets for chronic kidney disease in non diabetic adults. *Cochrane Database Syst Rev* 2009;(3):CD001892
2. Kidney Disease: Improving Global Outcomes. Chapter 3: management of progression and complications of CKD. KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney Int Suppl* 2013;3:73–90
3. American Diabetes Association. Microvascular complications and foot care: *Standards of Medical Care in Diabetes—2018*. *Diabetes Care* 2018;41(Suppl. 1):S105–S118
4. Lea JP, Nicholas SB. 10. Diabetes mellitus and hypertension: key risk factors for kidney disease. *J Natl Med Assoc* 2002;94(Suppl.):7S–15S
5. Feinman RD, Pogozelski WK, Astrup A, et al. Dietary carbohydrate restriction as the first approach in diabetes management: critical review and evidence base. *Nutrition* 2015;31:1–13

