



The At-Risk Diabetic Foot: Time to Focus on Prevention

Diabetes Care 2022;45:e144–e145 | <https://doi.org/10.2337/dc22-1111>

Robert G. Frykberg,^{1,2}
Loretta Vileikyte,^{3,4}
Andrew J.M. Boulton,^{3,4} and
David G. Armstrong⁵

As we approach the end of the coronavirus disease 2019 (COVID-19) pandemic, it has become clear that delayed care has had deleterious consequences for those with diabetes-related complications of the lower extremities. We have witnessed increasing numbers of patients presenting to emergency rooms with untreated diabetic foot ulcers (DFU), infections, and even gangrene, very often resulting from a delay in seeking care (1). Too many such patients consequently required urgent hospitalization and emergent surgery to control limb-threatening infections. While we recognize that foot-level amputations are often limb-sparing (and sometimes lifesaving) operations, we also know that many such outcomes can be prevented by early intervention. Furthermore, we also recognize that by preventing that initial or recurrent DFU, we can disrupt the causal pathway leading to limb loss (2). Accordingly, we must now focus on the prevention of such complications if we are going to reduce the global burden of diabetic lower extremity complications.

Recognizing the importance of preventing amputation in neuropathic individuals, the Health Resources and Services Administration (HRSA) developed their Lower Extremity Amputation Prevention (LEAP) program in 1992 (3). Although initially focused on people with Hansen disease, LEAP was also intended to apply to those with any condition that results in loss of protective sensation in the feet. Accordingly, the larger population of people with diabetes has become an

important focus of this prevention program. The five-step LEAP program consists of 1) annual foot screening, 2) patient education, 3) daily self-inspection, 4) footwear selection, and 5) management of simple (preulcerative) foot problems. Indeed, these important prevention components have long been embraced within diabetic foot guidelines put forth by several national and international organizations. These include the American Diabetes Association, the International Working Group on the Diabetic Foot, and many others. The Veterans Health Administration similarly incorporated the principles embodied in the LEAP program through their Preservation-Amputation Care and Treatment (PACT) program, first established in 1993. This was later renamed the Preventing Amputations in Veterans Everywhere (PAVE) program in 2012. A critical component of this program is the requirement for an annual foot examination for all patients with diabetes.

In its original position statement on preventive foot care, the ADA recommended annual foot screenings, including biomechanical, structural, dermatologic, neurologic, and vascular assessments, for all individuals with diabetes. This was further expanded upon in the 2008 publication that stressed the importance of a comprehensive foot examination and commensurate risk assessment (4). The introduction of “smart” technologies, where plantar foot temperatures and pressures can be monitored and feedback

provided in real time via audio and visual alerts, is indeed promising. It represents not only a paradigm shift in DFU risk screening and monitoring, but, crucially, it also transforms foot care education. The saying that “a picture is worth 1,000 words” is particularly relevant to this patient population, for whom symptoms and signs cannot be relied on when conveying messages about DFU risk. By allowing people to visualize their personal DFU risk, digital technologies are likely to enhance patients’ active participation in DFU prevention (5).

While we have made great strides in healing DFU, our progress in preventing them is debatable. As noted above, we have long had guidelines directing appropriate prevention recommendations as well technologies allowing for remote patient monitoring. It is therefore incumbent upon all practitioners managing patients with diabetes to focus on preventative protocols as much as on treatment protocols when considering diabetic foot disorders. While not as glamorous or financially rewarding as treating extant diabetic foot complications, our efforts must indeed be redirected toward preventing such complications.

Just as recommended retinal examinations have decreased the incidence of diabetes-related blindness over the last several decades, we propose that the requirement for annual comprehensive diabetic foot examinations be implemented as a measure of good clinical practice. The Healthcare Effectiveness Data and

¹Diabetic Foot Consultants, Phoenix, AZ

²Midwestern University, Glendale, AZ

³University of Manchester, Manchester, U.K.

⁴University of Miami, Miami, FL

⁵Keck School of Medicine of University of Southern California, Los Angeles, CA

Corresponding author: Robert G. Frykberg, rgfdpm@diabeticfoot.net

Received 13 July 2022 and accepted 17 July 2022

© 2022 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <https://www.diabetesjournals.org/journals/pages/license>.

Information Set (HEDIS) measures, commonly used to ensure appropriate evaluations of insured individuals and with commensurate rewarding for good clinical practices, currently track parameters for retinal examinations and measurement of HbA_{1c}, blood pressure, etc. However, there are currently no established HEDIS guidelines for annual comprehensive diabetic foot examinations. We believe that the growing incidence of diabetic foot disorders in our populations and their associated morbidities dictate the need for greater attention to such complications of diabetes. We therefore recommend and propose that a formal HEDIS measure for annual comprehensive diabetic foot examination, according to the principles embodied in the original LEAP initiative, be widely implemented in global health care systems.

By focusing on early detection of risk factors and/or early onset of diabetic foot lesions, we can more effectively reduce the incidence of potentially life-changing lower extremity complications of diabetes.

Duality of Interest. R.G.F. is a consultant to Arche Healthcare. No other potential conflicts of interest relevant to this article were reported.

Author Contributions. All authors contributed to manuscript preparation. All authors have read and agreed to the published version of the manuscript. R.G.F. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

1. Caruso P, Longo M, Signoriello S, et al. Diabetic foot problems during the COVID-19

pandemic in a tertiary care center: the emergency among the emergencies. *Diabetes Care* 2020;43:e123–e124

2. Armstrong DG, Boulton AJM, Bus SA. Diabetic foot ulcers and their recurrence. *N Engl J Med* 2017;376:2367–2375

3. Health Resources and Services Administration. Lower extremity amputation prevention (LEAP). Health Resources and Services Administration, 2022. Accessed 28 May 2022. Available from <https://www.hrsa.gov/hansens-disease/leap>

4. Boulton AJ, Armstrong DG, Albert SF, et al.; American Diabetes Association; American Association of Clinical Endocrinologists. Comprehensive foot examination and risk assessment: a report of the task force of the Foot Care Interest Group of the American Diabetes Association, with endorsement by the American Association of Clinical Endocrinologists. *Diabetes Care* 2008; 31:1679–1685

5. Boulton AJM, Armstrong DG, Londahl M, et al. *New Evidence-Based Therapies for Complex Diabetic Foot Wounds*. Arlington, VA, American Diabetes Association, 2022