



Glucagon-Like Peptide 1 Receptor Agonists, Carotid Atherosclerosis, and Cardiovascular Outcomes

Guillaume Marquis-Gravel and
Jean-Claude Tardif

Diabetes Care 2021;44:1252–1253 | <https://doi.org/10.2337/dci20-0076>

Glucagon-like peptide 1 receptor agonists (GLP-1 RA) represent an integral part of the arsenal used in clinical practice to improve cardiovascular outcomes in patients with diabetes. Large-scale randomized controlled trials have shown that liraglutide, dulaglutide, albiglutide, and semaglutide all reduce the risk of cardiovascular events in patients with diabetes and either established atherosclerotic cardiovascular disease (ASCVD) or high-risk characteristics (1–4). In contrast, exenatide and lixisenatide improved glycemic control but did not have a sizable impact on cardiovascular events in clinical trials (5,6), suggesting that glycemic control and cardiovascular outcomes are at least partly uncoupled. A meta-analysis including 42,920 participants from five randomized trials demonstrated that GLP-1 RA as a drug class are associated with a significant 13% reduction in the risk of the composite of myocardial infarction, stroke, or cardiovascular death in patients with diabetes and established ASCVD (7). GLP-1 RA are recommended in patients with diabetes and established ASCVD or at high risk of ASCVD (8). They are also indicated as second-line therapy in patients without ASCVD (or without indicators of high risk) who do not meet treatment goals with lifestyle modifications and metformin alone,

and they are preferred to insulin if injectable therapy is needed (8).

Although the impact of GLP-1RA on cardiovascular outcomes has been shown to be directly correlated with the reduction in glycated hemoglobin (9), pleiotropic effects of agents from this class have been documented, including reduction of systolic and diastolic blood pressure (10), body weight (11), and vascular inflammation (12), in addition to improvement of endothelial function (13). However, the impact of GLP-1 RA on atherosclerotic plaque volume and composition had not yet been investigated.

As reported in this issue of *Diabetes Care*, Koska et al. (14) examined whether the GLP-1 RA exenatide modifies carotid plaque volume and composition in patients with type 2 diabetes. They conducted a placebo-controlled, double-blind, pragmatic randomized trial in which 163 participants were randomly allocated to receive either exenatide 2 mg ($n = 109$) or placebo ($n = 54$) subcutaneously once weekly. Patients with a high carotid atherosclerosis burden at baseline were enrolled, with the majority presenting calcified plaques and lipid-rich necrotic cores. Plaque volume and composition were measured using serial carotid MRI at baseline, 9 months, and 18 months of follow-up. Despite a mean reduction in glycated hemoglobin of 0.55% with once-weekly exenatide compared with placebo,

there was no significant difference in the change over time in plaque volume between study groups. Plaque composition was also not modified by once-weekly exenatide, without significant differences detected in changes over time in the dimensions of calcified plaques, lipid-rich necrotic core plaques, and fibrous caps between the two treatment arms. Results were consistent among key prespecified subgroups and after excluding participants not completely adhering to the protocol.

The lack of benefit of once-weekly exenatide on carotid plaque volume and composition observed in this study is consistent with its modest effect on the primary composite end point of death from cardiovascular causes, nonfatal myocardial infarction, or nonfatal stroke in the Exenatide Study of Cardiovascular Event Lowering (EXSCEL) trial of 14,752 patients followed for a median of 3.2 years (hazard ratio 0.91, 95% CI 0.83–1.00, $P = 0.06$). Whether other GLP-1 RA associated with more marked cardiovascular benefits (liraglutide, dulaglutide, albiglutide, and semaglutide) would yield different results on carotid MRI is not known. Nevertheless, a previous study without a placebo group found that liraglutide, a GLP-1 RA that demonstrated cardiovascular benefits (1), significantly reduced carotid intima-media thickness (15). In the Harmony Outcomes trial, the reduction of cardiovascular risk with albiglutide was

Montreal Heart Institute, Université de Montréal, Montréal, Quebec, Canada

Corresponding author: Jean-Claude Tardif, jean-claude.tardif@icm-mhi.org

© 2021 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <https://www.diabetesjournals.org/content/license>.

See accompanying article, p. 1385.

greater than would have been expected given the degree of glycated hemoglobin reduction, suggesting that the cardioprotective effects are not exclusively linked to glycemic control (3,9). In the current study, a correlation was observed between reductions of glycated hemoglobin and plaque volume in the exenatide group, but the significance of this observation is uncertain given the lack of overall effect on changes in carotid MRI end points compared with placebo.

The increase in heart rate of 9 bpm at 3 months of treatment with once-weekly exenatide compared with placebo was more marked than that reported in clinical trials of GLP-1 RA associated with greater reductions in cardiovascular event rates (1,4,11). In light of the prognostic value of resting heart rate on cardiovascular outcomes and its link with experimental atherosclerosis (16), the heart rate-elevating effect of once-weekly exenatide might explain in part the disappointing carotid imaging results in the current study.

The study by Koska et al. (14) needs to be carefully interpreted in light of the relatively small sample size and the high rate (25%) of incomplete imaging procedures at 18 months of follow-up. The latter may be of importance given that patients who withdrew early had lower BMI and triglyceride values than those who completed the study. Given that some clinical trials of GLP-1 RA have shown limited differences in outcomes within the first 2 years but greater separation of cumulative cardiovascular event curves thereafter, the 18-month follow-up duration in the current study might also have represented a limitation. Finally, the lack of effect of once-weekly exenatide on carotid plaque volume and composition

observed in this study does not exclude a potential effect on coronary atherosclerosis. Notwithstanding these limitations, the study by Koska et al. suggests that the cardiovascular benefits of GLP-1 RA might not be entirely due to structural changes in atherosclerotic plaques. In order to definitely answer that question, additional placebo-controlled randomized trials involving imaging of carotid and coronary arteries should be conducted with those GLP-1 RA that have shown more pronounced benefits on cardiovascular outcomes.

Duality of Interest. J.-C.T. reports grants from Amarin, AstraZeneca, Ceapro, DalCor, Esperion, Ionis, Pfizer, and RegenXBio; honoraria from AstraZeneca, DalCor, HLS Therapeutics, Pharmascience, and Servier; and minor equity interest in DalCor. G.M.G. reports grants from Bayer. No other potential conflicts of interest relevant to this article were reported.

References

1. Marso SP, Daniels GH, Brown-Frandsen K, et al.; LEADER Steering Committee; LEADER Trial Investigators. Liraglutide and cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2016; 375:311–322
2. Gerstein HC, Colhoun HM, Dagenais GR, et al.; REWIND Investigators. Dulaglutide and cardiovascular outcomes in type 2 diabetes (REWIND): a double-blind, randomised placebo-controlled trial. *Lancet* 2019;394:121–130
3. Hernandez AF, Green JB, Janmohamed S, et al.; Harmony Outcomes committees and investigators. Albiglutide and cardiovascular outcomes in patients with type 2 diabetes and cardiovascular disease (Harmony Outcomes): a double-blind, randomised placebo-controlled trial. *Lancet* 2018;392:1519–1529
4. Marso SP, Bain SC, Consoli A, et al.; SUSTAIN-6 Investigators. Semaglutide and cardiovascular outcomes in patients with type 2 diabetes. *N Engl J Med* 2016;375:1834–1844
5. Holman RR, Bethel MA, Mentz RJ, et al.; EXSCEL Study Group. Effects of once-weekly exenatide on cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2017;377:1228–1239
6. Pfeffer MA, Claggett B, Diaz R, et al.; ELIXA Investigators. Lixisenatide in patients with type 2 diabetes and acute coronary syndrome. *N Engl J Med* 2015;373:2247–2257
7. Zelniker TA, Wiviott SD, Raz I, et al. Comparison of the effects of glucagon-like peptide receptor agonists and sodium-glucose cotransporter 2 inhibitors for prevention of major adverse cardiovascular and renal outcomes in type 2 diabetes mellitus. *Circulation* 2019;139:2022–2031
8. American Diabetes Association. 9. Pharmacologic approaches to glycemic treatment: *Standards of Medical Care in Diabetes—2021*. *Diabetes Care* 2021;44(Suppl. 1):S111–S124
9. Zweck E, Roden M. GLP-1 receptor agonists and cardiovascular disease: drug-specific or class effects? *Lancet Diabetes Endocrinol* 2019; 7:89–90
10. Wang B, Zhong J, Lin H, et al. Blood pressure-lowering effects of GLP-1 receptor agonists exenatide and liraglutide: a meta-analysis of clinical trials. *Diabetes Obes Metab* 2013;15:737–749
11. Robinson LE, Holt TA, Rees K, Randeve HS, O'Hare JP. Effects of exenatide and liraglutide on heart rate, blood pressure and body weight: systematic review and meta-analysis. *BMJ Open* 2013;3:e001986
12. Drucker DJ. The cardiovascular biology of glucagon-like peptide-1. *Cell Metab* 2016;24: 15–30
13. Koska J, Sands M, Burciu C, et al. Exenatide protects against glucose- and lipid-induced endothelial dysfunction: evidence for direct vasodilation effect of GLP-1 receptor agonists in humans. *Diabetes* 2015;64:2624–2635
14. Koska J, Migrino RQ, Chan KC, Cooper-Cox K, Reaven PD. The effect of exenatide once weekly on carotid atherosclerosis in individuals with type 2 diabetes: an 18-month randomized placebo-controlled study. *Diabetes Care* 2021; 44:1385–1392
15. Rizzo M, Rizvi AA, Patti AM, et al. Liraglutide improves metabolic parameters and carotid intima-media thickness in diabetic patients with the metabolic syndrome: an 18-month prospective study. *Cardiovasc Diabetol* 2016; 15:162
16. Fox K, Borer JS, Camm AJ, et al.; Heart Rate Working Group. Resting heart rate in cardiovascular disease. *J Am Coll Cardiol* 2007;50: 823–830