



RESPONSE TO COMMENT ON DUARTE ET AL.

Systematic Review and Network Meta-analysis of Neurostimulation for Painful Diabetic Neuropathy.

Diabetes Care 2022;45:2466–2475

Diabetes Care 2023;46:e113–e114 | <https://doi.org/10.2337/dci23-0008>

Rui V. Duarte,^{1,2} Sarah Nevitt,¹
Sue Copley,³ Michelle Maden,¹
Cecile C. de Vos,^{4,5} Rod S. Taylor,⁶ and
Sam Eldabe³

Sharan et al. (1) and Sadeghirad et al. (2) raise important points regarding the data sets used in our article (3) and whether a network meta-analysis (NMA) is appropriate. We appreciate the opportunity to respond.

We agree that it would have been preferable for our NMA to follow an intention-to-treat (ITT) approach. Although we are aware of the publicly available ITT analysis for the study of high-frequency spinal cord stimulation (SCS), ITT analysis was only available for 50% pain reduction at the 3-month follow-up. Comparable outcome data were not available for the studies of low-frequency SCS at the 3-month follow-up. Therefore, we were unable to compare the studies based on an ITT approach for any outcome. The results and conclusion are based on the available data.

No long-term data are available that are appropriate to inform comparisons in the NMA, since crossover was allowed at 6-month follow-up in the included trials.

Furthermore, Sadeghirad et al. (2) highlight the results of another subgroup analysis. The results of the primary comparison of SCS to placebo/sham showed significant improvements in pain (pooled mean difference -1.15 [95% CI -1.75 to -0.55 , $P = 0.001$]) (4). This review,

however, was conducted in a different population.

The probabilities that each treatment is the “best” (i.e., the largest effect size) (5) for each outcome are presented in Supplementary Table 1, and the footnote explains that these are calculated in the NMA. Importantly, our analysis shows that both forms of SCS provide better outcomes than conventional medical management (CMM) in a population with painful diabetic neuropathy (PDN).

The choice between fixed-effects and random-effects NMA was made a priori based on the anticipated similarity of population, trial design, outcomes, etc., for the research question. A fundamental issue for these analyses is that only a small number of trials with small sample sizes have been performed, as SCS currently is not commonly used in PDN treatment. We agree with Sadeghirad et al. (2) about power in NMA; however, it will likely be many years before 10 trials are published with the “optimal information size.” Do Sadeghirad et al. suggest we wait until then to do an NMA? Until a direct head-to-head comparison is available, indirect comparison, despite its limitations, is currently the only evidence available to inform clinician choice.

We have acknowledged and downgraded all outcomes due to imprecision, and the certainty of evidence was downgraded further for 50% reduction in pain because although the outcome of 50% pain reduction is based on pain intensity score, the dichotomization into $<50\%$ and $\geq 50\%$ pain reduction resulted in very small patient numbers in the CMM groups. The same does not apply to the evaluation of pain intensity as a continuous outcome.

In conclusion, we consider that the methods, interpretation, and conclusions made are justified and supported by the results presented. As we state in our article, there are limitations in the evidence base of SCS for patients with PDN, and a head-to-head trial of LF-SCS and HF-SCS would provide greater clarity on which intervention provides better outcomes for patients with PDN (3).

Duality of Interest. R.V.D. has received consultancy fees from Mainstay Medical, Medtronic, and Saluda Medical, all unrelated to this work. He is an employee of Saluda Medical. R.S.T. has received consultancy fees from Medtronic, Nevro Corp., and Saluda Medical, all unrelated to this work. He is due to serve on a Medtronic advisory board on

¹Liverpool Reviews and Implementation Group, University of Liverpool, Liverpool, U.K.

²Saluda Medical Pty. Ltd., Artarmon, New South Wales, Australia

³Department of Pain Medicine, The James Cook University Hospital, Middlesbrough, U.K.

⁴Department of Neurology and Neurosurgery, Medisch Spectrum Twente, Enschede, the Netherlands

⁵Centre for Pain Medicine, Erasmus MC, University Medical Centre, Rotterdam, the Netherlands

⁶Medical Research Council/Chief Scientist Office Social and Public Health Sciences Unit and Robertson Centre for Biostatistics, Institute of Health and Well Being, University of Glasgow, Glasgow, U.K.

Corresponding author: Rui V. Duarte, rui.duarte@liverpool.ac.uk

© 2023 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <https://www.diabetesjournals.org/journals/pages/license>.

SCS for PDN and is a co-investigator for the Comparison of 10 kHz SCS Combined With CMM to CMM Alone in the Treatment of Neuropathic Limb Pain (SENZA-PDN) trial. S.E. has received consultancy fees from Abbott, Boston Scientific Corp., Mainstay Medical, and Medtronic, all unrelated to this work. He has received department research funding from the National Institute of Health Research, Medtronic, and Nevro Corp. No other potential conflicts of interest relevant to this article were reported.

References

1. Sharan AD, Hatheway JA, Murphy M, et al. Comment on Duarte et al. Systematic review and network meta-analysis of neurostimulation for painful diabetic neuropathy. *Diabetes Care* 2022; 45:2466–2475 (Letter). *Diabetes Care* 46:e110–e111
2. Sadeghirad B, Wang X, Busse JW. Comment on Duarte et al. Systematic review and network meta-analysis of neurostimulation for painful diabetic neuropathy. *Diabetes Care* 2022;45: 2466–2475 (Letter). *Diabetes Care* 46:e112
3. Duarte RV, Nevitt S, Copley S, et al. Systematic review and network meta-analysis of neurostimulation for painful diabetic neuropathy. *Diabetes Care* 2022;45:2466–2475
4. Duarte RV, Nevitt S, McNicol E, et al. Systematic review and meta-analysis of placebo/sham controlled randomised trials of spinal cord stimulation for neuropathic pain. *Pain* 2020;161:24–35
5. Mbuagbaw L, Rochweg B, Jaeschke R, et al. Approaches to interpreting and choosing the best treatments in network meta-analyses. *Syst Rev* 2017;6:79