

Music as Medicine: The Therapeutic Potential of Music for Acute Stroke Patients

Charlene Supnet, PhD
April Crow, RN
Sonja Stutzman, PhD
DaiWai Olson, RN, PhD, CCRN

Nurses caring for patients with acute stroke are likely to administer both music and medication with therapeutic intent. The administration of medication is based on accumulated scientific evidence and tailored to the needs of each patient. However, the therapeutic use of music is generally based on good intentions and anecdotal evidence. This review summarizes and examines the current literature regarding the effectiveness of music in the treatment of critically ill patients and the use of music in neurologically injured patients. The rationale for hypothesis-driven research to explore therapeutic music intervention in acute stroke is compelling. (*Critical Care Nurse*. 2016;36[2]:e1-e7)

“Music hath charms to soothe the savage breast.”
W. Congreve, *The Mourning Bride*

Music has long been used in health care for its healing powers and has given rise to the practice of music therapy. The American Music Therapy Association defines music therapy as the “clinical and evidence-based use of music interventions to accomplish individualized goals within a therapeutic relationship by a credentialed professional who has completed an approved music therapy program.”¹ Board-certified music therapists employ customized music therapy interventions based on the best evidence from published literature, their training and expertise, and the needs of the patient.¹ Music therapy is powerful, physically noninvasive, and can mediate positive outcomes when

CE 1.0 hour

This article has been designated for CE contact hour(s). The evaluation tests your knowledge of the following objectives:

1. Differentiate between music as a therapeutic intervention and music therapy
2. List strategies to promote the use of music as a therapeutic intervention
3. Identify gaps in the current literature that require nursing research

To complete evaluation for CE contact hour(s) for test #C1623, visit www.cconline.org and click the “CE Articles” button. No CE test fee for AACN members. This test expires on April 1, 2019.

The American Association of Critical-Care Nurses is an accredited provider of continuing nursing education by the American Nurses Credentialing Center’s Commission on Accreditation. AACN has been approved as a provider of continuing education in nursing by the State Boards of Registered Nursing of California (#01036) and Louisiana (#LSBN12).

©2016 American Association of Critical-Care Nurses doi: <http://dx.doi.org/10.4037/ccn2016413>

interventions are directed to reduce pain, anxiety, stress, and depression.²⁻⁸ Music is often employed as an ambient or adjunct tool during medical procedures and for chronic pain management. However, no consensus has been reported regarding protocols for listening to music, its effectiveness in certain procedures, and which type of music is most effective. Furthermore, distinct differences are apparent between music-supported therapy (use of musical instruments in motor rehabilitation therapy),⁹ music as a therapeutic intervention (eg, music listening to affect vital signs, changes in analgesia),¹⁰⁻¹² and formal music therapy, which can be provided only by a music therapist.¹ Nonetheless, nurses and family members often play music with therapeutic intent in the presence of the patient.¹³

Following acute stroke, patients with severe brain injuries may benefit from mechanical ventilation to prevent secondary injury by providing airway protection and to ensure adequate oxygenation. Mechanical ventilation may also facilitate short-term goals to reduce brain swelling with the use of sedative, osmotic, or vasoactive medication therapy to reduce intracranial pressure¹⁴ and augment blood pressure.¹⁵⁻¹⁷ Patients with acute stroke who are receiving mechanical ventilation are often given sedative and hypnotic medications to prevent self-harm, facilitate nursing, and minimize their memory of the unpleasant experience of being intubated.¹⁸ However, oversedation of acute stroke patients may affect neurological performance on assessment examinations and result in inaccurate measures of

progress or deterioration. Care providers are constantly balancing the use of sedatives to keep patients comfortable with the need for minimal impact on their neurological state. Therefore, identifying nonpharmacotherapeutic interventions that help reduce anxiety and stress would markedly help staff caring for acute stroke patients to obtain a more precise assessment of their status, which could lead to the administration of appropriate interventions, decreased length of stay, and promotion of long-term recovery.^{19,20}

Music reduces postoperative pain, agitation, and narcotic requirements for critically ill cardiovascular and medical-surgical patients.²¹⁻²⁶ Although playing music has been associated with lower anxiety, pain, and agitation in patients undergoing mechanical ventilation,^{3,5,7,27-30} few studies have addressed the effect of music on patients with acute stroke in critical care.³¹⁻³³ The goal of this review is to explore the utility of music intervention as a means to influence outcomes in acute stroke patients.

Music and the Environment of Care

Nurses have long been recognized for their role in maintaining the environment of care³⁴ and establishing the therapeutic milieu to provide calm and foster optimal healing.³⁵ Milieu therapy, which includes controlling the levels of audio and visual stimulation the patient receives, promotes rest and sleep in neurocritically ill patients.³⁶⁻³⁸ A recent neurocritical care practice survey¹³ indicates that nurses and physicians often play music or turn the television on or off with therapeutic intent, despite a lack of best practice guidelines. Nevertheless, it has been demonstrated that music therapy can affect electroencephalographic brainwave activity, which is closely related to cerebral brain function.^{39,40} Playing music with a beat of roughly 60 to 66 beats per minute, such as classical baroque music,⁴¹ helps produce alpha waves in the brain⁴² and a state of relaxation, alertness, and meditation. Music was described in 1946 as an intervention that could alleviate pain and anxiety related to being in an intensive care unit (ICU) and reduce length of stay and sleep deprivation by providing a more patient-friendly ICU environment.⁴³ Furthermore, research on the use of relaxing music in the operating room, ICU, and critical care areas has been done before.^{21,40,44}

Chlan and Tracy⁴⁵ reported that allowing critically ill patients to choose the music played for intervention and providing uninterrupted listening periods gave patients

Authors

Charlene Supnet is an experienced basic/clinical neuroscience researcher and writer for the Department of Neurology and Neurotherapeutics and the Neuroscience Nursing Research Center, University of Texas Southwestern Medical Center, Dallas, Texas.

April Crow is a nurse in the inpatient rehabilitation unit of Zale Lipshy University Hospital, Dallas, Texas.

Sonja Stutzman is the clinical research manager for the Neuroscience Nursing Research Center, University of Texas Southwestern Medical Center. Additionally, she coordinates several acute studies in the neuroscience intensive care unit.

DaiWai Olson is an associate professor and director of the Neuroscience Nursing Research Center at University of Texas Southwestern.

Corresponding author: DaiWai Olson, RN, PhD, CCRN, UT Southwestern Medical Center, 5323 Harry Hines Boulevard, CS07.104, Dallas, TX 75390-8897 (e-mail: daiwai.olson@utsouthwestern.edu).

To purchase electronic or print reprints, contact the American Association of Critical-Care Nurses, 101 Columbia, Aliso Viejo, CA 92656. Phone, (800) 899-1712 or (949) 362-2050 (ext 532); fax, (949) 362-2049; e-mail, reprints@aacn.org.

an enhanced sense of control. Music therapy delivered by trained music therapists was associated with lower stress, as measured by lower apical heart rates and elevated peripheral temperatures, and lower incidence of complications compared with no music in patients with acute myocardial infarction.²² However, a recent Cochrane review noted that the majority of trials that used music as the main intervention did not use a trained music therapist.²⁶ Good²³ reported that playing music for patients was associated with a reduction in postoperative pain. In a review article examining music as a therapeutic intervention in the ICU, Chlan²⁹ reported that a majority of patients who receive mechanical ventilation experience some stress and anxiety, and although the use of music has the potential to benefit these patients by reducing anxiety, evidence to support music as a therapeutic intervention for a diverse group of patients is inadequate.

In a Cochrane review of music therapy for acquired brain injury published in 2010, 7 studies of music therapy for stroke patients were considered (N = 184).³¹ Notably, the participants in these studies were described as not having acute stroke and not being cared for in the ICU.³¹ A 2014 Cochrane review of music therapy for patients receiving mechanical ventilation reported on 14 studies.¹² Of these, 1 study⁴⁶ included only 3 patients with neurological injury and showed no benefit of listening to music for those patients. However, the review indicated that on average, music listening had a large and clinically significant effect on anxiety, causing a reduction of 1.1 standard deviation units more than in the standard care group (95% CI, -1.75 to -0.47, $P < .001$).¹² Because the meta-analysis performed revealed that music listening may reduce physiological responses, including the respiratory rate and systolic blood pressure, in addition to reducing anxiety in critically ill patients receiving mechanical ventilation, the authors recommend that music listening be offered as a stress management intervention.¹² However, they acknowledge that further studies including music therapists and using a randomized clinical trial design are needed to determine if and how music influences the environment of care.¹²

Music and Critically Ill Patients

A growing body of literature examines the utility of playing music to reduce anxiety and stress during critical illness.^{3,21,22,26,27,32} The use of music reduced postoperative pain sensation and distress 2 days after the procedure in

89% of patients who had undergone abdominal surgery, compared with control patients who did not have music played for them.²³ Studies that used music interventions in patients with myocardial infarction have consistently shown that music induces anxiety-reducing effects, with a mean anxiety reduction of 5.87 units on a scale of 20 to 80 units ($P < .001$).²⁶ One study⁴⁷ examined the effect of a 20-minute uninterrupted music intervention, versus 20 minutes of uninterrupted rest, in both intubated and nonintubated patients. Music intervention significantly decreased heart rate ($P < .05$), systolic blood pressure ($P < .05$), respiratory rate ($P < .05$), bispectral index ($P < .01$), agitation measured by the Richmond Agitation-Sedation Scale ($P < .05$), and pain measured by the Numerical Rating Scale ($P < .01$) in both intubated and nonintubated patients compared with patients who got rest only.⁴⁷ Thus, the use of music has been demonstrated to reduce postoperative agitation and narcotic requirements for critically ill cardiovascular and medical-surgical patients.^{23,26} A recent multicenter randomized clinical trial provides additional evidence that playing music is associated with lower anxiety, sedation intensity, and sedation frequency compared with usual care and noise ablation in patients undergoing mechanical ventilation.²⁷ However, the study excluded patients who could not provide self-consent (ie, stroke patients receiving mechanical ventilation).²⁷ Music has been associated with decreased anxiety, pain, and agitation in patients receiving mechanical ventilation.^{3,5,7,27-30} However, few studies have examined music intervention for patients with acute stroke in critical care.³¹⁻³³

The use of music reduced postoperative pain and distress in 89% of patients after abdominal surgery.

Stroke patients transition early, and often, to receive rehabilitation.⁴⁸ Stroke patients are also at higher risk of emotional, cognitive, and consciousness disorders. Results from the Healthy Ageing and Intellectual Disabilities study in the Netherlands demonstrated that anxiety and depression were common following illness.⁴⁹ Moreover, anxiety and depression are both prevalent in rehabilitative care and may limit recovery.⁵⁰⁻⁵⁴ Anxiety manifests in 2 ways: emotional symptoms, such as fear and worry, and physical symptoms, such as dry mouth, dizziness, increased heart rate, sweating, rapid breathing,

Role of Music in Stroke Rehabilitation

Stroke patients transition early, and often, to receive rehabilitation.⁴⁸ Stroke patients are also at higher risk of emotional, cognitive, and consciousness disorders. Results from the Healthy Ageing and Intellectual Disabilities study in the Netherlands demonstrated that anxiety and depression were common following illness.⁴⁹ Moreover, anxiety and depression are both prevalent in rehabilitative care and may limit recovery.⁵⁰⁻⁵⁴ Anxiety manifests in 2 ways: emotional symptoms, such as fear and worry, and physical symptoms, such as dry mouth, dizziness, increased heart rate, sweating, rapid breathing,

and possibly fatigue.⁵⁵ Depression is a disorder that has been characterized by low mood or loss of interest in things that once one found enjoyable, both of which affect the activities of daily living.⁵⁶

It has been demonstrated that music-supported therapy can induce changes to the brain, or plasticity, and cause the activation of the auditory-motor circuits that improve motor skills after acute or chronic stroke.⁹ The analysis of structural magnetic resonance imaging data of patients who listened to music for 6 months after stroke revealed an increase in the gray matter volume and reorganization in the frontal areas of the brain (left and right superior frontal gyrus and the middle frontal gyrus) that enhanced the recovery of verbal memory, focused attention, and language skills.⁹ Emotional effects induced by music learning, performing, and listening could engage reward-learning networks and corresponding neurotransmitter systems^{57,58} that help to consolidate new information and increase the likelihood of voluntary practicing of new movements.

Music has shown promise as an adjunct to treatment, and specific elements of music have been linked to significantly higher rates of positive emotion.^{45,58} For example, patients given music therapy experienced lower anxiety and a higher level of satisfaction with care provided, compared with patients who did not receive music intervention.^{45,59} Music therapy improves executive functioning and eases the adjustment to rehabilitation in patients with traumatic brain injury.^{60,61} Recently, it was reported that music listening significantly enhanced

Music therapy may help to maintain a positive outlook in stroke patients who are undergoing rehabilitation therapy.

cognitive functioning in verbal memory


and focused attention ($P < .01$ and $P < .05$, respectively, mixed-model analysis of variance), and decreased depression and confusion (both $P < .10$, 1-way analysis of variance), or positive emotion, in patients after middle cerebral artery stroke compared with control participants.⁶² It is often difficult to maintain a positive attitude when experiencing sudden life-changing events, which is especially true for patients who are emergently admitted to the ICU. Improving patients' mood is also important because it may affect the patient-care partner relationship following discharge.⁶³ In a study of music and patients' mood that used a single subject design, Magee and Davidson⁶ used the Profile of Mood States instrument

in a cohort of 14 patients (4 with stroke, 5 with multiple sclerosis, and 5 with traumatic brain injury) and reported that composed-anxious, energetic-tired, and agreeable-hostile mood states were significantly improved from before to after a music therapy intervention. Thus, it is reasonable to consider that music therapy may also help to maintain a positive outlook in other neurologically injured stroke patients who are undergoing rehabilitation therapy.

Is Music Beneficial to the Recovery of Acute Stroke Patients?

Following primary brain injury caused by a cerebrovascular event (eg, stroke), the focus of care during the acute phase is the prevention of secondary brain injury.⁶⁴ Improving and maintaining adequate cerebral perfusion is a chief concern. Therefore, severely injured acute stroke patients may benefit from mechanical ventilation to protect the airways and ensure adequate aeration of the lungs and oxygenation of the blood. Mechanical ventilation may also facilitate short-term goals to reduce brain swelling that may involve the use of sedative, osmotic, or vasoactive medication therapy to reduce intracranial pressure¹⁴ and increase blood pressure.¹⁵⁻¹⁷ Additionally, patients may require a host of supportive therapies such as nutritional support and antibiotic therapy. Prevention of complications associated with an ICU stay (eg, deep vein thrombus prophylaxis) is also a key factor in acute phase recovery.⁶⁵⁻⁶⁸

Patients with acute stroke who are undergoing mechanical ventilation are often given sedative and hypnotic medications to prevent self-harm, facilitate nursing and medical care, and blunt their memory of what is an unpleasant experience.¹⁸ Simultaneously, stroke patients require frequent neurological examinations to monitor for deterioration. Thus, one must ensure that any change in findings on the patient's neurological examination is not due, in whole or in part, to medication therapy (eg, oversedation causing a decreased level of consciousness). This situation creates a conundrum wherein staff must balance the use of sedative and hypnotic medications with the need to minimize the use of medications that could affect the findings on the neurological examination.⁶⁹ Hence, although it is true that anxiolysis and sedation can be accomplished solely by the use of medications, nonpharmacotherapeutic interventions that reduce anxiety have the potential to reduce length of



stay and posttraumatic stress disorder, decrease the cost of care, and promote long-term recovery.^{19,20} Given that music, as therapy, is effective in reducing agitation in a diverse patient population, the potential exists that the therapeutic use of music may have similar benefit for patients with cerebrovascular illness (see Figure). If beneficial, this low-cost, low-technology therapy may be used to reduce the use of sedatives and help in resolving the sedation assessment conundrum.

Although advantageous to the recovery of critically ill patients, the use of music as an intervention has not been studied extensively in acute stroke patients. However, a variety of independent studies provide insight regarding how music may benefit critically ill, acute stroke patients. Listening to music has been associated with neuronal recovery.^{33,70,71} Soothing music was associated with improved self-reported sleep quality in a study of 28 ICU patients.²⁵ Participants who listened to 45 minutes of music at nocturnal sleep time had shorter stage N2 non-rapid-eye-movement sleep, where brain waves become slower and more synchronized, and longer N3 (deeper) sleep in the first 2 hours of nocturnal sleep, improved self-reported sleep quality, and significantly lower heart rates compared with the control group ($P < .003$).²⁵ During anesthesia for surgical procedures,

music was beneficial for recovery and reducing the stress response.^{72,73} Postoperative patients exposed to music in combination with therapeutic suggestions required less rescue analgesia, experienced more effective analgesia, and at discharge were less fatigued than were control participants.⁷² In a randomized controlled trial designed to evaluate the influence of intraoperative and postoperative music therapy on stress and the immune response, researchers reported a significantly greater decrease in the level of cortisol, a stress hormone, in the urine of the patients who had music therapy after 2 hours in the postanesthesia care unit.⁷³ Furthermore, the postoperative music group required significantly less morphine than did the control group, but the level of immunoglobulin A, blood glucose, blood pressure, heart rate, and oxygen saturation did not differ significantly between the groups.⁷³ The protocol of 1 study⁷⁴ ensured that 99% of patients identified with delirium were exposed to music administered by a nurse daily. However, no attempt was made to measure the impact of this intervention on patients' outcomes.⁷⁴

Interest in music as a therapeutic intervention in the ICU has been steadily increasing. Thus far, research studies rarely include patients with acute stroke. Moreover, while the effect of playing music for patients in the ICU has resulted in what would be considered favorable outcomes, such as decreased stress, anxiety, and pain, formal examinations have failed to measure the impact of music interventions on neurocritically ill patients' outcomes.³ Despite the lack of an evidence-based consensus for music as therapy for critically ill stroke patients, music interventions are being implemented every day by nurses, physicians, and family members who are trying to optimize care.^{12,13} The Coma Cue-Response framework provides a rationale for determining which interventions aid in recovery from brain injury and prevention of secondary brain injury, as well as providing an understanding of those interventions.⁶⁴ If an intervention, such as playing music, listening to music, or music therapy actually affects patients' outcomes, then that impact must be determined. Only upon doing so can the effect of the particular intervention be controlled as we explore supplementary interventions.

Future Directions and Considerations

Given recent studies that highlight the power of non-pharmacological music intervention in reducing anxiety

in critically ill ICU patients,^{27,29,31} it is reasonable to test the translation of music as an intervention in the acute stroke population. Future studies should be designed with rigor using music interventions that are known to be effective in other populations, such as music therapy. Future goals should be to determine the effectiveness of the music intervention, identify which music genre or type should be used for which indication, dose (duration of exposure to music), and frequency of dosage. Results from these studies will help to empirically determine if music intervention can reduce anxiety, pain, and ultimately sedative exposure, in acute stroke patients who are critically ill or undergoing rehabilitation. If music is proven to be an effective intervention for acute stroke patients, resulting in improved outcomes, including reduced ICU stay and decreased recovery time, then individualized music therapy may be incorporated into the standard of care for stroke patients. **CCN**

Financial Disclosures
None reported.

eLetters

Now that you've read the article, create or contribute to an online discussion about this topic using eLetters. Just visit www.cconline.org and select the article you want to comment on. In the full-text or PDF view of the article, click "Responses" in the middle column and then "Submit a response."

dotmore

To learn more about music therapy, read "Does Music Ease Pain and Anxiety in the Critically Ill?" by Chlan and Halm in the *American Journal of Critical Care*, November 2013;22:528-532. Available at www.ajconline.org.

References

- American Music Therapy Association. Definition and Quotes about Music Therapy: What is Music Therapy? 2014. <http://www.musictherapy.org/about/quotes/>. Accessed January 2016.
- Gutgsell KJ, Schluchter M, Margevicius S, et al. Music therapy reduces pain in palliative care patients: a randomized controlled trial. *J Pain Symptom Manage*. 2013;45(5):822-831.
- Davis T, Jones P. Music therapy: decreasing anxiety in the ventilated patient—a review of the literature. *Dimens Crit Care Nurs*. 2012;31(3):159-166.
- Comeaux T, Comeaux T. The effect of complementary music therapy on the patient's postoperative state anxiety, pain control, and environmental noise satisfaction. *Medsurg Nurs*. 2013;22(5):313-318.
- Korhan EA, Khorshid L, Uyar M. The effect of music therapy on physiological signs of anxiety in patients receiving mechanical ventilatory support. *J Clin Nurs*. 2011;20(7-8):1026-1034.
- Magee WL, Davidson JW. The effect of music therapy on mood states in neurological patients: a pilot study. *J Music Ther*. 2002;39(1):20-29.
- Hunter BC, Oliva R, Sahler OJ, Gaisser D, Salipante DM, Arezina CH. Music therapy as an adjunctive treatment in the management of stress for patients being weaned from mechanical ventilation. *J Music Ther*. 2010;47(3):198-219.
- Walworth D, Rumana CS, Nguyen J, Jarred J. Effects of live music therapy sessions on quality of life indicators, medications administered and hospital length of stay for patients undergoing elective surgical procedures for brain. *J Music Ther*. 2008;45(3):349-359.
- Rodriguez-Fornells A, Rojo N, Amengual JL, Ripolles P, Altenmuller E, Munte TF. The involvement of audio-motor coupling in the music-supported therapy applied to stroke patients. *Ann N Y Acad Sci*. 2012;1252:282-293.
- Ikonomidou E, Rehnstrom A, Naesh O. Effect of music on vital signs and postoperative pain. *AORN J*. 2004;80(2):269-274, 277-268.
- Bae I, Lim HM, Hur MH, Lee M. Intra-operative music listening for anxiety, the BIS index, and the vital signs of patients undergoing regional anesthesia. *Complement Ther Med*. 2014;22(2):251-257.
- Bradt J, Dileo C. Music interventions for mechanically ventilated patients. *Cochrane Database Syst Rev*. 2014;12:CD006902.
- Olson DM, Batjer HH, Zanders ML, Harrison K, Suarez JI. Therapeutic use of music and television in neurocritical care: a practice survey. *J Holist Nurs*. 2015;Apr 6:pii. 0898010115577974.
- Broderick J, Connolly S, Feldmann E, et al. Guidelines for the management of spontaneous intracerebral hemorrhage in adults: 2007 update: a guideline from the American Heart Association/American Stroke Association Stroke Council, High Blood Pressure Research Council, and the Quality of Care and Outcomes in Research Interdisciplinary Working Group. *Stroke*. 2007;38(6):2001-2023.
- Seo W, Oh H. Comparisons of acute physiological parameters influencing outcome in patients with traumatic brain injury and hemorrhagic stroke. *Worldviews Evid Based Nurs*. 2009;6(1):36-43.
- Christensen MC, Mayer S, Ferran JM. Quality of life after intracerebral hemorrhage: results of the Factor Seven for Acute Hemorrhagic Stroke (FAST) trial. *Stroke*. 2009;40(5):1677-1682.
- Zhang Y, Reilly KH, Tong W, et al. Blood pressure and clinical outcome among patients with acute stroke in Inner Mongolia, China. *J Hypertens*. 2008;26(7):1446-1452.
- Jacobi J, Fraser GL, Coursin DB, et al. Clinical practice guidelines for the sustained use of sedatives and analgesics in the critically ill adult. *Crit Care Med*. 2002;30(1):119-141.
- Treggiari MM, Romand JA, Yanez ND, et al. Randomized trial of light versus deep sedation on mental health after critical illness. *Crit Care Med*. 2009;37(9):2527-2534.
- van Rijk MC, Wittekamp BH, Tjan DH, van Zanten AR. Posttraumatic stress disorder-related symptoms after critical care: the role of sedation and family. *Crit Care Med*. 2009;37(5):1831-1833.
- White JM. State of the science of music interventions: critical care and perioperative practice. *Crit Care Nurs Clin North Am*. 2000;12(2):219-225.
- Guzzetta CE. Effects of relaxation and music therapy on patients in a coronary care unit with presumptive acute myocardial infarction. *Heart Lung*. 1989;18(6):609-616.
- Good M. A comparison of the effects of jaw relaxation and music on postoperative pain. *Nurs Res*. 1995;44(1):52-57.
- Chlan LL. Music therapy as a nursing intervention for patients supported by mechanical ventilation. *AACN Clin Issues*. 2000;11(1):128-138.
- Su C-P, Lai H-L, Chang E-T, Yiin L-M, Perng S-J, Chen P-W. A randomized controlled trial of the effects of listening to non-commercial music on quality of nocturnal sleep and relaxation indices in medical intensive care unit. *J Adv Nurs*. 2013;69(6):1377-1389.
- Bradt J, Dileo C, Potvin N. Music for stress and anxiety reduction in coronary heart disease patients. *Cochrane Database Syst Rev*. 2013;12:CD006577.
- Chlan LL, Weinert CR, Heiderscheid A, et al. Effects of patient-directed music intervention on anxiety and sedative exposure in critically ill patients receiving mechanical ventilatory support: a randomized clinical trial. *JAMA*. 2013;309(22):2335-2344.
- Han L, Li JP, Sit JW, Chung L, Jiao ZY, Ma WG. Effects of music intervention on physiological stress response and anxiety level of mechanically ventilated patients in China: a randomised controlled trial. *J Clin Nurs*. 2010;19(7-8):978-987.
- Chlan L. A review of the evidence for music intervention to manage anxiety in critically ill patients receiving mechanical ventilatory support. *Arch Psychiatric Nurs*. 2009;23(2):177-179.
- Chlan LL, Engeland WC, Savik K. Does music influence stress in mechanically ventilated patients? *Intensive Crit Care Nurs*. 2013;29(3):121-127.
- Bradt J, Magee WL, Dileo C, Wheeler BL, McGiloway E. Music therapy for acquired brain injury. *Cochrane Database Syst Rev*. 2010(7):CD006787.
- Sarkamo T, Ripolles P, Vepsäläinen H, et al. Structural changes induced by daily music listening in the recovering brain after middle cerebral artery stroke: a voxel-based morphometry study. *Frontiers Human Neurosci*. 2014;8:245.
- Särkämö T, Soto D. Music listening after stroke: beneficial effects and potential neural mechanisms. *Ann N Y Acad Sci*. 2012;1252(1):266-281.

34. Nightingale F, Skretkovicz V. *Florence Nightingale's Notes on Nursing*. London, England: Scutari Press; 1992.
35. Meehan TC. The careful nursing philosophy and professional practice model. *J Clin Nurs*. 2012;21(19-20):2905-2916.
36. Olson DM, Borel CO, Laskowitz DT, Moore DT, McConnell ES. Quiet time: a nursing intervention to promote sleep in neurocritical care units. *Am J Crit Care*. 2001;10(2):74-78.
37. Dennis CM, Lee R, Woodard EK, Szalaj JJ, Walker CA. Benefits of quiet time for neuro-intensive care patients. *J Neurosc Nurs*. 2010;42(4):217-224.
38. Wuchner SS, Bakas T, Adams G, Buelow J, Cohn J. Nursing interventions and assessments for aneurysmal subarachnoid hemorrhage patients: a mixed methods study involving practicing nurses. *J Neurosci Nurs*. 2012;44(4):177-185.
39. Fachner J, Gold C, Erkkila J. Music therapy modulates fronto-temporal activity in rest-EEG in depressed clients. *Brain Topogr*. 2013;26(2):338-354.
40. Yang CY, Chen CH, Chu H, et al. The effect of music therapy on hospitalized psychiatric patients' anxiety, finger temperature, and electroencephalography: a randomized clinical trial. *Biol Res Nurs*. 2012;14(2):197-206.
41. Mornhinweg GC, Voignier RR. Music for sleep disturbance in the elderly. *J Holist Nurs*. 1995;13(3):248-254.
42. Rauscher FH, Shaw GL, Ky KN. Music and spatial task performance. *Nature*. 1993;365(6447):611.
43. Dunton WR Jr. Recreation and music therapy. *Occup Ther Rehabil*. 1946;25(6):247-252.
44. Thaut MH. Music as therapy in early history. *Progr Brain Res*. 2015;217:143-158.
45. Chlan L, Tracy MF. Music therapy in critical care: indications and guidelines for intervention. *Crit Care Nurse*. 1999;19(3):35-41.
46. Beaulieu-Boire G, Bourque S, Chagnon F, Chouinard L, Gallo-Payet N, Lesur O. Music and biological stress dampening in mechanically-ventilated patients at the intensive care unit ward-a prospective interventional randomized crossover trial. *J Crit Care*. 2013;28(4):442-450.
47. Jaber S, Bahloul H, Guetin S, Chanques G, Sebbane M, Eledjam JJ. [Effects of music therapy in intensive care unit without sedation in weaning patients versus non-ventilated patients]. *Ann Fr Anesth Reanim*. 2007;26(1):30-38.
48. O'Connor SE. Nursing and rehabilitation: the interventions of nurses in stroke patient care. *J Clin Nurs*. 1993;2(1):29-34.
49. Hermans H, Evenhuis HM. Factors associated with depression and anxiety in older adults with intellectual disabilities: results of the healthy ageing and intellectual disabilities study. *Int J Geriatr Psychiatry*. 2013;28(7):691-699.
50. Jolly K, Taylor RS, Lip GY, Stevens A. Home-based cardiac rehabilitation compared with centre-based rehabilitation and usual care: a systematic review and meta-analysis. *Int J Cardiol*. 2006;111(3):343-351.
51. Chemerinski E, Levine SR. Neuropsychiatric disorders following vascular brain injury. *Mt Sinai J Med*. 2006;73(7):1006-1014.
52. El Husseini N, Goldstein LB, Peterson ED, et al. Depression and antidepressant use after stroke and transient ischemic attack. *Stroke*. 2012;43(6):1609-1616.
53. Nuyen J, Spreuwenberg PM, Groenewegen PP, van den Bos GA, Schellevs FG. Impact of preexisting depression on length of stay and discharge destination among patients hospitalized for acute stroke: linked register-based study. *Stroke*. 2008;39(1):132-138.
54. Burvill PW, Johnson GA, Jamrozik KD, Anderson CS, Stewart-Wynne EG, Chakera TM. Anxiety disorders after stroke: results from the Perth Community Stroke Study. *Br J Psychiatry*. 1995;166(3):328-332.
55. Sousa KH, McDermott CM, Weiss J. The importance of assessing symptoms and symptom clusters in critical care units. *AACN Adv Crit Care*. 2011;22(3):275-282.
56. *Diagnostic and Statistical Manual of Mental Disorders (DSM)-5*. Arlington, VA: American Psychiatric Association; 2013.
57. Camara E, Rodriguez-Fornells A, Munte TF. Functional connectivity of reward processing in the brain. *Front Hum Neurosci*. 2008;2:19.
58. Salimpoor VN, Benovoy M, Larcher K, Dagher A, Zatorre RJ. Anatomically distinct dopamine release during anticipation and experience of peak emotion to music. *Nature Neurosci*. 2011;14(2):257-262.
59. Chang SC, Chen CH. Effects of music therapy on women's physiologic measures, anxiety, and satisfaction during cesarean delivery. *Res Nurs Health*. 2005;28(6):453-461.
60. Thaut MH, Gardiner JC, Holmberg D, et al. Neurologic music therapy improves executive function and emotional adjustment in traumatic brain injury rehabilitation. *Ann N Y Acad Sci*. 2009;1169:406-416.
61. Hegde S. Music-based cognitive remediation therapy for patients with traumatic brain injury. *Front Neurol*. 2014;5:34.
62. Sarkamo T, Tervaniemi M, Laitinen S, et al. Music listening enhances cognitive recovery and mood after middle cerebral artery stroke. *Brain*. 2008;131(Pt 3):866-876.
63. Jones AL, Charlesworth JF, Hendra TJ. Patient mood and carer strain during stroke rehabilitation in the community following early hospital discharge. *Disabil Rehabil*. 2000;22(11):490-494.
64. Olson DM, Graffagnino C. Consciousness, coma, and caring for the brain-injured patient. *AACN Clin Issues*. 2005;16(4):441-455.
65. Georgiadis D, Schwarz S, Kollmar R, Baumgartner RW, Schwab S. Influence of inspiration:expiration ratio on intracranial and cerebral perfusion pressure in acute stroke patients. *Intensive Care Med*. 2002;28(8):1089-1093.
66. Bardutzky J, Georgiadis D, Kollmar R, Schwarz S, Schwab S. Energy demand in patients with stroke who are sedated and receiving mechanical ventilation. *J Neurosurg*. 2004;100(2):266-271.
67. Bardutzky J, Georgiadis D, Kollmar R, Schwab S. Energy expenditure in ischemic stroke patients treated with moderate hypothermia. *Intensive Care Med*. 2004;30(1):151-154.
68. Douds GL, Hellkamp AS, Olson DM, et al. Venous thromboembolism in the Get With the Guidelines-Stroke acute ischemic stroke population: incidence and patterns of prophylaxis. *J Stroke Cerebrovasc Dis*. 2014;23(1):123-129.
69. Olson DM, Graffagnino C, King K, Lynch JR. Toward solving the sedation-assessment conundrum: bispectral index monitoring and sedation interruption. *Crit Care Nurs Clin North Am*. 2005;17(3):257-267.
70. Zatorre R, McGill J. Music, the food of neuroscience? *Nature*. 2005;434(7031):312-315.
71. Peretz I, Zatorre RJ. Brain organization for music processing. *Annu Rev Psychol*. 2005;56:89-114.
72. Nilsson U, Rawal N, Unestahl LE, Zetterberg C, Unosson M. Improved recovery after music and therapeutic suggestions during general anaesthesia: a double-blind randomised controlled trial. *Acta Anaesthesiol Scand*. 2001;45(7):812-817.
73. Nilsson U, Unosson M, Rawal N. Stress reduction and analgesia in patients exposed to calming music postoperatively: a randomized controlled trial. *Eur J Anaesthesiol*. 2005;22(2):96-102.
74. Skrobik Y, Ahern S, Leblanc M, Marquis F, Awissi DK, Kavanagh BP. Protocolized intensive care unit management of analgesia, sedation, and delirium improves analgesia and subsyndromal delirium rates. *Anesth Analg*. 2010;111(2):451-463.