

AACN Practice Alert

Pulmonary Artery/Central Venous Pressure Monitoring in Adults

Scope and Impact of the Problem

Pulmonary artery pressure (PAP), pulmonary artery occlusion pressure (PAOP), and central venous pressure (CVP) may aid in the differential diagnosis in pulmonary hypertension¹⁻⁴ and may be beneficial in complex shock states.⁵ Less invasive hemodynamic methods, in conjunction with the patient's medical history, may also be useful in the differential diagnosis of cardiogenic shock.⁶

The appropriate use of pulmonary artery catheter (PAC)-guided therapy is associated with decreased mortality in patients with refractory heart failure/cardiogenic shock⁷ but is not associated with improved outcomes in patients with less severe heart failure. In trauma patients, PAC-guided therapy may benefit older, more severely injured patients, but use of a PAC is not associated with improved outcomes in other trauma populations.^{8,9} Use of a PAC is not associated with improved outcomes in other populations, including high-risk cardiothoracic surgery¹⁰ and general intensive care unit (ICU) patients, although many studies are more than 20 years old.¹¹

The addition of CVP monitoring to guided therapy is not associated with improved outcomes in patients with severe sepsis and septic shock.¹²⁻¹⁶ The PAOP and CVP are not accurate for identifying which patients will respond to a fluid bolus with an increase in stroke volume, and alternative methods such as functional

AACN Levels of Evidence

- Level A** Meta-analysis of quantitative studies or metasynthesis of qualitative studies with results that consistently support a specific action, intervention, or treatment (including systematic review of randomized controlled trials)
- Level B** Well-designed, controlled studies with results that consistently support a specific action, intervention, or treatment
- Level C** Qualitative studies, descriptive or correlational studies, integrative reviews, systematic reviews, or randomized controlled trials with inconsistent results
- Level D** Peer-reviewed professional and organizational standards with the support of clinical study recommendations
- Level E** Multiple case reports, theory-based evidence from expert opinions, or peer-reviewed professional organizational standards without clinical studies to support recommendations
- Level M** Manufacturer's recommendations only

hemodynamic indicators should be used.¹⁷⁻¹⁹ Although the CVP has been identified as a fixed target for resuscitation, debate remains about whether CVP, if used at all, should more appropriately be considered a stopping point.^{20,21}

Use of PACs has decreased, particularly in nonsurgical ICUs.²² This decreased use indicates a need for alternative training methods, such as simulation, to maintain proficiency in PAP monitoring if continued use of PACs is anticipated.

Expected Practice

1. Monitor for complications during catheter insertion (arrhythmias), identify patients at increased risk for pulmonary artery rupture/infarction,



- and implement preventive actions to decrease risk for catheter-induced pulmonary artery rupture or infarction. [level D]
2. Verify the accuracy of the invasive pressure monitoring system by performing a square waveform test at the beginning of each shift and any time the system has been disturbed (eg, when obtaining blood samples). [level D]
 3. Obtain PAP/PAOP/CVP with the patient supine with the head of bed (HOB) elevated between 0° and 60°; with the patient in 20°, 30°, or 90° lateral position with the HOB at 0°; or with the patient prone. Allow the patient to stabilize for 5 to 15 minutes after a position change. Note that a longer stabilization period may be needed if the patient is prone. [level B]
 4. Level and reference the transducer's air/fluid interface to the phlebostatic axis (fourth intercostal space half the diameter of the chest) with the patient supine or prone, or with the patient on his or her side (provided that the correct angle-specific reference is verified by using a laser or carpenter's level before PAP/PAOP/CVP measurements). [level B]
 5. Obtain PAP/PAOP/CVP measurements from a graphic (analog) tracing at end-expiration. [level B] If an analog method is not available, use the stop-cursor method with concurrent airway tracing. [level B] Any automated transfer of hemodynamic data from the monitor to the electronic health record should be verified.
 6. With airway pressure release ventilation (APRV), patients breathe spontaneously around the upper pressure limit; thus, the airway pressure tracing is used to identify spontaneous end-expiration, which occurs immediately before the release of airway pressure and the initiation of inspiration. [level C]
 7. If a patient is actively exhaling or there is marked respiratory excursion (>10-15 mm Hg), measure the PAOP/CVP at a point halfway between the peak and the nadir pressures. [level C]
 8. Use a simultaneous electrocardiographic (ECG) tracing to assist with proper PAP/PAOP/CVP waveform identification. [level C]
 9. PACs can be safely withdrawn and removed by competent registered nurses. [level D]
 10. Accurate alternatives to measuring CVP include measuring peripheral venous pressure (PVP); PVP measurements are obtained from a catheter in the dorsum of the hand or forearm, or from a peripherally inserted central catheter (PICC). [level C]
 11. Continuity from the peripheral venous catheter tip to the central circulation should be verified by checking for an increase in the PVP in response to circumferential occlusion of the arm or leg proximal to the catheter or in response to a sustained inspiratory breath or Valsalva maneuver. [level C]

Supporting Evidence

Monitor for Complications During Catheter Insertion and Use

1. Complications associated with pulmonary artery catheterization are most commonly associated with central venous access; however, approximately 20% of patients experience transient arrhythmias during insertion of a PAC. Right-sided and complete heart block (most likely in patients with preexisting left bundle branch block) occur in 0.3% to 3.8% of patients. Life-threatening complications (pulmonary artery rupture, pulmonary infarction) have been reported in 0.3% to 3% of patients.^{11,23,24}
2. Contraindications for placement of a PAC include presence of a left bundle branch block (risk for complete heart failure), right-sided cardiac mass, tricuspid or pulmonic valve endocarditis, or presence of a mechanical tricuspid or pulmonic valve.²³ Severe tricuspid regurgitation may prevent catheter advancement.²³
3. Caution should be taken in placement and PAOP measurements in patients at increased risk for pulmonary artery rupture (eg, age >60 years, pulmonary hypertension, improper catheter position [catheter insertion >55 cm], cardiopulmonary bypass, and anticoagulation).²³ In addition, incorrect balloon inflation (eg, overwedging or forced inflation) and failure to

deflate the balloon increase the risk for pulmonary artery rupture and thrombosis. In patients at increased risk for pulmonary artery rupture, the end-diastolic pressure in the pulmonary artery should be measured rather than PAOP.

System Assessment to Ensure Accurate Measurement

1. Correct preparation of the pressure monitoring system is required to ensure accurate pressure measurements. The square waveform test, or dynamic response test, determines the ability of the transducer system to correctly reflect invasive pressures.^{25,26} The dynamic response is affected by system problems, such as air bubbles in the tubing, excessive tubing length, loose connections, and catheter patency. Removal of microbubbles during system setup results in an “adequate” or “optimal” system in more than 95% of cases.^{26,27} Any of these problems can affect the accuracy of PAP/PAOP/CVP measurements and must be corrected before pressure measurement. Perform the square waveform test on initial setup, at least once each shift, after opening the catheter system (eg, for zeroing, collecting a blood sample, or changing tubing), and whenever the PAP/PAOP/CVP waveform appears to be damped or distorted.²⁸
2. The addition of a blood conservation device or a needleless access port adversely affects the dynamic response characteristics of invasive pressure monitoring systems.^{26,29,30} The use of a damping device may improve the dynamic response characteristics of the system under these conditions.^{31,32} There are no standards that address the use of blood conservation devices on central catheters, but the effect of devices on the dynamic response characteristics should be considered in any decision.
3. Consider the following changes in PAPs as clinically significant (ie, not reflective of the normal variability in PAP): change in pulmonary artery systolic pressure greater than 4 to 7 mm Hg, change in pulmonary artery end-diastolic pressure (PAEDP) greater than 4 to 7 mm Hg, and change in PAOP greater than 4 mm Hg.³³⁻³⁵ All

hemodynamic values should be interpreted in the context of the patient’s condition. Changes in waveform configuration indicate a need for reassessment of the patient and support the need for documentation of actual waveforms in the patient’s record.

Positioning of Patients

1. Supine: Results of studies in a variety of populations of patients indicate that PAP/PAOP/CVP measurements are comparable when the patient is supine with the HOB elevated to any angle between 0° and 60°, or when the patient is in a 20°, 30°, or 90° lateral position with the HOB flat, as long as the correct angle-specific reference is used.³⁶ PAP/PAOP/CVP measurements obtained with patients in Trendelenburg or reverse Trendelenburg position are not comparable to measurements obtained with the patient supine.³⁷⁻⁴² The response to sitting with the legs in a dependent position is variable, and measurements made with patients in that position may not be comparable to measurements obtained with the patient supine.⁴³⁻⁴⁶ PAP, CVP, and cardiac index⁴⁷ values obtained with the patient in the seated legs-dependent position should be compared with measurements obtained with the patient supine with backrest elevation.
2. Prone: Patients may be placed prone as a part of therapy for acute respiratory distress syndrome (ARDS) or during surgical procedures. In patients with ARDS, if adequate time is allowed for stabilization (30-60 minutes) after repositioning, positioning patients prone does not cause any clinically significant differences in measurements of PAP, CVP, or cardiac output.⁴⁸⁻⁵⁴ However, variable responses in PAP, PAOP, CVP, and cardiac index may occur depending on whether the patient is a fluid responder or nonresponder.⁵⁵⁻⁵⁷
 - a. In studies done to evaluate the effects of prone positioning of patients on hemodynamic indices, the midaxillary line or the midanteroposterior diameter of the chest has been used as the reference point, although no study has been done to validate this

reference point.⁴⁸⁻⁵⁴ A reference point 5 cm vertical distance below the sternal angle may be more accurate with the patient prone.⁵⁷

3. Rotating beds: No studies have been done in patients in automated proning beds. Studies are needed to describe the effects of the prone/lateral rotation with the bed flat and the prone/reverse Trendelenburg position on hemodynamic indices.
4. Reliable measurements of cardiac output have been obtained only with patients positioned supine with the backrest at 20° or 45° and with patients prone. Clinically significant changes in cardiac output may occur in the 20° lateral position, which may limit concurrent measurements of PAP/PAOP/CVP and cardiac output. Because responses to a given position vary from patient to patient, use a standardized approach to evaluate each patient's PAP/PAOP/CVP compared with measurements obtained with the patient in the flat, supine position.

Waveform Assessment and Interpretation

1. Correct and consistent waveform interpretation is required to ensure accurate pressure measurements. A simultaneous ECG should be used to facilitate correct PAP/PAOP/CVP waveform measurement⁵⁸⁻⁶⁰ and should be read from an analog tracing or the stop cursor method, if an analog tracing is not available.⁶¹⁻⁶³
2. Digital readouts should not be used because they reflect pressures obtained throughout respiration and may be significantly different from end-expiratory pressures.⁶⁴⁻⁶⁶
3. Registered nurses who demonstrate competency can safely withdraw and/or remove PACs. Before incorporating withdrawing and/or removing PACs into nursing practice, verify that it is in your state's scope of practice for registered nurses.⁶⁷

Alternative Sites for Monitoring

1. Peripheral venous pressure (PVP), which is obtained from a catheter in the dorsum of the hand, forearm, or lower extremity, may be an accurate alternative to CVP. In a variety of populations of patients receiving mechanical

ventilation and spontaneously breathing patients (neurosurgical, laparoscopic, colorectal, cardiac, kidney and hepatic transplant, burns),⁶⁸⁻⁷⁵ and a variety of positions (supine, Fowler, Trendelenburg, prone),^{76,77} the PVP was higher than the CVP by approximately 2 or 3 mm Hg. Research in the postoperative and ICU setting is limited. In all positions, continuity from the catheter tip to central circulation must be confirmed by evaluation of an increase in the PVP in response to circumferential occlusion of the arm or leg proximal to the catheter,^{70,78} a sustained inspiratory breath,^{76,78} or a Valsalva maneuver.^{71,77} If continuity is absent (no increase in PVP in response to test), the PVP cannot be used and the catheter should be assessed for patency.^{71,78} The arm and shoulder should be placed in a neutral position, if possible.⁷⁹

2. Femoral venous pressure-CVP. Femoral (inferior vena caval or iliac venous) venous pressure agrees with the CVP if abdominal pressure is low.⁸⁰⁻⁸² If abdominal pressure (eg, bladder pressure) is increased (≥ 15 mm Hg), the femoral pressure will correlate with abdominal pressure, but will not agree with CVP.⁸³⁻⁸⁵ If using femoral venous/inferior vena caval pressure, confirm that intra-abdominal pressure is less than 15 mm Hg. The difference between CVP and femoral venous pressure is not related to catheter type (introducer, multilumen)⁸⁰ but is affected by HOB elevation (recommended to position the patient supine-flat or $<30^\circ$ HOB elevation during pressure measurements).⁸⁶
3. The CVP can be obtained from a PICC. A continuous flow device (3 mL/h) should be used. The PICC should be referenced at the phlebostatic axis.⁸⁷⁻⁹¹ A shallow slope following a fast flush suggests PICC occlusion.⁸⁸ The CVP obtained from PICCs that support high-pressure infusions such as infusions of radiographic contrast media is also accurate.⁸⁹

Effect of Ventilation on Measurement

1. In patients with pulmonary hypertension, a PAOP greater than 15 mm Hg is an adequate predictor of a left ventricular end-diastolic

pressure (LVEDP) exceeding 15 mm Hg; although absolute values of PAOP and LVEDP may differ 3 to 7 mm Hg on average.⁹² In patients with precapillary pulmonary hypertension (mean PAP \geq 25 mm Hg with normal left ventricular function), the PAOP may be an overestimate of the LVEDP, especially in obese patients or patients with chronic obstructive pulmonary disease.⁹³⁻⁹⁵ When measuring the PAOP, use end-expiratory analog measurement⁶⁶ and ensure complete occlusion without overwedging (eg, PAOP < PAEDP; atrial pressure waveform, free flow within catheter to confirm the tip is not impacted on the vessel wall, highly oxygenated blood [capillary] is obtained from distal port with balloon inflated).^{96,97}

2. Pressure support ventilation (PSV): During PSV, use the proximal airway pressure to identify end-expiration, as the type of ventilation (spontaneous, pressure control) varies depending on the level of pressure support being provided. In one study,⁶⁴ conversion from pressure control ventilation to a spontaneous respiratory pattern occurred at PSV of about 12 cm H₂O.
3. Airway pressure release ventilation (APRV): The effect of APRV on hemodynamic measurements has been evaluated in only 1 study (animal model).⁹⁸ The PAP and left atrial pressure agreed at continuous positive airway pressures from 10 to 35 cm H₂O. Both the PAOP and left atrial pressure increased as the airway pressure increased, which suggests a need to correct cardiac pressures for increased airway pressure (> 10 cm H₂O). No research on this topic has been done in humans; however, patients on APRV breathe spontaneously around the upper pressure limit; thus, the airway pressure tracing may be used to identify spontaneous end-expiration, which occurs immediately before the release of airway pressure and the initiation of inspiration.
4. Active exhalation/large respiratory swings. Debate persists on how to interpret PAP/PAOP in patients with active exhalation or large respiratory excursions (eg, respiratory-induced fluctuation in PAOP >10-15 mm Hg). Under

these conditions, end-expiratory pressures are overestimates of the true PAP/PAOP.⁹⁹ With active exhalation/large respiratory excursions, the PAP/PAOP should be consistently read at the midpoint between the peak and the nadir pressures⁹⁹ or averaged over 3 ventilatory cycles.¹⁰⁰

Implementation/Organizational Support for Practice

Always identify and mark the phlebostatic axis or appropriate angle-specific reference.

Measure PAP/PAOP/CVP with the patient supine (HOB elevated between 0° and 60°), in lateral position (20°, 30°, or 90°), or prone.

Perform a square waveform test at appropriate intervals as determined by your unit protocol.

Read pressures from an analog recording at end-expiration. However, if analog recording is not available, use a stop-cursor method with a concurrent ECG and respiratory waveform to identify end-expiration.

Confirm accuracy of hemodynamic data transferred from the monitor to the electronic health record and provide documentation of waveform recordings to allow evaluation of changes in waveform patterns.

Need More Information or Help?

1. Contact a clinical practice specialist for additional information: go to www.aacn.org then select Practice Resource Network.
2. AACN Hemodynamic Management Pocket Reference Card. Product 400751. <http://www.aacn.org/dm/Products/ProductDetails.aspx?productid=400751&prodtyp=product&subsystem=ORD>.
3. Lough ME. *Hemodynamic Monitoring: Evolving Technologies and Clinical Practice*. St Louis, MO: Elsevier; 2016.

*Original Author: Elizabeth Bridges, RN, PhD, CCNS
May 2004*

*Contributing Authors: Lia Barros, RN, BSN, CCRN, Elizabeth Bridges, RN, PhD, CCNS, Marie Cockerham, RN, BSN, CCRN, Sheryl Greco, RN, MN, CCRN, Farah Herrera, RN, BSN, CCRN, and Nicole Solvang, RN, BSN CCRN
December 2015*

Reviewed and approved by the AACN Clinical Resources Task Force, December 2015

Financial Disclosures
None reported.

References

- Vincent JL, Rhodes A, Perel A, et al. Clinical review: update on hemodynamic monitoring—a consensus of 16. *Crit Care*. 2011;15:229.
- Leuchte HH, Baezner CJ, Baumgartner RA, Mernitz P, Neurohr C, Behr J. Acute hemodynamic responses to supplemental oxygen and their prognostic implications in pulmonary hypertension. *Respiration*. 2013;85:400-407.
- Grignola JC. Hemodynamic assessment of pulmonary hypertension. *World J Cardiol*. 2011;3:10-17.
- Maron BA. Hemodynamics should be the primary approach to diagnosing, following, and managing pulmonary arterial hypertension. *Can J Cardiol*. 2015;31:515-520.
- De Backer D, Fagnoul D, Herpain A. The role of invasive techniques in cardiopulmonary evaluation. *Curr Opin Crit Care*. 2013;19:228-233.
- Cooper HA, Najafi AH, Ghafourian K, et al. Diagnosis of cardiogenic shock without the use of a pulmonary artery catheter. *Eur Heart J Acute Cardiovasc Care*. 2015;4:88-95.
- Sotomi Y, Sato N, Kajimoto K, et al. Impact of pulmonary artery catheter on outcome in patients with acute heart failure syndromes with hypotension or receiving inotropes: from the ATTEND Registry. *Int J Cardiol*. 2014;172:165-172.
- Friese RS, Shafi S, Gentilello LM. Pulmonary artery catheter use is associated with reduced mortality in severely injured patients: a National Trauma Data Bank analysis of 53,312 patients. *Crit Care Med*. 2006;34:1597-1601.
- Barmparas G, Inaba K, Georgiou C, et al. Swan-Ganz catheter use in trauma patients can be reduced without negatively affecting outcomes. *World J Surg*. 2011;35:1809-1817.
- Chiang Y, Hosseinian L, Rhee A, Itagaki S, Cavallaro P, Chikwe J. Questionable benefit of the pulmonary artery catheter after cardiac surgery in high-risk patients. *J Cardiothorac Vasc Anesth*. 2015;29:76-81.
- Rajaram SS, Desai NK, Kalra A, et al. Pulmonary artery catheters for adult patients in intensive care. *Cochrane Database Syst Rev*. 2013;2:CD003408.
- ProCESS Investigators, Yealy DM, Kellum JA, et al. A randomized trial of protocol-based care for early septic shock. *N Engl J Med*. 2014;370:1683-1693.
- ARISE Investigators, ANZICS Clinical Trials Group, Peake SL, et al. Goal-directed resuscitation for patients with early septic shock. *N Engl J Med*. 2014;371:1496-1506.
- Mouncey PR, Osborn TM, Power GS, et al. Trial of early, goal-directed resuscitation for septic shock. *N Engl J Med*. 2015;372:1301-1311.
- Angus DC, Barnato AE, Bell D, et al. A systematic review and meta-analysis of early goal-directed therapy for septic shock: the ARISE, ProCESS and ProMISe Investigators. *Intensive Care Med*. 2015;41(9):1549-1560.
- Rusconi AM, Bossi I, Lampard JG, Szava-Kovats M, Bellone A, Lang E. Early goal-directed therapy vs usual care in the treatment of severe sepsis and septic shock: a systematic review and meta-analysis. *Intern Emerg Med*. 2015;10(6):731-743.
- Cherpanath TG, Aarts LP, Groeneveld JA, Geerts BF. Defining fluid responsiveness: a guide to patient-tailored volume titration. *J Cardiothorac Vasc Anesth*. 2014;28:745-754.
- Marik PE, Cavallazzi R. Does the central venous pressure predict fluid responsiveness? An updated meta-analysis and a plea for some common sense. *Crit Care Med*. 2013;41:1774-1781.
- Sondergaard S, Parkin G, Aneman A. Central venous pressure: we need to bring clinical use into physiological context. *Acta Anaesthesiol Scand*. 2015;59:552-560.
- Pinsky MR, Kellum JA, Bellomo R. Central venous pressure is a stopping rule, not a target of fluid resuscitation. *Crit Care Resusc*. 2014;16:245-246.
- Polderman KH, Varon J, Marik PE. Fluid management decisions should not be guided by fixed central venous pressure targets. *Am J Emerg Med*. 2015;33:1311.
- Gershengorn HB, Wunsch H. Understanding changes in established practice: pulmonary artery catheter use in critically ill patients. *Crit Care Med*. 2013;41:2667-2676.
- Evans DC, Doraiswamy VA, Prosciak MP, et al. Complications associated with pulmonary artery catheters: a comprehensive clinical review. *Scand J Surg*. 2009;98:199-208.
- Practice guidelines for pulmonary artery catheterization: an updated report by the American Society of Anesthesiologists Task Force on Pulmonary Artery Catheterization. *Anesthesiology*. 2003;99:988-1014.
- Bridges E, Middleton R. Direct arterial vs oscillometric monitoring of blood pressure: stop comparing and pick one (a decision-making algorithm). *Crit Care Nurse*. 1997;17:58-72.
- Bridges E, Evers KG, Schmelz J, Woods S, Harrington E. Invasive pressure monitoring at altitude. *Crit Care Med*. 2005;33:A13.
- Bridges EJ. Pulmonary artery pressure monitoring: when, how, and what else to use. *AACN Adv Crit Care*. 2006;17:286-303.
- Gardner R. Direct blood pressure measurement: dynamic response requirements. *Anesthesiology*. 1981;54:227-236.
- Melamed R, Johnson K, Pothen B, Sprenkle MD, Johnson PJ. Invasive blood pressure monitoring systems in the ICU: influence of the blood-conserving device on the dynamic response characteristics and agreement with noninvasive measurements. *Blood Press Monit*. 2012;17:179-183.
- Gore S, Middleton R, Bridges E. Analysis of an algorithm to guide decision making regarding direct and oscillometric blood pressure measurement [abstract]. *Am J Resp Crit Care Med*. 1995;151:A331.
- Fujiwara S, Kawakubo Y, Mori S, Tachihara K, Toyoguchi I, Yokoyama T. Effect of Planecta and ROSE on the frequency characteristics of blood pressure-transducer kits. *J Clin Monit Comput*. 2015;29:681-689.
- Fujiwara S, Tachihara K, Mori S, et al. Effect of using a Planecta port with a three-way stopcock on the natural frequency of blood pressure transducer kits [published online October 14, 2015]. *J Clin Monit Comput*. doi:10.1007/s10877-015-9795-3.
- Nemens EJ, Woods SL. Normal fluctuations in pulmonary artery and pulmonary capillary wedge pressure in acutely ill patients. *Heart Lung*. 1982;11:393-398.
- Moser D, Woo M. Normal fluctuation in pulmonary artery pressure and cardiac output in patients with severe left ventricular dysfunction [abstract]. *Am J Crit Care*. 1996;5:236.
- Bridges EJ, Woods SL, Brengelmann GL, Mitchell P, Laurent-Bopp D. Effect of the 30 degree lateral recumbent position on pulmonary artery and pulmonary artery wedge pressures in critically ill adult cardiac surgery patients. *Am J Crit Care*. 2000;9:262-275.
- Rauen CA, Makic MB, Bridges E. Evidence-based practice habits: transforming research into bedside practice. *Crit Care Nurse*. 2009;29:46-59.
- Yoneda G, Katagiri S, Yamamoto M. Reverse Trendelenburg position is a safer technique for lowering central venous pressure without decreasing blood pressure than clamping of the inferior vena cava below the liver. *J Hepatobiliary Pancreat Sci*. 2015;22:463-466.
- Mekis D, Kamenik M. Influence of body position on hemodynamics in patients with ischemic heart disease undergoing cardiac surgery. *Wien Klin Wochenschr*. 2010;122(suppl 2):59-62.
- Lestar M, Gunnarsson L, Lagerstrand L, Wiklund P, Odeberg-Wernerman S. Hemodynamic perturbations during robot-assisted laparoscopic radical prostatectomy in 45 degrees Trendelenburg position. *Anesth Analg*. 2011;113:1069-1075.
- Rosendal C, Markin S, Hien MD, Motsch J, Roggenbach J. Cardiac and hemodynamic consequences during capnoperitoneum and steep Trendelenburg positioning: lessons learned from robot-assisted laparoscopic prostatectomy. *J Clin Anesth*. 2014;26:383-389.
- van Lieshout JJ, Harms MP, Pott F, Jenstrup M, Secher NH. Stroke volume of the heart and thoracic fluid content during head-up and head-down tilt in humans. *Acta Anaesthesiol Scand*. 2005;49:1287-1292.
- Reuter DA, Felbinger TW, Schmidt C, et al. Trendelenburg positioning after cardiac surgery: effects on intrathoracic blood volume index and cardiac performance. *Eur J Anaesthesiol*. 2003;20:17-20.
- Woods SL, Mansfield LW. Effect of body position upon pulmonary artery and pulmonary capillary wedge pressures in noncritically ill patients. *Heart Lung*. 1976;5:83-90.
- Fournier P, Mensch-Dechene J, Ranson-Bitker B, Valladares W, Lockhart A. Effects of sitting up on pulmonary blood pressure, flow, and volume in man. *J Appl Physiol Respir Environ Exerc Physiol*. 1979;46:36-40.
- Thadani U, Parker JO. Hemodynamics at rest and during supine and sitting bicycle exercise in normal subjects. *Am J Cardiol*. 1978;41:52-59.
- Thadani U, West RO, Mathew TM, Parker JO. Hemodynamics at rest and during supine and sitting bicycle exercise in patients with coronary artery disease. *Am J Cardiol*. 1977;39:776-783.
- Rader C, Nelson M, Sobek C, et al. Cardiac index based on measurements obtained in a bedside chair and in bed. *Am J Crit Care*. 2011;20:210-215.
- Blanch I, Mancebo J, Perez M, et al. Short-term effects of prone position in critically ill patients with acute respiratory distress syndrome. *Intensive Care Med*. 1997;23:1033-1039.
- Joliet P, Bulpa P, Chevrolet JC. Effects of the prone position on gas exchange and hemodynamics in severe acute respiratory distress syndrome. *Crit Care Med*. 1998;26:1977-1985.
- Vollman K, Bander J. Improved oxygenation using a prone positioner in patients with acute respiratory distress syndrome. *Intensive Care Med*. 1996;22:1105-1111.
- Hering R, Wrigge H, Vorwerk R, et al. The effects of prone positioning on intraabdominal pressure and cardiovascular and renal function in patients with acute lung injury. *Anesth Analg*. 2001;92:1226-1231.
- Hering R, Vorwerk R, Wrigge H, et al. Prone positioning, systemic hemodynamics, hepatic indocyanine green kinetics, and gastric intramucosal energy balance in patients with acute lung injury. *Intensive Care Med*. 2002;28:53-58.

53. Matejovic M, Rokyta R, Jr., Radermacher P, Krouzicky A, Sramek V, Novak I. Effect of prone position on hepato-splanchnic hemodynamics in acute lung injury. *Intensive Care Med.* 2002;28:1750-1755.
54. Borelli M, Lampati L, Vascotto E, Fumagalli R, Pesenti A. Hemodynamic and gas exchange response to inhaled nitric oxide and prone positioning in acute respiratory distress syndrome patients. *Crit Care Med.* 2000;28:2707-2712.
55. Jozwiak M, Teboul JL, Anguel N, et al. Beneficial hemodynamic effects of prone positioning in patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med.* 2013;188:1428-1433.
56. Wu CY, Lee TS, Chan KC, Jeng CS, Cheng YJ. Does targeted pre-load optimisation by stroke volume variation attenuate a reduction in cardiac output in the prone position. *Anaesthesia.* 2012;67:760-764.
57. Magder S. Is all on the level? Hemodynamics during supine versus prone ventilation. *Am J Respir Crit Care Med.* 2013;188:1390-1391.
58. Lipp-Ziff E, Kawanishi D. A technique for improving the accuracy of the pulmonary artery diastolic pressure as an estimate of left ventricular end-diastolic pressure. *Heart Lung.* 1991;20:107-115.
59. Magder S. How to use central venous pressure measurements. *Curr Opin Crit Care.* 2005;11:264-270.
60. Bridges EJ. Monitoring pulmonary artery pressures: just the facts. *Crit Care Nurse.* 2000;20:59-78.
61. Johnson MK, Schumann L. Comparison of three methods of measurement of pulmonary artery catheter readings in critically ill patients. *Am J Crit Care.* 1995;4:300-307.
62. Lundstedt JL. Comparison of methods of measuring pulmonary artery pressure. *Am J Crit Care.* 1997;6:324-332.
63. Ahrens TS, Schallom L. Comparison of pulmonary artery and central venous pressure waveform measurements via digital and graphic measurement methods. *Heart Lung.* 2001;30:26-38.
64. Wong FW. Where is end expiration? Measuring PAWP when the patient is on pressure support ventilation. *Dynamics (Pembroke, Ont).* 2010;21:11-16.
65. Pasion E, Good L, Tizon J, et al. Evaluation of the monitor cursor-line method for measuring pulmonary artery and central venous pressures. *Am J Crit Care.* 2010;19:511-521.
66. Ryan JJ, Rich JD, Thiruvoipati T, Swamy R, Kim GH, Rich S. Current practice for determining pulmonary capillary wedge pressure predisposes to serious errors in the classification of patients with pulmonary hypertension. *Am Heart J.* 2012;163:589-594.
67. Epping K, Hammer B. Removal of a pulmonary artery catheter in the presence of implanted leads. *Crit Care Nurse.* 2012;32:69-71.
68. Amar D, Melendez JA, Zhang H, Dobres C, Leung DH, Padilla RE. Correlation of peripheral venous pressure and central venous pressure in surgical patients. *J Cardiothorac Vasc Anesth.* 2001;15:40-43.
69. Kim SH, Park SY, Cui J, et al. Peripheral venous pressure as an alternative to central venous pressure in patients undergoing laparoscopic colorectal surgery. *Br J Anaesth.* 2011;106:305-311.
70. Munis JR, Bhatia S, Lozada LJ. Peripheral venous pressure as a hemodynamic variable in neurosurgical patients. *Anesth Analg.* 2001;92:172-179.
71. Bombardieri AM, Beckman J, Shaw P, Girardi FP, Ma Y, Memtsoudis SG. Comparative utility of centrally versus peripherally transduced venous pressure monitoring in the perioperative period in spine surgery patients. *J Clin Anesth.* 2012;24:542-548.
72. Tugrul M, Camci E, Pembeci K, Al-Darsani A, Telci L. Relationship between peripheral and central venous pressures in different patient positions, catheter sizes, and insertion sites. *J Cardiothorac Vasc Anesth.* 2004;18:446-450.
73. Hadimioglu N, Ertug Z, Yegin A, Sanli S, Gurkan A, Demirbas A. Correlation of peripheral venous pressure and central venous pressure in kidney recipients. *Transplant Proc.* 2006;38:440-442.
74. Hoftman N, Braunfeld M, Hoftman G, Mahajan A. Peripheral venous pressure as a predictor of central venous pressure during orthotopic liver transplantation. *J Clin Anesth.* 2006;18:251-255.
75. Sherif L, Joshi VS, Ollapally A, Jain P, Shetty K, Ribeiro KS. Peripheral venous pressure as a reliable predictor for monitoring central venous pressure in patients with burns. *Indian J Crit Care Med.* 2015;19:199-202.
76. Tobias JD. Measurement of central venous pressure from a peripheral intravenous catheter in the prone position during spinal surgery. *South Med J.* 2009;102:256-259.
77. Memtsoudis SG, Jules-Elyse K, Girardi FP, et al. Correlation between centrally versus peripherally transduced venous pressure in prone patients undergoing posterior spine surgery. *Spine (Phila Pa 1976).* 2008;33:E643-E647.
78. Cox P, Johnson J, Tobias J. Measurement of central venous pressure from a peripheral intravenous catheter in the lower extremity. *South Med J.* 2005;98:698-702.
79. Stoneking L, Deluca LA Jr, Fiorello AB, Munzer B, Baker N, Denninghoff KR. Alternative methods to central venous pressure for assessing volume status in critically ill patients. *J Emerg Nurs.* 2014;40:115-123.
80. Lee BK, Lee HY, Jeung KW, Jung YH, Lee GS. Estimation of central venous pressure using inferior vena caval pressure from a femoral endovascular cooling catheter. *Am J Emerg Med.* 2013;31:240-243.
81. Boone BA, Kirk KA, Tucker N, Gunn S, Forsythe R. Iliac venous pressure estimates central venous pressure after laparotomy. *J Surg Res.* 2014;191:203-207.
82. Walsh JT, Hildick-Smith DJ, Newell SA, Lowe MD, Satchithananda DK, Shapiro LM. Comparison of central venous and inferior vena caval pressures. *Am J Cardiol.* 2000;85:518-520, A11.
83. Ait-Oufella H, Boelle PY, Galbois A, et al. Comparison of superior vena cava and femoriliac vein pressure according to intra-abdominal pressure. *Ann Intensive Care.* 2012;2:21.
84. Groombridge CJ, Duplooy D, Adams BD, Paul E, Butt W. Comparison of central venous pressure and venous oxygen saturation from venous catheters placed in the superior vena cava or via a femoral vein: the numbers are not interchangeable. *Crit Care Resusc.* 2011;13:151-155.
85. Markou N, Grigorakos L, Myriantefs P, et al. Venous pressure measurements in the superior and inferior vena cava: the influence of intra-abdominal pressure. *Hepatogastroenterology.* 2004;51:51-55.
86. Pacheco S, Machado MN, Amorim RC, et al. Central venous pressure in femoral catheter: correlation with superior approach after heart surgery. *Rev Bras Cir Cardiovasc.* 2008;23:488-493.
87. Yun JY, Park SH, Cho DS, Jeung HJ, Lee SA, Seo SJ. Comparison of the central venous pressure from internal jugular vein and the pressure measured from the peripherally inserted antecubital central catheter (PICCP) in liver transplantation recipients. *Korean J Anesthesiol.* 2011;61:281-287.
88. Black IH, Blosser SA, Murray WB. Central venous pressure measurements: peripherally inserted catheters versus centrally inserted catheters. *Crit Care Med.* 2000;28:3833-3836.
89. Latham HE, Rawson ST, Dwyer TT, Patel CC, Wick JA, Simpson SQ. Peripherally inserted central catheters are equivalent to centrally inserted catheters in intensive care unit patients for central venous pressure monitoring. *J Clin Monit Comput.* 2012;26:85-90.
90. McLemore EC, Tessier DJ, Rady MY, et al. Intraoperative peripherally inserted central venous catheter central venous pressure monitoring in abdominal aortic aneurysm reconstruction. *Ann Vasc Surg.* 2006;20:577-581.
91. Alansari M, Hijazi M. Central venous pressure from peripherally inserted central catheters correlates well with that of centrally inserted central catheters [abstract]. *Chest.* 2004;126:873s.
92. Halpern SD, Taichman DB. Misclassification of pulmonary hypertension due to reliance on pulmonary capillary wedge pressure rather than left ventricular end-diastolic pressure. *Chest.* 2009;136:37-43.
93. Bitar A, Selej M, Bolad I, Lahm T. Poor agreement between pulmonary capillary wedge pressure and left ventricular end-diastolic pressure in a veteran population. *PLoS One.* 2014;9:e87304.
94. LeVarge BL, Pomerantsev E, Channick RN. Reliance on end-expiratory wedge pressure leads to misclassification of pulmonary hypertension. *Eur Respir J.* 2014;44:425-434.
95. de Oliveira RK, Ferreira EV, Ramos RP, et al. Usefulness of pulmonary capillary wedge pressure as a correlate of left ventricular filling pressures in pulmonary arterial hypertension. *J Heart Lung Transplant.* 2014;33:157-162.
96. Leatherman JW, Shapiro RS. Overestimation of pulmonary artery occlusion pressure in pulmonary hypertension due to partial occlusion. *Crit Care Med.* 2003;31:93-97.
97. Morris AH, Chapman RH, Gardner RM. Frequency of technical problems encountered in the measurement of pulmonary artery wedge pressure. *Crit Care Med.* 1984;12:164-170.
98. Slim AM, Martinho S, Slim J, Davenport E, Castillo-Rojas LM, Shry EA. The effect of airway pressure release ventilation on pulmonary catheter readings: specifically pulmonary capillary wedge pressure in a swine model [published online February 24, 2011]. *Anesthesiol Res Pract.* 2011;2011:371594. doi:10.1155/2011/371594.
99. Hoyt JD, Leatherman JW. Interpretation of the pulmonary artery occlusion pressure in mechanically ventilated patients with large respiratory excursions in intrathoracic pressure. *Intensive Care Med.* 1997;23:1125-1131.
100. Boerrigter BG, Waxman AB, Westerhof N, Vonk-Noordegraaf A, Systrom DM. Measuring central pulmonary pressures during exercise in COPD: how to cope with respiratory effects. *Eur Respir J.* 2014;43:1316-1325.