

COMMENTS AND RESPONSES

**Comment on:
Beverly et al. Do
Older Adults Aged
60–75 Years
Benefit From
Diabetes
Behavioral
Interventions?
Diabetes Care
2013;36:1501–
1506**

We wish to support Beverly et al. in confirming the efficacy of small-group education in older patients with diabetes (1). However, their suggestion that the majority of education programs are designed for younger/more recently diagnosed patients and that minimal attention has been paid to the development of successful interventions for older adults with diabetes is at conflict with much recent (2), and less recent (3), literature.

We have been running small-group education in individuals with type 2 and

type 1 diabetes for many years and have reported favorable lifestyle changes, improved health behaviors, and lower HbA_{1c} in both (4,5). We concur that, apparently, better results are observed when education is offered to older patients. Possibly, this is because interventions are centered on lifestyle issues whereas a more technical approach, including carbohydrate counting and self-monitoring, is necessary to improve clinical outcomes in individuals with type 1 diabetes (5). In either case, continuous reinforcement over the years is key to sustained change and improvement.

Health professionals are to realize that, similarly to glucose-lowering agents, education should be a lifelong treatment if it is to display its value as an instrument of change and self-care. One-off or even relatively short-term interventions do not provide lasting benefits and will lead to the flawed conclusion that education is ineffective.

MARINA TRENTO, MEDSCI, MBA¹
PIETRO PASSERA, MD¹
FRANCO CAVALLO, MD²
MASSIMO PORTA, MD, PHD¹

From the ¹Laboratory of Clinical Pedagogy, Department of Internal Medicine, University of Turin, Turin, Italy; and the ²Department of Public Health and Microbiology, University of Turin, Turin, Italy. Corresponding author: Marina Trento, marina.trento@unito.it.
DOI: 10.2337/dc13-0241

© 2013 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Acknowledgments—No potential conflicts of interest relevant to this article were reported.

References

1. Beverly EA, Fitzgerald S, Sitnikov L, Ganda OP, Caballero AE, Weinger K. Do older adults aged 60–75 years benefit from diabetes behavioral interventions? *Diabetes Care* 2013;36:1501–1506
2. Steinsbekk A, Rygg LØ, Lisulo M, Rise MB, Fretheim A. Group based diabetes self-management education compared to routine treatment for people with type 2 diabetes mellitus. A systematic review with meta-analysis. *BMC Health Serv Res* 2012;12:213
3. Norris SL, Engelgau MM, Narayan KM. Effectiveness of self-management training in type 2 diabetes: a systematic review of randomized controlled trials. *Diabetes Care* 2001;24:561–587
4. Trento M, Gamba S, Gentile L, et al.; ROMEO Investigators. Rethink Organization to improve Education and Outcomes (ROMEO): a multicenter randomized trial of lifestyle intervention by group care to manage type 2 diabetes. *Diabetes Care* 2010;33:745–747
5. Trento M, Borgo E, Kucich C, et al. Quality of life, coping ability, and metabolic control in patients with type 1 diabetes managed by group care and a carbohydrate counting program. *Diabetes Care* 2009;32:e134