

OBSERVATIONS

# No Difference in Vitamin D Levels Between Children Newly Diagnosed With Type 1 Diabetes and Their Healthy Siblings: A 13-Year Nationwide Danish Study

The causes of the worldwide increase in type 1 diabetes (T1D) in children are still largely unknown. In Denmark, the increase in incidence is steep at 3.4% annually and does not appear to be leveling out (1). Lower levels of 25-hydroxyvitamin D [25(OH)D] have been found in patients with newly diagnosed T1D compared with healthy control subjects, implying that 25(OH)D might play a role in the pathogenesis of T1D (2,3). We aimed to elucidate the possible association between low levels of 25(OH)D and T1D by measuring 25(OH)D levels in children with newly diagnosed T1D and their healthy siblings in Denmark across a 13-year study period (1997–2009).

We included 1,803 children (907 T1D patients and 896 siblings) in the data analyses. The children were aged 0–18 years (mean [SEM] 10.6 [0.13] years for patients and 9.9 [0.12] years for siblings). Of the 1,803 children included, 859 were females and 944 were males. Siblings were matched with the T1D patients for age, sex, month of sampling, and sample year.

Blood was sampled within 3 months after onset.

Vitamin D status was measured as serum 25(OH)D by high-performance liquid chromatography, and parathyroid hormone (PTH) level was measured with Siemens Immulite 2000. Table 1 shows the geometric mean and range of 25(OH)D, PTH, and calcium levels.

In the univariate model, 25(OH)D levels were 2.6% (95% CI –3.2 to 8.9%,  $P = 0.38$ ) higher in patients than in siblings, and in the adjusted model, levels were –5.2% (–14.9 to 3.4%,  $P = 0.22$ ) lower in patients. When sample year was forced as a linear variable, there was no sign of a decrease in 25(OH)D per year in patients or siblings, whereas when year was tested as a categorical variable in a joint model of patients and siblings, 25(OH)D varied significantly by year, with the highest levels in 1997 and in 2004–2006 ( $P = 0.02$ ). The PTH levels were not significantly different in patients and siblings in either the univariate analysis (–7.0% [–15.0 to 1.8%],  $P = 0.12$ ) or the adjusted model (–5.3% [–13.5 to 3.7%],  $P = 0.24$ ).

The present findings are somewhat contradictory to previous studies of 25(OH)D levels in patients with newly onset T1D (2,3) and years after (4). In contrast with other studies, we compared patients with siblings as a way to eliminate genetic confounding but which could result in overmatching on environmental factors (e.g., exposure to sunlight, vitamin supplementation, and exercise). The present study is larger than prior studies, where 25(OH)D levels were found to be 15–43% lower in patients (2–4). PTH is a valuable indicator of vitamin D status, and because PTH levels mirrored 25(OH)D levels in both patients and siblings, it supports the main finding.

In summary, the results show year-to-year variation in 25(OH)D levels, probably as a result of change in the “hours of bright sunshine” (5), and no association between 25(OH)D levels and T1D around onset in children and adolescents. The role of 25(OH)D levels in utero or in early childhood and a later risk for developing T1D needs further investigation.

STEFFEN U. THORSEN, MD<sup>1</sup>  
 HENRIK B. MORTENSEN, DMSC<sup>1</sup>  
 BENDIX CARSTENSEN, MSC<sup>2</sup>  
 MOGENS FENGER, DMSC<sup>3</sup>  
 BETINA H. THUESEN, MSC, PHD<sup>4</sup>  
 LISE LOTTE HUSEMOEN, MSC, PHD<sup>4</sup>  
 REGINE BERGHOLDT, DMSC<sup>5</sup>  
 CAROLINE BRORSSON, MSC<sup>6</sup>  
 FLEMMING POCIOT, MD<sup>2,5,7</sup>  
 ALLAN LINNEBERG, MD, PHD<sup>4</sup>  
 JANNET SVENSSON, MD, PHD<sup>1</sup>

From the <sup>1</sup>Herlev University Hospital, Herlev, Denmark; the <sup>2</sup>Steno Diabetes Center, Gentofte, Denmark; the <sup>3</sup>Department of Clinical Biochemistry and Molecular Biology, Hvidovre Hospital, Hvidovre, Denmark; the <sup>4</sup>Research Centre for Prevention and Health, Glostrup University Hospital, Glostrup, Denmark; the <sup>5</sup>Hagedorn Research Institute, Gentofte, Denmark; the <sup>6</sup>Glostrup Research Institute, Glostrup University Hospital, Glostrup, Denmark; and the <sup>7</sup>Department of Biomedical Science, University of Copenhagen, Copenhagen, Denmark.

Corresponding author: Steffen U. Thorsen, s.u.thorsen@gmail.com.

DOI: 10.2337/dc13-0342

© 2013 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

**Acknowledgments**—No potential conflicts of interest relevant to this article were reported.

S.U.T. and J.S. researched data and wrote the manuscript. H.B.M., B.H.T., L.L.H., R.B., F.P., and A.L. contributed to the discussion and reviewed and edited the manuscript. B.C. and C.B. researched data and reviewed and edited the manuscript. M.F. reviewed and edited the manuscript. J.S. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

- Svensson J, Lyngaae-Jørgensen A, Carstensen B, Simonsen LB, Mortensen HB; Danish Childhood Diabetes Registry. Long-term trends in the incidence of type 1 diabetes in Denmark: the seasonal variation changes

Table 1—Association between 25(OH)D, PTH, and total calcium levels and case subject status by sex

	All (n = 1,803)	Sibling		Patient	
		Female (n = 423)	Male (n = 473)	Female (n = 436)	Male (n = 471)
Age (years)	10.26 (0.09)	10.58 (0.19)	10.71 (0.18)	9.42 (0.18)	10.29 (0.17)
25(OH)D (nmol/L)	57.77	58.00	54.02	57.00	61.82
Q1; Q3	37.07; 86.64	36.99; 89.00	36.17; 80.65	37.16; 85.00	38.25; 90.75
PTH (pmol/L)	0.73	0.82	0.74	0.75	0.64
Q1; Q3	0.16; 1.51	0.16; 1.61	0.16; 1.51	0.16; 1.41	0.16; 1.47
Calcium (mmol/L)	2.54	2.49	2.52	2.56	2.57
Q1; Q3	2.40; 2.66	2.29; 2.64	2.36; 2.64	2.47; 2.68	2.43; 2.68

Data are mean (SEM) or median unless otherwise indicated.

- over time. *Pediatr Diabetes* 2009;10:248–254
2. Pozzilli P, Manfrini S, Crinò A, et al.; IMDIAB group. Low levels of 25-hydroxyvitamin D3 and 1,25-dihydroxyvitamin D3 in patients with newly diagnosed type 1 diabetes. *Horm Metab Res* 2005;37:680–683
  3. Borkar VV, Devidayal, Verma S, Bhalla AK. Low levels of vitamin D in North Indian children with newly diagnosed type 1 diabetes. *Pediatr Diabetes* 2010;11:345–350
  4. Cooper JD, Smyth DJ, Walker NM, et al. Inherited variation in vitamin D genes is associated with predisposition to autoimmune disease type 1 diabetes. *Diabetes* 2011;60:1624–1631
  5. Cappelen J. The climate of Denmark—key climatic figures, 1990-99 [Internet], 2000. Copenhagen, Denmark, Danish Meteorological Institute. Available from <http://www.dmi.dk/dmi/tr00-08.pdf>. Accessed 1 August 2012