



No Sign of Proliferative Retinopathy in 15 Patients With Permanent Neonatal Diabetes With a Median Diabetes Duration of 24 Years

Diabetes Care 2014;37:e181–e182 | DOI: 10.2337/dc14-0471

Dario Iafusco,¹ Silvana Salardi,² Giovanni Chiari,³ Sonia Toni,⁴ Ivana Rabbone,⁵ Roberta Pesavento,⁶ Bruno Pasquino,⁷ Antonella de Benedictis,⁸ Giulio Maltoni,² Carlo Colombo,⁹ Lucia Russo,¹⁰ Ornella Massa,¹⁰ Maurizio Sudano,¹¹ Francesco Cadario,¹² Massimo Porta,¹³ Fabrizio Barbetti,^{9,10} and the Early Onset Diabetes Study Group of the Italian Society of Pediatric Endocrinology and Diabetology (ISPED)*

The knowledge about the long-term consequences of diabetes with onset in the neonatal period is scanty. We investigated the impact of long-standing diabetes (>15 years) on the retina of 10 patients with permanent neonatal diabetes mellitus (PNDM) (diabetes diagnosis within 6 months of birth) associated with mutations of *GCK*, *KCNJ11*, *INS*, or *ABCC8* genes and of two parents carrying an *INS* gene mutation diagnosed with diabetes in their childhood (1,2) (Table 1, patients 1–12). Eye complications were also evaluated in three patients with diabetes onset within 1 year of age and negative for type 1A diabetes autoantibodies (2) (Table 1, patients 13–15).

The mean age at diagnosis of diabetes of patients with PNDM was 7 weeks (patients 1–9 and 12), and median duration of diabetes of the entire group was 24 years (\pm 10.2 SD; range 16–47 years). Six patients had diabetes for more than 30 years.

Fundus photography, performed according to EURODIAB recommendations

(3), was available for 12 patients and read independently by two experts (M.P., A.d.B.) who did not have access to clinical data. For the remaining three patients, reports of fundus oculi or fluorescein angiography were evaluated. Because the patients are transferred to adult diabetes clinic when they reach the age of 18 years, information on recent HbA_{1c} was obtained in some occasions directly from the patient and not from medical records.

The results obtained were compared with those published by our study group (4) describing 105 patients with diabetes duration of 20 years, 53 of which had diabetes onset before puberty.

Proliferative retinopathy was reported in none of the patients. Eight individuals (53%) with diabetes duration between 16 and 24 years had no sign of diabetic retinopathy (DR), five were judged to have mild DR, one had moderate DR with possible initial signs of diabetic macular edema (DME), and one had DME. All

patients with mild or moderate DR had diabetes for more than 30 years, but one. No difference was observed between the five patients with *INS* mutations (on insulin therapy) or the six patients with K_{ATP} channel (*KCNJ11/ABCC8*) mutations (four patients were weaned from insulin as adults and were currently on glyburide therapy) (Table 1).

These results compare well with those of our previous work (4), in which we observed that in patients with prepubertal onset of type 1A diabetes, 60% showed no DR after a mean duration of diabetes of 20 years. This subgroup had a better outcome than patients with pubertal or peripubertal onset of type 1A diabetes, who showed only 29% of cases free of DR and a significant number of cases with severe DR.

This low prevalence of severe DR in patients with PNDM of long duration is in contrast with that reported in other forms of monogenic diabetes with onset at puberty and beyond, such as *HNF1A*-maturity-

¹Department of Pediatrics, Second University of Naples, Naples, Italy

²Department of Pediatrics, University of Bologna, Bologna, Italy

³Regional Center for Children and Adolescents With Diabetes, University Hospital, Parma, Italy

⁴Pediatric Diabetology Unit, Meyer Children Hospital, Florence, Italy

⁵Department of Pediatrics, University of Turin, Turin, Italy

⁶Pediatric Unit, Boldrini Hospital, Thiene, Italy

⁷Department of Pediatrics, Regional Hospital, Bolzano, Italy

⁸Eye Clinic, Second University of Naples, Naples, Italy

⁹Department of Experimental Medicine and Surgery, University of Tor Vergata, Rome, Italy

¹⁰Bambino Gesù Children's Hospital, Istituto di Ricovero e Cura a Carattere Scientifico, Rome, Italy

¹¹Diabetes Clinic, Urbino City Hospital, Urbino, Italy

¹²Department of Pediatrics, University of Eastern Piedmont, Novara, Italy

¹³Department of Medical Sciences, University of Turin, Turin, Italy

Corresponding author: Fabrizio Barbetti, fabrizio.barbetti@uniroma2.it.

*A complete list of the members of the Early Onset Diabetes Study Group of the Italian Society of Pediatric Endocrinology and Diabetology (ISPED) can be found in the Supplementary Data online.

© 2014 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered.

Table 1—Clinical and genetic features of PNDM patients with a median diabetes duration of 24 years

Patient no.	Mutation	Age (years)/sex	Diabetes duration (years)	Therapy	HbA _{1c} % (mmol/mol) on INS	HbA _{1c} % (mmol/mol) on SU	DR
1	<i>INS</i> -R89C	18/F	18	INS	8.4 (68)	—	No
2	<i>INS</i> -LB39Y40delinsH	19/F	19	INS	7.3 (56)	—	No
3	<i>INS</i> -Y108X	20/F	20	INS	7.9 (63)	—	No
4	<i>ABCC8</i> -V324M/W688R	20/F	20	SU since 2010	7.1 (54)	6.4 (46)	No
5	<i>KCNJ11</i> -K170N	20/F	20	SU since 2008	8.3 (67)	6.0 (42)	No
6	<i>GCK</i> -T228M	22/F	22	INS	8.0 (64)	—	No
7	<i>ABCC8</i> -A355T	24/M	24	INS	9.2 (77)	—	No
8	<i>KCNJ11</i> -R201C	33/M	33	INS	8.9 (74)	—	Moderate, initial DME (?)
9	<i>KCNJ11</i> -G53D	33/F	33	SU since 2006	8.0 (64)	6.4 (46)	Mild
10	<i>INS</i> -R89C	39/M	35	INS	7.2 (55)	—	DME (initial)
11	<i>INS</i> -R89C	48/F	47	INS	10 (86)	—	Mild
12	<i>ABCC8</i> -L213P	48/M	48	SU since 2011	8.4 (68)	6.5 (48)	Mild
13	Unknown	17/M	16	INS	9.1 (76)	—	Mild
14	Unknown	25/M	24	INS	8 (64)	—	No
15	Unknown	44/M	43	INS	7.2 (55)	—	Mild

The interval (years) in which HbA_{1c} values were available for calculation of the mean(s) in the table are listed below; those patients with an observation period equal or longer than 10 years are in bold. **Patient 1 = 2000–2013; Patient 2 = 1994–2013; Patient 3 = 1994–2014;** Patient 4 INS = 2006–2009, SU = 2010–2013; Patient 5 INS = 2005–2007, SU = 2008–2013; Patient 6 = 2006–2013; **Patient 7 = 1990–2014; Patient 8 = 1990–2014; Patient 9 INS = 2004–2005, SU = 2006–2013;** Patient 10 = 2011–2012; **Patient 11 = 2000–2013; Patient 12 INS = 2004–2010, SU = 2011–2013;** Patient 13 = 2010–2012; Patient 14 = 2010–2012; Patient 15 = 2010–2012. INS, insulin; F, female; M, male; SU, sulfonylurea.

onset diabetes of the young 3, where the percentage of patients with proliferative DR is as high as in type 2 diabetes (5).

We acknowledge that a limitation of our study is the small group of patients considered. Nevertheless, these results support the notion that onset of diabetes before puberty has a lesser impact on DR than in patients with other monogenic forms of the disease.

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

Author Contributions. D.I., S.S., G.C., S.T., I.R., R.P., B.P., A.d.B., G.M., C.C., L.R., O.M., M.S.,

F.C., and M.P. researched data. F.B. wrote the manuscript. F.B. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

- Colombo C, Porzio O, Liu M, et al.; Early Onset Diabetes Study Group of the Italian Society of Pediatric Endocrinology and Diabetes (ISPED). Seven mutations in the human insulin gene linked to permanent neonatal/infancy-onset diabetes mellitus. *J Clin Invest* 2008;118:2148–2156
- Russo L, lafusco D, Brescianini S, et al.; ISPED Early Diabetes Study Group. Permanent

diabetes during the first year of life: multiple gene screening in 54 patients. *Diabetologia* 2011;54:1693–1701

3. Aldington SJ, Kohner EM, Meuer S, Klein R, Sjølie AK. Methodology for retinal photography and assessment of diabetic retinopathy: the EURODIAB IDDM complications study. *Diabetologia* 1995;38:437–444

4. Salardi S, Porta M, Maltoni G, et al.; Diabetes Study Group of the Italian Society of Paediatric Endocrinology and Diabetology. Infant and toddler type 1 diabetes: complications after 20 years' duration. *Diabetes Care* 2012;35:829–833

5. Isomaa B, Henricsson M, Lehto M, et al. Chronic diabetic complications in patients with MODY3 diabetes. *Diabetologia* 1998;41:467–473