

COMMENTS AND RESPONSES

Response to Comment on: Hermida et al. Influence of Time of Day of Blood Pressure-Lowering Treatment on Cardiovascular Risk in Hypertensive Patients With Type 2 Diabetes. Diabetes Care 2011;34: 1270-1276

We thank Rossen and Hansen (1) for their interest in our article (2). The participants represented a consecutive series of patients fulfilling the exclusion/inclusion criteria who were recruited from those referred to the hospital for evaluation by ambulatory blood pressure monitoring (ABPM) mainly to confirm or reject the diagnosis of hypertension inferred by clinic cuff blood pressure measurement. Diagnosis of hypertension in our trial was based on accepted ABPM criteria, i.e., an awake systolic blood pressure (SBP)/diastolic blood pressure (DBP) mean $\geq 135/85$ mmHg and/or an asleep SBP/DBP mean $\geq 120/70$ mmHg (3). Despite a recommended target goal for clinic SBP/DBP in diabetes of 130/80 mmHg, current guidelines do not provide separate ABPM threshold values for patients with and without diabetes (3). Patients with diabetes in our trial can be considered newly diagnosed hypertensive because a previous ABPM evaluation was not available for any participant before recruitment.

Patients were randomly assigned to one of two monotherapy treatment time groups, either upon awakening (control group) or at bedtime (experimental group). Assignment of participants to treatment time groups was done according to the order of recruitment, following allocation tables constructed for each allowed medication by a computerized random number generator. If patients were uncontrolled

based on ABPM criteria after 3 months of therapy, additional medications could be added in keeping with current clinical practice. Additional medications were not randomized according to treatment time. At the final evaluation, the percentages of patients in the experimental group ingesting 1, 2, or ≥ 3 medications at bedtime were 56.5, 25.0, and 18.5%, respectively. As previously stated (2), the sample size of our trial is limited to derive conclusions from the comparison between classes of medications and their combinations on the benefits, in terms of cardiovascular risk reduction, of bedtime treatment. Extensive reviews on the morning/evening treatment time differences in asleep blood pressure regulation of different classes of hypertension medications are already available (4).

CI for event rates can be directly calculated from the numbers of events and participants, plus the median duration of follow-up. As stated (2), the rate of total events was significantly higher in the control (54.2 [95% CI 43.4–65.1]) than in the experimental group (19.8 [12.1–27.4]; $P < 0.001$). The rate of major events was also significantly higher in the control group (17.6 [10.6–24.5] vs. 5.2 [1.1–9.2]; $P < 0.001$) (2).

With respect to the results described in Fig. 2 of our article (2), hypertension treatment, and not just the severity of disease, markedly influenced achieved blood pressure. As such, our statements and conclusions on the different shape of the relationship with cardiovascular risk for achieved clinic blood pressure and asleep blood pressure mean seem adequate (2). Data from the baseline ABPM evaluation indicated that cardiovascular risk increased exponentially with progressively increasing asleep blood pressure mean. Baseline clinic blood pressure was not a significant predictor of outcome in the survival model including the asleep blood pressure mean as a confounding variable (5). Finally, analyses of changes in blood pressure during follow-up revealed a significant reduction in cardiovascular risk associated with the decrease in asleep blood pressure mean during follow-up, independently of changes in clinic or awake blood pressure mean (2,5).

**RAMÓN C. HERMIDA, PHD
DIANA E. AYALA, MD, MPH, PHD
ARTEMIO MOJÓN, PHD
JOSÉ R. FERNÁNDEZ, PHD**

From the Bioengineering and Chronobiology Laboratories, University of Vigo, Campus Universitario, Vigo, Spain.

Corresponding author: Ramón C. Hermida, rhermida@uvigo.es.

DOI: 10.2337/dc11-1611

© 2011 by the American Diabetes Association.

Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Acknowledgments—This independent investigator-promoted research was supported by unrestricted grants from Ministerio de Ciencia e Innovación (SAF2006-6254-FEDER; SAF2009-7028-FEDER); Consellería de Presidencia, Relacións Institucionais e Administración Pública, Secretaría Xeral de Investigación e Desenvolvemento, Xunta de Galicia (PGIDIT03-PXIB-32201PR); Consellería de Economía e Industria, Dirección Xeral de Investigación e Desenvolvemento, Xunta de Galicia (INCITE07-PXI-322003ES; INCITE08-E1R-322063ES; INCITE09-E2R-322099ES; 09CSA018322PR); and Vicerrectorado de Investigación, University of Vigo.

No potential conflicts of interest relevant to this article were reported.

References

- Rossen NB, Hansen KW. Comment on: Hermida et al. Influence of time of day of blood pressure-lowering treatment on cardiovascular risk in hypertensive patients with type 2 diabetes. *Diabetes Care* 2011; 34:1270–1276 (Letter). *Diabetes Care* 2011; 34:e184. DOI: 10.2337/dc11-1514
- Hermida RC, Ayala DE, Mojón A, Fernández JR. Influence of time of day of blood pressure-lowering treatment on cardiovascular risk in hypertensive patients with type 2 diabetes. *Diabetes Care* 2011;34:1270–1276
- Mancia G, De Backer G, Dominiczak A, et al.; Management of Arterial Hypertension of the European Society of Hypertension; European Society of Cardiology. 2007 guidelines for the management of arterial hypertension: The Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *J Hypertens* 2007;25:1105–1187
- Smolensky MH, Hermida RC, Ayala DE, Tiseo R, Portaluppi F. Administration-time-dependent effects of blood pressure-lowering medications: basis for the chronotherapy of hypertension. *Blood Press Monit* 2010;15: 173–180
- Hermida RC, Ayala DE, Mojón A, Fernández JR. Decreasing sleep-time blood pressure determined by ambulatory monitoring reduces cardiovascular risk. *J Am Coll Cardiol* 2011;58:1165–1173