
 COMMENTS AND
 RESPONSES

Comment on: Raz et al. Personalized Management of Hyperglycemia in Type 2 Diabetes: Reflections From a Diabetes Care Editors' Expert Forum. Diabetes Care 2013;36:1779-1788

Not all existing therapies for type 2 diabetes provide similar glycemic control for every affected patient; with a few notable exceptions, how different responses to therapy relate to specific variations within identifiable subgroups of patients remained, and remains, largely unexplored.

Also on the basis of current uncertainty, the *Diabetes Care* Editors' Expert Forum delivered their reflections to help physicians personalize diabetes care (1). Apart from the many given reasons for the need to decode the joint position statement by the American Diabetes Association and the European Association for the Study of Diabetes (EASD) (2), the list could also include recognition of the difficulty of implementing effective personalized therapy in a clinical setting, the need to enlarge consensus, and the panoply of diabetes drugs, which may also imply the "personalization of a niche" for each different antihyperglycemic agent within the 12 drug classes on the U.S. market. However, multiple treatment

guidelines, algorithms, and goals periodically released to improve guidance may also enhance uncertainty (3).

Personalized medicine should be based on evidence rather than clinical impression; unfortunately, it still lacks scientific evidence. For example, personalization of HbA_{1c} target for diabetic individuals is paramount as the aggressiveness of any therapy is ultimately based on how low the target is set. A good example of this challenge is the recent INTERVAL trial, which aims to assess the feasibility of setting and achieving individualized targets in elderly (aged 70 years or older) type 2 diabetic patients (4). Although investigators from seven European countries were free to set individualized treatment targets on the basis of age, baseline HbA_{1c}, comorbidities, and frailty status, the mean investigator-defined individualized HbA_{1c} targets were around 7.0%, substantially lower than expected for that elderly population. To our knowledge, this is the only controlled study trying to apply the philosophy of personalized medicine in type 2 diabetes.

Human beings tend to revert to the familiar: given the paucity of pragmatic aid (for example, Web-driven algorithms that estimate the target with simple parameters or tailor pharmacological therapy on the basis of clinical features [5]), personal decisions tend to be conservative and uniform to what is familiar and known. A lesson learned is that physicians need practical help to feel safer with consistent recommendations.

DARIO GIUGLIANO, MD, PHD¹
 KATHERINE ESPOSITO, MD, PHD²

From the ¹Department of Medical, Surgical, Neurological, Metabolic Sciences, and Geriatrics, Second University of Naples, Naples, Italy; and the ²Department of Clinical and Experimental Medicine and Surgery, Second University of Naples, Naples, Italy.

Corresponding author: Dario Giugliano, dario.giugliano@unina2.it.

DOI: 10.2337/dc13-1427

© 2013 by the American Diabetes Association.

Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Acknowledgments—No potential conflicts of interest relevant to this article were reported.

.....

References

1. Raz I, Riddle MC, Rosenstock J, et al. Personalized management of hyperglycemia in type 2 diabetes: reflections from a *Diabetes Care* Editors' Expert Forum. *Diabetes Care* 2013;36:1779-1788
2. Inzucchi SE, Bergenstal RM, Buse JB, et al.; American Diabetes Association (ADA); European Association for the Study of Diabetes (EASD). Management of hyperglycemia in type 2 diabetes: a patient-centered approach: position statement of the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care* 2012;35:1364-1379
3. Esposito K, Ceriello A, Giugliano D. Does personalized diabetology overcome clinical uncertainty and therapeutic inertia in type 2 diabetes? *Endocrine* 2013;44:343-345
4. Strain WD, Lukashevich V, Kothny W, Hoellinger MJ, Paldanius PM. Individualised treatment targets for elderly patients with type 2 diabetes using vildagliptin add-on or lone therapy (INTERVAL): a 24 week, randomised, double-blind, placebo-controlled study. *Lancet* 2013;382:409-416
5. Italian Association of Medical Diabetologists (AMD). Personalization in type 2 diabetes [Internet], 2013. Available from http://www.aemmedi.it/algorithmi_en_2013. Accessed 17 June 2013