

Lusha Xiang, Peter N. Mittwede, and Robert L. Hester



COMMENT ON SATO ET AL.

Improving Type 2 Diabetes Through a Distinct Adrenergic Signaling Pathway Involving mTORC2 That Mediates Glucose Uptake in Skeletal Muscle. *Diabetes* 2014;63:4115–4129

Diabetes 2014;63:e20–e21 | DOI: 10.2337/db14-1187

Sato et al. (1) showed that β_2 -adrenoreceptor-mediated phosphorylation of mammalian target of rapamycin complex 2 (mTORC2) in skeletal muscles increased GLUT4 translocation to the plasma membrane and thus increased cellular glucose uptake. In diabetes and obesity, an upregulated G-protein-coupled receptor kinase, which inhibits G-protein-coupled receptors, has been reported to contribute to insulin resistance in skeletal muscle (2). In support of this, the current study demonstrated an increased peripheral glucose uptake following β_2 -adrenoreceptor activation in diabetic rats and obese mice (1). However, the statement that “this signaling pathway provides new opportunities for the treatment of type 2 diabetes” (1) may be debatable because the impacts of systemic β_2 -adrenoreceptor activation on glucose control and hemodynamic changes are not considered.

Increased sympathetic nervous system activity (exercise and the fight or flight response, as Sato et al. mentioned) is associated not only with increased glucose uptake by skeletal muscle but also with increased hepatic glycogenolysis and/or glucagon release, likely leading to an increase in plasma glucose levels. Plasma glucose and glucagon levels can increase significantly during exercise despite increased glucose uptake in exercising muscle. We recently demonstrated that traumatic injury rapidly increased plasma glucose levels via β_2 -adrenoreceptor-dependent glycogenolysis in the liver. Treatment with a selective β_2 -adrenoreceptor antagonist blunted the

hepatic glycogenolysis and the increase in plasma glucose levels (3). Thus, following systemic β_2 -adrenoreceptor activation, the beneficial effect from increased peripheral glucose uptake in skeletal muscle may be overridden by simultaneous increases in hepatic glycogenolysis and/or glucagon release. Notably, due to insulin resistance, obese rats exhibited exacerbated post-trauma hyperglycemia as compared with lean Zucker rats (3). Therefore, the net increase in plasma glucose following β_2 -adrenoreceptor activation could be more profound in obesity and type 2 diabetes.

Sato et al. (1) showed that systemic treatment with β_2 -adrenoreceptor agonist improved glucose tolerance in diabetic rats and obese mice. However, whether this is due to the same mechanism (mTORC2 pathway) as demonstrated in their *in vitro* experiments is uncertain. For example, β_2 -adrenoreceptor activation also causes arteriolar dilation and capillary recruitment in skeletal muscle and thus increases tissue perfusion and (insulin-mediated) glucose uptake (4). Although the current study eloquently provided evidence for β_2 -adrenoreceptor-mediated glucose uptake in skeletal muscle via the cAMP-mTORC2-GLUT4 pathway at the cellular level, future studies are needed to determine the interactions between cellular and systemic glucose responses following β_2 -adrenoreceptor activation.

Funding. This work was supported by National Institutes of Health (P20GM104357, HL-51971, and HL-89581), American Heart Association

(12SDG12050525 and 14PRE17810005), and a Basic Science Research Grant from the Orthopaedic Trauma Association.

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

References

1. Sato M, Dehvari N, Öberg AI, et al. Improving type 2 diabetes through a distinct adrenergic signaling pathway involving mTORC2 that mediates glucose uptake in skeletal muscle. *Diabetes* 2014;63:4115–4129
2. Garcia-Guerra L, Nieto-Vazquez I, Vila-Bedmar R, et al. G protein-coupled receptor kinase 2 plays a relevant role in insulin resistance and obesity. *Diabetes* 2010;59:2407–2417
3. Xiang L, Lu S, Mittwede PN, Clemmer JS, Husband GW, Hester RL. β 2 adrenoreceptor blockade improves early post-trauma hyperglycemia and pulmonary injury in obese rats. *Am J Physiol Heart Circ Physiol* 2014;307:H621–H627
4. Baron AD, Tarshoby M, Hook G, et al. Interaction between insulin sensitivity and muscle perfusion on glucose uptake in human skeletal muscle: evidence for capillary recruitment. *Diabetes* 2000;49:768–774