

Did the great dying of life take 700 k.y.? Evidence from global astronomical correlation of the Permian-Triassic boundary interval

Chunju Huang^{1,2}, Jinnan Tong³, Linda Hinnov², and Zhong Qiang Chen³

¹Key Laboratory of Tectonics and Petroleum Resources of Ministry of Education, China University of Geosciences, Wuhan 430074, PR China

²Department of Earth and Planetary Sciences, Johns Hopkins University, Baltimore, Maryland 21218, USA

³State Key Laboratory of Biogeology and Environmental Geology, China University of Geosciences, Wuhan 430074, PR China

We thank Rampino and Kaiho (2012) for their Comment on our Geology paper (Huang et al., 2011). They questioned our definition of a mass extinction interval (MEI) for the Permian-Triassic (P-Tr) boundary interval. We define the MEI as the stratigraphic interval from the start of the *Neogondolella meishanensis* conodont zone, or its equivalents at the onset of the negative $\delta^{13}\text{C}$ excursion (ME_{start}), to the base of the *Isarcicella isarcica* conodont zone and increasing $\delta^{13}\text{C}$ values (ME_{end}).

At the Meishan (China) section, the *N. meishanensis* zone starts at the base of Bed 25, which is widely accepted as the main mass extinction horizon. Thus, we treat this horizon as ME_{start} . Kaiho et al. (2009) proposed the main mass extinction to be within an interval of 1.2 cm near the top of Bed 24e based on examination of continuously sampled microfossils. However, quantitative analysis, e.g., evaluation of the Signor-Lipps effect, shows that the main mass extinction falls at the top of Bed 24e or the base of Bed 25 (Jin et al., 2000; Song et al., 2009; Shen et al., 2011). The *I. isarcica* zone starts at the base of Bed 29 (Jiang et al., 2007). An extinction horizon was recognized in the base of Bed 28 (Yin et al., 2007; Song et al., 2009). However, Bed 28 is a volcanic ash bed, representing a very short duration, and is not preserved in P-Tr boundary sections worldwide. To enable recognition of this extinction horizon and its equivalent in other sections, we used the base of the *I. isarcica* zone as a marker of ME_{end} . We also considered MEI durations based on both options (GSA Data Repository item 2011231). Also, the “cycles” in our eccentricity count at Meishan included Beds 25 and 28, which are not true cycles, but ash beds—hence likely “instantaneous” events.

For the Gartnerkofel core (Austria) discussed by Rampino and Kaiho, we considered two definitions of the MEI (Huang et al., 2011). Biostratigraphic correlation places ME_{start} at 231 m (base of the *Hindeodus latidentatus* zone) and ME_{end} at 188.6 m (base of *I. isarcica*) for a MEI thickness of 42.4 m (Schönlaub, 1991). Alternatively, $\delta^{13}\text{C}$ profile correlation places ME_{start} at 224.97 m and ME_{end} at 182 m. Either scheme gives the same duration.

Our definition of an MEI focuses not only on the main mass extinction event in the Late Permian, but also considers the immediate aftermath and the succeeding extinctions through the P-Tr boundary. Rampino and Kaiho consider only the Late Permian mass extinction event, which we have termed here as the main mass extinction. Recent studies indicate that abundant Permian fossils occur above the main extinction event horizon (Chen et al., 2005; Song et al., 2009; Korchagin, 2011), and most finally disappear at Bed 28. The most distinctive change in benthic community structures did not occur in the main mass extinction event, but coincided with the later demise of the temporary survivors (Chen et al., 2010). Consequently, P-Tr biotic extinctions and ecosystem collapse are recorded in the MEI. We show that the MEI can be correlated worldwide with biostratigraphic and $\delta^{13}\text{C}$ profile correlations (Huang et al., 2011).

Permian fossils may occur only in an interval of 0.3 m to 1 m above

the main mass extinction horizon in the Italian sections (Rampino and Kaiho, 2012). However, abundant Permian foraminifers extend up to the *I. staeschei* zone at Meishan (Song et al., 2009; Korchagin, 2011). If our MEI definition is considered, combined conodont biostratigraphy and $\delta^{13}\text{C}$ patterns suggest a much broader extinction interval than suggested by Rampino and Kaiho.

We therefore believe that the P-Tr mass extinctions may be accounted for by a series of critical events rather than a single catastrophe. We also note that an eccentricity minimum occurs ~200 k.y. prior to the onset of the MEI (Huang et al., 2011), which could have influenced or paced the impending extinctions. We favor a new campaign to measure cyclostratigraphy at high resolution across these and other biostratigraphically constrained sections worldwide to help solve the timing problems of the P-Tr mass extinctions.

ACKNOWLEDGMENT

This research was supported by the Program for New Century Excellent Talents in University (NCET-11-0723) and the Fundamental Research Funds for the Central Universities, China University of Geosciences (Wuhan) (007-G1323521118).

REFERENCES CITED

- Chen, Z.Q., Kaiho, K., and George, A.D., 2005, Early Triassic recovery of the brachiopod faunas from the end-Permian mass extinction: A global review: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 224, p. 270–290, doi:10.1016/j.palaeo.2005.03.037.
- Chen, Z.Q., Tong, J., Liao, Z.T., and Chen, J., 2010, Structural changes of marine communities over the Permian-Triassic transition: Ecologically assessing the end-Permian mass extinction and its aftermath: *Global and Planetary Change*, v. 73, p. 123–140, doi:10.1016/j.gloplacha.2010.03.011.
- Huang, C., Tong, J., Hinnov, L., and Chen, Z.Q., 2011, Did the great dying of life take 700 k.y.? Evidence from a global astronomical correlation of the Permian-Triassic boundary interval: *Geology*, v. 39, p. 779–782, doi:10.1130/G32126.1.
- Jiang, H., Lai, X., Luo, G., Aldridge, R., Zhang, K., and Wignall, P.B., 2007, Restudy of conodont zonation and evolution across the P/T boundary at Meishan section, Changxing, Zhejiang, China: *Global and Planetary Change*, v. 55, p. 39–55, doi:10.1016/j.gloplacha.2006.06.007.
- Jin, Y.G., Wang, Y., Wang, W., Shang, Q.H., Cao, C.Q., and Erwin, D.H., 2000, Pattern of marine mass extinction near the Permian-Triassic boundary in southern China: *Science*, v. 289, p. 432–436, doi:10.1126/science.289.5478.432.
- Kaiho, K., Chen, Z.Q., and Sawada, K., 2009, Possible causes for a negative shift in the stable carbon isotope ratio before, during and after the end-Permian mass extinction in Meishan, South China: *Australian Journal of Earth Sciences*, v. 56, p. 799–808, doi:10.1080/08120090903002615.
- Korchagin, O.A., 2011, Foraminifers in the global stratotype (GSSP) of the Permian-Triassic boundary (Bed 27, Meishan, South China): *Stratigraphy and Geological Correlation*, v. 19, p. 160–172, doi:10.1134/S0869593811020055.
- Rampino, M.R., and Kaiho, K., 2012, Did the great dying of life take 700 k.y.? Evidence from a global astronomical correlation of the Permian-Triassic boundary interval: *Geology*, e267, doi:10.1130/G32821C.1.
- Schönlaub, H.P., 1991, The Permian-Triassic of the Gartnerkofel-1 core (Carnic alps, Austria): Conodont biostratigraphy: *Abhandlungen der Geologischen Bundesanstalt*, v. 45, p. 79–98.
- Shen, S.Z., and 21 others, 2011, Calibrating the end-Permian mass extinction: *Science*, v. 334, p. 1367–1372, doi:10.1126/science.1213454.
- Song, H., Tong, J., and Chen, Z.Q., 2009, Two episodes of foraminiferal extinction near the Permian-Triassic boundary at the Meishan section, South China: *Australian Journal of Earth Sciences*, v. 56, p. 765–773, doi:10.1080/08120090903002599.
- Yin, H., Feng, Q., Lai, X., Baud, A., and Tong, J., 2007, The protracted Permian-Triassic crisis and multi-episode extinction around the Permian-Triassic boundary: *Global and Planetary Change*, v. 55, p. 1–20, doi:10.1016/j.gloplacha.2006.06.005.