

Carbon sequestration on Mars

Christopher S. Edwards¹ and Bethany L. Ehlmann^{2,3}

¹Astrogeology Science Center, U.S. Geological Survey, Flagstaff, Arizona 86001, USA

²California Institute of Technology, 1200 E. California Boulevard, MC 150-21, Pasadena, California 91125, USA

³Jet Propulsion Laboratory/California Institute of Technology, Pasadena, California 91109, USA

Martian atmospheric pressure has important implications for the past and present habitability of the planet, including the timing and causes of environmental change. The ancient Martian surface is strewn with evidence for early water bound in minerals (e.g., Ehlmann and Edwards, 2014) and recorded in surface features such as large catastrophically created outflow channels (e.g., Carr, 1979), valley networks (Hynek et al., 2010; Irwin et al., 2005), and crater lakes (e.g., Fassett and Head, 2008). Using orbital spectral data sets coupled with geologic maps and a set of numerical spectral analysis models, Edwards and Ehlmann (2015) constrained the amount of atmospheric sequestration in early Martian rocks and found that the majority of this sequestration occurred prior to the formation of the early Hesperian/late Noachian valley networks (Fassett and Head, 2011; Hynek et al., 2010), thus implying the atmosphere was already thin by the time these surface-water-related features were formed.

As Lee et al. (2016) rightly highlight in their Comment, the meteorite record provides important constraints on Martian carbon sequestration processes. The example used in the discussion by Edwards and Ehlmann of the possibility of deep diffuse carbonate alteration—rather than precipitation of carbonate in discrete rock deposits like those analyzed—was selected based on measurements of carbonate in dust, *in-situ* mission data, and meteorites. That is, a 1 km global crustal layer of 1 vol.% carbonate-bearing materials, similar to the abundances in the Nakhallite-class Lafayette meteorite (Changela and Bridges, 2010; Nyquist et al., 2001; Tomkinson et al., 2013; Wright et al., 1992), if formed from waters in contact with the atmosphere, would lead to sequestration of ~500 mbar CO₂.

To sequester significantly more carbon than this from the Martian atmosphere requires an assumption that typical Martian rocks and soils have vol% carbonate several factors greater than that observed in Martian meteorites. Abundances as high as this would be more readily detectable to remote sensing and landed missions, but are so far observed only rarely or at abundances of a few percent or less (e.g., Leshin et al., 2013; Niles et al., 2013) except in an isolated olivine-rich region of the Columbia Hills, Gusev crater (Morris, 2010). Thus, a 1–5 vol% carbonate range over a large volume of crust is a reasonable estimate, which leads to a maximum of ~1–2 bars sequestered since valley network formation, under most environmental scenarios, a value consistent with isotopic evidence and known geologic processes (e.g., Hu et al., 2015), including newly measured escape processes (Jakosky, 2015). Interestingly, the period prior to valley network formation likely had additional carbon sequestration by different processes to remove early or pre-Noachian atmosphere (Wray et al., 2016). Future landed missions are considering visiting a site with in-place carbonate-bearing materials (<http://marsnext.jpl.nasa.gov>; Hand, 2015), possibly returning samples from them, a prospect which would provide a key suite of petrologic and isotopic data to address this most important question about Mars atmospheric and climate evolution: the fate of the atmosphere.

REFERENCES CITED

- Carr, M. H., 1979, Formation of martian flood features by release of water from confined aquifers: *Journal of Geophysical Research*, v. 84, B6, p. 2995–3007.
- Changela, H.G., and Bridges, J.C., 2010, Alteration assemblages in the nakhlites: Variation with depth on Mars: *Meteoritics & Planetary Science*, v. 45, p. 1847–1867, doi:10.1111/j.1945-5100.2010.01123.x.
- Edwards, C.S., and Ehlmann, B.L., 2015, Carbon sequestration on Mars: *Geology*, v. 43, p. 863–866, doi:10.1130/G36983.1.
- Ehlmann, B.L., and Edwards, C.S., 2014, Mineralogy of the Martian Surface: *Annual Review of Earth and Planetary Sciences*, v. 42, p. 291–315, doi:10.1146/annurev-earth-060313-055024.
- Fassett, C.I., and Head, J.W., 2008, The timing of martian valley network activity: Constraints from buffered crater counting: *Icarus*, v. 195, p. 61–89, doi:10.1016/j.icarus.2007.12.009.
- Fassett, C.I., and Head, J.W., 2011, Sequence and timing of conditions on early Mars: *Icarus*, v. 211, p. 1204–1214, doi:10.1016/j.icarus.2010.11.014.
- Hand, E., 2015, Scientists tap ancient river deltas and hot springs as promising targets for 2020 rover: *Science News*, doi:10.1126/science.aac8981.
- Hu, R., Kass, D.M., Ehlmann, B.L., and Yung, Y.L., 2015, Tracing the fate of carbon and the atmospheric evolution of Mars: *Nature Communications*, v. 6, doi:10.1038/ncomms10003.
- Hynek, B.M., Beach, M., and Hoke, M.R.T., 2010, Updated global map of Martian valley networks and implications for climate and hydrologic processes: *Journal of Geophysical Research*, v. 115, E09008, doi:10.1029/2009JE003548.
- Irwin, R.P., Craddock, R.A., and Howard, A.D., 2005, Interior channels in Martian valley networks: Discharge and runoff production: *Geology*, v. 33, p. 489–492, doi:10.1130/G21333.1.
- Jakosky, B.M., 2015, MAVEN Explores the Martian Upper Atmosphere: *Science*, v. 350, p. 643, doi:10.1126/science.aad3443.
- Lee, M. R., Tomkinson, T., Mark, D. F., and Smith, C. L., 2016, Carbon sequestration on Mars: Comment: *Geology*, v. 44, p. xxx, doi:10.1130/G37617C.1
- Leshin, L.A., et al., 2013, Volatile, isotope, and organic analysis of martian fines with the Mars Curiosity rover: *Science*, v. 341, doi:10.1126/science.1238937.
- Morris, R.V., et al., 2010, Identification of Carbonate-Rich Outcrops on Mars by the Spirit Rover: *Science*, v. 329, p. 421–424, doi: 10.1126/science.1189667.
- Niles, P.B., Catling, D.C., Berger, G., Chassefiere, E., Ehlmann, B.L., Michalski, J.R., Morris, R., Ruff, S.W., and Sutter, B., 2013, Geochemistry of carbonates on Mars: Implications for climate history and nature of aqueous environments: *Space Science Reviews*, v. 174, p. 301–328, doi:10.1007/s11214-012-9940-y.
- Nyquist, L. E., Bogard, D. D., Shih, C.-Y., Greshake, A., Stoffler, D., and Eugster, O., 2001, Ages and geologic histories of martian meteorites: *Chronology and Evolution of Mars*, v. 96, p. 105–164, doi: 10.1023/A:1011993105172.
- Tomkinson, T., Lee, M. R., Mark, D. F., and Smith, C. L., 2013, Sequestration of Martian CO₂ by mineral carbonation: *Nature Communications*, v. 4, doi:10.1038/ncomms3662
- Wray, J.J., Murchie, S.L., Bishop, J.L., Ehlmann, B.L., Milliken, R.E., Wilhelm, M.B., Seelos, K.D., and Chojnacki, M., 2016, Orbital evidence for more widespread carbonate-bearing rocks on Mars: *Journal of Geophysical Research: Planets*, doi:10.1002/2015JE004972.
- Wright, I.P., Grady, M.M., and Pillinger, C.T., 1992, Chassigny and the nakhlites: Carbon-bearing components and their relationship to martian environmental conditions: *Geochimica et Cosmochimica Acta*, v. 56, p. 817–826, doi:10.1016/0016-7037(92)90100-W.