

COMMENTS AND RESPONSES

Response to Comment on: Wheeler et al. Macronutrients, Food Groups, and Eating Patterns in the Management of Diabetes: A Systematic Review of the Literature, 2010. Diabetes Care 2012;35:434-445

We appreciate the comments from Drs. Esposito and Giugliano (1) concerning our systematic review (2). Each issue will be addressed as outlined in their letter.

Wrong statement about results of published articles (reference 104): We acknowledge that the results were misquoted in the systematic review and appreciate the identification of the error. The correct data have been substituted in the online version of the article.

Reporting of one triplicate study (references 105–107): The three publications include different patients from the same hospital, different meals consumed, and, to some extent, different methodology used. The authors (personal communication with Smaragdi Antonopoulou, PhD on 22 February 2012) have confirmed that the publications are three separate studies.

Missing published studies: The two other clinical trial references provided (3,4) do meet the inclusion criteria for the systematic review and should have been included in the review. However, because both of these studies had a retention rate of <80%, according to our systematic review procedure, they would not have been included in the key summary for each topic area. Further, the diet that significantly improved A1C over the control diet in the

three-arm study by Elhayany et al. was a combination of a Mediterranean diet and a moderately low-carbohydrate diet (35% carbohydrate, 45% fat); the traditional Mediterranean diet (50–55% carbohydrate, 30% fat) did not demonstrate an advantage for A1C over the control diet.

Two other studies provided—a cross-sectional study (5) and a systematic review (6)—also meet our inclusion criteria and should have been included in the review. However, neither would have affected the summary statement. The cross-sectional study would not have been included in the summary as it did not compare the Mediterranean diet to any other type of eating pattern, and the systematic review did not add any additional studies that fit our inclusion criteria.

To clarify, our review does not suggest that the Mediterranean diet should be avoided in patients with diabetes; we simply state that “there appears to be no advantage in using the Mediterranean-style eating pattern compared with other eating patterns for glycemic control.” From our review, we found that several different eating patterns, including the Mediterranean diet, show promise for achieving glycemic control. There was, however, not one particular eating pattern with definitive evidence as being better than the others. This leaves health care providers the opportunity to work with individuals to adopt eating patterns honoring their own culture, food preferences, and food availability while still achieving glycemic control.

MADelyn L. WHEELER, MS, RD, FADA, CD¹
 STEPHANIE A. DUNBAR, MPH, RD²
 LINDSAY M. JAACKS, BS³
 WAHIDA KARMALLY, DRPH, RD, CDE, CLS, FNLA⁴
 ELIZABETH J. MAYER-DAVIS, MSPH, PHD, RD⁵
 JUDITH WYLIE-ROSETT, EDD, RD⁶
 WILLIAM S. YANCY JR., MD, MHS⁷

From ¹Nutritional Computing Concepts, Zionsville, Indiana; ²Medical Affairs, American Diabetes Association, Alexandria, Virginia; the ³School of Public Health, Nutritional Epidemiology, The University of North Carolina, Chapel Hill, Chapel Hill, North Carolina; the ⁴Irving Institute for Clinical and Translational Research, Columbia University, New York, New York; the ⁵Department of Nutrition, The University of North Carolina,

Chapel Hill, Chapel Hill, North Carolina; the ⁶Department of Epidemiology and Population Health, Albert Einstein College of Medicine, Bronx, New York; and the ⁷Division of General Internal Medicine, Duke University School of Medicine, Durham, North Carolina.

Corresponding author: Stephanie A. Dunbar, sdunbar@diabetes.org.

DOI: 10.2337/dc12-0585

© 2012 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Acknowledgments—W.K. has reported being a member of the Research Committee for the American Pistachio Growers. No other potential conflicts of interest relevant to this article were reported.

References

- Esposito K, Giugliano D. Comment on: Wheeler et al. Macronutrients, food groups, and eating patterns in the management of diabetes: a systematic review of the literature, 2010. *Diabetes Care* 2012;35:434-445 (Letter). *Diabetes Care* 2012;35:e51. DOI: 10.2337/dc12-0247
- Wheeler ML, Dunbar SA, Jaacks LM, et al. Macronutrients, food groups, and eating patterns in the management of diabetes: a systematic review of the literature, 2010. *Diabetes Care* 2012;35:434-445
- Elhayany A, Lustman A, Abel R, Attal-Singer J, Vinker S. A low carbohydrate Mediterranean diet improves cardiovascular risk factors and diabetes control among overweight patients with type 2 diabetes mellitus: a 1-year prospective randomized intervention study. *Diabetes Obes Metab* 2010;12:204-209
- Shai I, Schwarzfuchs D, Henkin Y, et al.; Dietary Intervention Randomized Controlled Trial (DIRECT) Group. Weight loss with a low-carbohydrate, Mediterranean, or low-fat diet. *N Engl J Med* 2008;359:229-241
- Esposito K, Maiorino MI, Di Palo C, Giugliano D; Campanian Postprandial Hyperglycemia Study Group. Adherence to a Mediterranean diet and glycaemic control in type 2 diabetes mellitus. *Diabet Med* 2009;26:900-907
- Esposito K, Maiorino MI, Ceriello A, Giugliano D. Prevention and control of type 2 diabetes by Mediterranean diet: a systematic review. *Diabetes Res Clin Pract* 2010;89:97-102

Downloaded from <http://diabetesjournals.org/care/article-pdf/35/6/e52/61385/e52.pdf> by guest on 24 September 2022