

## OBSERVATIONS

## Diagnosis of Gestational Diabetes Mellitus in China

After the publication of the Hyperglycemia and Adverse Pregnancy Outcome (HAPO) study (1), the International Association of the Diabetes and Pregnancy Study Groups (IADPSG) (2) formulated consensus guidelines for the testing and diagnosis of gestational diabetes mellitus (GDM). These have been adopted by the American Diabetes Association (3). The Ministry of Health (MOH) of China also recommended that new testing and diagnostic criteria based on the IADPSG guidelines (4) be effective beginning 1 December 2011. The Chinese criteria agree with the American Diabetes Association that a diagnostic 75-g oral glucose tolerance test (OGTT) should be performed between the 24th and 28th week of gestation for all pregnant women not previously known to have diabetes. A diagnosis of GDM can be made if one or more of the following glucose levels are elevated: fasting

$\geq 5.1$  mmol/L, 1 h  $\geq 10.0$  mmol/L, and 2 h  $\geq 8.5$  mmol/L.

This is the one-step approach that the MOH has recommended to well-resourced medical institutions. However, in China a formal OGTT may be difficult to implement in low-resourced rural areas. Therefore, and different from the American Diabetes Association, the MOH has recommended a two-step approach under these circumstances. A fasting plasma glucose can be used as a screening tool to reduce the number of OGTTs required. If the fasting plasma glucose is  $\geq 5.1$  mmol/L, GDM can be diagnosed and if  $< 4.4$  mmol/L, GDM is unlikely. Women with a result of  $\geq 4.4$  and  $\leq 5.0$  mmol/L will still require a glucose tolerance test. This approach will approximately halve the number of glucose tolerance tests required (5).

WEI-WEI ZHU, MD  
HUI-XIA YANG, MD

From the Department of Obstetrics and Gynecology, Peking University First Hospital, Beijing, China.  
Corresponding author: Hui-xia Yang, yanghuixia@bjmu.edu.cn.

DOI: 10.2337/dc12-2624

© 2013 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

**Acknowledgments**—No potential conflicts of interest relevant to this article were reported.

H.-x.Y. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

### References

1. Metzger BE, Lowe LP, Dyer AR, et al.; HAPO Study Cooperative Research Group. Hyperglycemia and adverse pregnancy outcomes. *N Engl J Med* 2008;358:1991–2002
2. Metzger BE, Gabbe SG, Persson B, et al.; International Association of Diabetes and Pregnancy Study Groups Consensus Panel. International association of diabetes and pregnancy study groups recommendations on the diagnosis and classification of hyperglycemia in pregnancy. *Diabetes Care* 2010;33:676–682
3. American Diabetes Association. Standards of medical care in diabetes—2011. *Diabetes Care* 2011;34(Suppl. 1):S11–S61
4. Yang HX; Medical Service Specialty Standard Committee of Ministry of Health. Diagnostic criteria for gestational diabetes mellitus (WS 331-2011). *Chin Med J (Engl)* 2012;125:1212–1213
5. Zhu WW, Fan L, Yang HX, et al. Fasting plasma glucose at 24–28 weeks to screen gestational diabetes: new evidence from China. *Diabetes Care*. 27 March 2013 [Epub ahead of print]