



RESPONSE TO COMMENT ON SHAH ET AL.

## Cardiovascular Complications and Mortality After Diabetes Diagnosis for South Asian and Chinese Patients: A Population-Based Cohort Study. *Diabetes Care* 2013;36:2670–2676

*Diabetes Care* 2014;37:e80 | DOI: 10.2337/dc13-2836

We wish to thank Tillin et al. (1) for their comments on our study (2). We agree that the absence of information on smoking status from population-level health care administrative data limits our ability to adjust for this important cardiovascular risk factor, and that this and other unmeasured risk factors could confound the relationships we found between ethnicity and cardiovascular complications or mortality. However, as noted in the article, we conducted sensitivity analyses to see how imbalanced an unmeasured confounder would have to be to explain the observed results. A risk factor that quadrupled the risk for mortality would have to have a prevalence of 5% in the South Asian population and 30% in the European population to explain the observed reduced hazard for mortality. Compared with this hypothetical risk factor, cigarette smoking is both less potent as a risk factor and less imbalanced in prevalence in the Ontario population (3,4); hence, it is unlikely that it explains the striking difference in mortality observed in our study.

The discrepancy between our results and those of Tillin et al. in the SABRE study (5) highlights an important message often lost in studies of minority

populations: broad ethnic categories such as South Asians are not homogeneous, and there may be important differences in diabetes and other cardiovascular risk factors within members of these groups (6,7). The differing migration histories to the U.K. and to Canada mean that the South Asian populations in each country have different regional origins, generation status, cultural and religious traditions, socioeconomic position, and acculturation and integration after migration. Hence, it should not be surprising that their health status and disease risk are also different. Comparing and contrasting research findings from different countries can shed light onto the biological, behavioral, socioeconomic, and health care system differences between different immigrant populations to better understand their health risks.

**Duality of Interest.** No potential conflicts of interest relevant to this article were reported.

### References

1. Tillin T, Sattar N, Forouhi N, Hughes AD, Chaturvedi N. Comment on Shah et al. Cardiovascular complications and mortality after diabetes diagnosis for South Asian and

Baiju R. Shah,<sup>1,2,3</sup> J. Charles Victor,<sup>1</sup> Maria Chiu,<sup>1,2</sup> Jack V. Tu,<sup>1,2,3</sup> Sonia S. Anand,<sup>4</sup> Peter C. Austin,<sup>1,2</sup> Douglas G. Manuel,<sup>1,5,6</sup> and Janet E. Hux<sup>1,2,3</sup>

Chinese patients: a population-based cohort study. *Diabetes Care* 2013;36:2670–2676 (Letter). *Diabetes Care* 2014;37:e78–e79. DOI: 10.2337/dc13-2490

2. Shah BR, Victor JC, Chiu M, et al. Cardiovascular complications and mortality after diabetes diagnosis for South Asian and Chinese patients: a population-based cohort study. *Diabetes Care* 2013;36:2670–2676
3. Jha P, Ramasundarahettige C, Landsman V, et al. 21st-century hazards of smoking and benefits of cessation in the United States. *N Engl J Med* 2013;368:341–350
4. Chiu M, Austin PC, Manuel DG, Tu JV. Cardiovascular risk factor profiles of recent immigrants vs long-term residents of Ontario: a multi-ethnic study. *Can J Cardiol* 2012;28:20–26
5. Tillin T, Forouhi NG, McKeigue PM, Chaturvedi N; SABRE Study Group. Southall And Brent REvisited: Cohort profile of SABRE, a UK population-based comparison of cardiovascular disease and diabetes in people of European, Indian Asian and African Caribbean origins. *Int J Epidemiol* 2012;41:33–42
6. Creatore MI, Moineddin R, Booth G, et al. Age- and sex-related prevalence of diabetes mellitus among immigrants to Ontario, Canada. *CMAJ* 2010;182:781–789
7. Bhopal R, Unwin N, White M, et al. Heterogeneity of coronary heart disease risk factors in Indian, Pakistani, Bangladeshi, and European origin populations: cross sectional study. *BMJ* 1999;319:215–220

<sup>1</sup>Institute for Clinical Evaluative Sciences, Toronto, Ontario, Canada

<sup>2</sup>University of Toronto, Toronto, Ontario, Canada

<sup>3</sup>Department of Medicine, Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada

<sup>4</sup>McMaster University, Hamilton, Ontario, Canada

<sup>5</sup>University of Ottawa, Ottawa, Ontario, Canada

<sup>6</sup>Ottawa Hospital Research Institute, Ottawa, Ontario, Canada

Corresponding author: Baiju R. Shah, baiju.shah@ices.on.ca.

© 2014 by the American Diabetes Association. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.