


REVIEW

Surface ocean-lower atmospheric processes in the Indian Ocean: Current understanding, knowledge gaps, and future directions

Ashwini Kumar^{1,*} , Susann Tegtmeier², Sheryl Oliveira Fernandes³, Haimanti Biswas⁴, Imran Girach⁵, M. K. Roxy⁶, Siby Kurian⁷, Christa A. Marandino⁸, V. V. S. S. Sarma⁹, and Damodar M. Shenoy⁷

Our understanding of surface ocean and lower atmosphere processes in the Indian Ocean (IO) region shows significant knowledge gaps mainly due to the paucity of observational studies. The IO basin is bordered by landmasses and an archipelago on 3 sides with more than one-quarter of the global population dwelling along these coastal regions. Therefore, interactions between dynamical and biogeochemical processes at the ocean-atmosphere interface and human activities are of particular importance here. Quantifying the impacts of changing oceanic and atmospheric processes on the marine biogeochemical cycle, atmospheric chemistry, ecosystems, and extreme events poses a great challenge. A comprehensive understanding of the links between major physical, chemical, and biogeochemical processes in this region is crucial for assessing and predicting local changes and large-scale impacts. The IO is one of the SOLAS (Surface Ocean-Lower Atmosphere Study) cross-cutting themes as summarized in its implementation strategy. This article attempts to compile new scientific results over the past decade focusing on SOLAS relevant processes within the IO. Key findings with respect to monsoon and air-sea interactions, oxygen minimum zones, ocean biogeochemistry, atmospheric composition, upper ocean ecosystem, and interactions between these components are discussed. Relevant knowledge gaps are highlighted, with a goal to assist the development of future IO research programs. Furthermore, we provided several recommendations to conduct interdisciplinary research to advance our understanding on the land-ocean-atmospheric interaction in the IO.

Keywords: SOLAS, Indian Ocean, Air-sea interaction, Oxygen minimum zone, Anthropogenic pollutants, Biogeochemistry

1. Introduction

The Indian Ocean (IO) is unique, when compared to the Atlantic and Pacific, due to being bordered by landmasses and an archipelago on 3 sides and the seasonally reversing winds and ocean currents (Hood et al., 2015). The IO is becoming increasingly important in terms of global and regional relevance as signals of climate change emerge rapidly in this basin with the adjacent coastlines harboring one-

third of the global population (Cai et al., 2014; Doyle, 2018; Hermes et al., 2019; Jyoti et al., 2023). Observations reveal that the IO is consistently warming, with some areas showing accelerated warming, leading to extreme weather events (Dhame et al., 2020; Cai et al., 2021), such as cyclones (Deshpande et al., 2021; Singh and Roxy, 2022), storm surges (Dube et al., 2009; Shaji et al., 2014; Needham et al., 2015), and floods (Roxy et al., 2017; Ankur et al.,

¹ Geological Oceanography Division, CSIR-National Institute of Oceanography, Dona Paula, Goa, India

² Institute of Space and Atmospheric Studies, University of Saskatchewan, Saskatoon, Canada

³ School of Earth, Ocean and Atmospheric Sciences, Goa University, Goa, India

⁴ Biological Oceanography Division, CSIR-National Institute of Oceanography, Dona Paula, Goa, India

⁵ Space Applications Centre, Indian Space Research Organisation, Ahmedabad, Gujarat, India

⁶ Indian Institute of Tropical Meteorology, Ministry of Earth Sciences, Pune, Maharashtra, India

⁷ Chemical Oceanography Division, CSIR-National Institute of Oceanography, Dona Paula, Goa, India

⁸ GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

⁹ National Institute of Oceanography, Regional Centre, Visakhapatnam, Andhra Pradesh, India

* Corresponding author:

Emails: ashwinikumajha@gmail.com; ashwinik@nio.org




Figure 1. Schematic of dynamical and biogeochemical processes in and above the Indian Ocean relevant for Surface Ocean-Lower Atmosphere Study (SOLAS) science.

2020; Zhou et al., 2021). Apart from its influence on the global climate system, the IO has a major effect on millions of people who live around its shores and depend on its resources. In particular, the 2 semi-enclosed basins in the North Indian Ocean (NIO), namely, the Arabian Sea (AS) and the Bay of Bengal (BoB), are ecologically and socioeconomically vital as they provide marine living and nonliving resources to the IO rim countries and generating a significant revenue for them. With a confirmed steady warming of the NIO and increasing atmospheric temperatures, a steady decline in the marine primary production (due to increased oligotrophy) is a worrisome possibility (Roxy et al., 2015). This may have severe biogeochemical consequences for the rim countries by impacting the entire food chain and fishery resources. Moreover, continued deoxygenation and its intensification over large areas of the NIO could lead to massive fish killing events and shifts in fish migration habits affecting the income of fishery-dependent coastal communities (Techera, 2018; Gangal et al., 2023). With rapidly emerging concepts and implementation of blue economy (including elements like fisheries, aquaculture, industry, shipping, and tourism), the IO is likely to experience increases in marine pollution and habitat degradation leading to changes in the ecosystem and biogeochemical characteristics. In view of this development, it is crucial to assess and understand the complex and interactive processes of the physical and biogeochemical components at the air–sea interface of this oceanic basin.

Despite several international efforts made in the past, through programs such as, the International Indian Ocean Expedition (IIOE), Geochemical Section Study (GEOSECS), Joint Global Ocean Flux Study (JGOFS), World Ocean Circulation Experiments (WOCE), and Geotraces, the IO region is still remained as one of the most poorly sampled regions among the world's oceans, with various scientific gaps (Beal et al., 2020). Owing to this, the International Indian Ocean Expedition-2 (IIOE-2; Hood et al., 2016), a decadal review of the Indian Ocean Observing System (IndOOS-2; Beal et al., 2020), monsoon mission and repetition of WOCE transects are being carried out by international scientific community to improve our understanding of the IO and its role in the Earth climate. Such an improved understanding of physical,

biogeochemical, and ecological processes also has strong relevance for the sustainable development of marine resources and environmental stewardship of the IO and surrounding coastal regions.

The Surface Ocean-Lower Atmosphere Study (SOLAS) program came into existence 2 decades ago to address some of the key surface water lower atmosphere coupled interactions and processes (Law et al., 2013). **Figure 1** displays a variety of processes, covered under the SOLAS domain that are relevant for the IO. The characteristic IO monsoon, which is a seasonal reversal of wind patterns (south-westerly during May–September and north-easterly during December–February), plays an important role in modulating, not only atmospheric dynamics and composition but also surface water currents and deep-water processes (Shankar et al., 2002; Phillips et al., 2021). One of the unique features of the IO atmosphere is the seasonally contrasting chemical regimes, which are controlled by anthropogenic emissions and natural mineral dust from the surrounding land masses (Lelieveld et al., 2001; Aswini et al., 2022; Tegtmeier et al., 2022) together with the large-scale monsoon circulations (McCreary et al., 2013). Air–sea exchange couples the atmospheric composition to biogeochemical processes in the surface IO and impacts elemental cycling (Kumar et al., 2020; Sarma et al., 2021). Another unique feature is the formation of vast oxygen minimum zones (OMZs) in the NIO (Naqvi et al., 2006; Sarma et al., 2020b; Gupta et al., 2021).

In the recent years, efforts have been made to review and synthesize various physical and biogeochemical processes that occur within the IO. Phillips et al. (2021) summarized new studies related to ocean–atmosphere coupling, highlighting the importance of small-scale processes on large-scale circulations, interactions between physical and biological processes as well as interaction between the surface and deep ocean. Vinayachandran et al. (2021) reviewed various features of upwelling along the coastal region of the IO, including underlying mechanisms and the impact on ecosystems. A review of the OMZ in the NIO and associated implications was carried out by Rixen et al. (2020) and, in particular, for the BoB by Sridevi and Sarma (2020) and Sarma and Udaya Bhaskar (2018). Tegtmeier

et al. (2022) provided a comprehensive review of the atmospheric gas phase composition over the IO and its coupling to trace gas emissions from the land and ocean.

This article aims to synthesize new knowledge accumulated over the past decade focusing on SOLAS relevant processes within the IO. Important research questions and knowledge gaps that SOLAS science can help to address are highlighted in Section 2, to assist the development of future IO research programs. Furthermore, we discuss interdisciplinary efforts required to advance our understanding of this oceanic basin with possible future recommendations in Section 3. The last section summarizes this synthesis by highlighting important results in recent years over the IO.

2. Recent progress and knowledge gaps

2.1. Monsoon and air–sea interactions

Ocean–atmosphere interactions are crucial for understanding the dynamics of the seasonally reversing monsoon system, and recent studies have advanced our understanding of these interactions. Monsoon variability can be driven thermodynamically through the exchange of heat and moisture that regulate the formation of deep convective clouds and rainfall or dynamically through the exchange of momentum and changes in circulation due to winds and ocean currents (Seager et al., 2010). The IO is the breeding ground of the intraseasonal variability of the monsoon, particularly the northward propagating monsoon intraseasonal oscillations (MISO) in summer and the eastward propagating Madden–Julian oscillations (MJO) in winter. The MISO and MJO modulate rainfall and cyclogenesis in South Asia and throughout the global tropics and evolve in response to the ocean–atmosphere interactions in the tropical IO (Roxy et al., 2013; Sharmila et al., 2013; Beal et al., 2020). The 2 oscillations induce sea surface temperature (SST) variations that are larger in the IO than in the Pacific Ocean, and incorporating the ocean–atmosphere interactions associated with these SST variations can improve the tropical rainfall predictability (DeMott et al., 2015).

While the ocean–atmosphere interactions that drive the intraseasonal variability have been extensively investigated and documented, the role of ocean dynamics needs to be better understood. For example, the BoB receives substantial freshwater through monsoon rainfall and river runoff, leading to a strong salinity stratification that causes a thin mixed layer and thick barrier layer. This stratification suppresses vertical ocean mixing, thereby affecting the monsoon intraseasonal SST variability and associated convection (Sengupta et al., 2016). Furthermore, initiation and early development of MJO events are influenced by ocean–atmospheric processes in the Seychelles–Chagos Thermocline Ridge (SCTR), where ocean dynamics and processes play a significant role (West et al., 2020). However, these regions with complex ocean–atmosphere interactions, such as the BoB, the SCTR, and the Indonesian through-flow have only scarce high-resolution observations, limiting our understanding of the key processes associated with the monsoon intraseasonal variability and curtailing model predictability. In this context, noble observations can help to quantify the effects of physical processes on dissolved gases and to separate

biological from physical impacts on constituents like oxygen, nitrogen, and carbon (Emerson et al., 2012).

Large gradients in nutrients were observed below and above stratified waters with higher concentration below than above. In such cases, nutrient diffusion through vertical eddy mixing is very important. While vertical eddy mixing has been used as a constant at the rate of 10^5 or 10^7 in the past, it can now be measured using noble gases (Hamme et al., 2017). Because noble gases have no biological function and respond only to physical processes, their measurements can be used to quantify these physical processes and estimate associated nutrient.

The SST variability is also an essential factor impacting monsoon variability on interannual and interdecadal-to-climate timescales. The Indian Ocean Dipole (IOD), a mode of variability characterized by the contrast in SSTs between the western and eastern IO, is a major driver of regional monsoon variability (Ashok et al., 2001) and influences the biogeochemical and ecosystem signatures (Parvathi et al., 2017). The IOD also plays a crucial role in the teleconnection between the El Niño Southern Oscillation (ENSO) and the monsoon (Cherchi et al., 2021). When positive IOD events (characterized by warmer SSTs in the western IO and cooler SSTs in the eastern IO) co-occur with El Niño events, the IO conditions act to counter El Niño's drought-inducing subsidence by enhancing moisture convergence over the Indian subcontinent, resulting in an average monsoon season (Ummenhofer et al., 2011; Chowdary et al., 2015). The IOD–monsoon relationship has strengthened in recent decades due to the nonuniform warming of the IO, while the ENSO–monsoon relationship has weakened (Cherchi et al., 2021). Meanwhile, the long-term monotonic warming of the IO with respect to the Indian subcontinent is altering the regional land–ocean thermal contrast, weakening the monsoon circulation and rainfall on interdecadal timescales (Roxy et al., 2015; Wang et al., 2021). These large-scale modes of climate variability (e.g., IOD, ENSO) influence the local atmospheric circulation which is coupled with the ocean, thereby modulating the ocean–lower atmospheric processes and biogeochemistry in the IO.

2.2. Oxygen minimum zones

The NIO, comprising the AS and the BoB, is characterized by low oxygen conditions below the subsurface layers (Wyrski, 1971; Sen Gupta and Naqvi, 1984). The western boundary upwelling system (off Somalia and Oman) during summer, and convective mixing during winter, supply nutrients to the surface waters resulting in high productivity in the AS (Lachkar et al., 2023). Conversely, upwelling in the BoB is suppressed by the presence of low-saline waters (Sarma et al., 2015), and primary productivity is supported through riverine input (Krishna et al., 2016) and mesoscale eddies (Prasanna Kumar, 2004; Sarma et al., 2019). Based on the work carried out till now, we sum up existing results in a map depicting the OMZ in the NIO (**Figure 2**), which includes the open ocean perennial nitrite-bearing waters of the AS, the coastal seasonal hypoxic zone along the west and east coast of India and few locations in the BoB. Nitrogen loss in the OMZ can happen either via classical denitrification (Ward et al., 2008; Ward et al., 2009) or




Figure 2. The Arabian Sea (AS) and the Bay of Bengal (BoB) both house low oxygen zones in the subsurface waters. The blue contour line demarcates $0.2 \mu\text{M}$ of secondary nitrite in the AS oxygen minimum zones (OMZs; Naqvi, 1991). The pink arrows show the seasonal changes in the AS OMZ as reported by Rixen et al. (2014) and Shenoy et al. (2020), whereas the red arrow indicates the northward expansion as reported by Lachkar et al. (2021). The shaded region shows the seasonal hypoxic zone in the Western Indian shelf (Naqvi et al., 2000) and Eastern Indian Shelf (Sarma et al., 2013a; Sarma et al., 2013b). The red star demarcates the station where secondary nitrite was detected by Bristow et al. (2017), and the blue squares are stations where Toyoda et al. (2023) observed high concentrations of N_2O in the BoB OMZ.

anammox (Jensen et al., 2011). The majority of the N-loss in the AS has been attributed to denitrification (Ward et al., 2009; Bulow et al., 2010) contributing to one-third of the global N-loss to the atmosphere (Naqvi et al., 2006; Ward et al., 2009; Jensen et al., 2011).

The intense OMZ and nitrite-bearing zone occurs in the northeastern AS associated with high primary production in the overhead and sinking fluxes to the depth (Naqvi, 1991; Morrison et al., 1999). While several theories have been put forth to explain this anomaly (McCreary et al., 2013 and references therein; Acharya and Panigrahi, 2016), through circulation, eddies, and transport of detritus from intense upwelling zone (off Oman) to east, using numerical models, there are no observational evidences are available. Recently, Sarma et al. (2020a) highlighted the role of cross-shelf transport of organic matter from the northwestern Indian shelf to the OMZ supporting its intensification using the backscatter data collected by Argo floats. However, the potential mechanism involved in transporting them and quantitative estimation of organic matter transported to the OMZ and its fate are unknown. Recent work has also shown the importance of AS high-saline water mass for supplying excess organic carbon and its possible role in sustaining the OMZ (Shetye et al., 2021). The interannual variability in OMZ is partly controlled by phytoplankton community composition and primary productivity due to ENSO events and biophysical

coupling (Vidya and Kurian, 2018; Kurian et al., 2020; Chndrasekhararao et al., 2022).

While the first demarcation of the nitrite-bearing zone was carried out by Naqvi (1991), the first observation on the seasonality was reported by de Souza et al. (1996). Recent studies have also reported seasonal variations in the western (Rixen et al., 2014) and southern end (Shenoy et al., 2020) of the nitrite-bearing zone (Figure 2). Apart from the perennial OMZs in the NIO, the western continental shelf of India (WCSI) houses the world's largest seasonal hypoxic zone covering $180,000 \text{ km}^2$ during the southwest monsoon (SWM) (Figure 2). Earlier studies indicated that hypoxic conditions along the WCSI might have intensified due to eutrophication mainly induced by anthropogenic input (Naqvi et al., 2000). However, recent work suggested that the formation of the seasonal hypoxic-anoxic zone is a natural process initiated through upwelling of deoxygenated waters during the summer monsoon (Gupta et al., 2021). Using time-series data and modeling, Parvathi et al. (2017) demonstrated that a positive IOD prevents the formation of coastal anoxia, while a negative or neutral IOD are among the essential conditions for its formation.

Anoxic conditions in the water column cause accumulation of hydrogen sulfide (Naqvi et al., 2000) and production of high concentrations of radiatively important gases namely, nitrous oxide (N_2O ; Naqvi et al., 2000), methane (CH_4 ; Shirodkar et al., 2018), and dimethylsulphide (DMS;

Bepari et al., 2020). Massive N-loss has been reported from the WCSI during the SWM (Naqvi et al., 2006; Bardhan and Naqvi, 2020; Sarkar et al., 2020) and accumulation of N_2O has been attributed to denitrification in the water column and sediments (Sudheesh et al., 2016), and also to chemolithoautotrophic denitrification (Pratihary et al., 2023). The emission of higher concentrations of radiatively important gases in this region has the potential to significantly contribute and modulate greenhouse gas budgets in the lower atmosphere. Thus, it is important to understand their long-term changes and feedback with processes at the air–sea interface.

Long-term variations of the strength of the OMZ and the denitrification in the AS have been debated in the literature and contrary statements can be found. Based on the available measured nonsystematic time-series dissolved oxygen (DO) data for more than 4 decades (1959–2004), Banse et al. (2014) reported a marked seasonality in the AS OMZ with higher oxygen concentrations during the northeast (NE) and spring intermonsoon than the SWM. The mean annual decrease in oxygen levels during NE and SWM seasons were -0.2 and -0.08 μM , respectively. Interestingly, the dataset also showed a statistically significant decrease of DO over 4 decades between 15°N and 20°N , but an opposing trend was noticed in the north of 21°N (Banse et al., 2014).

Based on the regional model, Lachkar et al. (2021) reported an expansion of the OMZ and attributed to the widespread warming of the sea surface resulting in a reduction in the ventilation of subsurface and intermediate layers (**Figure 2**). Recent observations in the Qatar exclusive economic zone revealed a decline in oxygen concentrations in the Persian (Arabian) Gulf over the past 4 decades accompanied by an expansion of seasonal near-bottom hypoxia (Al-Ansari et al., 2015; Lachkar et al., 2022). In contrast to regional models, the global models did not identify significant changes in the OMZ in the past several decades (Breitburg et al., 2018) and proposed the need for more observations to confirm the same. In recent years, modeling studies have revealed that the AS OMZ could shrink in the future due to an increased supply of DO from its southern boundary (Vallivattathillam et al., 2023).

The expansion of the OMZ can significantly influence nitrogen cycling, in particular the emission of nitrous oxide, which requires further evaluation. At the same time, decreasing oxygen concentrations in the northern AS have led to a regime shift in the pelagic ecosystem with dinoflagellates now dominating a usually diatom-dominated area (Rixen et al., 2020). As dinoflagellates are known to produce more DMS, it is paramount to assess the impact of such a regime shift on the production of DMS and their contribution to atmospheric aerosols from the region. The future of the OMZ intensity and expansion in the AS is largely unpredictable (Lachkar et al., 2023) in contrast to the other OMZs due to knowledge gaps with respect to ventilation, bacterial respiration, and supply of oxygen through water masses in the models. The biogeochemical feedback mechanisms such as iron inputs, N_2 fixation, and denitrification are neither well understood nor well represented in the models (Ulloa et al., 2012; Oschlies et al., 2019; Moffett and Landry, 2020; Wallmann et al., 2022). The biological

production is one of the important processes that control the OMZ and its changes in the past and future were contrastingly represented. For instance, a decline in the primary production in the AS was reported due to warming and weakening of upwelling (Roxy et al., 2016; Kwiatkowski et al., 2020), whereas insignificant trends in net primary production were observed (Sridevi et al., 2023) except in the southwestern basin during summer, due to the compensation of nutrients through the atmospheric deposition. The variations in the circulation, including speed and direction of currents, are equally important on formation of the OMZ. Therefore, significant work needs to be carried out through time-series observations, moorings, and repetition of transect observation to understand behavior of OMZ and its possible changes in the future due to climate change.

The OMZ in the BoB is as intense as the AS OMZ, albeit it does not host a perennial nitrite-bearing zone (Sarma et al., 2016; Bristow et al., 2017). Recent DO data measured from the Argo-based sensors revealed significant spatial variability of OMZ's boundary and its thickness in the BoB (Udaya Bhaskar et al., 2021). The lower and upper boundaries of the OMZ were observed between 100–800 m and 60–200 m, respectively, with a thickness of 80–650 m in the BoB (Udaya Bhaskar et al., 2021). Associated with salinity stratification a thicker OMZ was found in the northern BoB when compared to the southern BoB. A thick and intense OMZ was found in the northwestern BoB region (with oxygen levels <1.5 μM) associated with strong salinity stratification and high primary production in contrast to the NE BoB region with higher oxygen concentrations (2.5 μM). Like in the AS, cross-shelf transport of organic matter was observed in the northern BoB that further supports the OMZ (Udaya Bhaskar et al., 2021).

Though earlier studies reported low primary production (Gauns et al., 2005) in the BoB, recent measurements of dissolved organic carbon (DOC) exudation rates indicate total primary production (particulate organic carbon production and DOC exudation) is almost same as in the AS (Rao et al., 2021). Such high primary production in the BoB is supported by organic nutrients (Sarma et al., 2021; Rao and Sarma, 2022). The weaker OMZ in the BoB was attributed to the ballast of organic matter in association with mineral particles (Ittekkot et al., 1992). Recent measurements using STOX (switchable trace oxygen) sensors suggested that oxygen levels in the BoB are at nano molar levels (Bristow et al., 2017) suggesting that ballast of organic matter may not be potential reason and OMZ is not weaker in the BoB. This is also supported by measurement by Kurian et al. (2023), which reported the sinking carbon fluxes at 430 m and found that only $<5\%$ of the primary production is sinking to the depth below. Though such low oxygen levels were observed (Bristow et al., 2017), maintaining such low oxygen levels for longer times may be unlikely due to the presence of anticyclonic eddies that ventilate oxygen from surface waters to the OMZ (Sarma et al., 2018; Sarma and Udaya Bhaskar, 2018). On the other hand, a recent study by Toyoda et al. (2023) reported an extensive accumulation of N_2O in the OMZ of the BoB and attribute to the archeal ammonia oxidation

and denitrification in the micro-environment induced by riverine sinking particles during the monsoon and also raise the concern that the bay could be a source of N_2O to the atmosphere. The cross-shelf transport of organic matter (Udaya Bhaskar et al., 2021) may also bring sedimentary N_2O to the offshore region, however, there were no evidences to support. Considering the importance of air–sea interactions for the atmospheric gaseous budget, there is a need for sustained observations of climatically important gases in the low oxygen zones of the NIO.

2.3. Ocean biogeochemistry, air–sea exchange, and carbon cycling

The IO, particularly the NIO, is influenced by atmospheric forcing (monsoons), aerosol deposition, riverine outflow, as well as anthropogenic interferences. Associated interactions are complex and significantly impact surface water biogeochemical processes which include primary productivity, phytoplankton composition, and carbon dynamics. Recent observations and studies in the IO, that are highlighted in this section, significantly improved our understanding of the relevant biogeochemical processes.

The ocean–atmospheric exchange of CO_2 , as well as changes in the surface ocean carbon system parameters are one of the least studied topics for the IO (Valsala et al., 2020; Chakraborty et al., 2021; Valsala et al., 2021), in particular for the southern IO (Valsala et al., 2012). The BoB is characterized by a high spatial and seasonal variability of CO_2 levels and the rate of ocean acidification (Sarma et al., 2012; Sarma et al., 2015; Sarma et al., 2021). Experimental studies assessing the impact of increasing CO_2 levels on phytoplankton communities in the BoB revealed that coastal communities show resilience to a short-term increase in CO_2 with enhanced carbon fixation rates and particulate organic carbon accumulation (Biswas et al., 2011; Biswas et al., 2012; Biswas et al., 2017). While the BoB acts as a sink for CO_2 , the AS is a perennial source of CO_2 to the atmosphere due to upwelling and winter convective mixing (Valsala et al., 2013; Valsala and Maksyutov, 2013).

The prevalence of oligotrophy, in addition to change in pH, could potentially change the phytoplankton community and particularly low dissolved silicate availability, which could be one of the key players in a community shift. A series of experimental studies (Biswas et al., 2011; Biswas et al., 2012; Biswas et al., 2017; Shaik et al., 2017; Sharma et al., 2022a; Sharma et al., 2022b; Sharma et al., 2022c) have been published in the past years addressing the impacts of ocean acidification on natural phytoplankton communities. From all these studies, it is clear that the responses of natural phytoplankton communities to short-term and fast changes in pH/p CO_2 were highly diverse and species-specific. Moreover, the phytoplankton living close to the coastlines of both, the BoB and the AS, were more resilient and benefited from increased CO_2 supply. Some species of diatoms like *Thalassiosira* (Biswas et al., 2012; Biswas et al., 2017) and *Chaetoceros* (Shaik et al., 2017; Sharma et al., 2022a) exhibited higher resilience and growth rates indicating that they could be the species for future oceans. This higher resilience also

explains their bloom development in the coastal waters as they adapt to a wide level of fluctuations in seawater pH even within a diel period.

The AS acts as a source of CO_2 and usually shows higher levels in many places, hence could act as a natural laboratory for ocean acidification experiments. In most of the experiments in the coastal (Sharma et al., 2022a) and shelf waters (Sharma et al., 2022c), as well as in the open ocean (Sharma et al., 2022b; Sharma et al., 2023), many diatoms showed enhanced growth in response to increasing CO_2 supply suggesting their pre-acclimatization to such conditions. Some genera/species did not change their growth rates at variable pH/p CO_2 levels indicating higher resilience (Sharma et al., 2022b). Multiple stressor studies, which include other abiotic factors like atmospheric dust, and trace metals (e.g., copper and zinc) showed that some species are resilient even under such combined stress conditions indicating their high potential to cope with increasing pollution levels. It should be noted that some pennate diatoms with the potential to produce toxins were prominent in many cases, and such community shifts could have some adverse biogeochemical consequences. In the open ocean region of the BoB, such studies have not been carried out and the responses of the plankton community are largely unknown. In this region, low salinity values along with increasing CO_2 may result in different outcomes compared to the AS, where high salinity persists. Nevertheless, the available results could be used for ecological modeling on a regional scale for predicting future responses.

Atmospheric extreme events, such as cyclones, depressions, and eddies as well as large-scale atmospheric variability such as the IOD and ENSO, significantly impact ocean biogeochemical processes in the NIO on shorter timescales of a few days to months. These events modify upper ocean mixing causing variations in nutrient input, temperature, primary production, and CO_2 fluxes (e.g., Sarma et al., 2015; Sarma et al., 2019). For instance, a cyclone can churn up the upper ocean and bring nutrients and CO_2 -rich waters to the surface, which may enhance the CO_2 flux to the atmosphere. It can also lead to the absorption of CO_2 due to enhanced primary production, which would lead to a much smaller net impact of a tropical cyclone on CO_2 dynamics than believed previously (Lévy et al., 2012). Overall, previous investigations only covered snapshots of the full picture and therefore the average net impact of atmospheric extreme events is still unclear. The IO being a hub for such events, which are expected to increase in the future due to climate change (Swapna et al., 2022), underscores the need to study the impact of cyclones on biogeochemical processes in this basin.

In addition to decreasing pH, the increasing trend in SSTs (particularly in the mid-western AS; Roxy et al., 2015) has negatively impacted the overall primary productivity due to an enhanced stratification and restricted mixing that suppresses nutrient supply to the euphotic layers. In line with this, a comprehensive study covering a larger part of the IO from Dalpadado et al. (2021) confirmed an increasing warming trend in the last decade. Recently, Sridevi et al. (2023) noticed a decrease in net primary

production in the southwestern AS and southeastern BoB. The absence of a significant trend in the northern AS and BoB may be due to an increased deposition of anthropogenic aerosols (Sarma et al., 2022) compensating for the deficit of nutrient input through vertical mixing caused by stratification (Sridevi et al., 2023).

Based on observations over the last decade, several researchers have highlighted frequent occurrences of heterotrophic dinoflagellate *Noctiluca* blooms during the NE monsoon concluding that it might occur during all seasons (do Rosário Gomes et al., 2014 and references therein). The occurrence of *Noctiluca* blooms is not new to the AS (Subrahmanyam, 1954), however, their seasonal occurrences were not well captured in previous studies. A comparison of the phytoplankton community observed during 1990 (Sawant and Madhupratap, 1996) in the boreal winter (December and January) with recent data collected during February, a community shift (diatoms to dinoflagellate) due to anthropogenic activities was reported (do Rosário Gomes et al., 2014). The isotopic studies using carbon and nitrogen did not reveal the contribution of anthropogenic inputs to the AS during winter (Sarma et al., 2019). Some studies have confirmed the succession of the bloom development from diatom to *Noctiluca* (Lotliker et al., 2018; Lakshmi et al., 2021; Sridevi and Sarma, 2022). Other studies, however, found contradictory results showing diatom-dominated phytoplankton bloom during summer monsoon comparable to earlier JGOFS data in the central (Chowdhury et al., 2021; Silori et al., 2021) and eastern AS (Albin et al., 2022; Chowdhury and Biswas, 2023). Two recent studies on the phytoplankton community structure in these regions highlighted the key role of the strength of monsoon winds in governing the phytoplankton community and cell size, which in turn control the intensity of the carbon export flux (Chowdhury et al., 2021; Chowdhury and Biswas, 2023). Importantly, the contribution of nanoplanktonic coccolithophores to the total phytoplankton community from the western Indian shelf waters has also been highlighted recently (Chowdhury et al., 2022; Shetye et al., 2022). As these calcifiers significantly modulate the calcium carbonate pump, their role in carbon transport to the sediment needs to be evaluated.

A special feature of the IO is relatively high atmospheric deposition of aerosols which controls the surface biological processes (Banerjee and Kumar, 2016; Bange et al., 2024 and references therein). Over the last 2 decades, the northern IO experienced the globally highest rate of aerosol increase (Lotliker et al., 2018; Lakshmi et al., 2021; Sridevi and Sarma, 2021; Yadav et al., 2021; Sridevi and Sarma, 2022). Several cruise-based campaigns have been undertaken in the NIO (Tindale and Pease, 1999; Kumar and Sarin, 2010; Srinivas and Sarin, 2013; Aswini et al., 2020; Bikkina et al., 2020; Panda et al., 2022 and references therein) to evaluate the chemical composition of aerosols and their possible sources. These studies highlighted the dominance of mineral dust (derived from surrounding arid/semi-arid and desert regions) and anthropogenic aerosols (derived from fossil fuel

combustion and biomass burning) in both basins (Bange et al., 2024).

Large-scale mineral dust transport from the Arabian Peninsula and Northeast Africa has been reported during the SWM over the AS (Suresh et al., 2021; Aswini et al., 2022) while anthropogenic-rich aerosols are dominant in both basins during the NE monsoon when the continental outflow is maximum (Kumar et al., 2010; Bikkina et al., 2020; Nayak et al., 2022). Based on measured reactive nitrogen (inorganic nitrogen, IN) in aerosols and anthropogenic aerosol optical depth, atmospheric input of IN to the AS (1.7 TgN per year) and BoB (0.9 TgN per year) were estimated, contributing together approximately 30% of the total external sources of nitrogen to the NIO (Sarma et al., 2022). The contribution of atmospheric sources of total nitrogen may be much higher if organic nitrogen is also taken into account. Organic components (nitrogen, phosphorus) of aerosols over the IO are almost unknown in terms of their quantities as well as temporal variations. An important knowledge gap is the currently unconstrained macronutrient budget and its impact on the surface water biogeochemistry. The new production, supported by external sources of nitrogen (mainly IN), contributes about 23% and 53% of export production to the AS and BoB, respectively (Sarma et al., 2022).

Long-term studies in the northwestern coastal BoB revealed that rapid acidification of coastal waters has occurred due to the deposition of atmospheric aerosols (Sarma et al., 2015; Sarma et al., 2021). Very few studies have reported on the trace metal distribution over the IO, particularly from the NIO (Srinivas and Sarin, 2013; Panda et al., 2022 and references therein). Relatively high dry deposition fluxes of Fe, Mn, and Cu (757 , 38.8 , and $1.4 \mu\text{g m}^{-2} \text{day}^{-1}$, respectively) to the AS surface waters with high fractional solubility were detected during SWM campaigns (Panda et al., 2022). These observations, although being only a snapshot, provide new insight to the solubility behavior of trace elements and thus underscore the need for more observations at varying spatial and temporal scale. In recent years, pyrogenic Fe, which is mainly derived from biomass burning emissions and other anthropogenic activities, has been found to contribute significantly to the soluble Fe budget (Hamilton et al., 2022). In case of the IO and particularly the BoB, emissions from the agricultural residue burning from the Indo-Gangetic Plains could be a potential supplier of soluble Fe (Kumar et al., 2010). Therefore, more observations need to be initiated over the Indo-Gangetic Plains, focusing on the quantification of mineral dust associated trace metals and their solubility as well as radiogenic isotope signature of dust. Recent dust flux estimates over the NIO (Panda et al., 2024) show a good agreement with previous model-based flux estimates by Grand et al. (2015) ($2.5 \pm 0.5 \text{ g m}^{-2} \text{ year}^{-1}$), Jickells et al. (2005) ($2\text{--}5 \text{ g m}^{-2} \text{ year}^{-1}$), and Mahowald et al. (2005) ($3 \pm 1 \text{ g m}^{-2} \text{ year}^{-1}$). However, dust flux over the equatorial IO region is overestimated by a factor 5 ($2.5 \pm 1.8 \text{ g m}^{-2} \text{ year}^{-1}$) to 2 ($1.1 \pm 0.1 \text{ g m}^{-2} \text{ year}^{-1}$) compared to model estimates ($0.5 \pm 0.08 \text{ g m}^{-2} \text{ year}^{-1}$; Grand et al., 2015). The latter are impacted by model uncertainties and the large variability of aerosol

composition and solubility (Grand et al., 2015) highlighting the need for measurements to evaluate the model performance for the IO. In the absence of real-time aerosol deposition data along with accurate river discharge data, biogeochemical models fail to reproduce actual CO₂ fluxes (Sarma et al., 2023).

2.4. Greenhouse gases and pollution over the IO

Trace gases and their air–sea exchange play an important role for the IO regarding atmospheric chemistry, oceanic processes, chemistry–climate interactions, and stratospheric composition. Therefore, their spatial and temporal distributions, in both the lower atmosphere and upper ocean, as well as the parameters that control their distributions and fluxes must be understood. Synergies of in situ measurements (shipborne, balloon borne, and aircraft based), remote sensing observations (satellite and ground-based), and modeling (regional and global) have improved our understanding of pollutants and greenhouse gases (GHGs) over the IO and associated atmospheric chemistry and dynamics (Tegtmeier et al., 2022; Bange et al., 2024 and reference therein). In this section, the main characteristics of various pollutants, GHGs, and short-lived gases are discussed together with current knowledge gaps. As an illustrative example, surface CO and CH₄ distributions during the winter and post-monsoon seasons are based on 2018 (Girach et al., 2020) and 2010 (Mallik et al., 2013) ship-based measurements, respectively; recent (2019) satellite observations (MOPITT—Measurements of Pollution in the Troposphere; GOSAT—Greenhouse Gases Observing Satellite); and CAMS (Copernicus Atmosphere Monitoring Service) model reanalysis are shown in **Figure 3**. Pollutants and GHGs exhibit relatively higher concentrations over the northern IO as compared to the southern or equatorial IO, as seen here for CO and CH₄, primarily due to transport from continental sources, that is, fossil fuel and biomass burning (Girach et al., 2014; Girach and Nair, 2014; Chandra et al., 2016; Sahu et al., 2019). For CO, higher levels exist not only at the surface (up to 450 ppbv) but also in the upper boundary layer (up to 300 ppbv at 0.75 or 1.5 km) over the southeast BoB during winter owing to the influence of biomass burning over Southeast Asia (Srivastava et al., 2012). Similar to CO, the variability of surface O₃ is relatively well known. The latter is driven by various factors such as outflow from South and Southeast Asia, en route and in situ photochemical production, losses due to HO_x and halogen chemistry, vertical transport, boundary layer processes, and cyclone-induced stratospheric intrusion (e.g., Mallik et al., 2013; Das et al., 2016a; Das et al., 2016b; Girach et al., 2017; Ajayakumar et al., 2019; Mahajan et al., 2019a; Mahajan et al., 2019b; Girach et al., 2020b; Inamdar et al., 2020; Mahajan et al., 2021; Soni et al., 2022).

For most trace gases, only a few in situ observations are available over the IO. One example is NO₂, for which higher mixing ratios have been found over coastal regions (approximately 1.2 ppbv) and shipping lanes (up to 4 ppbv) compared to background values of 0.05–0.2 ppbv (Takashima et al., 2012; Georgoulias et al., 2020). Satellite observed NO₂ over international shipping lanes has been demonstrated to be a tracer of global economic changes

(de Ruyter de Wildt et al., 2012). Maritime shipping is known to emit also other pollutants including formaldehyde (HCHO) which has been found to be nearly twice as high along the major IO shipping lanes when compared to background levels (Gopikrishnan and Kuttippurath, 2021). Another example of limited in situ data is the pollutant SO₂, for which the few available observations suggest higher levels over the northern IO with prominent enhancements (0.05–0.15 DU) around the coastal regions and over volcanoes (Sarma et al., 2012; Chutia et al., 2022). Such elevated SO₂ abundances near active volcanoes could play a role in forming the recently discovered OH minimum in the eastern BoB during boreal spring (Kuttippurath et al., 2023).

Air–sea exchange impacts many trace gases over the IO such as CH₄, CO₂, N₂O, and volatile organic compounds (VOCs). Upwelling over the northern IO brings CH₄ (diffused from underlying sediments) up to the surface, which subsequently leads to sea-to-air CH₄ flux (Naqvi et al., 2010) in addition to strong continental sources. An important characteristic of atmospheric CH₄ is the seasonal variation of approximately 150 ppbv in the boundary layer (Kavitha et al., 2018), with model simulations suggesting that these are not present above 750 hPa (Guha et al., 2018). Overall CH₄ belongs to the group of rather understudied gases over the IO with surface satellite observations being uncertain (Girach et al., 2018) partially due to frequent cloudy conditions. For CO₂, another important GHG, variability in surface ocean partial pressure over the IO is mainly driven by SSTs with minor influence from the dynamics of dissolved inorganic carbon (Valsala and Murtugudde, 2015). The increasing level of CO₂ over the IO can change the intensity of IOD which has pronounced impact worldwide, including on the Indian summer monsoon (An et al., 2022).

Trace gas emissions from oceanic ecosystems are a major contributor to atmospheric VOCs over the IO, especially in the AS where high biological productivity prevails mainly due to upwelling and the influence of dust deposition (Singh and Ramesh, 2015; Guieu et al., 2019). The emissions through biogeochemical processes and sea–air exchange are influenced by incoming solar radiation, SST, wind speed, and microbial sources. Elevated levels of isoprene (i.e., the most abundant natural VOC) and alkenes are linked with higher chlorophyll-*a* (proxy for algae biomass) and phytoplankton species in seawater over the AS, which is an intense OMZ (Tripathi et al., 2020). According to the latest DMS climatology (Hulswar et al., 2022), the IO is one of the worldwide hotspots for DMS fluxes throughout all seasons. An interesting aspect of this updated climatology is that the DMS seawater concentrations and subsequent air–sea fluxes are less than those reported in the previous climatology (Lana et al., 2011), highlighting the uncertainty resulting from the IO being under sampled for DMS.

Relatively more O₃ and CO observations are available, however, surface observations of NO, NO₂, SO₂, NH₃, CH₄, CO₂, isoprene, monoterpenes, and other VOCs are very limited. Hence new in situ measurements and subsequent extensive studies combining observations with modeling exercises and satellite retrievals are highly desirable to




Figure 3. Spatial distribution of an air pollutant, CO, and a greenhouse gas, CH₄, during pre-winter and post-monsoon season over the Indian Ocean, respectively. The datasets are based on in situ observations (Girach et al., 2020 [a], Mallik et al., 2013 [d]), satellite observations (MOPITT [b], GOSAT [e]) and reanalysis output (CAM5 [c], [f]). Shown are the surface or boundary layer mixing ratios, except figure [e] which is a tropospheric average.

address some of the following knowledge gaps: (a) Radical concentrations, O₃ formation, and biogeochemical fluxes of O₃-precursors; (b) Modifications of fluxes of CO₂ and hydrocarbons including VOCs by marine heatwaves (Mignot et al., 2022); (c) Chemical transformation of DMS to SO₂; (d) Gas-to-particle conversions (e.g., SO₂ to sulfate aerosols) and heterogeneous chemistry; (e) Deposition of pollutants to the sea surface and subsequent impact on the production/consumption of other trace gases in the sea surface and their air–sea exchange. Finally, long-term trends of many gases over the IO are not well known and often only poorly understood. For instance, despite increasing CO emissions as seen in the inventories,

satellite, and in situ measurements suggest a declining trend in the surface CO (0.7% year⁻¹) over the northern IO (Worden et al., 2013; Girach et al., 2020a). Challenges of improved satellite retrievals of pollutants and GHGs over the ocean, especially under frequent cloudy conditions, need to be addressed for better characterizations of short- and long-term variability.

3. Interdisciplinary efforts and future recommendations to address the processes linked with SOLAS

The IO plays a key role in regional as well as global climate and is one of the most interesting areas to understand

ocean–atmospheric interactions that have a pivotal role in biological productivity. The positive trend in SSTs of the IO with pronounced spatial variability (Roxy et al., 2014 and references therein) leads to several questions about the future responses of the biological systems and fisheries, as well as the climatic feedback. Since a significant area of the IO, particularly the NIO, fall within exclusive economic zones, a strong collaboration among the IO rim countries would be beneficial for answering these questions.

One of the major impediments in this research area is the limited IO observing system with poor instrument coverage over space and time. This shortcoming could be addressed by increasing the number of instruments that measure critical parameters including nutrients, pH, oxygen, and fluorescence, such as Bio-Argo floats with an enhanced number of sensors. Argo floats deployed in the southern IO have added to our understanding of seasonal/interannual variability of environmental variables such as subsurface ocean temperature, salinity, and chlorophyll (Masuda, 2020; Prakash et al., 2020), however, there is a dearth of observational data such as on nutrients, their transformation mechanisms, interactions between the abiotic and biotic components in the region, as well as in situ production and air–sea exchange of climate-relevant gases. Therefore, more continuous collaborative efforts are required from the observational scientist to generate such comprehensive datasets. These datasets could be used for basic research on biogeochemical aspects and for the evaluation or as boundary conditions for regional general circulation or biogeochemical models. Combining such new observations with recent advances in atmospheric dynamics and modeling can improve our understanding of the impact of extreme events on ocean biogeochemistry.

Quantifying ocean–atmosphere interactions at interannual to interdecadal timescales, understanding their influences on regional monsoon patterns, and attributing them to natural variability or anthropogenic forcing is an ongoing challenge, mainly due to the lack of high-resolution long-term observations of dynamical variables (Beal et al., 2020). Further, widely used SST datasets can use statistical methods to fill gaps between sporadic observations for constructing ocean temperature products, without explicitly involving subsurface oceanic processes that are crucial for representing the IO variability at interannual and climate timescales (Yang et al., 2020). Hence, comprehensive observations that better-resolve upper ocean processes and numerical models that incorporate these processes are critical for an improved understanding of ocean–atmospheric coupling and prediction of monsoon variability. Long-term monitoring and high-resolution modeling of the critical processes of the near-surface ocean, including diurnal mixed layer and barrier layer variability are highly recommended future exercises to meet the need for improved monsoon forecasting.

Considering the present trends of global change, it is pertinent to understand whether the AS-OMZ has expanded and intensified over the last few decades, and whether the BoB-OMZ will turn into a denitrifying zone in the future. The influence of the increasing peripheral human population on the cycling of carbon, nitrogen, and

sulfur and the resultant impact on the ecosystem needs to be understood. Precise measurements of DO (using STOX sensors) are required to understand its seasonal variability and the redox transformations of nitrogen, sulfur, iron, and iodine as well as links to the sustenance of OMZ. As microbes play an important role in redox transformations, in-depth understanding of microbial ecology, including transcriptomics of the OMZ, needs to be a key element of future research. Studies in coastal regions focusing on the impact of anoxic and sulphidic conditions on primary and secondary producers are needed to understand the potential impact on fisheries. All these studies will require a dedicated platform for long-term observations and comparisons with other OMZs around the World.

Numerical models do not use real-time river discharge data to simulate biogeochemical processes in the BoB resulting in significant differences in the processes observed and simulated (Sarma et al., 2023). Trends in primary productivity are known to be impacted by the supply of nutrients from the atmosphere, submarine groundwater, and nitrogen fixation (Sridevi et al., 2023). Dissolved organic nutrients, DOC exudation, atmospheric deposition of nutrients, and the role of eddies for the supply of nutrients and oxygen ventilation are often not included in global models. Therefore, prediction of future changes of the OMZs or any other biogeochemical processes are questionable and significant improvements in the numerical models are warranted.

Biogeochemical and biological processes in the IO can also be altered by atmospheric deposition of particles, air-sea exchange of trace gases, as well as extraneous input of nutrients and toxic substances leading to potential climate feedbacks. Dedicated collaborations between atmospheric scientists studying aerosol-trace gas interactions and oceanic biogeochemists are needed to better understand the impacts of atmospheric pollution, dust and nutrients on the IO's biogeochemistry, and trace gas cycling. Considering relatively short lifetime of aerosols and other trace gases, large number of ship-based campaigns on varying seasons with large spatial converge should be undertaken for collection of aerosols as well as in situ and/or on-board measurements of atmospheric species. Model output of trace gases and aerosols over the IO should be compared with the observations in order to evaluate the representation of associated processes in regional and global models. Trace gas emissions, their conversions to aerosols and impact on clouds as well as feedbacks on ocean biogeochemistry are extremely important. This calls for extensive collaborations among researchers working on microphysics of aerosols as well as clouds, chemistry of trace gases, and biogeochemistry of the ocean.

The sources and distribution of macronutrients associated with organic phases and their role in biological processes are hardly known in the IO region except for a few reports (Shetye et al., 2019; Rao et al., 2021). Recent biogeochemical models have shown a substantial increase in nitrogen fixation when organic nutrients were considered (Hamilton et al., 2022). Thus, emphasis should be given toward the measurement of organic components

(nitrogen, phosphorus) in aerosols in the IO region. Aerosol pH is another important factor which plays a major role for the availability of nutrients from aerosol deposition. Thus, the quantification of aerosol acidity along with nutrients are equally important to assess the relative roles of various processes (source, transport, and chemical transformation) affecting nutrient abundance over the IO. In general, aerosol pH is estimated using thermodynamic equilibrium models (e.g., ISORROPIA), which require the particulate and gas phase composition of ambient atmosphere as input parameters (Shukla et al., 2023). Simultaneous measurements of aerosol chemical species and trace gases are highly recommended as they will provide more robust information on the aerosol pH and other factors controlling nutrient availability over the IO region. Another important source of uncertainty in the atmospheric nutrient supply are the deposition rates and velocities associated with varying particle sizes. Measurements of the latter over the Ocean are difficult and prone to problems of temporal and spatial variability. A promising approach is the use of natural radionuclides (^7Be) delivered to the ocean from the atmosphere. Simultaneous measurement of ^7Be in seawater and aerosols will be helpful in constraining the deposition velocities and reducing uncertainties in nutrient flux estimations over the IO.

Microcosm experiments conducted to evaluate the impact of atmospheric aerosols on coastal and open ocean waters revealed a variety of changes in their biogeochemical features including surface water pH, primary productivity, and phytoplankton composition (Kumari et al., 2021; Kumari et al., 2022; Sharma et al., 2022b). Several published studies indicated the significant impact of aerosols on surface ocean biogeochemistry, while most of the data on aerosol composition come from field surveys that provide only snapshot information. The impact of aeolian dust supply on the surface ocean biogeochemical processes, including primary productivity, is important on daily to seasonal timescale. More insight can be gained via micro and mesocosm experiments, in particular at regional scales. However, only a very few experiments have been reported for the IO region. Efforts should be put into a higher number of laboratory and onboard simulation experiments to identify processes and ecology parameters that impact surface water biogeochemical processes.

Within the IO, the equatorial and southern part of the IO are the least explored regions in terms of observational and modeling studies related to SOLAS processes. A series of comprehensive measurements and research initiatives are crucial in order to unravel the complex interaction of physical, chemical, and biological processes in this remote area. Understanding the biogeochemistry of the Southern IO is not only essential for advancing our knowledge of global ocean dynamics but also for constraining the global biogeochemical cycles of carbon as well as macro- and micronutrients. It is imperative that we prioritize scientific exploration and measurement campaigns in this region to address the critical gaps in our understanding of its ecosystems and their interactions with the Earth's climate system.

4. Summary

This article provides a synthesis of observational and modeling results for the IO region related to SOLAS processes and obtained over the last 10 years. The significant increase in the number of observational as well as modeling studies found for this time frame has contributed to a better understanding of the atmosphere-oceanic coupled processes in this basin. Some of the key findings are summarized as follows:

1. Mesoscale events like IOD play a crucial role for the teleconnection between the ENSO and the Indian monsoon. When positive IOD events co-occur with El Niño events, the IO conditions act to counter El Niño's drought-inducing subsidence by enhancing moisture convergence over the Indian subcontinent, resulting in an average monsoon season.
2. One of the niche areas of the IO is the seasonal and perennial OMZs in the AS and possibly in the BoB. Modeling and observational results from the last decade suggest an expansion of the OMZ at the northern part of the AS, which is attributed to the widespread surface warming causing a reduction in the ventilation of subsurface and intermediate layers. The seasonal hypoxic-anoxic zone in the WCSI is primarily driven by the natural process of upwelling of deoxygenated waters during the summer monsoon. Oxygen data from the Argo-based sensors has shed new light on the seasonal variability of low oxygen zones in the BoB, which are mainly controlled by stratification, primary and export productions, organic matter decomposition, and eddy-driven mixing.
3. Ocean acidification experiments in the NIO revealed that the responses of natural phytoplankton to short-term and fast changes in pH/pCO₂ are highly diverse. Species-specific responses were found with phytoplankton living close to the coastline being more resilient and benefiting from enhanced acidification. In addition, the impact of decreasing pH with increasing SSTs on primary productivity is significant in the southwestern AS and southeastern BoB, but not in the northern parts of the 2 basins. The missing trend in the northern parts may be attributed to an increased deposition of anthropogenic aerosols compensating for the deficit of nutrients input through vertical mixing caused by stratification.
4. Significant advancements have been made in measurements of aeolian nutrient supply over the IO, particularly in the NIO, which may be considered as baseline information necessary for the regional and global scale models. The real-time flux measurements are far off from the model simulated output, particularly in the equatorial and southern part of the IO, suggesting the need for a large number of campaign-based observations in future.
5. Atmospheric pollutants and GHGs exhibit relatively higher concentrations over the northern IO as compared to the southern or equatorial IO primarily due

to transport from continental sources, that is, fossil fuel and biomass burning.

- Trace gases like SO₂ show enhanced abundances near coastal and volcanic regions, while some pollutants (e.g., HCHO, NO₂) display clearly increased concentrations along the major IO shipping lanes. Recent observations highlighted that IO is one of the worldwide hotspots for DMS fluxes throughout all seasons.

Despite all efforts over the past decade, there are significant knowledge gaps and major research question that need to be addressed in the near future. Although there is a plethora of studies for the NIO, almost negligible work has been reported for the southern part of the IO. More observational studies in the southern region of the IO should be a major research focus for the future. In addition, long-term dedicated, observational time-series of physical, chemical, and biological parameters in the IO are necessary for addressing some of the most pressing research questions. These observations should be undertaken in a collaborative effort from all Rim countries surrounding the IO, which is important for their economic and sustainable development.

Data accessibility statement

All data used in this review article are published and the source of data have been indicated through article citation. All data are available in public domain.

Acknowledgments

The authors (AK, ST, and SOF) thank Lisa and Cliff for inviting them as lead authors and providing useful suggestions/comments during the paper preparation. The authors thank the 3 anonymous reviewers and editor for their valuable comments/suggestions, which helped in improving this article. AK thanks Garima Shukla for assisting in formatting of the manuscript and references. NIO contribution number is 7254.

Funding

AK and VVSSS thank the Department of Science and Technology, Government of India, for partial funding under the Core research Grant (File no. CRG/2021/003307) during the study period. This publication resulted in part from support from the U.S. National Science Foundation (Grant OCE-1840868) to the Scientific Committee on Oceanic Research (SCOR).

Competing interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this article.

Author contributions

All authors have approved the submitted version for publication: AK, ST, and SOF drafted the chapter outline and each one coordinated writing for different sections. MKR wrote Section 2.1. SOF coordinated writing of Section 2.2 with inputs from DMS, SK, and VVSSS. AK coordinated writing of Section 2.3 with inputs from HB and VVSSS. ST coordinated writing of Section 2.4 with inputs from IG

and CAM. AK and ST wrote Abstract, Introduction, Conclusion, Section 3 with input from CAM, IG, and other coauthors. All the coauthors have commented during revision of the manuscript.

References

- Acharya, SS, Panigrahi, MK.** 2016. Eastward shift and maintenance of Arabian Sea oxygen minimum zone: Understanding the paradox. *Deep Sea Research Part I: Oceanographic Research Papers* **115**: 240–252. DOI: <http://dx.doi.org/10.1016/j.dsr.2016.07.004>.
- Ajayakumar, RS, Nair, PR, Girach, IA, Sunilkumar, SV, Muhsin, M, Chandran, PS.** 2019. Dynamical nature of tropospheric ozone over a tropical location in Peninsular India: Role of transport and water vapour. *Atmospheric Environment* **218**(D1): 117018.
- Al-Ansari, EM, Rowe, G, Abdel-Moati, M, Yigiterhan, O, Al-Maslamani, I, Al-Yafei, M, Al-Shaikh, I, Upstill-Goddard, RC.** 2015. Hypoxia in the Central Arabian gulf Exclusive Economic Zone (EEZ) of Qatar during summer season. *Estuarine, Coastal and Shelf Science* **159**: 60–68. DOI: <http://dx.doi.org/10.1016/j.ecss.2015.03.022>.
- Albin, KJ, Jyothibabu, R, Alok, KT, Santhikrishnan, S, Sarath, S, Sudheesh, V, Sherin, CK, Balachandran, KK, Asha Devi, CR, Gupta, GVM.** 2022. Distinctive phytoplankton size responses to the nutrient enrichment of coastal upwelling and winter convection in the eastern Arabian Sea. *Progress in Oceanography* **203**: 102779. DOI: <http://dx.doi.org/10.1016/j.pocean.2022.102779>.
- An, S-I, Park, H-J, Kim, S-K, Shin, J, Yeh, S-W, Kug, J-S.** 2022. Intensity changes of Indian Ocean dipole mode in a carbon dioxide removal scenario. *npj Climate and Atmospheric Science* **5**(1): 1–8. DOI: <http://dx.doi.org/10.1038/s41612-022-00246-6>.
- Ankur, K, Busireddy, NKR, Osuri, KK, Niyogi, D.** 2020. On the relationship between intensity changes and rainfall distribution in tropical cyclones over the North Indian Ocean. *International Journal of Climatology* **40**(4): 2015–2025.
- Ashok, K, Guan, Z, Yamagata, T.** 2001. Impact of the Indian Ocean dipole on the relationship between the Indian monsoon rainfall and ENSO. *Geophysical Research Letters* **28**(23): 4499–4502. DOI: <http://dx.doi.org/10.1029/2001GL013294>.
- Aswini, MA, Kumar, A, Das, SK.** 2020. Quantification of long-range transported aeolian dust towards the Indian peninsular region using satellite and ground-based data—A case study during a dust storm over the Arabian Sea. *Atmospheric Research* **239**: 104910. DOI: <http://dx.doi.org/10.1016/j.atmosres.2020.104910>.
- Aswini, MA, Tiwari, S, Singh, U, Kurian, S, Patel, A, Gunthe, SS, Kumar, A.** 2022. Aeolian dust and sea salt in marine aerosols over the Arabian Sea during the southwest monsoon: Sources and spatial variability. *ACS Earth and Space Chemistry* **6**(4): 1044–1058. DOI: <http://dx.doi.org/10.1021/acsearthspacechem.1c00400>.

- Banerjee, P, Kumar, SP.** 2016. ENSO modulation of inter-annual variability of dust aerosols over the North-west Indian Ocean. *Journal of Climate* **29**(4): 1287–1303. DOI: <http://dx.doi.org/10.1175/JCLI-D-15-0039.1>.
- Bange, H, Arévalo-Martínez, DL, Bikkina, S, Marandino, CA, Sarin, M, Tegtmeier, S, Valsala, V.** 2024. Chapter 14: Air-sea exchange and its impacts on biogeochemistry in the Indian Ocean, in Ummenhofer, CC, Hood, RR eds., *The Indian Ocean and its role in the global climate system*. Amsterdam, the Netherlands: Elsevier. DOI: <https://doi.org/10.1016/B978-0-12-822698-8.00010-X>.
- Banase, K, Naqvi, SWA, Narvekar, PV, Postel, JR, Jayakumar, DA.** 2014. Oxygen minimum zone of the open Arabian Sea: Variability of oxygen and nitrite from daily to decadal timescales. *Biogeosciences* **11**(8): 2237–2261. DOI: <http://dx.doi.org/10.5194/bg-11-2237-2014>.
- Bardhan, P, Naqvi, SWA.** 2020. Nitrogen and carbon cycling over the western continental shelf of India during seasonal anoxia: A stable isotope approach. *Journal of Marine Systems* **207**(38): 103144.
- Beal, LM, Vialard, J, Roxy, MK, Li, J, Andres, M, Annamalai, H, Parvathi, V.** 2020. A road map to IndOOS-2: Better observations of the rapidly warming Indian Ocean. *Bulletin of the American Meteorological Society* **101**(11): E1891–E1913.
- Bepari, KF, Shenoy, DM, Rao, AC, Kurian, S, Gauns, MU, Naik, BR, Naqvi, SWA.** 2020. Dynamics of dimethylsulphide and associated compounds in the coastal waters of Goa, west coast of India. *Journal of Marine Systems* **207**(C11): 103228.
- Bikkina, P, Sarma, VVSS, Kawamura, K, Bikkina, S, Kunwar, B, Sherin, CK.** 2020. Chemical characterization of wintertime aerosols over the Arabian Sea: Impact of marine sources and long-range transport. *Atmospheric Environment* **239**: 117749. DOI: <http://dx.doi.org/10.1016/j.atmosenv.2020.117749>.
- Biswas, H, Alexander, C, Yadav, K, Ramana, VV, Prasad, VR, Acharyya, T, Babu, PR.** 2011. The response of a natural phytoplankton community from the Godavari River estuary to increasing CO₂ concentration during the pre-monsoon period. *Journal of Experimental Marine Biology and Ecology* **407**(2): 284–293.
- Biswas, H, Gadi, SD, Ramana, VV, Bharathi, MD, Priyan, RK, Manjari, DT, Kumar, MD.** 2012. Enhanced abundance of tintinnids under elevated CO₂ level from coastal Bay of Bengal. *Biodiversity and Conservation* **21**(5): 1309–1326.
- Biswas, H, Shaik, AUR, Bandyopadhyay, D, Chowdhury, N.** 2017. CO₂ induced growth response in a diatom dominated phytoplankton community from SW Bay of Bengal coastal water. *Estuarine, Coastal and Shelf Science* **198**(Part A): 29–42.
- Breitburg, D, Levin, LA, Oschlies, A, Grégoire, M, Chavez, FP, Conley, DJ, Garçon, V, Gilbert, D, Gutiérrez, D, Isensee, K, Jacinto, GS, Limburg, KE, Montes, I, Naqvi, SWA, Pitcher, GC, Rabalais, NN, Roman, MR, Rose, KA, Seibel, BA, Telszewski, M, Yasuhara, M, Zhang, J.** 2018. Declining oxygen in the global ocean and coastal waters. *Science* **359**(6371). DOI: <http://dx.doi.org/10.1126/science.aam7240>.
- Bristow, LA, Callbeck, CM, Larsen, M, Altabet, MA, Dekaezemacker, J, Forth, M, Gauns, M, Glud, RN, Kuypers, MMM, Lavik, G, Milucka, J, Naqvi, SWA, Pratihary, A, Revsbech, NP, Thamdrup, B, Treusch, AH, Canfield, D.** 2017. N₂ production rates limited by nitrite availability in the Bay of Bengal oxygen minimum zone. *Nature Geoscience* **10**(1): 24–29. DOI: <http://dx.doi.org/10.1038/ngeo2847>.
- Bulow, SE, Rich, JJ, Naik, HS, Pratihary, AK, Ward, BB.** 2010. Denitrification exceeds anammox as a nitrogen loss pathway in the Arabian Sea oxygen minimum zone. *Deep Sea Research Part I: Oceanographic Research Papers* **57**(3): 384–393.
- Cai, W, Santoso, A, Wang, G, Weller, E, Wu, L, Ashok, K, Yamagata, T.** 2014. Increased frequency of extreme Indian Ocean dipole events due to greenhouse warming. *Nature* **510**(7504): 254–258.
- Cai, W, Yang, K, Wu, L, Huang, G, Santoso, A, Ng, B, Yamagata, T.** 2021. Opposite response of strong and moderate positive Indian Ocean dipole to global warming. *Nature Climate Change* **11**(1): 27–32.
- Chakraborty, K, Valsala, V, Bhattacharya, T, Ghosh, J.** 2021. Seasonal cycle of surface ocean pCO₂ and pH in the northern Indian Ocean and their controlling factors. *Progress in Oceanography* **198**: 102683. DOI: <http://dx.doi.org/10.1016/j.pocean.2021.102683>.
- Chandra, N, Lal, S, Venkataramani, S, Patra, PK, Sheel, V.** 2016. Temporal variations of atmospheric CO₂ and CO at Ahmedabad in western India. *Atmospheric Chemistry and Physics* **16**(10): 6153–6173. DOI: <http://dx.doi.org/10.5194/acp-16-6153-2016>.
- Cherchi, A, Terray, P, Ratna, SB, Sankar, S, Sooraj, KP, Behera, S.** 2021. Chapter 8—Indian Ocean dipole influence on Indian summer monsoon and ENSO: A review, in Chowdary, J, Parekh, A, Gnanaseelan, C eds., *Indian summer monsoon variability*. Cambridge, MA: Elsevier: 157–182. DOI: <http://dx.doi.org/10.1016/B978-0-12-822402-1.00011-9>.
- Chndrasekhararao, AV, Kurian, S, Vidya, PJ, Gauns, M.** 2022. Seasonal and inter-annual variability of chemotaxonomic marker pigments in the north-eastern Arabian Sea. *Deep Sea Research Part I: Oceanographic Research Papers* **179**: 103679. DOI: <http://dx.doi.org/10.1016/j.dsr.2021.103679>.
- Chowdary, JS, Bandgar, AB, Gnanaseelan, C, Luo, J-J.** 2015. Role of tropical Indian Ocean air–sea interactions in modulating Indian summer monsoon in a coupled model. *Atmospheric Science Letters* **16**(2): 170–176. DOI: <http://dx.doi.org/10.1002/asl.2561>.
- Chowdhury, M, Biswas, H.** 2023. A coherent status of summer monsoon phytoplankton communities (2017–2018) along the Western Indian continental

- shelf: Implications for fisheries. *Science of the Total Environment* **878**: 162963.
- Chowdhury, M, Biswas, H, Mitra, A, Silori, S, Sharma, D, Bandyopadhyay, D, Shaik, AUR, Fernandes, V, Narvekar, J.** 2021. Southwest monsoon-driven changes in the phytoplankton community structure in the Central Arabian Sea (2017–2018): After two decades of JGOFs. *Progress in Oceanography* **197**: 102654.
- Chowdhury, M, Biswas, H, Sharma, D, Silori, S, Winter, A.** 2022. Distribution of extant coccolithophores from the northwest continental shelf of India during the summer monsoon. *Phycologia* **61**(3): 284–298.
- Chutia, L, Ojha, N, Girach, I, Pathak, B, Sahu, LK, Sarangi, C, Flemming, J, da Silva, A, Bhuyan, PK.** 2022. Trends in sulfur dioxide over the Indian sub-continent during 2003–2019. *Atmospheric Environment* **284**: 119189. DOI: <http://dx.doi.org/10.1016/j.atmosenv.2022.119189>.
- Dalpadado, P, Arrigo, KR, van Dijken, GL, Gunasekara, SS, Ostrowski, M, Bianchi, G, Sperfeld, E.** 2021. Warming of the Indian Ocean and its impact on temporal and spatial dynamics of primary production. *Progress in Oceanography* **198**: 102688. DOI: <http://dx.doi.org/10.1016/j.pocean.2021.102688>.
- Das, SS, Ratnam, MV, Uma, KN, Patra, AK, Subrahmanyam, KV, Girach, IA, Ramkumar, G.** 2016b. Stratospheric intrusion into the troposphere during the tropical cyclone Nilam (2012). *Quarterly Journal of the Royal Meteorological Society* **142**(698): 2168–2179.
- Das, SS, Ratnam, MV, Uma, KN, Subrahmanyam, KV, Girach, IA, Patra, AK, Ramkumar, G.** 2016a. Influence of tropical cyclones on tropospheric ozone: Possible implications. *Atmospheric Chemistry and Physics* **16**(8): 4837–4847.
- DeMott, CA, Klingaman, NP, Woolnough, SJ.** 2015. Atmosphere-ocean coupled processes in the Madden-Julian oscillation. *Reviews of Geophysics* **53**(4): 1099–1154. DOI: <http://dx.doi.org/10.1002/2014RG000478>.
- de Ruyter de Wildt, M, Eskes, H, Boersma, KF.** 2012. The global economic cycle and satellite-derived NO₂ trends over shipping lanes. *Geophysical Research Letters* **39**(1). DOI: <http://dx.doi.org/10.1029/2011GL049541>.
- de Souza, SN, Kumar, MD, Sardesai, S, Sarma, VVSS, Shirodkar, PV.** 1996. Seasonal variability in oxygen and nutrients in the Central and Eastern Arabian Sea. *Current Science* **71**(11): 847–851.
- Deshpande, M, Singh, VK, Ganadhi, MK, Roxy, MK, Emmanuel, R, Kumar, U.** 2021. Changing status of tropical cyclones over the North Indian Ocean. *Climate Dynamics* **57**(3): 3545–3567.
- Dhame, S, Taschetto, AS, Santoso, A, Meissner, KJ.** 2020. Indian Ocean warming modulates global atmospheric circulation trends. *Climate Dynamics* **55**(1): 2053–2073.
- do Rosário Gomes, H, Goes, JI, Matondkar, SP, Buskey, EJ, Basu, S, Parab, S, Thoppil, P.** 2014. Massive outbreaks of *Noctiluca scintillans* blooms in the Arabian Sea due to spread of hypoxia. *Nature Communications* **5**(1): 4862.
- Doyle, T.** 2018. Blue economy and the Indian Ocean rim. *Journal of the Indian Ocean Region* **14**(1): 1–6.
- Dube, SK, Jain, I, Rao, AD, Murty, TS.** 2009. Storm surge modelling for the Bay of Bengal and Arabian Sea. *Natural Hazards* **51**(1): 3–27.
- Emerson, S, Ito, T, Hamme, RC.** 2012. Argon supersaturation indicates low decadal-scale vertical mixing in the ocean thermocline. *Geophysical Research Letters* **39**(18).
- Gangal, M, Suri, V, Arthur, R.** 2023. How well does Indian fisheries policy engage with fisheries biology? Exploring the science-policy interface of coastal capture fisheries along the west coast of India. *Marine Policy* **156**: 105796.
- Gauns, M, Madhuratap, M, Ramaiah, N, Jyothibabu, R, Fernandes, V, Paul, JT, Prasanna Kumar, S.** 2005. Comparative accounts of biological productivity characteristics and estimates of carbon fluxes in the Arabian Sea and the Bay of Bengal. *Deep Sea Research Part II: Topical Studies in Oceanography* **52**: 2003–2017.
- Georgoulias, AK, Boersma, KF, van Vliet, J, Zhang, X, van der, AR, Zanis, P, de Laat, J.** 2020. Detection of NO₂ pollution plumes from individual ships with the TROPOMI/S5P satellite sensor. *Environmental Research Letters* **15**(12): 124037. DOI: <http://dx.doi.org/10.1088/1748-9326/abc445>.
- Girach, IA, Nair, PR.** 2014. On the vertical distribution of carbon monoxide over Bay of Bengal during winter: Role of water vapour and vertical updrafts. *Journal of Atmospheric and Solar-Terrestrial Physics* **117**: 31–47.
- Girach, IA, Nair, PR, Ojha, N, Sahu, LK.** 2020a. Tropospheric carbon monoxide over the Northern Indian Ocean during winter: Influence of inter-continental transport. *Climate Dynamics* **54**(11): 5049–5064.
- Girach, IA, Nair, VS, Babu, SS, Nair, PR.** 2014. Black carbon and carbon monoxide over Bay of Bengal during W_ICARB: Source characteristics. *Atmospheric Environment* **94**: 508–517.
- Girach, IA, Ojha, N, Nair, PR, Pozzer, A, Tiwari, YK, Kumar, KR, Lelieveld, J.** 2017. Variations in O₃, CO, and CH₄ over the Bay of Bengal during the summer monsoon season: Shipborne measurements and model simulations. *Atmospheric Chemistry and Physics* **17**(1): 257–275.
- Girach, IA, Ojha, N, Nair, PR, Tiwari, YK, Kumar, KR.** 2018. Variations of trace gases over the Bay of Bengal during the summer monsoon. *Journal of Earth System Science* **127**(1): 15. DOI: <http://dx.doi.org/10.1007/s12040-017-0915-y>.
- Girach, IA, Tripathi, N, Nair, PR, Sahu, LK, Ojha, N.** 2020b. O₃ and CO in the South Asian outflow over the Bay of Bengal: Impact of monsoonal dynamics

- and chemistry. *Atmospheric Environment* **233**(D19): 117610.
- Gopikrishnan, GS, Kuttippurath, J.** 2021. A decade of satellite observations reveal significant increase in atmospheric formaldehyde from shipping in Indian Ocean. *Atmospheric Environment* **246**: 118095.
- Grand, MM, Measures, CI, Hatta, M, Hiscock, WT, Buck, CS, Landing, WM.** 2015. Dust deposition in the eastern Indian Ocean: The ocean perspective from Antarctica to the Bay of Bengal. *Global Biogeochemical Cycles* **29**(3): 357–374.
- Guha, T, Tiwari, YK, Valsala, V, Lin, X, Ramonet, M, Mahajan, A, Datye, A, Kumar, KR.** 2018. What controls the atmospheric methane seasonal variability over India? *Atmospheric Environment* **175**: 83–91. DOI: <http://dx.doi.org/10.1016/j.atmosenv.2017.11.042>.
- Guieu, C, Al Azhar, M, Aumont, O, Mahowald, N, Levy, M, Ethé, C, Lachkar, Z.** 2019. Major impact of dust deposition on the productivity of the Arabian Sea. *Geophysical Research Letters* **46**(12): 6736–6744. DOI: <http://dx.doi.org/10.1029/2019GL082770>.
- Gupta, GVM, Jyothibabu, R, Ramu, CV, Reddy, AY, Balachandran, KK, Sudheesh, V, Kumar, S, Chari, NVHK, Bepari, KF, Marathe, PH, Reddy, BB, Vijayan, AK.** 2021. The world's largest coastal deoxygenation zone is not anthropogenically driven. *Environmental Research Letters* **16**(5): 054009. DOI: <http://dx.doi.org/10.1088/1748-9326/abe9eb>.
- Hamilton, DS, Perron, MM, Bond, TC, Bowie, AR, Buchholz, RR, Guieu, C, Mahowald, NM.** 2022. Earth, wind, fire, and pollution: Aerosol nutrient sources and impacts on ocean biogeochemistry. *Annual Review of Marine Science* **14**: 303–330.
- Hamme, RC, Emerson, SR, Severinghaus, JP, Long, MC, Yashayaev, I.** 2017. Using noble gas measurements to derive air-sea process information and predict physical gas saturations. *Geophysical Research Letters* **44**(19): 9901–9909.
- Hermes, JC, Masumoto, Y, Beal, LM, Roxy, MK, Vialard, J, Andres, M, Annamalai, H, Behera, S, D'Adamo, N, Doi, T, Feng, M, Han, W, Hardman-Mountford, N, Hendon, H, Hood, R, Kido, S, Lee, C, Lee, T, Lengaigne, M, Li, J, Lumpkin, R, Navaneeth, KN, Milligan, B, McPhaden, MJ, Ravichandran, M, Shinoda, T, Singh, A, Sloyan, B, Stratton, PG, Subramanian, AC, Thurston, S, Tozuka, T, Ummenhofer, CC, Unnikrishnan, AS, Venkatesan, R, Wang, D, Wiggert, J, Yu, L, Yu, W.** 2019. A sustained ocean observing system in the Indian Ocean for climate related scientific knowledge and societal needs. *Frontiers in Marine Science* **6**: 355. DOI: <https://doi.org/10.3389/fmars.2019.00355>.
- Hood, RR, Bange, HW, Beal, L, Beckley, LE, Burkill, P, Cowie, GL, D'Adamo, N, Ganssen, G, Hendon, H, Hermes, J, Honda, M, McPhaden, M, Roberts, M, Singh, S, Urban, E, Yu, W.** 2015. *Science plan of the second International Indian Ocean Expedition (IIOE-2): A basin-wide research program*. Newark, DE: Scientific Committee on Oceanic Research.
- Hood, RR, Urban, ER, McPhaden, MJ, Su, D, Raes, E.** 2016. The 2nd International Indian Ocean Expedition (IIOE-2): Motivating new exploration in a poorly understood basin. *Limnology and Oceanography Bulletin* **25**(4): 117–124.
- Hulswar, S, Simó, R, Galí, M, Bell, TG, Lana, A, Inamdar, S, Halloran, PR, Manville, G, Mahajan, AS.** 2022. Third revision of the global surface seawater dimethyl sulfide climatology (DMS-Rev3). *Earth System Science Data* **14**(7): 2963–2987. DOI: <http://dx.doi.org/10.5194/essd-14-2963-2022>.
- Inamdar, S, Tinel, L, Chance, R, Carpenter, LJ, Sabu, P, Chacko, R, Mahajan, AS.** 2020. Estimation of reactive inorganic iodine fluxes in the Indian and Southern Ocean marine boundary layer. *Atmospheric Chemistry and Physics* **20**(20): 12093–12114.
- Ittekkot, V, Haake, B, Bartsch, M, Nair, RR, Ramaswamy, V.** 1992. Organic carbon removal in the sea: The continental connection. *Geological Society, London, Special Publications* **64**(1): 167–176. DOI: <http://dx.doi.org/10.1144/gsl.sp.1992.064.01.11>.
- Jensen, MM, Lam, P, Revsbech, NP, Nagel, B, Gaye, B, Jetten, MS, Kuypers, MM.** 2011. Intensive nitrogen loss over the Omani Shelf due to anammox coupled with dissimilatory nitrite reduction to ammonium. *The ISME Journal* **5**(10): 1660–1670. DOI: <http://dx.doi.org/10.1038/ismej.2011.44>.
- Jickells, TD, An, ZS, Andersen, KK, Baker, AR, Bergametti, G, Brooks, N, Torres, R.** 2005. Global iron connections between desert dust, ocean biogeochemistry, and climate. *Science* **308**(5718): 67–71.
- Jyoti, J, Swapna, P, Krishnan, R.** 2023. North Indian Ocean Sea level rise in the past and future: The role of climate change and variability. *Global and Planetary Change* **228**(502–505): 104205.
- Kavitha, M, Nair, PR, Girach, IA, Aneesh, S, Sijikumar, S, Renju, R.** 2018. Diurnal and seasonal variations in surface methane at a tropical coastal station: Role of mesoscale meteorology. *Science of the Total Environment* **631–632**: 1472–1485. DOI: <http://dx.doi.org/10.1016/j.scitotenv.2018.03.123>.
- Krishna, KS, Ismaiel, M, Srinivas, K, Rao, DG, Mishra, J, Saha, D.** 2016. Sediment pathways and emergence of Himalayan source material in the Bay of Bengal. *Current Science* **110**(3): 363–372. Available at <https://www.jstor.org/stable/24906781>. Accessed January 23, 2023.
- Kumar, A, Sarin, MM.** 2010. Atmospheric water-soluble constituents in fine and coarse mode aerosols from high-altitude site in Western India: Long-range transport and seasonal variability. *Atmospheric Environment* **44**(10): 1245–1254. DOI: <http://dx.doi.org/10.1016/j.atmosenv.2009.12.035>.
- Kumar, A, Sarin, MM, Srinivas, B.** 2010. Aerosol iron solubility over Bay of Bengal: Role of anthropogenic sources and chemical processing. *Marine Chemistry* **121**(1): 167–175. DOI: <http://dx.doi.org/10.1016/j.marchem.2010.04.005>.
- Kumar, A, Suresh, K, Rahaman, W.** 2020. Geochemical characterization of modern aeolian dust over the

- Northeastern Arabian Sea: Implication for dust transport in the Arabian Sea. *Science of the Total Environment* **729**: 138576.
- Kumari, VR, Neeraja, B, Rao, DN, Ghosh, VRD, Rajula, GR, Sarma, VVSS.** 2022. Impact of atmospheric dry deposition of nutrients on phytoplankton pigment composition and primary production in the coastal Bay of Bengal. *Environmental Science and Pollution Research* **29**(54): 82218–82231. DOI: <http://dx.doi.org/10.1007/s11356-022-21477-3>.
- Kumari, VR, Yadav, K, Sarma, VVSS, Dileep Kumar, M.** 2021. Acidification of the coastal Bay of Bengal by aerosols deposition. *Journal of Earth System Science* **130**(4): 1–13.
- Kurian, S, Chndrasekhararao, AV, Vidya, PJ, Shenoy, DM, Gauns, M, Uskaikar, H, Aparna, SG.** 2020. Role of oceanic fronts in enhancing phytoplankton biomass in the eastern Arabian Sea during an oligotrophic period. *Marine Environmental Research* **160**: 105023. DOI: <http://dx.doi.org/10.1016/j.marenvres.2020.105023>.
- Kurian, S, Shenoy, DM, Akhil, VP, Kessarkar, PM, Gauns, M, Shetye, SS, Kabeer, M, Vijayan, AP, Methar, A, Karapurkar, S, Chndrasekhararao, AV, Naqvi, SWA.** 2023. Spatial and seasonal variations in the particulate sinking flux in the Bay of Bengal. *Progress in Oceanography* **211**: 102983.
- Kuttippurath, J, Maishal, S, Anjaneyan, P, Sunanda, N, Chakraborty, K.** 2023. Recent changes in atmospheric input and primary productivity in the North Indian Ocean. *Heliyon* **9**(7): e17940.
- Kwiatkowski, L, Torres, O, Bopp, L, Aumont, O, Chamberlain, M, Christian, JR, Dunne, JP, Gehlen, M, Ilyina, T, John, JG, Lenton, A, Li, H, Lovenduski, NS, Orr, JC, Palmieri, J, Santana-Falcón, Y, Schwinger, J, Séférian, R, Stock, CA, Tagliabue, A, Takano, Y, Tjiputra, J, Toyama, K, Tsujino, H, Watanabe, M, Yamamoto, A, Yool, A, Ziehn, T, Dunne, JP, Gehlen, M, Ilyina, T, John, JG, Lenton, A, Li, H, Lovenduski, NS, Orr, JC, Palmieri, J, Santana-Falcón, Y, Schwinger, J, Séférian, R, Stock, CA, Tagliabue, A, Takano, Y, Tjiputra, J, Toyama, K, Tsujino, H, Watanabe, M, Yamamoto, A, Yool, A, Ziehn, T.** 2020. Twenty-first century ocean warming, acidification, deoxygenation, and upper-ocean nutrient and primary production decline from CMIP6 model projections. *Biogeosciences* **17**(13): 3439–3470. DOI: <http://dx.doi.org/10.5194/bg-17-3439-2020>.
- Lachkar, Z, Lévy, M, Hailegeorgis, D, Vallivattathillam, P.** 2023. Differences in recent and future trends in the Arabian Sea oxygen minimum zone: Processes and uncertainties. *Frontiers in Marine Science* **10**: 1122043.
- Lachkar, Z, Mehari, M, Al Azhar, M, Lévy, M, Smith, S.** 2021. Fast local warming is the main driver of recent deoxygenation in the Northern Arabian Sea. *Biogeosciences* **18**(20): 5831–5849.
- Lachkar, Z, Mehari, M, Lévy, M, Paparella, F, Burt, JA.** 2022. Recent expansion and intensification of hypoxia in the Arabian Gulf and its drivers. *Frontiers in Marine Science* **9**: 1616. DOI: <http://dx.doi.org/10.3389/fmars.2022.891378>.
- Lakshmi, RS, Prakash, S, Lotliker, AA, Baliarsingh, SK, Samanta, A, Mathew, T, Chatterjee, A, Sahu, BK, Nair, TMB.** 2021. Physicochemical controls on the initiation of phytoplankton bloom during the winter monsoon in the Arabian Sea. *Scientific Reports* **11**(1): 13448. DOI: <http://dx.doi.org/10.1038/s41598-021-92897-3>.
- Lana, A, Bell, TG, Simó, R, Vallina, SM, Ballabrera-Poy, J, Kettle, AJ, Dachs, J, Bopp, L, Saltzman, ES, Stefels, J, Johnson, JE, Liss, PS.** 2011. An updated climatology of surface dimethylsulfide concentrations and emission fluxes in the global ocean. *Global Biogeochemical Cycles* **25**(1). DOI: <http://dx.doi.org/10.1029/2010GB003850>.
- Law, CS, Brévière, E, de Leeuw, G, Garçon, V, Guieu, C, Kieber, DJ, Konradowitz, S, Paulmier, A, Quinn, PK, Saltzman, ES, Stefels, J, von Glasow, R.** 2013. Evolving research directions in Surface Ocean–Lower Atmosphere (SOLAS) science. *Environmental Chemistry* **10**(1): 1–16. DOI: <http://dx.doi.org/10.1071/EN12159>.
- Lelieveld, J, Crutzen, PJ, Ramanathan, V, Andreae, MO, Brenninkmeijer, CAM, Campos, T, Cass, GR, Dickerson, RR, Fischer, H, de Gouw, JA, Hansel, A, Jefferson, A, Kley, D, de Laat, ATJ, Lal, S, Lawrence, MG, Lobert, JM, Mayol-Bracero, O, Mitra, AP, Novakov, T, Oltmans, SJ, Prather, KA, Reiner, T, Rodhe, H, Scheeren, HA, Sikka, D, Williams, J.** 2001. The Indian Ocean experiment: Widespread air pollution from South and Southeast Asia. *Science* **291**(5506): 1031–1036.
- Lévy, M, Lengaigne, M, Bopp, L, Vincent, EM, Madec, G, Ethé, C, Kumar, D, Sarma, VVSS.** 2012. Contribution of tropical cyclones to the air-sea CO₂ flux: A global view. *Global Biogeochemical Cycles* **26**(2). DOI: <http://dx.doi.org/10.1029/2011GB004145>.
- Lotliker, AA, Baliarsingh, SK, Trainer, VL, Wells, ML, Wilson, C, Udaya Bhaskar, TVS, Samanta, A, Shahimol, SR.** 2018. Characterization of oceanic *Noctiluca* blooms not associated with hypoxia in the Northeastern Arabian Sea. *Harmful Algal Bloom* **74**: 46–57. DOI: <http://dx.doi.org/10.1016/j.hal.2018.03.008>.
- Mahajan, AS, Li, Q, Inamdar, S, Ram, K, Badia, A, Saiz-Lopez, A.** 2021. Modelling the impacts of iodine chemistry on the Northern Indian Ocean marine boundary layer. *Atmospheric Chemistry and Physics* **21**(11): 8437–8454.
- Mahajan, AS, Tinel, L, Hulswar, S, Cuevas, CA, Wang, S, Ghude, S, Lopez, AS.** 2019a. Observations of iodine oxide in the Indian Ocean marine boundary layer: A transect from the tropics to the high latitudes. *Atmospheric Environment: X* **1**: 100016.
- Mahajan, AS, Tinel, L, Sarkar, A, Chance, R, Carpenter, LJ, Hulswar, S, Vinayachandran, PN.** 2019b. Understanding iodine chemistry over the northern

- and equatorial Indian Ocean. *Journal of Geophysical Research: Atmospheres* **124**(14): 8104–8118.
- Mahowald, NM, Baker, AR, Bergametti, G, Brooks, N, Duce, RA, Jickells, TD, Tegen, I.** 2005. Atmospheric global dust cycle and iron inputs to the ocean. *Global Biogeochemical Cycles* **19**(4).
- Mallik, C, Lal, S, Venkataramani, S, Naja, M, Ojha, N.** 2013. Variability in ozone and its precursors over the Bay of Bengal during post monsoon: Transport and emission effects. *Journal of Geophysical Research: Atmospheres* **118**(17): 10–190.
- Masuda, S.** 2020. Determining subsurface oceanic changes in the Indian sector of the Southern Ocean using Argo float data. *Polar Science* **23**: 1873–9652.
- McCreary, JP, Yu, Z, Hood, RR, Vinayachandran, PN, Furue, R, Ishida, A, Richards, KJ.** 2013. Dynamics of the Indian-Ocean oxygen minimum zones. *Progress in Oceanography* **112–113**: 15–37. DOI: <http://dx.doi.org/10.1016/j.pocean.2013.03.002>.
- Mignot, A, von Schuckmann, K, Landschützer, P, Gasparin, F, van Gennip, S, Perruche, C, Lamouroux, J, Amm, T.** 2022. Decrease in air-sea CO₂ fluxes caused by persistent marine heatwaves. *Nature Communications* **13**(1): 4300. DOI: <http://dx.doi.org/10.1038/s41467-022-31983-0>.
- Moffett, JW, Landry, MR.** 2020. Grazing control and iron limitation of primary production in the Arabian Sea: Implications for anticipated shifts in southwest monsoon intensity. *Deep Sea Research Part II: Topical Studies in Oceanography* **179**: 104687.
- Morrison, JM, Codispoti, LA, Smith, SL, Wishner, K, Flagg, C, Gardner, WD, Gaurin, S, Naqvi, SWA, Manghnani, V, Prosperie, L, Gundersen, JS.** 1999. The oxygen minimum zone in the Arabian Sea during 1995. *Deep Sea Research Part II: Topical Studies in Oceanography* **46**: 1903–1931.
- Naqvi, SWA, Bange, HW, Farías, L, Monteiro, PMS, Scranton, MI, Zhang, J.** 2010. Marine hypoxia/anoxia as a source of CH₄ and N₂O. *Biogeosciences* **7**(7): 2159–2190.
- Naqvi, SWA, Jayakumar, DA, Narvekar, PV, Naik, H, Sarma, V, D'souza, W, Joseph, S, George, MD.** 2000. Increased marine production of N₂O due to intensifying anoxia on the Indian continental shelf. *Nature* **408**(6810): 346–349.
- Naqvi, SWA, Naik, H, Pratihary, A, D'Souza, W, Narvekar, PV, Jayakumar, DA, Devol, AH, Yoshinari, T, Saino, T.** 2006. Coastal versus open-ocean denitrification in the Arabian Sea. *Biogeosciences* **3**: 621–633.
- Naqvi, WA.** 1991. Geographical extent of denitrification in the Arabian Sea in relation to some physical processes. *Oceanologica Acta* **14**(3): 281–290.
- Nayak G, Kumar, A, Bikkina, S, Tiwari SS, Sheteye, SK, Sudheer, A.** 2022. Carbonaceous aerosols and their light absorption properties over the Bay of Bengal during continental outflow. *Environmental Science: Processes & Impacts* **24**(1): 72–88. DOI: <http://dx.doi.org/10.1039/D1EM00347J>.
- Needham, HF, Keim, BD, Sathiaraj, D.** 2015. A review of tropical cyclone-generated storm surges: Global data sources, observations, and impacts. *Reviews of Geophysics* **53**(2): 545–591.
- Oschlies, A, Koeve, W, Landolfi, A, Kähler, P.** 2019. Loss of fixed nitrogen causes net oxygen gain in a warmer future ocean. *Nature Communications* **10**(1): 2805.
- Panda, PP, Aswini, MA, Bhatt, P, Srimuruganandam, B, Peketi, A, Kumar, A.** 2022 Jul 15. Bioactive trace elements' composition and their fractional solubility in aerosols from the Arabian Sea during the southwest monsoon. *ACS Earth and Space Chemistry* **6**(8): 1969–1981. DOI: <http://dx.doi.org/10.1021/acsearthspacechem.2c00067>.
- Panda, PP, Shukla, G, Kumar, A, Aswini, MA, Kaushik, A, Nayak, G, Matta, VM.** 2024. Atmospheric deposition of mineral dust and associated nutrients over the Equatorial Indian Ocean. *Science of The Total Environment* **915**: 169779.
- Parvathi, V, Suresh, I, Lengaigne, M, Ethé, C, Vialard, J, Levy, M, Neetu, S, Aumont, O, Resplandy, L, Naik, H.** 2017. Positive Indian Ocean dipole events prevent anoxia along the west coast of India. *Biogeosciences* **14**(6): 1541–1559.
- Phillips, HE, Tandon, A, Furue, R, Hood, R, Ummenhofer, CC, Benthuyssen, JA, Wiggert, J.** 2021. Progress in understanding of Indian Ocean circulation, variability, air–sea exchange, and impacts on biogeochemistry. *Ocean Science* **17**(6): 1677–1751.
- Prakash, P, Prakash, S, Ravichandran, M, Udaya Bhaskar, TVS, Anil Kumar, N.** 2020. Seasonal evolution of chlorophyll in the Indian sector of the Southern Ocean: Analyses of bio-argo measurements. *Deep Sea Research Part II: Topical Studies in Oceanography* **178**: 104791.
- Prasanna Kumar, S.** 2004. Intrusion of the Bay of Bengal water into the Arabian Sea during winter monsoon and associated chemical and biological response. *Geophysical Research Letters* **31**(15): L15304. DOI: <http://dx.doi.org/10.1029/2004GL020247>.
- Pratihary, A, Lavik, G, Naqvi, SWA, Shirodkar, G, Sarkar, A, Marchant, H, Ohde, T, Shenoy, D, Kurian, S, Uskaikar, H, Kuypers, MM.** 2023. Sulphidic event modulates the nitrogen cycling over the Eastern Arabian Sea shelf: Evidence for chemolithoautotrophic denitrification causing massive N loss. *Progress in Oceanography* **217**: 103075.
- Rao, DN, Chopra, M, Rajula, GR, Durgadevi, DSL, Sarma, VVSS.** 2021. Release of significant fraction of primary production as dissolved organic carbon in the Bay of Bengal. *Deep Sea Research Part I: Oceanographic Research Papers* **168**(1): 103445.
- Rao, DN, Sarma, VVSS.** 2022. Accumulation of dissolved organic carbon and nitrogen in the photic zone in the nitrogen-depleted waters of the Bay of Bengal. *Marine Chemistry* **239**: 104074.
- Rixen, T, Baum, A, Gaye, B, Nagel, B.** 2014. Seasonal and interannual variations in the nitrogen cycle in the Arabian Sea. *Biogeosciences* **11**(20): 5733–5747. DOI: <http://dx.doi.org/10.5194/bg-11-5733-2014>.

- Rixen, T, Cowie, G, Gaye, B, Goes, J, do Rosário Gomes, H, Hood, RR, Lachkar, Z, Schmidt, H, Segsneider, J, Singh, A.** 2020. Reviews and syntheses: Present, past, and future of the oxygen minimum zone in the northern Indian Ocean. *Biogeosciences* **17**(23): 6051–6080. DOI: <http://dx.doi.org/10.5194/bg-17-6051-2020>.
- Roxy, M, Tanimoto, Y, Preethi, B, Terray, P, Krishnan, R.** 2013. Intraseasonal SST-precipitation relationship and its spatial variability over the tropical summer monsoon region. *Climate Dynamics* **41**(1): 45–61. DOI: <http://dx.doi.org/10.1007/s00382-012-1547-1>.
- Roxy, MK, Ghosh, S, Pathak, A, Athulya, R, Mujumdar, M, Murtugudde, R, Rajeevan, M.** 2017. A threefold rise in widespread extreme rain events over Central India. *Nature Communications* **8**(1): 708.
- Roxy, MK, Modi, A, Murtugudde, R, Valsala, V, Panickal, S, Prasanna Kumar, S, Ravichandran, M, Vichi, M, Levy, M.** 2016. A reduction in marine primary productivity driven by rapid warming over the tropical Indian Ocean. *Geophysical Research Letters* **43**(2): 826–833. DOI: <http://dx.doi.org/10.1002/2015GL066979>.
- Roxy, MK, Ritika, K, Terray, P, Masson, S.** 2014. The curious case of Indian Ocean warming. *Journal of Climate* **27**(22): 8501–8509. DOI: <http://dx.doi.org/10.1175/JCLI-D-14-00471.1>.
- Roxy, MK, Ritika, K, Terray, P, Murtugudde, R, Ashok, K, Goswami, BN.** 2015. Drying of Indian subcontinent by rapid Indian Ocean warming and a weakening land-sea thermal gradient. *Nature Communications* **6**(1): 7423. DOI: <http://dx.doi.org/10.1038/ncomms8423>.
- Sahu, LK, Tripathi, N, Sheel, V, Ojha, N.** 2019. The influence of local meteorology and convection on carbon monoxide distribution over Chennai. *Journal of Earth System Science* **128**(5): 119. DOI: <http://dx.doi.org/10.1007/s12040-019-1156-z>.
- Sarkar, A, Naqvi, SWA, Lavik, G, Pratihary, A, Naik, H, Shirodkar, G, Kuypers, MM.** 2020. Massive nitrogen loss over the western Indian continental shelf during seasonal anoxia: Evidence from isotope pairing technique. *Frontiers in Marine Science* **7**: 678.
- Sarma, V, Krishna, MS, Rao, VD, Viswanadham, R, Kumar, NA, Kumari, TR, Gawade, L, Ghatkar, S, Tari, A.** 2012. Sources and sinks of CO₂ in the west coast of Bay of Bengal. *Tellus B: Chemical and Physical Meteorology* **64**(1): 10961.
- Sarma, V, Krishna, MS, Srinivas, TNR, Kumari, VR, Yadav, K, Kumar, MD.** 2021. Elevated acidification rates due to deposition of atmospheric pollutants in the coastal Bay of Bengal. *Geophysical Research Letters* **48**(16): e2021GL095159.
- Sarma, V, Patil, JS, Shankar, D, Anil, AC.** 2019. Shallow convective mixing promotes massive *Noctiluca scintillans* bloom in the Northeastern Arabian Sea. *Marine Pollution Bulletin* **138**: 428–436.
- Sarma, V, Paul, YS, Vani, DG, Murty, VSN.** 2015. Impact of river discharge on the coastal water pH and pCO₂ levels during the Indian Ocean Dipole (IOD) years in the Western Bay of Bengal. *Continental Shelf Research* **107**: 132–140.
- Sarma, V, Sridevi, B, Metzl, N, Patra, PK, Lachkar, Z, Chakraborty, K, Chandra, N.** 2023. Air-sea fluxes of CO₂ in the Indian Ocean between 1985 and 2018: A synthesis based on observation-based surface CO₂, hindcast and atmospheric inversion models. *Global Biogeochemical Cycles* **37**(5): e2023GB007694.
- Sarma, V, Vivek, R, Rao, DN, Ghosh, VRD.** 2020a. Severe phosphate limitation on nitrogen fixation in the Bay of Bengal. *Continental Shelf Research* **205**: 104199.
- Sarma, VVSS, Bhaskar, TVSU, Kumar, JP, Chakraborty, K.** 2020b. Potential mechanisms responsible for occurrence of core oxygen minimum zone in the north-eastern Arabian Sea. *Deep Sea Research Part I: Oceanographic Research Papers* **165**: 103393. DOI: <http://dx.doi.org/10.1016/j.dsr.2020.103393>.
- Sarma, VVSS, Jagadeesan, L, Dalabehera, HB, Rao, DN, Kumar, GS, Durgadevi, DS, Priya, MMR.** 2018. Role of eddies on intensity of oxygen minimum zone in the Bay of Bengal. *Continental Shelf Research* **168**: 48–53.
- Sarma, VVSS, Krishna, MS, Viswanadham, R, Rao, GD, Rao, VD, Sridevi, B, Kumar, BSK, Prasad, VR, Subbaiah, ChV, Acharyya, T, Bandopadhyay, D.** 2013a. Intensified oxygen minimum zone on the western shelf of Bay of Bengal during summer monsoon: Influence of river discharge. *Journal of Oceanography* **69**(1): 45–55.
- Sarma, VVSS, Rao, GD, Viswanadham, R, Sherin, CK, Salisbury, J, Omand, MM, Stafford, KM.** 2016. Effects of freshwater stratification on nutrients, dissolved oxygen, and phytoplankton in the Bay of Bengal. *Oceanography* **29**(2): 222–231.
- Sarma, VVSS, Sridevi, B, Kumar, A, Bikkina, S, Kumari, VR, Bikkina, P, Rao, VD.** 2022. Impact of atmospheric anthropogenic nitrogen on new production in the Northern Indian Ocean: Constrained based on satellite aerosol optical depth and particulate nitrogen levels. *Environmental Science: Processes & Impacts* **24**(10): 1895–1911.
- Sarma, VVSS, Sridevi, B, Maneesha, K, Sridevi, T, Naidu, SA, Prasad, VR, Venkataramana, V, Acharya, T, Bharati, MD, Subbaiah, ChV, Kiran, BS, Reddy, NPC, Sarma, VV, Sadhuram, Y, Murty, TVR.** 2013b. Impact of atmospheric and physical forcings on biogeochemical cycling of dissolved oxygen and nutrients in the coastal Bay of Bengal. *Journal of Oceanography* **69**(2): 229–243.
- Sarma, VVSS, Udaya Bhaskar, TVS.** 2018. Ventilation of oxygen to oxygen minimum zone due to anticyclonic eddies in the Bay of Bengal. *Journal of Geophysical Research: Biogeosciences* **123**(7): 2145–2153.
- Sawant, S, Madhupratap, M.** 1996. Seasonality and composition of phytoplankton in the Arabian Sea. *Current Science* **71**: 869–873.
- Seager, R, Naik, N, Vecchi, GA.** 2010. Thermodynamic and dynamic mechanisms for large-scale changes in the hydrological cycle in response to global warming. *Journal of Climate* **23**(17): 4651–4668.

- Sen Gupta, R, Naqvi, SWA.** 1984. Chemical oceanography of the Indian Ocean, north of the equator. *Deep Sea Research Part A. Oceanographic Research Papers* **31**: (6–8): 671–706.
- Sengupta, D, Bharath Raj, GN, Ravichandran, M, Sree Lekha, J, Papa, F.** 2016. Near-surface salinity and stratification in the North Bay of Bengal from moored observations. *Geophysical Research Letters* **43**(9): 4448–4456. DOI: <http://dx.doi.org/10.1002/2016GL068339>.
- Shaik, AUR, Biswas, H, Surendra Babu, N, Reddy, NPC, Ansari, ZA.** 2017. Investigating the impacts of treated effluent discharge on coastal water health (Visakhapatnam, SW coast of Bay of Bengal, India). *Environmental Monitoring and Assessment* **189**(12): 1–15.
- Shaji, C, Kar, SK, Vishal, T.** 2014. Storm surge studies in the North Indian Ocean: A review. *Indian Journal of Geo-Marine Sciences* **43**(2): 125–147.
- Shankar, D, Vinayachandran, PN, Unnikrishnan, AS.** 2002. The monsoon currents in the North Indian Ocean. *Progress in Oceanography* **52**(1): 63–120.
- Sharma, D, Biswas, H, Bandyopadhyay, D.** 2022a. Simulated ocean acidification altered community composition and growth of a coastal phytoplankton assemblage (South West Coast of India, Eastern Arabian Sea). *Environmental Science and Pollution Research* **29**(13): 19244–19261.
- Sharma, D, Biswas, H, Chowdhury, M, Silori, S, Pandey, M, Ray, D.** 2023. Phytoplankton community shift in response to experimental Cu addition at the elevated CO₂ levels (Arabian Sea, winter monsoon). *Environmental Science and Pollution Research* **30**(3): 7325–7344.
- Sharma, D, Biswas, H, Panda, PP, Chowdhury, M, Silori, S, Pandey, M, Kumar, A.** 2022b. Atmospheric dust addition under elevated CO₂ restructured phytoplankton community from the Arabian Sea: A microcosm approach. *Marine Chemistry* **247**(4): 104183.
- Sharma, D, Biswas, H, Silori, S, Bandyopadhyay, D, Shaik, AUR.** 2022c. Phytoplankton growth and community shift over a short-term high-CO₂ simulation experiment from the southwestern shelf of India, Eastern Arabian Sea (summer monsoon). *Environmental Monitoring and Assessment* **194**(8): 581.
- Sharmila, S, Pillai, PA, Joseph, S, Roxy, M, Krishna, RPM, Chattopadhyay, R, Abhilash, S, Sahai, AK, Goswami, BN.** 2013. Role of ocean–atmosphere interaction on northward propagation of Indian summer Monsoon Intra-Seasonal Oscillations (MISO). *Climate Dynamics* **41**(5): 1651–1669. DOI: <http://dx.doi.org/10.1007/s00382-013-1854-1>.
- Shenoy, DM, Suresh, I, Uskaikar, H, Kurian, S, Vidya, PJ, Shirodkar, G, Gauns, MU, Naqvi, SWA.** 2020. Variability of dissolved oxygen in the Arabian Sea Oxygen Minimum Zone and its driving mechanisms. *Journal of Marine Systems* **204**: 103310. DOI: <http://dx.doi.org/10.1016/j.jmarsys.2020.103310>.
- Shetye, SS, Kurian, S, Naik, H, Gauns, M, Chndrasekhararao, AV, Kumar, A, Naik, B.** 2019. Variability of organic nitrogen and its role in regulating phytoplankton in the Eastern Arabian Sea. *Marine Pollution Bulletin* **141**(11): 550–560.
- Shetye, SS, Kurian, S, Vidya, PJ, Gauns, M, Shenoy, DM, Aparna, SG, Nandakumar, K, Karapurkar, SG.** 2021. Total organic carbon and its role in oxygen utilization in the eastern Arabian Sea. *Marine Pollution Bulletin* **163**: 111939. DOI: <http://dx.doi.org/10.1016/j.marpolbul.2020.111939>.
- Shetye, SS, Nandakumar, K, Kurian, S, Gauns, M, Shenoy, DM, Naik, H, Karapurkar, SG.** 2022. Organic carbon dynamics in the continental shelf waters of the Eastern Arabian Sea. *Environmental Monitoring and Assessment* **194**(10): 716.
- Shirodkar, G, Naqvi, SWA, Naik, H, Pratihary, AK, Kurian, S, Shenoy, DM.** 2018. Methane dynamics in the shelf waters of the west coast of India during seasonal anoxia. *Marine Chemistry* **203**: 55–63.
- Shukla, G, Sudheer, AK, Gunthe, SS, Beig, G, Kumar, A.** 2023. Seasonal variability in fine particulate matter water content and estimated pH over a coastal region in the Northeast Arabian Sea. *Atmosphere* **14**(2): 259.
- Silori, S, Sharma, D, Chowdhury, M, Biswas, H, Bandyopadhyay, D, Shaik, AUR, Cardinal, D, Mandeng-Yogo, M, Narvekar, J.** 2021. Contrasting phytoplankton and biogeochemical functioning in the eastern Arabian Sea shelf waters recorded by carbon isotopes (SW monsoon). *Marine Chemistry* **232**: 103962.
- Singh, A, Ramesh, R.** 2015. Environmental controls on new and primary production in the Northern Indian Ocean. *Progress in Oceanography* **131**: 138–145.
- Singh, VK, Roxy, MK.** 2022. A review of ocean–atmosphere interactions during tropical cyclones in the North Indian Ocean. *Earth-Science Reviews* **226**: 103967.
- Soni, M, Verma, S, Mishra, MK, Mall, RK, Payra, S.** 2022. Estimation of particulate matter pollution using WRF-Chem during dust storm event over India. *Urban Climate* **44**: 101202.
- Sridevi, B, Sabira, S, Sarma, VVSS.** 2023. Impact of ocean warming on net primary production in the Northern Indian Ocean: Role of aerosols and freshening of surface ocean. *Environmental Science and Pollution Research* **30**(18): 53616–53634.
- Sridevi, B, Sarma, V.** 2021. Role of river discharge and warming on ocean acidification and pCO₂ levels in the Bay of Bengal. *Tellus B: Chemical and Physical Meteorology* **73**(1): 1–20.
- Sridevi, B, Sarma, VVSS.** 2020. A revisit to the regulation of oxygen minimum zone in the Bay of Bengal. *Journal of Earth System Science* **129**: 107.
- Sridevi, B, Sarma, VVSS.** 2022. Enhanced atmospheric pollutants strengthened winter convective mixing and phytoplankton blooms in the Northern Arabian Sea. *Journal of Geophysical Research: Biogeosciences* **127**(10): e2021JG006527. DOI: <http://dx.doi.org/10.1029/2021JG006527>.

- Srinivas, B, Sarin, MM.** 2013. Atmospheric dry-deposition of mineral dust and anthropogenic trace metals to the Bay of Bengal. *Journal of Marine Systems* **126**: 56–68. DOI: <http://dx.doi.org/10.1016/j.jmarsys.2012.11.004>.
- Srivastava, S, Lal, S, Venkataramani, S, Guha, I, Subrahmanyam, DB.** 2012. Airborne measurements of O₃, CO, CH₄ and NMHCs over the Bay of Bengal during winter. *Atmospheric Environment* **59**: 597–609.
- Subrahmanyam, R.** 1954 Mar. A new member of the Euglenineae, Protoeuglena Noctilucae gen. et sp. nov., occurring in Noctiluca miliaris suriray, causing green discoloration of the sea off Calicut. *Proceedings of the Indian Academy of Sciences* **39**(3): 118–127.
- Sudheesh, V, Gupta, GVM, Sudharma, KV, Naik, H, Shenoy, DM, Sudhakar, M, Naqvi, SWA.** 2016. Upwelling intensity modulates N₂O concentrations over the western Indian shelf. *Journal of Geophysical Research: Oceans* **121**(12): 8551–8565.
- Suresh, K, Singh, U, Kumar, A, Karri, D, Peketi, A, Ramaswamy, V.** 2021. Provenance tracing of long-range transported dust over the Northeastern Arabian Sea during the southwest monsoon. *Atmospheric Research* **250**: 105377. DOI: <http://dx.doi.org/10.1016/j.atmosres.2020.105377>.
- Swapna, P, Sreeraj, P, Sandeep, N, Jyoti, J, Krishnan, R, Prajeesh, AG, Ayantika, DC, Manmeet, S.** 2022. Increasing frequency of extremely severe cyclonic storms in the North Indian Ocean by anthropogenic warming and southwest monsoon weakening. *Geophysical Research Letters* **49**(3): e2021GL094650. DOI: <http://dx.doi.org/10.1029/2021GL094650>.
- Takashima, H, Irie, H, Kanaya, Y, Syamsudin, F.** 2012. NO₂ observations over the western Pacific and Indian Ocean by MAX-DOAS on *Kaiyo*, a Japanese research vessel. *Atmospheric Measurement Techniques* **5**(10): 2351–2360. DOI: <http://dx.doi.org/10.5194/amt-5-2351-2012>.
- Techera, EJ.** 2018. Supporting blue economy agenda: Fisheries, food security and climate change in the Indian Ocean. *Journal of the Indian Ocean Region* **14**(1): 7–27.
- Tegtmeier, S, Marandino, C, Jia, Y, Quack, B, Mahajan, AS.** 2022. Atmospheric gas-phase composition over the Indian Ocean. *Atmospheric Chemistry and Physics* **22**(10): 6625–6676. DOI: <http://dx.doi.org/10.5194/acp-22-6625-2022>.
- Tindale, NW, Pease, PP.** 1999. Aerosols over the Arabian Sea: Atmospheric transport pathways and concentrations of dust and sea salt. *Deep Sea Research Part II: Topical Studies in Oceanography* **46**(8–9): 1577–1595. DOI: [http://dx.doi.org/10.1016/S0967-0645\(99\)00036-3](http://dx.doi.org/10.1016/S0967-0645(99)00036-3).
- Toyoda, S, Terajima, K, Yoshida, N, Yoshikawa, C, Makabe, A, Hashihama, F, Ogawa, H.** 2023. Extensive accumulation of nitrous oxide in the oxygen minimum zone in the Bay of Bengal. *Global Biogeochemical Cycles* **37**(9): e2022GB007689. DOI: <http://dx.doi.org/10.1029/2022GB007689>.
- Tripathi, N, Sahu, LK, Singh, A, Yadav, R, Patel, A, Patel, K, Meenu, P.** 2020. Elevated levels of biogenic nonmethane hydrocarbons in the marine boundary layer of the Arabian Sea during the intermonsoon. *Journal of Geophysical Research: Atmospheres* **125**(22): e2020JD032869.
- Udaya Bhaskar, TV, Sarma, VVSS, Pavan Kumar, J.** 2021. Potential mechanisms responsible for spatial variability in intensity and thickness of oxygen minimum zone in the Bay of Bengal. *Journal of Geophysical Research: Biogeosciences* **126**(6): e2021JG006341. DOI: <http://dx.doi.org/10.1029/2021JG006341>.
- Ulloa, O, Canfield, DE, DeLong, EF, Letelier, RM, Stewart, FJ.** 2012. Microbial oceanography of anoxic oxygen minimum zones. *Proceedings of the National Academy of Sciences* **109**(40): 15996–16003.
- Ummenhofer, CC, Gupta, AS, Li, Y, Taschetto, AS, England, MH.** 2011. Multi-decadal modulation of the El Niño–Indian monsoon relationship by Indian Ocean variability. *Environmental Research Letters* **6**(3): 034006. DOI: <http://dx.doi.org/10.1088/1748-9326/6/3/034006>.
- Vallivattathillam, P, Lachkar, Z, Lévy, M.** 2023. Shrinking of the Arabian Sea oxygen minimum zone with climate change projected with a downscaled model. *Frontiers in Marine Science* **10**: 1123739.
- Valsala, V, Maksyutov, S.** 2013. Interannual variability of the air–sea CO₂ flux in the North Indian Ocean. *Ocean Dynamics* **63**(2): 165–178.
- Valsala, V, Maksyutov, S, Murtugudde, R.** 2012. A window for carbon uptake in the southern subtropical Indian Ocean. *Geophysical Research Letters* **39**(17).
- Valsala, V, Murtugudde, R.** 2015. Mesoscale and intra-seasonal air–sea CO₂ exchanges in the Western Arabian Sea during boreal summer. *Deep Sea Research Part I: Oceanographic Research Papers* **103**: 101–113.
- Valsala, V, Sreeush, MG, Anju, M, Sreenivas, P, Tiwari, YK, Chakraborty, K, Sijikumar, S.** 2021. An observing system simulation experiment for Indian Ocean surface pCO₂ measurements. *Progress in Oceanography* **194**: 102570. DOI: <http://dx.doi.org/10.1016/j.pocean.2021.102570>.
- Valsala, V, Sreeush, MG, Chakraborty, K.** 2020. The IOD impacts on the Indian Ocean carbon cycle. *Journal of Geophysical Research: Oceans* **125**(11): e2020JC016485.
- Valsala, V, Tiwari, YK, Pillai, P, Roxy, M, Maksyutov, S, Murtugudde, R.** 2013. Intraseasonal variability of terrestrial biospheric CO₂ fluxes over India during summer monsoons. *Journal of Geophysical Research: Biogeosciences* **118**(2): 752–769.
- Vidya, PJ, Kurian, S.** 2018. Impact of 2015–2016 ENSO on the winter bloom and associated phytoplankton community shift in the northeastern Arabian Sea. *Journal of Marine Systems* **186**: 96–104. DOI: <http://dx.doi.org/10.1016/j.jmarsys.2018.06.005>.
- Vinayachandran, PNM, Masumoto, Y, Roberts, MJ, Huggett, JA, Halo, I, Chatterjee, A, Amol, P, Gupta, GVM, Singh, A, Mukherjee, A, Prakash,**

- S, Beckley, LE, Raes, EJ, Hood, R.** 2021. Reviews and syntheses: Physical and biogeochemical processes associated with upwelling in the Indian Ocean. *Biogeosciences* **18**(22): 5967–6029. DOI: <http://dx.doi.org/10.5194/bg-18-5967-2021>.
- Wallmann, K, José, YS, Hopwood, MJ, Somes, CJ, Dale, AW, Scholz, F, Oschlies, A.** 2022. Biogeochemical feedbacks may amplify ongoing and future ocean deoxygenation: A case study from the Peruvian oxygen minimum zone. *Biogeochemistry* **159**(1): 45–67.
- Wang, B, Biasutti, M, Byrne, MP, Castro, C, Chang, C-P, Cook, K, Fu, R, Grimm, AM, Ha, K-J, Hendon, H, Kitoh, A, Krishnan, R, Lee, J-Y, Li, J, Liu, J, Moise, A, Pascale, S, Roxy, MK, Seth, A, Sui, C-H, Turner, A, Yang, S, Yun, K-S, Zhang, L, Zhou, T.** 2021. Monsoons climate change assessment. *Bulletin of the American Meteorological Society* **102**(1): E1–E19. DOI: <http://dx.doi.org/10.1175/BAMS-D-19-0335.1>.
- Ward, BB, Devol, AH, Rich, JJ, Chang, BX, Bulow, SE, Naik, H, Pratihary, A, Jayakumar, A.** 2009. Denitrification as the dominant nitrogen loss process in the Arabian Sea. *Nature* **461**(7260): 78–81. DOI: <http://dx.doi.org/10.1038/nature08276>.
- Ward, BB, Tuit, CB, Jayakumar, A, Rich, JJ, Moffett, J, Naqvi, SWA.** 2008. Organic carbon, and not copper, controls denitrification in oxygen minimum zones of the ocean. *Deep Sea Research Part I: Oceanographic Research Papers* **55**(12): 1672–1683.
- West, BJ, Han, W, Zhang, L, Li, Y.** 2020. The role of oceanic processes in the initiation of boreal winter intraseasonal oscillations over the Indian Ocean. *Journal of Geophysical Research: Oceans* **125**(2): e2019JC015426. DOI: <http://dx.doi.org/10.1029/2019JC015426>.
- Worden, HM, Deeter, MN, Frankenberg, C, George, M, Nichitiu, F, Worden, J, Aben, I, Bowman, KW, Clerbaux, C, Coheur, PF, de Laat, ATJ, Detweiler, R, Drummond, JR, Edwards, DP, Gille, JC, Hurtmans, D, Luo, M, Martínez-Alonso, S, Massie, S, Pfister, G, Warner, JX.** 2013. Decadal record of satellite carbon monoxide observations. *Atmospheric Chemistry and Physics* **13**(2): 837–850. DOI: <http://dx.doi.org/10.5194/acp-13-837-2013>.
- Wyrtki, K.** 1971. *Oceanographic atlas of the International Indian Ocean Expedition*. Washington, DC: National Science Foundation Publication.
- Yadav, K, Rao, VD, Sridevi, B, Sarma, VVSS.** 2021. Decadal variations in natural and anthropogenic aerosol optical depth over the Bay of Bengal: The influence of pollutants from Indo-Gangetic Plain. *Environmental Science and Pollution Research* **28**(39): 55202–55219. DOI: <http://dx.doi.org/10.1007/s11356-021-14703-x>.
- Yang, K, Cai, W, Huang, G, Wang, G, Ng, B, Li, S.** 2020. Oceanic processes in ocean temperature products key to a realistic presentation of positive Indian Ocean dipole nonlinearity. *Geophysical Research Letters* **47**(16): e2020GL089396. DOI: <http://dx.doi.org/10.1029/2020GL089396>.
- Zhou, ZQ, Xie, SP, Zhang, R.** 2021. Historic Yangtze flooding of 2020 tied to extreme Indian Ocean conditions. *Proceedings of the National Academy of Sciences* **118**(12): e2022255118.

How to cite this article: Kumar, A, Tegtmeier, S, Fernandes, SO, Biswas, H, Girach, I, Roxy, MK, Kurian, S, Marandino, CA, Sarma, VVSS, Shenoy, DM. 2024. Surface ocean-lower atmospheric processes in the Indian Ocean: Current understanding, knowledge gaps, and future directions. *Elementa: Science of the Anthropocene* 12(1). DOI: <https://doi.org/10.1525/elementa.2023.00041>

Domain Editor-in-Chief: Detlev Helmig, Boulder AIR LLC, Boulder, CO, USA

Associate Editor: Byron Blomquist, University of Colorado Boulder, Boulder, CO, USA

Knowledge Domain: Atmospheric Science

Part of an Elementa Special Feature: Boundary Shift: The Air-Sea Interface in a Changing Climate

Published: August 21, 2024 **Accepted:** May 9, 2024 **Submitted:** March 13, 2023

Copyright: © 2024 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.



Elem Sci Anth is a peer-reviewed open access journal published by University of California Press.

OPEN ACCESS The Open Access logo, consisting of the words "OPEN ACCESS" followed by a circular icon containing a stylized lock symbol.