

The impact of ocean acidification on reproduction and early development of marine organisms

Pauline M Ross¹, Laura Parker¹ and Wayne A O'Connor²

¹School of Natural Sciences, College of Health and Science, University of Western Sydney Locked Bag 1797, Penrith South DC, 1797 Australia Email: pm.ross@uws.edu.au

² Port Stephens Fisheries Centre, Taylors Beach, New South Wales, 2316 Australia

ABSTRACT

Elevations in atmospheric carbon dioxide (CO₂) are anticipated to acidify oceans because of fundamental changes in ocean chemistry created by CO₂ absorption from the atmosphere into the oceans in a process known as ocean acidification. Over the next century, elevated CO₂ is expected to cause a reduction in the pH of the surface ocean from 8.1 to 7.7 units and a reduction in carbonate ion (CO₃²⁻) concentration required for calcifying marine organisms. Of growing concern is the potential impact that this change in ocean chemistry will have on marine and estuarine organisms and ecosystems, particularly molluscs and echinoderms which are broadcast spawners with larvae that develop in seawater. Although fertilisation in molluscs and echinoderms appears to be robust to the effects of elevated CO₂, larval development is characterised by impacts on the rate of larval development through successive stages, larval survival and abnormality including the failure to produce shells and skeletons. Despite these trends, our current understanding of the biological consequences of an acidifying ocean over the next century is still dominated by large uncertainties. Some of the greatest gaps in our understanding is the synergistic impacts of elevated CO₂ with other environmental stressors such as increasing ocean temperature and changing salinity. Until we have a better picture from laboratory and field based experiments which investigate multiple stressors in a chronic way over multiple generations, we will be limited in predicting which mollusc or echinoderm species will be able to acclimate or adapt.

Key words: Ocean acidification, ocean warming, molluscs, echinoderms, climate change

Introduction

It is essential to predict the impact of ocean acidification on marine organisms and habitats in order to anticipate the severity and consequences of future climate change and create adaptive management plans. Doing this in reality, however, is difficult (Pörtner *et al.* 2004; Przeslawski *et al.* 2008) because studies investigating the response of marine organisms to ocean acidification are comparatively in their infancy (Kleypas *et al.* 1999; Turley *et al.* 2006; Gooding *et al.* 2009) and ultimately meta-analyses attempting to draw out main effects, albeit authoritative, are restricted to small sample sizes and limited by a paucity of data (Hendriks *et al.* 2010, Dupont *et al.* 2010a, Hendriks and Duarte 2010; Hofman *et al.* 2010; Kroeker *et al.* 2010; Byrne 2011). The potential rate of acidification to marine organisms and habitats is, however, great exceeding that experienced by marine organisms at any other time on the planet (Caldeira and Wickert 2003; Turley *et al.* 2006; Guinotte and Fabry 2008). To counteract this great risk that climate change and ocean acidification specifically poses to the biodiversity of marine habitats, more recently research has turned to the impact of ocean acidification on fertilisation and early larval developmental stages of marine organisms because these life history stages may be the most sensitive and vulnerable to elevated CO₂ (Kurihara 2008; Dupont and Thorndyke 2009) and any negative impact may

form a dead end in one life history stage and thus create a “bottleneck” for populations (Dupont *et al.* 2010a). In such a scenario all early life history stages of marine organisms are likely to be affected by ocean acidification, including egg and sperm development and production, fertilisation, cleavage, larval development, larval dispersal, settlement and post-settlement survival (Figure 1). Embryos and larvae in these early developmental stages have specific environmental needs (Thorson 1950) and are more likely to have acute responses to environmental stress as the first deposition of CaCO₃ of many marine groups is known to occur during the larval stage (Kurihara




Figure 1. Different life-cycle stages of benthic calcifiers, including reproduction, fertilization, planktonic larva, settlement, metamorphosis, juvenile and benthic adult stages, that are potentially affected in different manners by ocean acidification. With permission from Kurihara 2008

2008). Effects from stress at fertilisation may also have profound carry-over consequences for individual larvae, larval cohorts (Green *et al.* 2004; Kurihara 2008; Parker *et al.* 2009, 2010) and larval dispersal. Collectively, the effects of elevated CO₂ on larvae may cascade, directly influencing the distribution and abundance of settlers and post settlement survival, and thus shape the structure and function of marine ecosystems and adult marine populations (Gosselin and Qian 1997; Kurihara 2008; ESF 2009). Here, we review the current literature on the effects of ocean acidification on fertilisation and larval development of molluscs and echinoderms on which the majority of work has been done and highlight the current gaps in our understanding and the areas for future research.

Historically, studies investigating the responses of early life history stages of molluscs and echinoderms to acidic conditions in a range of molluscs and echinoderms including the oysters *Crassostrea virginica* and *Saccostrea glomerata*, clams *Mercenaria mercenaria* and *Tivela stultorum* and the giant scallop *Placopecten magellanicus*; sea urchins *Paracentrotus lividus* and; *Sphaerechinus granularis*, manipulated instead of manipulating pH by mineralisation at levels often far outside those predicted for marine environments as a response to elevated atmospheric CO₂ (Table 1; Calabrese and Davis 1966; Pagano *et al.* 1985a,b; Cipollaro *et al.* 1986; Alvarado-Alvarez *et al.* 1996; Desrosiers *et al.* 1996; Wilson and Hyne 1997). Desrosiers *et al.* (1996) found no reduction in fertilisation but did find that the speed of cleavage was slowed due to polyspermy. There were also negative impacts on fertilisation in bivalve molluscs (Alvarado-Alvarez *et al.* 1996), cleavage in echinoids (*Sphaerechinus granularis* Cipollaro *et al.* 1986; *Paracentrotus lividus* Pagano *et al.*, 1985 a and b), morphological development in bivalve molluscs (*Saccostrea glomerata* Wilson and Hyne 1997) and increased mortality, decreased growth and skeletal malformations in bivalve molluscs (*Crassostrea virginica* and *Mercenaria mercenaria* Calabrese and Davis 1966). Caution is however, required when using the results of such studies to predict potential impacts of ocean acidification on marine organisms and ecosystems given the levels of elevated CO₂ predicted by IPCC scenarios for the end of the century are considerably lower and current best practise methodologies recommend acidifying culture media seawater to most accurately reflect future carbonate chemistry by: i. directly aerating with CO₂ at required level; ii. directly adding HCO₃ and HCl in equal molar values or iii. Saturating seawater with CO₂ prior to adding to the culture media (Gattuso *et al.* 2010). Most recently, studies manipulating pH using best practise methodologies of Gattuso *et al.* (2010) and CO₂ scenarios for the end of the century (IPCC 2001; Caldeira and Wickett 2003, 2005; IPCC 2007) have found significant, non-significant and no effect of elevated CO₂ on fertilisation, larval development and growth of early developmental stages of marine organisms even when using the same or closely related species (Table 1). Nevertheless, emerging trends suggest that the impact of ocean acidification on marine organisms, particularly molluscs and echinoderms with planktotrophic larvae and feeding larval stages is more significant for larval development than for fertilisation.

This is because it is during the early larval stage that shell formation occurs and amorphous calcium carbonate (ACC) crystal skeleton is deposited, known to be 30 times more soluble than stable forms of aragonite and calcite secreted later in larval and adult life (Orr *et al.* 2005; Dupont *et al.* 2010b).

Fertilisation, Cleavage and Gastrulation

The impact of ocean acidification on fertilisation has been investigated in a range of marine invertebrates, mainly echinoderms and bivalves which are broadcast spawners releasing eggs and gametes into the water column. There are fewer studies on crustaceans, corals and fish (Table 1). The life history stages of broadcast spawners potentially differ in sensitivity with early life history stages predicted to be more sensitive than later life history stages (Byrne *et al.* 2009) forming a major hypotheses in the literature being used to explain the variable responses (Kroeker *et al.* 2010) with less reference to the influence of natural variation in eggs and larvae (Przeslawski and Webb 2009) which may lead to a variability in larval response. Other hypotheses proposed to explain the variable responses of fertilisation in echinoderms and molluscs to the impact of ocean acidification include the level of elevated CO₂ used, the synergistic impact of temperature, the environment from which organisms are sourced, and debates around polyandry versus single crosses and other experimental techniques and analyses.

Echinoderms

Generally it is thought that echinoderm fertilisation is relatively robust to elevated CO₂ and decreased pH at levels predicted for the year 2100 (Byrne *et al.* 2009, 2010a, b; Dupont *et al.* 2010a), but can be negatively impacted when pH levels predicted for 2300 and lower are used (Kurihara and Shirayama 2004 and Kurihara *et al.* 2004b) or if sperm concentration is low and limiting (Byrne *et al.* 2010b, Ericson *et al.* 2010; Reuter *et al.* 2011). However, contrasting results have been reported for the same species from the same geographic region (Havenhand *et al.* 2008; Byrne *et al.* 2009).

Earliest studies by Kurihara and Shirayama (2004) and Kurihara *et al.* (2004b) exposed sea urchins, *Hemicentrotus pulcherrimus* and *Echinometra mathaei*, to pCO₂ concentrations ranging from 360 – 10365 ppm (pH 8.1-6.8). At exposures of pH 7.6 and 7.7 (-0.4-0.3 less than control levels) there was no effect on fertilisation of *Hemicentrotus pulcherrimus* and *Echinometra mathaei*, but at pH 6.8 for *Hemicentrotus pulcherrimus* and 6.8-7.1 *Echinometra mathaei* (-0.5 to -0.2 less than that predicted for 2300), there were significant effects with reduced fertilisation success and reduced cleavage speed in both species of sea urchins (Kurihara and Shirayama 2004; Kurihara *et al.* 2004b) and a trend for decreasing fertilisation with decreasing pH between 7.3-7.4 (Table 1). These studies were criticised for using CO₂ concentrations outside that predicted even for 2300 so later studies focussed on fertilisation responses to realistic scenarios




Figure 2. Summary of CO₂ effects at different life cycle stages of benthic calcifiers under CO₂ concentrations that are expected to occur in the future ocean (380–2000 μatm pCO₂ / pH 8.2~7.3). Although the magnitude of CO₂ tolerance may differ between species and life stages, effects of high CO₂ are proposed for several different life stages, including reproduction, egg, cleavage, larva, settlement and adult stages. Used with permission from Kurihara 2008.

predicted for 2100. Havenhand *et al.* (2008) found significant reductions in sperm swimming speed, motility and fertilisation success in the sea urchin *Heliocidaris erythrogramma* using a decrease of 0.4 units (pH of 7.7, 1000 ppm, -0.4 units compared to the control pH of 8.1). There was a reduction of 11.7% in swimming speed of sperm, 16.3% reduction in sperm motility, a 24.9% reduction in fertilisation success after 3-4 hours exposure. In addition the number of eggs developing into cleaving embryos after 20 hours exposure was 20% less when treated at an elevated concentration of CO₂ (Table 1). Such reductions in fertilisation were not detected in the next series of studies on the same species. Even with the additional stressor of elevated temperature, Byrne *et al.* (2009) found no significant effect of elevated CO₂ on the fertilisation success, cleavage and gastrulation in the sea urchin, *H. erythrogramma*; sourced from the same geographical region as Havenhand *et al.* (2008), with a larger pH range of 7.6-8.2 over a shorter exposure period (2-3 hours for cleavage and 19-20 hours for gastrulation). There was, however, a trend for decreased cleavage at elevated concentrations of CO₂ and pH of 7.6-7.9 compared to the control (8.2) at the higher temperature of 26 °C (4 units above ambient). Overall they detected no significant synergistic effect of elevated CO₂ and temperature on cleavage and gastrulation but rather a significant decrease in cleavage and gastrulation with increased temperature alone.

Such contrasting results, even for identical species, could be due to differences in experimental methodology arising from single male and female crosses in Havenhand *et al.* (2008) versus multiple males and females in Byrne *et al.* (2009; 2010a and 2010b) and high fertilisation success

in controls (Havenhand *et al.*, 2008 62% versus Byrne *et al.*, 2010 a, b 80-90%). It remains unclear whether intra and inter specific differences are real or due to differences in experimental techniques (Byrne *et al.* 2010a; Ericson *et al.* 2010; Reuter *et al.* 2011) by the pooling of gametes from males and females (Byrne *et al.* 2009, 2010a, 2010b), creating arguments about polyandry versus single crosses and their influence on results and interpretation and or misinterpretations and measurements in fertilisation due to polyspermy. Whether polyandry or single crosses, even recent studies are based on small sample sizes of males and females (Byrne *et al.* 2009 2 males: 2 females; Byrne *et al.* 2010a 3 males: 3 females; Ericson *et al.* 2010, 1 male: 1 female) and additionally may suffer from pseudoreplication where containers are not factored as a source of variation in the analysis. This raises doubt over whether differences are representative because of low replication (Moulin *et al.* 2011). Value would be added if post-hoc tests (e.g. Byrne *et al.* 2009) were done. Indeed the experimental design of Havenhand *et al.* (2008) used 5-6 males, while Reuter *et al.* (2011) used 12 male-female pairs of replicates over eight serial sperm dilutions and emphasised that fertilisation success may be inaccurately estimated if fertilisation success is a measure of both fertilised and polyspermy embryos. Other authors agree stating that differences in experimental methodologies hamper our ability to make robust generalisations (Dupont *et al.* 2010a).

Single stressor studies also reduce our understanding of the impacts of ocean acidification. These have been replaced in recent times by synergistic studies investigating the effect of elevated CO₂ and temperature on fertilisation. Against expectation, elevated CO₂ has

not been found to reduce fertilisation nor temperature to enhance fertilisation (Byrne *et al.* 2010a). There was also no synergistic interaction between elevated CO₂ and temperature and no trend for reduced fertilisation with increased concentration of CO₂ over a range of sperm concentrations. In a further study by Byrne *et al.* (2010b), on a range of echinoderms including once again the echinoid, *Heliocidaris erythrogramma* and the closely related species, *Heliocidaris tuberculata*, *Tripneustes gratilla*, *Centrostephanus rogersii* and the asteroid, *Patiriella regularis*, using multiple male and female combinations, there was no significant effect of elevated CO₂ nor temperature on fertilisation. There was a trend, however, for decreased fertilisation in the subtidal sea urchin, *H. tuberculata*, prompting the comment that species characterising shallow coastal waters are perhaps more robust to fluctuations in their environment (Byrne *et al.* 2010b). Similarly, Martin *et al.* (2011) found high fertilisation rates (>97%) with no effect on fertilisation down to a low pH (7.25) for the widely distributed Mediterranean and North Atlantic sea urchin, *Paracentrotus lividus*.

Although, fertilisation remains robust when synergistic factors such as temperatures are combined with elevated CO₂ (Byrne *et al.* 2010 a, b) this may not occur when sperm is limiting (Ericson *et al.* 2010; Reuter *et al.* 2011). Similar to Byrne *et al.* (2009, 2010a, 2010b), Ericson *et al.* (2010) found no effect on fertilisation of elevated CO₂ and decreased pH at levels predicted for the year 2100 for the echinoid *Sterechinus neumayeri*, but they did find an effect of elevated CO₂ when sperm concentration was suboptimal. Sperm concentration appears to be a significant factor for organisms when other conditions are suboptimal. Reuter *et al.* (2011) found maximum fertilisation rates in the sea urchin, *Strongylocentrotus franciscanus*, were maintained only at increased sperm concentrations. A reduction in fertilisation efficiency related to sperm concentration raised at pCO₂ levels and pH 7.81 and 7.55 (800 and 1800 ppm, respectively) and the time of egg block to polyspermy was also significantly longer at pH 7.55 (Reuter *et al.* 2011).

Studies such as (Byrne *et al.* 2010a b) and more recent findings (Moulin *et al.* 2011) have led to the suggestion that robustness of fertilisation in echinoderms to ocean acidification may be because they can acclimate to the low pH conditions experienced in their environment. In addition, the natural process of fertilisation is characterised by low pH conditions, with an inhibitory effect on sperm motility (Havenhand *et al.* 2008) over ridden by jelly peptides on eggs and respiratory effects (Byrne *et al.* 2010b). Moulin *et al.* (2011) found when gametes of the temperate sea urchin, *Paracentrotus lividus*, originated from individuals sourced from a more acidic tidal pool (which recorded an overnight drop to pH 7.4) fertilisation rates were greater in the acidic pool compared to control pool. Although in general fertilisation and cleavage rates decreased in the acidic pool, below pH 7.6 (control pH 8.0). Moulin's study, however, compares an acidic with a control pool and thus any differences between the 2 pools, may be due to chance, rather than represent the responses of sea urchins in rock pools with lowered pH.

Such acute studies with low replication limit our ability to make robust conclusions. Acute studies need to be replaced with chronic multigenerational studies which test hypotheses related to acclimation and pre adaptive capacity as suggested in molluscs (Parker *et al.* 2011). Thus the evidence for the potential of echinoderms to acclimate to acidification requires further investigations to be done.

Molluscs

Similar to echinoderms, the methodology of early studies investigating fertilisation in molluscs exposed eggs and sperm to elevated CO₂ used pH concentrations outside the scenarios predicted for 2300 (Caldeira and Wickett 2003; Table 1). Yet in contrast to the echinoderms there was no reduction in swimming speed of sperm, sperm motility, fertilisation or cleavage even at these low pH levels. Kurihara *et al.* (2008a) exposed the mussel, *Mytilus galloprovincialis*, to elevated concentrations of pCO₂ (2000 ppm) with resultant pH of 7.4 (a reduction of 0.7 pH units compared to controls), and found no significant differences in fertilisation between controls and elevated CO₂.

Intraspecific differences in fertilisation responses to elevated CO₂ in molluscs once again make general conclusions difficult. There was no effect of elevated CO₂ on sperm speed, sperm motility or fertilisation in the Pacific oyster, *Crassostrea gigas* (pH 7.4-7.8 Kurihara *et al.* 2007; pH 7.8 Havenhand and Schlegel 2009), located in Japan and Sweden respectively, but there was reduced fertilisation at pH 7.83 for this same species in Australia (Parker *et al.* 2010). In an unpublished follow up study using almost identical methodologies, sperm concentrations and exposure levels; there was still no difference in fertilisation responses of populations of *C. gigas* in Japan and Sweden, but significant differences for populations of *C. gigas* in Australia (Parker *et al.* unpublished data). These contrasting contradictory fertilisation responses to elevated CO₂ and lowered pH within identical species of molluscs is characteristic of acute studies done on sea urchins (Byrne *et al.* 2009, 2010b; Havenhand *et al.* 2008). Yet perhaps because the results occur even when using identical methodology, they provide support for models explaining that differences within and among populations are caused by environmental or genetic differences, rather than experimental methodology.

Intra and interspecific differences in fertilisation responses have also been found between related species of oyster; the Sydney Rock oyster, *Saccostrea glomerata* and the Pacific oyster, *Crassostrea gigas*. There were significant differences in fertilisation rates at elevated CO₂ in *S. glomerata* (pH 7.9; Parker *et al.* 2009, 2010), and when responses of *S. glomerata* and *C. gigas* were compared, there was a significantly greater reduction in fertilisation in *S. glomerata* compared to the more robust *C. gigas* (Parker *et al.* 2010). When these acute experiments combined the additional stressor of suboptimal or elevated temperature, there was a synergistic impact with elevated CO₂ and correspondingly significantly reduced fertilisation (reduction in fertilisation of up to

26% and 51% for *C. gigas* and *S. glomerata* respectively at suboptimal temperature). In molluscs, intra and species specific differences make generalisations difficult and such variable results may mirror the fluctuations in pH in these estuarine habitats where these intertidal species are found. In the only study done on a subtidal species, where pH may fluctuate less, there was no effect on fertilisation of temperature or elevated CO₂, using concentrations of CO₂ expected for 2100 on the temperate abalone, *Haliotis coccoradiata* (Byrne *et al.* 2010b).

Recent meta-analyses of the overall effects of elevated CO₂ on fertilisation are restricted by the paucity of papers available with reductions in effect size below 2000 ppm pCO₂ for sea urchin embryos (-9%, 3 papers, 2 species; Hendriks *et al.*, 2010) and yet increased effects for bivalves (+2%, maximum 2 papers 2 species) and a cephalopod (*Sepia officinalis*, +98% Hendriks and Duarte 2010), although the ppm range was not specified. With such small databases being employed, generalisations on the overall effect on fertilisation for groups is not possible, leaving our understanding of the impacts of ocean acidification on fertilisation based on species specific studies done in sometimes different geographic regions where population and experimental similarities and differences cloud the conclusions. Despite these unanswered questions, authors are now in agreement that fertilisation in echinoderms may be relatively robust to near future ocean acidification (Dupont *et al.* 2010a, c; Hendriks *et al.* 2010; Hendriks and Duarte 2010). This cannot be supported for fertilisation of molluscs nor widely for other groups when there is still a paucity of data on the diversity of marine invertebrates (Byrne *et al.* 2010b) and where results are outside the context of the normal fluctuations of pH in their environment, which may provide some pre adaptive capacity (Clark *et al.* 2009; Moulin *et al.* 2011; Parker *et al.* 2011).

Larval development and growth

Even if it turns out that fertilisation is robust to near future predictions of ocean acidification, larval developmental stages may be more vulnerable (Dupont *et al.* 2010a). Many marine invertebrates start to calcify in their larval or juvenile stage (Kurihara 2008; Kroeker *et al.* 2010). Several studies have suggested that the early life history stages may be more sensitive to environmental perturbations (Kurihara *et al.* 2007; Kurihara 2008; Dupont *et al.* 2010a, c).

Echinoderms

Larval development has been shown to be more sensitive than fertilisation to elevated concentrations of atmospheric CO₂, particularly for echinoderm embryos and larvae, with delayed development, reduced survival and size and skeletal abnormalities mainly investigated in planktotrophic larvae (Table 1). The most dramatic impacts have been measured in the planktotrophic ecological keystone brittlestar, *Ophiothrix fragilis*, (Dupont *et al.* 2008). Larvae of *O. fragilis* grown in CO₂-acidified seawater at pH 7.7-7.9 suffered 100% mortality within 8 days, while control larvae showed 70% survival over

the same time. There was also a decrease in larval size with no larvae raised at pH 7.9 reaching the normal 8 pluteus arm stage and a high proportion of these larvae were either abnormal or had altered skeletal proportions and asymmetry during skeletogenesis. Further, there was a delay in development at low pH, with larvae at elevated CO₂ treatments taking longer to reach the same developmental stage (Dupont *et al.* 2008); for example, 50% of the control larvae were 6 armed after 5.42 days compared to 5.73 days at pH 7.9. Similarly after 2 days while abnormalities were completely absent in the control larvae, larvae at low pH with no other abnormalities had high proportions of asymmetry (25% at pH 7.9 and 32% at 7.7, Dupont *et al.* 2008).

These more recent studies support the findings of earlier studies reporting delayed development and abnormal pluteus morphology in the planktotrophic sea urchins, *Hemicentrotus pulcherrimus* and *Echinometra mathaei*, in which used a wider range of pH levels, some being within emission scenarios for the end of the century (Kurihara and Shirayama 2004; Kurihara *et al.* 2004b). In acute exposure experiments of three days, *H. pulcherrimus* larvae grown at pH 6.8 failed to develop arms and spicules or reach pluteus stage, consequently displaying high abnormality (Kurihara and Shirayama 2004). Larvae grown at pH 7.8 and 7.6 were smaller than controls and formed normal triangular and abnormally trapezium shapes respectively (Kurihara and Shirayama 2004; Figure 3). Using a similar pH range larval development in another temperate planktotrophic sea urchin, *Paracentrotus lividus*, was only affected below pH 7.4 with an increase in abnormal pluteus larva, although rod length was shorter only below pH 7.2 (Moulin *et al.* 2011). Similar to *H. pulcherrimus* (Kurihara and Shirayama 2004), at the acute pH of 6.8 all *P. lividus* embryos failed to develop to pluteus stage, displaying an abnormal bowl-shaped morphology (Moulin *et al.* 2011).

The negative responses to elevated CO₂ in planktotrophic larvae are not necessarily reflected across all echinoderms. Elevated CO₂, had no negative impact on survival or skeletal formation on the lecithotrophic larvae of the sea star, *Crossaster papposus* over the 38 day duration of the experiment. In fact there was a positive effect, on larvae which grew 2.8 times faster and juveniles 2.2 times faster at pH 7.7 compared to controls; 50% of larvae had progressed into the juvenile stage by day 28, whilst 95% of controls had developed rudimentary, but remained in the preceding settlement stage (Dupont *et al.* 2010b). In contrast, Havenhand *et al.* (2008; 2010b), that the percentage of eggs which developed into swimming larvae of the lecithotrophic sea urchin, *Heliocidaris erythrogramma*, after 24 hours was 25.9% lower than the control. Dupont *et al.* (2010b) suggested the need for further investigation of the impact of ocean acidification on lecithotrophic larvae, because the lecithotrophic cephalopod, *Sepia officinalis*, developed and calcified in low pH (Gutowska and Melzner 2009). These studies show that even amongst marine organisms with lecithotrophic development there can be variability in response.




Figure 3. Larval morphology in the sea urchins *Hemicentrotus pulcherrimus* (a,b,c) *Echinometra mathaei* (d,e,f), and the bivalves *Crassostrea gigas* (g, h, i) and *Mytilus galloprovincialis* (j, k), exposed to pCO₂ concentrations of 380 ppm (control, a, d, g, j), 1000 ppm (b, e, h) and 2000 ppm (c, f, i, k). Scale bars represent 50 μm. Figure from Kurihara (2008).

There is little evidence to suggest that a geographic distribution model explains the delayed development and skeletal abnormalities in echinoderm larvae. Clark *et al.*, (2009), tested the prediction that the calcification of larvae of polar species would be more negatively impacted than temperate and tropical species because calcification is less energetically favourable at higher latitudes, and found opposite to that predicted. The Antarctic species, *Sterechinus neumayeri*, was least affected by elevated CO₂ while there were variable yet similar impacts on sea urchin larvae distributed from the tropic to Antarctic regions. Lowering of the pH to 7.0, resulted in significantly reduced survival of tropical, *Triploneustes gratilla*, but did not significantly affect survival of the temperate species, *Pseudechinus huttoni* and *Evechinus chloroticus*, or the polar species, *Sterechinus neumayeri*. The main effect, however, was a reduction in calcification of pluteus larvae which ranged from a small and non-significant decrease of 3.9% in polar, *Sterechinus neumayeri*, to significant decreases of 13.8% in tropical, *Triploneustes gratilla*, 30.6% in temperate, *Evechinus chloroticus*, and 36.9% in temperate, *Pseudechinus huttoni*, with 0.4 pH units reduction (7.51-7.87) compared to controls (7.9-8.17) and a reduction

in growth for 3 out of the 4 species, but with no significant difference in *P. huttoni* (treatments pH 7.6-7.8, controls 8-8.2). Although acidified seawater did not change the morphology of larval skeletons among species. As found in other studies, there were fine scale differences in skeletal structures with an apparent loss of integrity of the surfaces of the skeletal rods for temperate, *E. chloroticus* (was pitted), *P. huttoni* (was eroded), contrasting with the smooth skeletal rods from larvae of tropical, *T. gratilla*, and polar, *S. neumayeri*, grown at ambient or reduced pH. Further, Ericson *et al.* (2010) found no difference among pH treatments (7, 7.3, 7.7 and ambient 8.0) in rate of development or visual abnormality in *Sterechinus neumayeri* larvae, the polar species used by Clark *et al.* (2009), through early development up to coeloblastula stage (day 2 post fertilisation). Significant differences among pH treatments first appeared at gastrulation (day 6) where embryos were 20% shorter, displayed no normal larval development, had rupturing of the gastrula epithelium and irregular morphology at pH 7, but no difference in abnormality levels or length among the other treatments (although at pH 7.3 some delayed development was apparent).

Multiple stressor experiments, combining elevated CO₂ and temperature are more rare, but Sheppard Brennan *et al.* (2010), testing the hypothesis that temperature enhancement would counteract the negative effect of acidification, found elevated CO₂ reduced calcification in the tropical sea urchin, *Triploneustes gratilla*, used by Clark *et al.* (2009). Varying temperatures (24 °C-control, 27 and 30°C) and pH levels (7.6, 7.8 and control 8.15), significantly increased larval growth as temperature increased until the thermal limit was reached (30°C), and decreased as pH decreased, although difference in post oral arm length (larval asymmetry) was sensitive to temperature, not pH. These increased growth rates with +3°C rise in temperature effectively reduced, but did not negate, adverse effects in larvae raised at low pH (Sheppard Brennan *et al.* 2010). The effect of elevated temperature and lowered pH on growth of *T. gratilla* larvae was similar to the findings of Clark *et al.* (2009), once again highlighting the species specific nature of responses to environmental stressors. Byrne *et al.* (2011a) in a similar experiment, but on the temperate species, *Heliocidaris erythrogramma*, found that both temperature and elevated CO₂ reduced the number of spines and percentage of juveniles developing, but that increased temperature reduced the overall negative effect of elevated CO₂.

The search to determine the vital physiological mechanisms and processes affected in marine invertebrates by elevated CO₂ during larval development is revealing that gene expression can be compromised at elevated atmospheric CO₂ levels and temperature in the sea urchins, *Strongylocentrotus franciscanus* and *Lytechinus pictus* (O'Donnell *et al.* 2009, 2010). Potential impairment of physiological response to heat stress by *S. franciscanus* larvae at the four-arm echinopluteus stage was observed at pCO₂ levels of 540 and 970 ppm (pH 7.98 and 7.87 respectively), with significant up-regulation and change in expression profile of the heat stress gene *hsp70*, resulting in peak gene expression occurring 2°C higher than controls (O'Donnell *et al.* 2009). Although 540 ppm pCO₂ (pH 7.87) had no effect on *Lytechinus pictus* raised to pluteus stage, at 970 ppm (pH 7.78) larvae were significantly smaller and had more triangular skeletons and shorter arms than controls, as well as down regulation of biomineralization, skeletogenesis and energy metabolism genes, whilst some acid-base and ion regulation genes were up-regulated (O'Donnell *et al.* 2010). Similarly Martin *et al.* (2011) found up-regulation of development and biomineralisation genes by factors of up to 26 at low pH.

In the absence of adequate adaptation or acclimation, such CO₂-induced sub lethal effects on *O. fragilis* and other echinoderms species may result in the disappearance of these organisms from the surface oceans within the next 50–100 years (Dupont *et al.* 2008). As studies accumulate there is some evidence that some species may be resilient. For example, although larvae of the purple sea urchin *Strongylocentrotus purpuratus* were significantly smaller at pH of 7.7 and 7.5 after 6 days of exposure to elevated CO₂, the difference was only 7–13% less than

the controls suggesting potential resilience (Yu *et al.* 2011). Further, the widely distributed Mediterranean and North Atlantic sea urchin, *Paracentrotus lividus*, was extremely resistant to low pH (down to 7.25 pH) with no effect on larval survival. Also although there was a significant delay in larval development, this had no impact on relative larval morphology or calcification (Martin *et al.* 2011). Overall, however, to date studies from a range of geographical regions show that sea urchin, brittlestar and seastar larvae may develop more slowly, be smaller, have skeletal abnormalities and have impaired physiological responses to environmental stressors under elevations in CO₂ which may be either ameliorated or exacerbated by temperature increases predicted for the end of the century. Much more research is required on larval physiology to determine whether there is an increase or decrease in metabolism (Parker *et al.* 2012).

Molluscs

The vulnerability to elevated CO₂ observed in echinoderms is also reported in studies on larvae of molluscs (Table 1). Although there was no effect on fertilisation, in response to reductions of pH to 7.4, veliger larvae of the Pacific oyster *Crassostrea gigas* were reduced in size and had shell malformations (Kurihara *et al.* 2007). When embryos of *Crassostrea gigas* were reared at elevated pCO₂ of 2268 ppm for 48 hours, there was no significant difference in development relative to the controls until the D veliger stage (Kurihara *et al.* 2007) and the onset of shell mineralization (Waller 1981; Hayakaze and Tanabe 1999; Kurihara *et al.* 2007). At the completion of the experiment, however, the embryos of *C. gigas* were found to have reduced rates of development, calcification and growth and increased rates of abnormal development, with only 5% of the CO₂-stressed embryos developing normally to D-stage larvae compared to 68% in the controls (Kurihara *et al.* 2007; Table 1, Figure 3.4). In embryos of the mussel *Mytilus galloprovincialis* reared at elevated pCO₂ of 2000 ppm (pH 7.4), there was also a delay in development visible once the mussels reached the trochophore larval stage (Kurihara *et al.* 2008a). This was followed by morphological shell abnormalities such as convex hinge and mantle protrusion, as well as a 26% and 20% reduction in shell height and length, respectively (Kurihara *et al.* 2008a; Table 1, Figure 3). Similarly, a 4.5±1.3% and 12±5.4% decrease in shell length and thickness respectively was observed in mussel larvae of *Mytilus edulis* raised at pH 7.8 (Gazeau *et al.* 2010). Those raised at pH 7.6 had a reduced shell length of 6±2.3%, overall size of D-veligers was reduced 12.7±0.9% and hatching rates were suppressed by 24±4% (Gazeau *et al.* 2010). Embryos of the gastropod mollusc, *Littorina obtusata*, raised at pH 7.6 also showed shell changes with shorter lateral, but longer spiral shell length (Ellis *et al.* 2009). Mean development time was statistically significantly longer (although only 0.45 day: 27.5±0.1 day for treatments; 26.9±0.05 for controls) and hatching rate was lower at pH 7.6 than in controls (pH 8.1).

Similarly, Parker *et al.* (2010) found reduced development, size and increased abnormality in D veliger larvae and spat for both *C. gigas* and *Saccostrea glomerata* to elevated $p\text{CO}_2$ (600, 750, 1000 ppm, mean pH 8.02, 7.95 and 7.83 respectively) compared to controls (375 ppm, pH 8.2) (Parker *et al.* 2009, 2010) as did Watson *et al.* (2009) for *S. glomerata*. When elevated CO_2 was combined with temperature, (18, 22, 26 and 30°C) differences increased at suboptimal temperatures (outside 26°C) and elevated CO_2 . The pediveligers of *S. glomerata* were more sensitive than spat to changes to CO_2 and temperature, but this reversed in *C. gigas* with newly metamorphosed spat showing greater sensitivity than pediveligers (Parker *et al.* 2010). This result is surprising given that the CaCO_3 polymorph (calcite) deposited by spat is less soluble than that deposited by the preceding pediveliger larval stage. Differences in responses between closely related species have been found in other studies. For example Miller *et al.* (2009) compared the response of veliger larval development exposed to a range of elevations of CO_2 on of *Crassostrea virginica* and *C. ariakensis*. Whereas *C. virginica* experienced a 16% decrease in shell area and a 42% reduction in calcium content, there was no impact on the larval development of *C. ariakensis*. Both species demonstrated net calcification and growth even when aragonite was undersaturated.

Given that the first effects of elevated CO_2 on the bivalve species were observed during the trochophore and D-veliger stage, corresponding with the first onset of shell mineralisation, it has been suggested that calcification during larval development will be one of the major processes affected by elevated CO_2 (Kurihara *et al.* 2007; Kurihara 2008), although other authors such as Kurihara *et al.* (2007) suggest that other processes such as protein synthesis and metabolism may also be affected. Rather than earlier stages being more sensitive than later (Beiras and His 1994) there is an absence of evidence of hierarchical sensitivity and more evidence of sensitivity of larval stages being species specific. Variable sensitivity of life stages to low pH has been found in the lecithotrophic abalone, *Haliotis rufescens*. Thermal tolerance was impaired at pH 7.87 compared to control (pH 8.05) for pretorsion and late veliger life stages, but not posttorsion and premetamorphic veligers (Zippay and Hofmann 2010). Supporting the findings of Zippay and Hoffman (2010), more recent studies on abalone have found that in response to elevated CO_2 that larvae of *Haliotis cocciradiata* failed to produce skeletons when reared at -0.4-0.6 pH less than controls (Byrne *et al.* 2011a) and *Haliotis kamtschatkana* had reduced survival, increased abnormalities in shell structure at elevated CO_2 , with larvae in the highest treatments of 1800 ppm CO_2 (-0.4-0.5 pH less than controls) either developed an abnormal shell or lacked a shell completely (Crim *et al.* 2011). Of those larvae that did survive to the end of the experiment, however, there was no effect on settlement whether larvae had or did not have a shell (Crim *et al.* 2011).

Shell abnormalities were observed in veliger larvae of the Mediterranean pteropod *Cavolinia inflexa* exposed to acidified seawater (Comeau *et al.* 2010). Reduced growth and flat, irregular shell shape was observed at pH

7.82, but at pH 7.61 shells were absent after 13 days due to dissolution, yet larvae displayed “normal” swimming action (Comeau *et al.* 2010). Pre-winter juveniles of the polar pteropod *Limacina helicina* also had more degraded shells when raised at or below pH 7.78 (Lischka *et al.* 2010). Shell diameter, degradation and increment were significantly affected by CO_2 , but not temperature, whereas the combination of elevated temperature and CO_2 increased mortality, yet with temperature the more dominant factor (Lischka *et al.* 2010). Survival, growth, rate of metamorphosis, size, shell thickness and hinge integrity decreased significantly with increasing CO_2 of the bay scallop *Argopecten irradians* and the hard clam *Mercenaria mercenaria* at $p\text{CO}_2$ levels of 250, 390, 750 and 1500 ppm (pH 8.17, 8.05, 7.8 and 7.5 respectively) (Talmage and Gobler 2010). Waldbusser *et al.* (2010) also revealed a decrease in calcification rate with reducing pH and organism size, indicating size dependency of pH effects of *Mercenaria* spp. when subjected to $p\text{CO}_2$ of 1120 and 1950 ppm (pH 7.64 and 7.41 respectively) with calcification rate differing between source populations signifying possible genotypic variation. Parker *et al.* (2011) also found significant variation in growth of spat among populations of the endemic Sydney Rock Oyster, *Saccostrea glomerata*. There was no effect of elevated CO_2 in mass and single pair mated family lines which had been bred for resistance to overwintering and QX disease. Indicating that there may be inherent genetic diversity and feeding efficiency among populations of species or environmental pre adaptive capacity which may ameliorate the impacts of ocean acidification on oyster populations (Parker *et al.* 2011).

In contrast to the other molluscs with mainly planktotrophic larvae, the lecithotrophic juvenile European cuttlefish, *Sepia officinalis*, showed no adverse growth or developmental effects when raised at $p\text{CO}_2$ of 4000 and 6000 ppm (pH 7.23 and 7.1 respectively) compared to controls at 628-705 ppm (pH 7.94-8.01) (Gutowska *et al.* 2008). In fact, at 6000 ppm significantly more CaCO_3 ($0.80 \pm 0.15\text{g}$) was accreted into cuttlebones than in controls ($0.71 \pm 0.15\text{g}$) (Gutowska *et al.* 2008). A later study by Lacoue-Labarthe *et al.* (2009) found eggs of the same species increased in weight as $p\text{CO}_2$ decreased without affecting hatchling size (900 and 1400 ppm; pH 7.85 and 7.6 respectively). There are also potential ecotoxicological consequences associated with elevated CO_2 . Lacoue-Labarthe *et al.* (2009) also found accumulation of metals varied: $^{110\text{m}}\text{Ag}$ increased and, ^{109}Cd decreased with decreasing pH, whereas ^{65}Zn accumulation was highest at pH 7.85, but lower at pH 7.6 than 8.1 (control). Similarly there was no effect on calcification rates, size or weight in juveniles of the grooved carpet clam *Ruditapes decussates* raised at $p\text{CO}_2$ of 1694 and 4245 ppm (pH 7.84 and 7.46 respectively), though mortality was reduced in acidified treatment (pH 7.46), possibly due to delayed reproductive development of clams preventing spawning in acidified treatments (Range *et al.* 2011) as an energy saving survival strategy. Decreased pH also had no effect on expression of two shell mineralisation genes *ap24* or *engrailed* at any developmental stage in the abalone *Haliotis rufescens* (Zippay and Hofmann 2010).

Meta analyses

The overall response of larval development of both echinoderms and molluscs to elevated CO₂ and associated decrease in pH appears highly variable, yet with a general negative effect. Indeed such generalisations are difficult because such a synthesis masks the complex interaction of species specific sensitivities during progressive life stages, early feeding strategies, latitudinal distribution, previous acclimation due to habitat and robustness of gene expression processes. Meta-analyses (Hendriks *et al.* 2010; Hendriks and Duarte 2010 and Kroeker *et al.* 2010) do not support the early life history stages being the most sensitive stages to ocean acidification compared to adults. Hendriks *et al.* (2010) found reduced larval growth (-16%) in sea urchins, but found adult growth significantly more sensitive (-62%), yet data was limited with only 2 papers and 2 species contributing to the sea urchin embryo and 1 paper and 2 species contributing to the adult data. A later exercise by Hendriks and Duarte (2010) consolidated this position concluding there was no difference in effect size of different life stages by pooled groups of organisms using a dataset restricted to 16 studies species in 2 studies. Yet in contrast Dupont *et al.* (2010a) found gametes and early developmental stages of echinoderms appeared “far more impacted” than adults relative to calcification, growth and survival although the number of experimental responses was heavily weighted towards larvae with 47, 10 and 17 responses for larvae, juveniles and adults respectively. These conflicting views on sensitivity of early life stages have been further addressed by a more extensive meta-analysis by Kroeker *et al.* (2010), on the effects of ocean acidification on life stages and biological processes of marine organisms based on 73 studies with 251 experimental results finding no variation in mean effect among life stages for different biological variables concluding “this synthesis did not support all the leading hypotheses for variation in response to ocean acidification (i.e., early developmental stages and more soluble CaCO₃ polymorphs are more susceptible to ocean acidification). We instead found the explanatory power of these hypotheses was specific to organisms within taxonomic groups. Such meta-analyses, although authoritative are based on a paucity of studies, particularly in relation to molluscs and echinoderms and it may be premature in their attempt to use them to support hypotheses related to overall trends.

Conclusion

Despite recent research, our current understanding of the biological consequences of an acidifying ocean over the next century is relatively in its infancy and there are still large unknowns. Increasingly studies are considering the synergistic effects of other stressors (i.e. temperature, nutrients, hypoxia). In order to fully understand the consequences of ocean acidification at the population and ecosystem level the synergistic or antagonistic impacts of environmental variables must be considered. The echinoderms and molluscs discussed in this review contribute significant ecological roles to the marine environment. In the case of molluscs, forming habitats and

providing ecosystem services such as water purification, as a food source for other organisms and producing large amounts of CaCO₃ (Comeau *et al.* 2010). Also molluscs and echinoderms have enormous economic value. In 2007 in the United States molluscs alone, accounted for 748 million USD of domestic ex-vessel revenues and in 2008, 13.1 million tonnes of shellfish were produced by aquaculture worldwide, with a market value of over 13 billion USD (Gazeau *et al.* 2010). In Japan sea urchin roe is worth over 320 million USD alone. Thus any negative impacts to fertilisation and larval development even if sub lethal may result in severe economic and ecological consequences. Over the coming decades, it is likely that a rising surface ocean CO₂ will also be accompanied by rising surface ocean temperatures, yet only a handful of studies have considered the synergistic impacts of both elevated CO₂ and temperature (Reynaud *et al.* 2003; Metzger *et al.* 2007; Anthony *et al.* 2008; Byrne *et al.* 2009; Gooding *et al.* 2009; Martin and Gattuso 2009; Munday *et al.* 2009a; Parker *et al.* 2009, 2010). Current models also remain largely based on acute experiments over single generations where the sudden drop in pH of 0.4 units does not mimic well the longer time frame over which this will occur (0.0044pH/yr Hendriks and Duarte 2010). Future experiments are needed to consider the potential for species to acclimate over long term perturbations (Parker *et al.* 2012) and not enough studies yet are investigating the variability in responses of organisms within and between populations (Parker *et al.* 2011; Waldbusser *et al.* 2010) perhaps explaining some of the variability behind contrasting findings, albeit with also contrasting methods (Byrne *et al.* 2009 with Havenhand *et al.* 2008). Those that have, are finding that differences do exist (Parker *et al.* 2011), and the potential to ameliorate climate change perhaps through environmentally induced plasticity (Bibby *et al.* 2008; Hofman *et al.* 2010). Further work is, however, required on the underlying physiological mechanisms which may allow for species adaptation or acclimation (Parker *et al.* 2012).

Negative results from short term studies on larvae make it difficult to extrapolate to long term impacts on marine environments, especially when these potentially could couple with positive effects on adults (increased production and robustness of propagules, Hendriks and Duarte 2010) and greater capacity to acclimate (Moulin *et al.* 2011; Parker *et al.* 2012). This gives further support for an entire life cycle environmental approach to measuring the impact of ocean acidification. Difficulty in extrapolation also occurs because laboratory studies and results are not necessarily repeatable in the field. Caution is required in translating between these two contexts, where in the laboratory all variables are held constant except the treatment and the opposite in the field where sources of variability are numerous and interactive (Connell 1975; Green 1979; Underwood 1997) with unforeseen consequences. In order to have a greater understanding of how organisms will respond, over the next century, to warming and acidifying oceans and the capacity for species to adapt, we must move beyond measuring acute responses and focus on identifying the underlying mechanisms involved in the sensitivity and/

or acclimation of marine species to elevated CO₂ levels (Pörtner 2008). The physiological, molecular and gene mechanisms involved in ocean acidification responses have also been relatively unexplored (European Science Foundation 2009) and have been identified as priorities

of future research (Orr et al. 2009; Doney et al. 2009; Pörtner 2008). Such research is important if we are to create the authoritative and robust meta-analyses and generalisations that we need to manage our marine environments (Hofman et al. 2010; Kroeker et al. 2010).

References

- Alvarado-Alvarez, R., Gould, M. C. and Stephano, J. L. 1996. Spawning, in vitro maturation, and changes in oocyte electrophysiology induced by serotonin in *Tiwela stultorum*. *Biological Bulletin* 190(3): 322-328.
- Anger, K. 1987. The DO threshold: a critical point in the larval development of decapod crustaceans. *Journal of Experimental Marine Biology and Ecology* 108(1): 15-30.
- Anlauf, H., D'Croz, L. and O'Dea, A. 2011. A corrosive concoction: The combined effects of ocean warming and acidification on the early growth of a stony coral are multiplicative. *Journal of Experimental Marine Biology and Ecology* 397(1): 13-20.
- Anthony, K. R. N., Kline, D. I., Diaz-Pulido, G., Dove, S. and Hoegh-Guldberg, O. 2008. Ocean acidification causes bleaching and productivity loss in coral reef builders. *Proceedings of the National Academy of Sciences of the United States of America* 105(45): 17442-17446.
- Arnold, K. E., Findlay, H. S., Spicer, J. I., Daniels, C. L. and Boothroyd, D. 2009. Effect of CO₂-related acidification on aspects of the larval development of the European lobster, *Homarus gammarus* (L.). *Biogeosciences* 6(8): 1747-1754.
- Beiras, R. and His, E. 1994. Effects of dissolved mercury on embryogenesis, survival, growth and metamorphosis of *Crassostrea gigas* oyster larvae. *Marine Ecology Progress Series* 113(1-2): 95-104.
- Bibby, R., Widdicombe, S., Parry, H., Spicer, J. and Pipe, R. 2008. Effects of ocean acidification on the immune response of the blue mussel *Mytilus edulis*. *Aquatic Biology* 2(1): 67-74.
- Byrne, M. 2011. Impact of ocean warming and ocean acidification on marine invertebrate life history stages: vulnerabilities and potential for persistence in a changing ocean. *Oceanography and Marine Biology – An Annual Review* 49: 1-42.
- Byrne, M., Ho, M., Selvakumaraswamy, P., Nguyen, H. D., Dworjanyn, S. A. and Davis, A. R. 2009. Temperature, but not pH, compromises sea urchin fertilization and early development under near-future climate change scenarios. *Proceedings of the Royal Society B: Biological Sciences* 276(1663): 1883-1888.
- Byrne, M., Ho, M., Wong, E., Selvakumaraswamy, P., Sheppard Brennand, H., Dworjanyn, S. A. and Davis, A. R. 2011a. Unshelled abalone and corrupted urchins, development of marine calcifiers in a changing ocean. *Proceedings of the Royal Society B: Biological Sciences* 278(1716): 2376-2383.
- Byrne, M., Selvakumaraswamy, P., Ho, M. A., Woolsey, E. and Nguyen, H. D. 2011b. Sea urchin development in a global change hotspot, potential for southerly migration of thermotolerant propagules. *Deep-Sea Research Part II: Topical Studies in Oceanography* 58(5): 712-719.
- Byrne, M., Soars, N. A., Ho, M. A., Wong, E., McElroy, D., Selvakumaraswamy, P., Dworjanyn, S. A. and Davis, A. R. 2010a. Fertilization in a suite of coastal marine invertebrates from SE Australia is robust to near-future ocean warming and acidification. *Marine Biology* 157(9): 2061-2069.
- Byrne, M., Soars, N.A., Selvakumaraswamy, P., Dworjanyn, S. A. and Davis, A. R. 2010b. Sea urchin fertilization in a warm, acidified and high pCO₂ ocean across a range of sperm densities. *Marine Environmental Research* 69(4): 234-239.
- Calabrese, A. and Davis, H. C. 1966. The pH tolerance of embryos and larvae of *Mercenaria mercenaria* and *Crassostrea virginica*. *Biological Bulletin* 131(3): pp. 427-436.
- Caldeira, K. and Wickett, M. 2005. Ocean model predictions of chemistry changes from carbon dioxide emissions to the atmosphere and ocean. *Journal of Geophysical Research C: Oceans* 110(9): 1-12.
- Caldeira, K. and Wickett, M. E. 2003. Anthropogenic carbon and ocean pH. *Nature* 425(6956): 365.
- Cigliano, M., Gambi, M. C., Rodolfo-Metalpa, R., Patti, F. P. and Hall-Spencer, J. M. 2010. Effects of ocean acidification on invertebrate settlement at volcanic CO₂ vents. *Marine Biology* 157(11): 2489-2502.
- Cipollaro, M., Corsale, G. and Esposito, A. 1986. Sublethal pH decrease may cause genetic damage to eukaryotic cell: A study on sea urchins and *Salmonella typhimurium*. *Teratogenesis Carcinogenesis and Mutagenesis* 6(4): 275-287.
- Clark, D., Lamare, M. and Barker, M. 2009. Response of sea urchin pluteus larvae (Echinodermata: Echinoidea) to reduced seawater pH: A comparison among a tropical, temperate, and a polar species. *Marine Biology* 156(6): 1125-1137.
- Comeau, S., Gorsky, G., Alliouane, S. and Gattuso, J. P. 2010. Larvae of the pteropod *Cavolinia inflexa* exposed to aragonite undersaturation are viable but shell-less. *Marine Biology* 157(10): 2341-2345.
- Connell, J. H. 1975. Some mechanisms producing structure in natural communities: a model and evidence from field experiments Pp. 460-490 in *Ecology and Evolution of Communities*, edited by M. Cody and J. Diamond. Harvard University Press, Cambridge, MA, USA.
- Crim, R. N., Sunday, J. M. and Harley, C. D. G. 2011. Elevated seawater CO₂ concentrations impair larval development and reduce larval survival in endangered northern abalone (*Haliotis kamtschatkana*). *Journal of Experimental Marine Biology and Ecology* 400(1-2): 272-277.
- Desrosiers, R. R., Désilets, J. and Dubé, F. 1996. Early developmental events following fertilization in the giant scallop *Placopecten magellanicus*. *Canadian Journal of Fisheries and Aquatic Sciences* 53(6): 1382-1392.
- Doney, S. C., Fabry, V. J., Feely, R. A. and Kleypas, J. A. 2009. Ocean acidification: the other CO₂ problem. *Annual Review of Marine Science* 1(1): 169-192.
- Dupont, S., Dorey, N. and Thorndyke, M. 2010a. What meta-analysis can tell us about vulnerability of marine biodiversity to ocean acidification? *Estuarine, Coastal and Shelf Science* 89(2): 182-185.
- Dupont, S., Havenhand, J., Thorndyke, W., Peck, L. and Thorndyke, M. 2008. Near-future level of CO₂-driven ocean acidification radically affects larval survival and development in the brittlestar *Ophiothrix fragilis*. *Marine Ecology Progress Series* 373: 285-294.
- Dupont, S., Lundve, B. and Thorndyke, M. 2010b. Near future ocean acidification increases growth rate of the lecithotrophic larvae and juveniles of the sea star *Crossaster papposus*. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution* 314 B(5): 382-389.

- Dupont, S., Ortega-Martínez, O. and Thorndyke, M. 2010c. Impact of near-future ocean acidification on echinoderms. *Ecotoxicology* 19(3): 449-462.
- Dupont, S. and Thorndyke, M. C. 2009. Impact of CO₂-driven ocean acidification on invertebrates early life-history - What we know, what we need to know and what we can do. *Biogeosciences Discussions* 6(2): 3109-3131.
- Egilsdottir, H., Spicer, J. I. and Rundle, S. D. 2009. The effect of CO₂ acidified sea water and reduced salinity on aspects of the embryonic development of the amphipod *Echinogammarus marinus* (Leach). *Marine Pollution Bulletin* 58(8): 1187-1191.
- Ellis, R. P., Bersey, J., Rundle, S. D., Hall-Spencer, J. M. and Spicer, J. I. 2009. Subtle but significant effects of CO₂ acidified seawater on embryos of the intertidal snail, *Littorina obtusata*. *Aquatic Biology* 5(1): 41-48.
- Ericson, J. A., Lamare, M. D., Morley, S. A. and Barker, M. F. 2010. The response of two ecologically important Antarctic invertebrates (*Sterechinus neumayeri* and *Parborlasia corrugatus*) to reduced seawater pH: effects on fertilisation and embryonic development. *Marine Biology* 157(12): 2689-2702.
- European Science Foundation 2009. Impacts of ocean acidification, Science Policy Briefing. 37: 1-12.
- Findlay, H. S., Kendall, M. A., Spicer, J. I. and Widdicombe, S. 2009. Future high CO₂ in the intertidal may compromise adult barnacle *Semibalanus balanoides* survival and embryonic development rate. *Marine Ecology Progress Series* 389: 193-202.
- Findlay, H. S., Kendall, M. A., Spicer, J. I. and Widdicombe, S. 2010a. Post-larval development of two intertidal barnacles at elevated CO₂ and temperature. *Marine Biology* 157(4): 725-735.
- Findlay, H. S., Kendall, M. A., Spicer, J. I. and Widdicombe, S. 2010b. Relative influences of ocean acidification and temperature on intertidal barnacle post-larvae at the northern edge of their geographic distribution. *Estuarine, Coastal and Shelf Science* 86(4): 675-682.
- Frommel, A. Y., Stiebens, V., Clemmesen, C. and Havenhand, J. 2010. Effect of ocean acidification on marine fish sperm (Baltic cod: *Gadus morhua*). *Biogeosciences Discussions* 7(4): 5859-5872.
- Gattuso, J. P., Gao, K., Lee, K., Rost, B. and Schulz, K. G. 2010. Approaches and tools to manipulate the carbonate chemistry Pp. 41-51 in *Guide to Best Practices for Ocean Acidification Research and Data Reporting*, edited by U. Riebesell, V. J. Fabry, L. Hansson and J. P. Gattuso. Publications Office of the European Union, Luxembourg.
- Gazeau, E., Gattuso, J. P., Dawber, C., Pronker, A. E., Peene, E., Peene, J., Heip, C. H. R. and Middelburg, J. J. 2010. Effect of ocean acidification on the early life stages of the blue mussel (*Mytilus edulis*). *Biogeosciences Discussions* 7(2): 2927-2947.
- Gooding, R. A., Harley, C. D. G. and Tang, E. 2009. Elevated water temperature and carbon dioxide concentration increase the growth of a keystone echinoderm. *Proceedings of the National Academy of Sciences of the United States of America* 106(23): 9316-9321.
- Gosselin, L. A. and Qian, P. Y. 1997. Juvenile mortality in benthic marine invertebrates. *Marine Ecology Progress Series* 146(1-3): 265-282.
- Green, M. A., Jones, M. E., Boudreau, C. L., Moore, R. L. and Westman, B. A. 2004. Dissolution mortality of juvenile bivalves in coastal marine deposits. *Limnology and Oceanography* 49(3): 727-734.
- Guinotte, J. M. and Fabry, V. J. 2008. Ocean acidification and its potential effects on marine ecosystems. *Annals of the New York Academy of Sciences* 1134: 320-342.
- Gutowska, M. A. and Melzner, F. 2009. Abiotic conditions in cephalopod (*Sepia officinalis*) eggs: Embryonic development at low pH and high pCO₂. *Marine Biology* 156(3): 515-519.
- Gutowska, M. A., Pörtner, H. O. and Melzner, F. 2008. Growth and calcification in the cephalopod *Sepia officinalis* under elevated seawater pCO₂. *Marine Ecology Progress Series* 373: 303-309.
- Hall-Spencer, J. M., Rodolfo-Metalpa, R., Martin, S., Ransome, E., Fine, M., Turner, S. M., Rowley, S. J., Tedesco, D. and Buia, M. C. 2008. Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. *Nature* 454(7200): 96-99.
- Hart, M. W. and Strathmann, R. R. 1995. Mechanisms and rates of suspension feeding Pp. 193-222 in *Ecology of Marine Invertebrate Larvae*, edited by L. McEdward. CRC Press, Boca Raton, FL, USA.
- Hauton, C., Tyrrell, T. and Williams, J. 2009. The subtle effects of sea water acidification on the amphipod *Gammarus locusta*. *Biogeosciences Discussions* 6(1): 919-946.
- Havenhand, J. N., Buttler, F. R., Thorndyke, M. C. and Williamson, J. E. 2008. Near-future levels of ocean acidification reduce fertilization success in a sea urchin. *Current Biology* 18(15).
- Havenhand, J. N. and Schlegel, P. 2009. Near-future levels of ocean acidification do not affect sperm motility and fertilization kinetics in the oyster *Crassostrea gigas*. *Biogeosciences* 6(12): 3009-3015.
- Hayakaze, E. and Tanabe, K. 1999. Early larval shell development in mytilid bivalve *Mytilus galloprovincialis*. *Venus* 58: 119-127.
- Hendriks, I. E. and Duarte, C. M. 2010. Ocean acidification: separating evidence from judgment - A reply to Dupont et al. *Estuarine, Coastal and Shelf Science* 89(2): 186-190.
- Hendriks, I. E., Duarte, C. M. and Álvarez, M. 2010. Vulnerability of marine biodiversity to ocean acidification: a meta-analysis. *Estuarine, Coastal and Shelf Science* 86(2): 157-164.
- Hofmann, G. E., Barry, J. P., Edmunds, P. J., Gates, R. D., Hutchins, D. A., Klinger, T. and Sewell, M. A. 2010. The effect of ocean acidification on calcifying organisms in marine ecosystems: an organism-to-ecosystem perspective. *Annual Review of Ecology, Evolution, and Systematics* 41: 127-147.
- Intergovernmental Panel on Climate Change 2001. The Scientific Basis: Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change, edited by J. T. Houghton, Y. Ding, D. J. Griggs, M. Noguer, P. J. van der Linden, X. Dai, K. Maskell and C. A. Johnson. Cambridge University Press, Cambridge, UK, New York, USA.
- Intergovernmental Panel on Climate Change 2007. Synthesis Report: Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, edited by R. K. Pachauri and A. Reisinger. Intergovernmental Panel on Climate Change, Geneva, Switzerland.
- Jokiel, P. L., Rodgers, K. S., Kuffner, I. B., Andersson, A. J., Cox, E. F. and Mackenzie, F. T. 2008. Ocean acidification and calcifying reef organisms: a mesocosm investigation. *Coral Reefs* 27(3): 473-483.
- Kikkawa, T., Ishimatsu, A. and Kita, J. 2003. Acute CO₂ tolerance during the early developmental stages of four marine teleosts. *Environmental toxicology* 18(6): 375-382.

- Kleypas, J. A., Buddemeier, R. W., Archer, D., Gattuso, J. P., Langdon, C. and Opdyke, B. N. 1999. Geochemical consequences of increased atmospheric carbon dioxide on coral reefs. *Science* 284(5411): 118-120.
- Kroeker, K. J., Kordas, R. L., Crim, R. N. and Singh, G. G. 2010. Meta-analysis reveals negative yet variable effects of ocean acidification on marine organisms. *Ecology Letters* 13(11): 1419-1434.
- Kurihara, H. 2008. Effects of CO₂-driven ocean acidification on the early developmental stages of invertebrates. *Marine Ecology Progress Series* 373: 275-284.
- Kurihara, H., Asai, T., Kato, S. and Ishimatsu, A. 2008a. Effects of elevated pCO₂ on early development in the mussel *Mytilus galloprovincialis*. *Aquatic Biology* 4(3): 225-233.
- Kurihara, H. and Ishimatsu, A. 2008. Effects of high CO₂ seawater on the copepod (*Acartia tsuensis*) through all life stages and subsequent generations. *Marine Pollution Bulletin* 56(6): 1086-1090.
- Kurihara, H., Kato, S. and Ishimatsu, A. 2007. Effects of increased seawater pCO₂ on early development of the oyster *Crassostrea gigas*. *Aquatic Biology* 1: 91-98.
- Kurihara, H., Matsui, M., Furukawa, H., Hayashi, M. and Ishimatsu, A. 2008b. Long-term effects of predicted future seawater CO₂ conditions on the survival and growth of the marine shrimp *Palaemon pacificus*. *Journal of Experimental Marine Biology and Ecology* 367(1): 41-46.
- Kurihara, H., Shimode, S. and Shirayama, Y. 2004a. Effects of raised CO₂ concentration on the egg production rate and early development of two marine copepods (*Acartia steueri* and *Acartia erythroa*). *Marine Pollution Bulletin* 49(9-10): 721-727.
- Kurihara, H., Shimode, S. and Shirayama, Y. 2004b. Sub-lethal effects of elevated concentration of CO₂ on planktonic copepods and sea urchins. *Journal of Oceanography* 60(4): 743-750.
- Kurihara, H. and Shirayama, Y. 2004. Effects of increased atmospheric CO₂ on sea urchin early development. *Marine Ecology Progress Series* 274: 161-169.
- Lacoue-Labarthe, T., Martin, S., Oberhänsli, F., Teyssié, J. L., Markich, S., Ross, J. and Bustamante, P. 2009. Effects of increased pCO₂ and temperature on trace element (Ag, Cd and Zn) bioaccumulation in the eggs of the common cuttlefish, *Sepia officinalis*. *Biogeosciences* 6(11): 2561-2573.
- Lischka, S., Büdenbender, J., Boxhammer, T. and Riebesell, U. 2010. Impact of ocean acidification and elevated temperatures on early juveniles of the polar shelled pteropod *Limacina helicina*: mortality, shell degradation, and shell growth. *Biogeosciences Discussions* 7(6): 8177-8214.
- Martin, S. and Gattuso, J. P. 2009. Response of Mediterranean coralline algae to ocean acidification and elevated temperature. *Global Change Biology* 15(8): 2089-2100.
- Martin, S., Richier, S., Pedrotti, M.-L., Dupont, S., Castejon, C., Gerakis, Y., Kerros, M.-E., Oberhänsli, F., Teyssié, J.-L., Jeffrey, R. and Gattuso, J.-P. 2011. Early development and molecular plasticity in the Mediterranean sea urchin *Paracentrotus lividus* exposed to CO₂-driven acidification. *Journal of Experimental Biology* 214(8): 1357-1368.
- Mayor, D. J., Matthews, C., Cook, K., Zuur, A. F. and Hay, S. 2007. CO₂-induced acidification affects hatching success in *Calanus finmarchicus*. *Marine Ecology Progress Series* 350: 91-97.
- McDonald, M. R., McClintock, J. B., Amsler, C. D., Rittschof, D., Angus, R. A., Orihuela, B. and Lutostanski, K. 2009. Effects of ocean acidification over the life history of the barnacle *Amphibalanus amphitrite*. *Marine Ecology Progress Series* 385: 179-187.
- Metzger, R., Sartoris, F. J., Langenbuch, M. and Pörtner, H. O. 2007. Influence of elevated CO₂ concentrations on thermal tolerance of the edible crab *Cancer pagurus*. *Journal of thermal biology* 32(3): 144-151.
- Miller, A. W., Reynolds, A. C., Sobrino, C. and Riedel, G. F. 2009. Shellfish face uncertain future in high CO₂ world: Influence of acidification on oyster larvae calcification and growth in estuaries. *PLoS ONE* 4(5).
- Morita, M., Suwa, R., Iguchi, A., Nakamura, M., Shimada, K., Sakai, K. and Suzuki, A. 2010. Ocean acidification reduces sperm flagellar motility in broadcast spawning reef invertebrates. *Zygote* 18(2): 103-107.
- Moulin, L., Catarino, A. I., Claessens, T. and Dubois, P. 2011. Effects of seawater acidification on early development of the intertidal sea urchin *Paracentrotus lividus* (Lamarck 1816). *Marine Pollution Bulletin* 62(1): 48-54.
- Munday, P. L., Crawley, N. E. and Nilsson, G. E. 2009a. Interacting effects of elevated temperature and ocean acidification on the aerobic performance of coral reef fishes. *Marine Ecology Progress Series* 388: 235-242.
- Munday, P. L., Dixon, D. L., Donelson, J. M., Jones, G. P., Pratchett, M. S., Devitsina, G. V. and Døving, K. B. 2009b. Ocean acidification impairs olfactory discrimination and homing ability of a marine fish. *Proceedings of the National Academy of Sciences of the United States of America* 106(6): 1848-1852.
- Munday, P. L., Donelson, J. M., Dixon, D. L. and Endo, G. G. K. 2009c. Effects of ocean acidification on the early life history of a tropical marine fish. *Proceedings of the Royal Society B: Biological Sciences* 276(1671): 3275-3283.
- O'Donnell, M. J., Hammond, L. M. and Hofmann, G. E. 2009. Predicted impact of ocean acidification on a marine invertebrate: elevated CO₂ alters response to thermal stress in sea urchin larvae. *Marine Biology* 156(3): 439-446.
- O'Donnell, M. J., Todgham, A. E., Sewell, M. A., Hammond, L. M., Ruggiero, K., Fanguie, N. A., Zippay, M. L. and Hofmann, G. E. 2009. Ocean acidification alters skeletogenesis and gene expression in larval sea urchins. *Marine Ecology Progress Series* 398: 157-171.
- Orr, J. C., Caldeira, K., Fabry, V., Gattuso, J. P., Haugan, P., Lehodey, P., Pantoja, S., Pörtner, H. O., Riebesell, U., Trull, T., Hood, M., Urban, E. and Broadgate, W. 2009. Research priorities for ocean acidification. Report from the second symposium on the ocean in a high-CO₂ world, Monaco, October 6-9, 2008, Convened by SCOR, IOC-UNESCO, IAEA, and IGBP.
- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R. G., Plattner, G. K., Rodgers, K. B., Sabine, C. L., Sarmiento, J. L., Schlitzer, R., Slater, R. D., Totterdell, I. J., Weirig, M. E., Yamanaka, Y. and Yool, A. 2005. Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature* 437(7059): 681-686.
- Pagano, G., Cipollaro, M. and Corsale, G. 1985a. pH-Induced changes in mitotic and developmental patterns in sea urchin embryogenesis. I. Exposure of embryos. *Teratogenesis Carcinogenesis and Mutagenesis* 5(2): 101-112.
- Pagano, G., Cipollaro, M. and Corsale, G. 1985b. pH-Induced changes in mitotic and developmental patterns in sea urchin embryogenesis. II. Exposure of sperm. *Teratogenesis Carcinogenesis and Mutagenesis* 5(2): 113-121.

- Parker, L. M., Ross, P. M. and O'Connor, W. A. 2009. The effect of ocean acidification and temperature on the fertilization and embryonic development of the Sydney rock oyster *Saccostrea glomerata* (Gould 1850). *Global Change Biology* 15(9): 2123-2136.
- Parker, L. M., Ross, P. M. and O'Connor, W. A. 2010. Comparing the effect of elevated pCO₂ and temperature on the fertilization and early development of two species of oysters. *Marine Biology* 157(11): 2435-2452.
- Parker, L. M., Ross, P. M. and O'Connor, W. A. 2011. Populations of the Sydney rock oyster, *Saccostrea glomerata*, vary in response to ocean acidification. *Marine Biology* 158(3): 689-697.
- Parker, L. M., Ross, P. M., O'Connor, W. A., Borysko, L., Raftos, D. A. and Portner, H. O. 2012. Adult exposure influences offspring response to ocean acidification in oysters. *Global Change Biology*. 18(1): 82-92.
- Pörtner, H. O. 2008. Ecosystem effects of ocean acidification in times of ocean warming: a physiologist's view. *Marine Ecology Progress Series* 373: 203-217.
- Pörtner, H. O. and Farrell, A. P. 2008. Ecology: physiology and climate change. *Science* 322(5902): 690-692.
- Pörtner, H. O., Langenbuch, M. and Reipschläger, A. 2004. Biological impact of elevated ocean CO₂ concentrations: lessons from animal physiology and earth history. *Journal of Oceanography* 60(4): 705-718.
- Przeslawski, R., Ah Yong, S., Byrne, M., Wörheide, G. and Hutchings, P. 2008. Beyond corals and fish: the effects of climate change on noncoral benthic invertebrates of tropical reefs. *Global Change Biology* 14(12): 2773-2795.
- Przeslawski, R. and Webb, A. R. 2009. Natural variation in larval size and developmental rate of the northern quahog *Mercenaria mercenaria* and associated effects on larval and juvenile fitness. *Journal of Shellfish Research* 28(3): 505-510.
- Range, P., Chicharo, M. A., Ben-Hamadou, R., Piló, D., Matias, D., Joaquim, S., Oliveira, A. P. and Chicharo, L. 2011. Calcification, growth and mortality of juvenile clams *Ruditapes decussatus* under increased pCO₂ and reduced pH: variable responses to ocean acidification at local scales? *Journal of Experimental Marine Biology and Ecology* 396(2): 177-184.
- Reuter, K. E., Lotterhos, K. E., Crim, R. N., Thompson, C. A. and Harley, C. D. G. 2011. Elevated pCO₂ increases sperm limitation and risk of polyspermy in the red sea urchin *Strongylocentrotus franciscanus*. *Global Change Biology* 17(1): 163-171.
- Reynaud, S., Leclercq, N., Romaine-Lioud, S., Ferrier-Pagès, C., Jaubert, J. and Gattuso, J. P. 2003. Interacting effects of CO₂ partial pressure and temperature on photosynthesis and calcification in a scleractinian coral. *Global Change Biology* 9(11): 1660-1668.
- Royal Society 2005. Ocean acidification due to increasing atmospheric carbon dioxide. Policy Document 12/05, The Royal Society, London. Pp 1-58.
- Sheppard Brennd, H., Soars, N., Dworjanyn, S. A., Davis, A. R. and Byrne, M. 2010. Impact of ocean warming and ocean acidification on larval development and calcification in the sea urchin *Tripneustes gratilla*. *PLoS ONE* 5(6): e11372.
- Strathmann, M. F. 1987. Opisthobranchia Pp. 268-302 in *Reproduction and Development of Marine Invertebrates of the Northern Pacific Coast*, edited by M. F. Strathmann. University of Washington Press, Seattle, WA.
- Suwa, R., Nakamura, M., Morita, M., Shimada, K., Iguchi, A., Sakai, K. and Suzuki, A. 2010. Effects of acidified seawater on early life stages of scleractinian corals (Genus *Acropora*). *Fisheries Science* 76(1): 93-99.
- Talmage, S. C. and Gobler, C. J. 2010. Effects of past, present, and future ocean carbon dioxide concentrations on the growth and survival of larval shellfish. *Proceedings of the National Academy of Sciences of the United States of America* 107(40): 17246-17251.
- Thorson, G. 1950. Reproduction and larval ecology of marine bottom invertebrates. *Biology Reviews* 25: 1-45.
- Todgham, A. E. and Hofmann, G. E. 2009. Transcriptomic response of sea urchin larvae *Strongylocentrotus purpuratus* to CO₂-driven seawater acidification. *Journal of Experimental Biology* 212(16): 2579-2594.
- Turley, C., Blackford, J. C., Widdicombe, S., Lowe, D., Nightingale, P. D. and Rees, A. P. 2006. Reviewing the impact of increased atmospheric CO₂ on oceanic pH and the marine ecosystem Pp. 65-70 in *Avoiding Dangerous Climate Change*, edited by H. J. Schellnhuber, W. Cramer, N. Nakicenovic, T. Wigley and G. Yohe. Cambridge University Press, Cambridge, UK.
- Underwood, A. J. 1997. *Experiments in Ecology: Their Logical Design and Interpretation Using Analysis of Variance*. Cambridge University Press, Cambridge.
- Waldbusser, G. G., Bergschneider, H. and Green, M. A. 2010. Size-dependent pH effect on calcification in post-larval hard clam *Mercenaria* spp. *Marine Ecology Progress Series* 417: 171-182.
- Waller, T. R. 1981. Functional morphology and development of veliger larvae of the European oyster, *Ostrea edulis* Linné. *Smithsonian Contributions to Zoology*: 328: 1-70.
- Walther, K., Anger, K. and Pörtner, H. O. 2010. Effects of ocean acidification and warming on the larval development of the spider crab *Hyas araneus* from different latitudes (54° vs. 79° N). *Marine Ecology Progress Series* 417: 159-170.
- Watson, S. A., Southgate, P. C., Tyler, P. A. and Peck, L. S. 2009. Early larval development of the Sydney rock oyster *Saccostrea glomerata* under near-future predictions of CO₂-driven ocean acidification. *Journal of Shellfish Research* 28(3): 431-437.
- Wilson, S. P. and Hyne, R. V. 1997. Toxicity of acid-sulfate soil leachate and aluminum to embryos of the Sydney rock oyster. *Ecotoxicology and Environmental Safety* 37(1): 30-36.
- Yu, P. C., Matson, P. G., Martz, T. R. and Hofmann, G. E. 2011. The ocean acidification seascape and its relationship to the performance of calcifying marine invertebrates: laboratory experiments on the development of urchin larvae framed by environmentally-relevant pCO₂/pH. *Journal of Experimental Marine Biology and Ecology* 400(1-2): 288-295.
- Zippay, M. L. and Hofmann, G. E. 2010. Effect of pH on gene expression and thermal tolerance of early life history stages of red abalone (*Haliotis rufescens*). *Journal of Shellfish Research* 29(2): 429-439.

Table 1. The effects of ocean acidification on the reproduction and early development of marine biota. n/k : not known; s: seconds; d: days; w: weeks; mo: month; *: effect only when in synergy with salinity; -:negative impact; =: no significant difference; +: positive impact

Organism or Species	Stage tested	CO ₂ system parameters		Manipulation method	Exposure length	Impact			Reference	
		pCO ₂ (ppm)	pH			CO ₂ /Mineral acid	Fertilization	Larval Development		Settlement/ Recruitment
Mollusca										
<i>Saccostrea glomerata</i> (Sydney rock oyster)	Egg, larva	600	7.9	Acid	48 h	-	-	-	Parker et al., (2009)	Reduced fertilisation, larval development & growth; increased abnormality & mortality
<i>Saccostrea glomerata</i> (Sydney rock oyster)	Spat	1000	7.84	CO ₂	4 d	=	=	=	Parker et al., (2011)	Reduced growth varied between selectively bred (~25%) & wild caught (~64%)
<i>Saccostrea glomerata</i> (Sydney rock oyster)	Larva	508-775	7.77-7.8	CO ₂	8 d	-	-	-	Watson et al., (2009)	Reduced growth; larval development; increased mortality & shell abnormality
<i>Saccostrea glomerata</i> (Sydney rock oyster)	Larva	n/k	6.75	Acid H ₂ SO ₄	48 h	-	-	-	Wilson & Hyne (1997)	pH ≤ 6.75 reduced development and increased abnormality of D-veliger larvae
<i>Saccostrea glomerata</i> (Sydney rock oyster)	Egg, larva	600	7.9	CO ₂	48 h	-	-	-	Parker et al., (2010)	Reduced fertilisation, larval development & growth; increased abnormality
<i>Crassostrea gigas</i> (Pacific oyster)	Egg, larva	600	7.9	CO ₂	48 h	-	-	-	Parker et al., (2011)	Reduced fertilisation, larval development & growth; increased abnormality
<i>Crassostrea gigas</i> (Pacific oyster)	Egg, larva	2268	7.4	CO ₂	48 h	=	=	=	Kurihara et al., (2007)	No effect on fertilisation; inhibition of shell synthesis; reduced larval size; increased abnormality
<i>Crassostrea gigas</i> (Pacific oyster)	Egg, sperm	1000 (n/k)	7.8	CO ₂	1 h	=	=	=	Havenhand & Schlegel (2009)	No effect on fertilisation, sperm speed or motility
<i>Crassostrea virginica</i> (Eastern oyster)	Larva	n/k	6.25	Acid	12 d	-	-	-	Calabrese & Davis (1966)	pH < 6.25 increased mortality rate pH < 6.75 decreased growth rate
<i>Crassostrea virginica</i> (Eastern oyster)	Larva	280, 380, 560, 800	8.16 8.06 7.91 7.76	CO ₂	30 d	-	-	-	Miller et al., (2009)	<560 pCO ₂ Growth significantly decreased. 800 pCO ₂ calcification reduced.
<i>Crassostrea ariakensis</i> (Surinooe oyster)	Larva	280, 380, 560, 800	8.17 8.08 7.92 7.79	CO ₂	32 d	=	=	=	Miller et al., (2009)	No effect on growth or calcification
<i>Mytilus galloprovincialis</i> (Mussel)	Egg, larva	2000	7.4	CO ₂	6 d	-	-	-	Kurihara et al., (2008a)	Development delayed at trochophore stage following onset of shell development; increased abnormality of larvae; 26% & 20% reduction in larval shell height and length, respectively after 6 d

Organism or Species	Stage tested	CO ₂ system parameters		Manipulation method	Exposure length	Impact			Reference
		pCO ₂ (ppm)	pH			Fertilization	Larval Development	Settlement/ Recruitment	
<i>Mytilus edulis</i> (Blue mussel)	Larva	1124-1929	7.8, 7.6	CO ₂	15 d	-	-	-	Gazeau et al., (2010)
<i>Tivela stultorum</i> (Pismo clam)	Egg	n/k	8.0	Acid???	n/k	-	-	-	Alvarado-Alvarez et al., (1996)
<i>Mercenaria mercenaria</i> (Hard clam)	Larva	n/k	6.25	Acid	10 d	-	-	-	Calabrese & Davis (1966)
<i>Littorina obtusata</i> (Intertidal gastropod)	Egg, larva	1100	7.6	CO ₂	24 d	-	-	-	Ellis et al., (2009)
<i>Cavolinia inflexa</i> (Opisthobranch gastropod/ pteropod)	Larva	857, 1713	7.82, 7.51	n/k	13 d	-	-	-	Comeau et al., (2010)
<i>Haliotis rufescens</i> (Abalone)	Larva	570, 990	7.97, 7.87	CO ₂	6 d	-	-	-	Zippay et al., (2010)
<i>Haliotis coccoradiata</i> (Abalone)	Larva	814-1828	7.9-7.6	CO ₂	2 h	=	=	=	Byrne et al., (2010a)
<i>Haliotis coccoradiata</i> (Abalone)	Larva, Juvenile	700, 2000	7.8, 7.6	CO ₂	4 d	-	-	-	Byrne et al., (2011a)
<i>Haliotis kamtschatkana</i> (Abalone)	Larva	800, 1800	8.07, 7.81	CO ₂	8 d	-	=	=	Crim et al., (2011)
<i>Sepia officinalis</i> (European cuttlefish)	Juvenile	4000, 6000	7.23, 7.1	CO ₂	6 w	+	-	-	Gutowska et al., (2008)
<i>Sepia officinalis</i> (Common cuttlefish)	Eggs	900, 1400	7.85, 7.6	CO ₂	65 d	+	-	-	Lacoue-Labarthe et al., (2009)

Organism or Species	Stage tested	CO ₂ system parameters		Manipulation method	Exposure length	Impact			Reference
		pCO ₂ (ppm)	pH			Fertilization	Larval Development	Settlement/ Recruitment	
<i>Mercenaria mercenaria</i> (Hard clam)	Larva	250	8.17	CO ₂ /Mineral acid	36 d	-	-	-	Survival, growth, rate of metamorphosis, size & shell thickness all decreased significantly with increasing CO ₂ , hinge integrity was also impacted. Talmage & Gobler (2010)
		390	8.05						
		750	7.8						
<i>Argopecten irradians</i> (Bay scallop)	Larva	1500	7.5	CO ₂	38 d	-	-	-	Survival, growth, rate of metamorphosis, size & shell thickness all decreased significantly with increasing CO ₂ , hinge integrity was also impacted. Talmage & Gobler (2010)
		250	8.17						
		390	8.05						
<i>Ruditapes decussates</i> (Grooved carpet clam)	Juvenile	1698	7.84	CO ₂	75 d	=	=	=	No effect on calcification rates, size or weight. Reduced mortality possibly due to absence of spawning. Range et al., (2011)
		4345	7.46						
<i>Mercenaria</i> spp. (Hard clam)	Post larval	1120	7.64	CO ₂	n/k	-	-	-	Calcification rate decreased with pH & organism size (size dependency of pH effects). Waldbusser et al., (2010)
<i>Limacina helicina</i> (Polar pteropod)	Pre-winter juveniles	230	8.21	CO ₂	29 d	-	-	-	Increased pCO ₂ (not temperature) reduced shell diameter, shell increment & increased shell degradation. pCO ₂ increased mortality, though temperature overriding factor. Lischka et al., (2010)
		350	8.06						
		750	7.78						
Echinodermata	Egg larva	1100	7.63	Both HCl & CO ₂	3 d	-	-	-	pH < 7.8 skeletal malformation, reduced larval size, fertilisation decreased with increasing pCO ₂ - significant at pH 6.8 (33-50%) Kurihara & Shirayama (2004), Kurihara et al., 2004b
		865-10365	7.8-6.8						
<i>Echinometra mathaeii</i> (Sea urchin)	Egg larva	865-10365	7.8-6.8	Both HCl & CO ₂	3 d	-	-	-	pH < 7.8 skeletal malformation, reduced larval size, Fertilisation decreased with increasing pCO ₂ - significant at pH 7.1. Shirayama (2004), Kurihara et al., 2004b
<i>Paracentrotus lividus</i> (Sea urchin)	Egg larva	n/k	7.5	Acid	48 d	-	-	-	pH < 7.5 skeletal malformation, pH < 7.7 morphological abnormality, Mitotic abnormality with decreasing pH Pagano et al., (1985a,b)
<i>Paracentrotus lividus</i> (Sea urchin)	Egg larva	704-6632	7.9-7.0	CO ₂	3 d	=	=	=	Development delayed to pluteus stage. <pH 7.25 morphology affected. < pH 7.5 overall length, post oral and antero-lateral arm length shorter. <pH7.9 rods shorter. Protein & biomineralisation genes upregulated at low pH. Martin et al., (2011)

Organism or Species	Stage tested	CO ₂ system parameters		Manipulation method	Exposure length	Impact			Reference
		pCO ₂ (ppm)	pH			CO ₂ /Mineral acid	Fertilization	Larval Development	
<i>Helicidaris erythrogramma</i> (Sea urchin)	Larva, juvenile	700	7.8	CO ₂	5 d		-		Byrne et al., (2011a)
<i>Helicidaris tuberculata</i> (Sea urchin)	Egg, larva	814-1828	7.9-7.6	CO ₂	2 h	=			Byrne et al., (2010a)
<i>Tripneustes gratilla</i> (Sea urchin)	Egg, larva	814-1828	7.9-7.6	CO ₂	2 h	=			Byrne et al., (2010a)
<i>Centrostephanus rogersii</i> (Sea urchin)	Egg, larva	1051-1828	7.8-7.6	CO ₂	2 h	=			Byrne et al., (2010a)
<i>Patriella regularis</i> (Sea star)	Egg, larva	1051-1828	7.8-7.6	CO ₂	2 h	=			Byrne et al., (2010a)
<i>Grossaster papposus</i> (Sea star)	Larva	930	7.7	CO ₂	38 d		+		Dupont et al., (2010b)
<i>Ophiothrix fragilis</i> (Brittlestar)	Larva	n/k	7.9, 7.7	CO ₂	8 d		-		Dupont et al., (2008)
<i>Evechinus chloroticus</i> (Sea urchin)	Larva	1320	7.7	CO ₂	13 d		-		Clark et al., (2009)
<i>Pseudechinus huttoni</i> (Sea urchin)	Larva	1282	7.7	CO ₂	9 d		-		Clark et al., (2009)
<i>Sterechinus neumayeri</i> (Sea urchin)	Larva	1380	7.6	CO ₂	7-17 d		-		Clark et al., (2009)
<i>Tripneustes gratilla</i> (Sea urchin)	Larva	1119	7.8	CO ₂	4 d		-		Clark et al., (2009)
<i>Tripneustes gratilla</i> (Sea urchin)	Larva	n/k	7.8, 7.6	n/k	5 d		-		Sheppard Brennan et al., (2010)
<i>Paracentrotus lividus</i> (Intertidal temperate sea urchin)	Egg, larva	310 8335	7.8-6.8	CO ₂	72 h	-	-		Moulin et al., (2010)

Organism or Species	Stage tested	CO ₂ system parameters		Manipulation method	Exposure length	Impact			Reference	
		pCO ₂ (ppm)	pH			Fertilization	Larval Development	Settlement/Recruitment		Comment
<i>Sterechinus neumayeri</i> (Sea urchin)	Egg, larva	1121, 2886, 5806	7.7, 7.3, 7.0	CO ₂	2h -6 d	=	-	-	Ericson et al., (2010)	
Arthropoda										
<i>Acartia erythraea</i> (Copepod)	Egg, larva	2360-10360	7.31-6.82	CO ₂	2 d	-	-	-	Kurihara et al., (2004a)	
<i>Acartia tsuensis</i> (Copepod)	Egg, larva	2380	7.31	CO ₂	9 d	=	=	=	Kurihara & Shimatsu (2008)	
<i>Calanus finmarchicus</i> (Copepod)	Egg	8000	6.95	CO ₂	72 h	-	-	-	Mayor et al., (2007)	
<i>Palaemon pacificus</i> (Shrimp)	Egg Juvenile	1000, 1900	7.89, 7.64	CO ₂	30 w, 15 w	-	-	-	Kurihara et al., (2008b)	
<i>Homarus gammarus</i> (European lobster)	Larva	1200	8.1	CO ₂	28 d	=	=	=	Arnold et al., (2009)	
<i>Echinogammarus marinus</i> Leach (Amphipod)	Egg	1900	7.5	CO ₂	Approx. 18 d	-*	-	-	Egisdottir et al., (2009)	
<i>Gammarus locusta</i> (Amphipod)	Juvenile	550, 980	7.8, 7.6	CO ₂	14-28 d	-	-	-	Hauton et al., (2009)	
<i>Hyas araneus</i> (Spider crab)	Larva	710 3000	n/k	CO ₂	n/k	-	-	-	Waither et al., (2010)	
Barnacles										
<i>Amphibalanus amphitrite</i> (Barnacle)	Larva; cyprid	n/k	7.4	CO ₂	5d	=	=	=	McDonald et al., (2009)	

Organism or Species	Stage tested	CO ₂ system parameters		Manipulation method	Exposure length	Impact			Reference	
		pCO ₂ (ppm)	pH			Fertilization	Larval Development	Settlement/ Recruitment		Comment
<i>Semibalanus balanoides</i> (Barnacle)	Egg	922	7.7	CO ₂	104 d	=			Reduced rate of embryonic development (19 day delay), but within natural rates	Findlay et al., (2009)
<i>Semibalanus balanoides</i> (Barnacle)	Post-larval	1000	7.7	CO ₂	30 d	=			No effect on growth rate, but reduced calcium content & survival at elevated temperature & CO ₂	Findlay et al., (2010a)
<i>Elminius modestus</i> (Barnacle)	Post-larval	1000	7.7	CO ₂	30 d	=			No effect on calcium content or survival; decrease in growth rate at elevated CO ₂	Findlay et al., (2010a)
<i>Semibalanus balanoides</i> (Barnacle)	Post-larval	1000 3000	7.7 7.3	CO ₂	20 d	=			pH 7.7 reduced growth & development; no effect on shell mineralisation	Findlay et al., (2010b)
Vertebrata										
<i>Pagrus major</i> (Silver seabream)	Egg, larva, juvenile	10000	6.2	CO ₂	24 h	-			Hatching and survival decreased with increased pCO ₂ & exposure; stage specific sensitivity	Kikkawa et al., (2003)
<i>Sillago japonica</i> (Japanese sillago)	Egg, larva, juvenile	10000	6.2	CO ₂	24 h	-			Hatching and survival decreased with increased pCO ₂ & exposure; stage specific sensitivity	Kikkawa et al., (2003)
<i>Amphiprion percula</i> Pomacentridae (Clown fish)	Egg, larva	1030	7.84	CO ₂	11 d	=			No effect on embryonic development, egg survival and hatching size; some parental pairs increased length up to 18% & weight by 52%	Munday et al., (2009b)
<i>Amphiprion percula</i> Pomacentridae (Clown fish)	Larva	1050, 1710	7.8, 7.6	CO ₂	11 d	-			pH 7.8 disruption, pH 7.6 total Loss of larval discriminatory ability for olfactory settlement cues at elevated pCO ₂	Munday et al., (2009a)
<i>Gadhus morhua</i> (Baltic cod)	Sperm	1400	7.55	CO ₂	n/k	=			No effect on sperm speed, rate of change of direction or percent motility.	Frommel et al., (2010)
Corals & communities										
<i>Pocillopora damicornis</i> (Coral)	Larva	745	n/k c.7.6-7.9	Acid 10% HCl	10 mo	=			No effect on recruitment (effects on adults)	Jokiel et al., (2008)
<i>Montipora capitata</i> (Coral)	Egg, sperm	745	n/k c.7.6-7.9	Acid 10% HCl	6 mo	=			No effect on gametes (effects on adults)	Jokiel et al., (2008)
<i>Acropora digitifera</i> (Coral)	Larva, Primary polyp	905-3585	7.31-7.64	CO ₂	10 d	=			pH 7.3 reduced primary polyp occupation area 14% compared to control; larvae mortality not significantly affected by pH	Suwa et al., (2010)
<i>Acropora tenuis</i> (Coral)	Larva	905-3585	7.31-7.64	CO ₂	7d	=			Mortality inconsistent- lower at pH 7.3 than in control & significantly lower at 7.3 (15.2%) than at 7.6 (38%)	Suwa et al., (2010)

Organism or Species	Stage tested	CO ₂ system parameters		Manipulation method	Exposure length	Impact			Reference
		pCO ₂ (ppm)	pH			CO ₂ /Mineral acid	Fertilization	Larval Development	
<i>Porites panamensis</i>	Larva Primary polyp	861-	7.81-	CO ₂	10-42 d	=	=	-	Anlauf et al., 2010
		1006	7.85						
<i>Cladocora caespitosa</i> (Temperate coral)	Polyp	700	7.86	CO ₂	1-10 mo	=			Rodolfo-Metalpa et al., (2010)
<i>Astrangia porculata</i> (Temperate coral)	Polyp	780	7.78	CO ₂	6 mo	-			Holcomb et al., (2010)
<i>Porites</i> sp (Long-lived temperate coral)	Polyp	1904	7.49	CO ₂	6-14 mo	-	-	-	Krief et al., (2010)
		3970	7.19						
<i>Stylophora pistillata</i> (Short lived temperate coral)	Polyp	1904	7.49	CO ₂	6-14 mo	-	-	-	Krief et al., (2010)
		3970	7.19						
<i>Acropora digitifera</i> (Coral)	Sperm	775-	7.77-	CO ₂	2 s	-	-	-	Morita et al., (2009)
		21100	6.55						