

The molecular signature of mediastinal large B-cell lymphoma differs from that of other diffuse large B-cell lymphomas and shares features with classical Hodgkin lymphoma

Kerry J. Savage, Stefano Monti, Jeffery L. Kutok, Giorgio Cattoretti, Donna Neuberg, Laurence de Leval, Paul Kurtin, Paola Dal Cin, Christine Ladd, Friedrich Feuerhake, Ricardo C. T. Aguiar, Sigui Li, Gilles Salles, Francoise Berger, Wen Jing, Geraldine S. Pinkus, Thomas Habermann, Riccardo Dalla-Favera, Nancy Lee Harris, Jon C. Aster, Todd R. Golub, and Margaret A. Shipp

Mediastinal large B-cell lymphoma (MLBCL) is a recently identified subtype of diffuse large B-cell lymphoma (DLBCL) that characteristically presents as localized tumors in young female patients. Although MLBCL has distinctive pathologic features, it clinically resembles the nodular sclerosis subtype of classical Hodgkin lymphoma (cHL). To elucidate the molecular features of MLBCL, we compared the gene expression profiles of newly diagnosed MLBCL and DLBCL and developed a classifier of these diseases. MLBCLs

had low levels of expression of multiple components of the B-cell receptor signaling cascade, a profile resembling that of Reed-Sternberg cells of cHL. Like cHLs, MLBCLs also had high levels of expression of the interleukin-13 (IL-13) receptor and downstream effectors of IL-13 signaling (Janus kinase-2 [JAK2] and signal transducer and activator of transcription-1 [STAT1]), tumor necrosis factor (TNF) family members, and TNF receptor-associated factor-1 (TRAF1). Increased expression of STAT1 and TRAF1 in ML-

BCL was confirmed by immunohistochemistry. Given the TRAF1 expression and known link to nuclear factor- κ B (NF- κ B), MLBCLs were also evaluated for nuclear translocation of c-REL protein. In almost all cases, c-REL was localized to the nucleus, consistent with activation of the NF- κ B pathway. These studies identify a molecular link between MLBCL and cHL and a shared survival pathway. (Blood. 2003;102:3871-3879)

© 2003 by The American Society of Hematology

Introduction

Diffuse large B-cell lymphomas (DLBCLs) are heterogeneous tumors with variable clinical presentations and responses to therapy.¹ At diagnosis, these aggressive B-cell malignancies involve a wide variety of nodal and extranodal disease sites; however, it is unclear whether tumors presenting in specific anatomic locations exhibit unique molecular features. One likely exception is the recently identified entity, primary mediastinal (thymic) large B-cell lymphoma (MLBCL), a subtype of DLBCL defined by a combination of clinical and pathologic features.^{2,3}

Unlike DLBCL, which commonly arises in elderly patients of both sexes, MLBCL typically presents in younger women. These patients have bulky mediastinal masses with frequent invasion of adjacent structures.² Although patients with MLBCL rarely have extrathoracic disease at diagnosis, extranodal sites including the central nervous system (CNS), liver, adrenals, and kidneys are often involved at relapse.⁴ MLBCLs typically consist of tumor cells with a pale cytoplasm and a diffuse growth pattern associated with variable degrees of sclerosis.^{3,5} However, there are no histologic features that reliably distinguish these tumors from DLBCLs that often involve mediastinal regional lymph nodes.

Although MLBCLs have rearranged immunoglobulin (Ig) genes and express B-cell lineage markers such as CD19, CD20, CD22, and CD79a (Ig α), these tumors do not express Ig.² In fact, the discordant expression of specific components of the B-cell receptor (BCR) (Ig $^-$ and CD79a $^+$) is a unique feature of MLBCLs.⁶ The anatomic location, morphologic appearance, and immunophenotypic signature of MLBCLs suggest that these tumors arise from thymic medullary B cells.^{2,7} MLBCLs and a subpopulation of normal thymic medullary B cells express MAL, an integral membrane protein and lipid raft component that is rarely found in DLBCL.⁸

MLBCLs have mutated, class-switched Ig genes without evidence of ongoing somatic mutation.⁹ These tumors also exhibit several characteristic genetic abnormalities, including gains of chromosomes 9p and 2p, and the associated *JAK2* (9p24) and *REL* (2p16) loci.^{10,11} In contrast to DLBCLs, MLBCLs do not exhibit *BCL2* rearrangements and rarely manifest translocations involving *BCL6*.¹²

Because MLBCL is currently considered a subtype of DLBCL (as opposed to a distinct disease), these tumors are treated similarly

From the Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA; Whitehead Institute/Massachusetts Institute of Technology (MIT) Center for Genome Research, Cambridge, MA; Department of Pathology, Brigham and Women's Hospital, Boston, MA; Institute for Cancer Genetics, Columbia University, New York, NY; Department of Biostatistical Science, Dana-Farber Cancer Institute, Boston, MA; Department of Pathology, Massachusetts General Hospital, Boston, MA; Department of Pathology, Mayo Clinic, Rochester, MN; Hematology Department, Centre Hospitalier Lyon-Sud, Lyon, France; Department of Pathology, Centre Hospitalier Lyon-Sud, Lyon, France; Division of Hematology and Department of Medicine, Mayo Clinic, Rochester, MN; Department of Pediatric Oncology, Dana-Farber Cancer Institute and Howard Hughes Medical Institute, Boston, MA.

Submitted June 9, 2003; accepted August 17, 2003. Prepublished online as

Blood First Edition Paper, August 21, 2003; DOI 10.1182/blood-2003-06-1841.

K.J.S. and S.M. contributed equally to this study. T.R.G. and M.A.S. contributed equally to this study.

The online version of the article contains a data supplement.

Reprints: Margaret A. Shipp, Dana-Farber Cancer Institute, 44 Binney St, Boston, MA 02115; e-mail: margaret_shipp@dfci.harvard.edu.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 U.S.C. section 1734.

© 2003 by The American Society of Hematology

with empiric adriamycin-containing combination chemotherapy regimens. Involved-field radiation therapy is often added to the site of bulky, localized disease.¹³ However, combined modality therapy is only partially effective, because more than 40% of MLBCL patients die of their disease.¹³ Further, involved-field radiation therapy has been associated with long-term side effects including secondary malignancies and cardiac dysfunction.¹⁴ For these reasons, additional insights into the molecular signature of MLBCL and potential rational treatment targets are critically needed.

To delineate unique features of MLBCL compared with DLBCL, we performed gene expression profiling of these 2 tumors and found them to be markedly different. Importantly, we also identified a molecular link between MLBCL and another lymphoid neoplasm with shared clinical features, classical Hodgkin lymphoma (cHL).

Patients, materials, and methods

Case selection and histologic classification

Frozen tumor specimens from newly diagnosed, previously untreated MLBCL patients (34) and DLBCL patients (176) were analyzed according to protocol approved by the institutional review board of the Dana Farber/Harvard Cancer Center. MLBCL tumor specimens were derived from mediastinal masses or contiguous nodal biopsies, and DLBCLs were all nodal tumor specimens. Primary MLBCLs were identified using clinical criteria (predominant mediastinal mass with or without local extension and no extrathoracic disease) and pathologic features. The histopathology and immunophenotype of each primary MBLCL were reviewed by expert hematopathologists to confirm the diagnosis. The diagnosis of each DLBCL was originally made by expert hematopathologists at the participating institutions.

Target cRNAs of oligonucleotide microarrays

Total RNA was extracted from each frozen tumor specimen, and biotinylated cRNAs were generated as previously described.^{15,16} Samples were hybridized overnight to Affymetrix U133A and U133B oligonucleotide microarrays (Affymetrix, Santa Clara, CA), which include probe sets from more than 44 000 genes. Arrays were subsequently developed with phycoerythrin-conjugated streptavidin (SAPE) and biotinylated antibody against streptavidin and scanned to obtain quantitative gene expression levels.¹⁵ The raw gene expression values were then scaled to account for differences in global chip intensity.

Gene expression analysis

The top 15 000 genes from the U133A and U133B Affymetrix chips were selected based on their ranking as measured by a median absolute deviation (MAD) variation filter across all samples. From within this 15 000-gene pool, genes correlating with the class distinction of interest (mediastinal = 1 versus nonmediastinal = 2) were identified by ranking them according to their signal-to-noise ratio (SNR). For a given gene g , $SNR(g) = (\bar{x}_1 - \bar{x}_2) / (s_1 + s_2)$, where \bar{x}_i and s_i denote, respectively, gene g 's sample mean and sample standard deviation within class $i = 1, 2$. Similar rankings were obtained by using the median in place of the mean or by using the t statistic. A Monte Carlo simulation of the permutation distribution of the SNRs was performed by permuting the sample labels indicating class membership ($n = 1000$); thereafter, the observed values in the data were compared with the 99th percentile of the permutation distribution (Supplemental Material: see the Supplemental Document link at the top of the online article on the *Blood* website).

MLBCL versus DLBCL classifier

The discriminatory power of the gene expression signature was evaluated by building classifiers for the MLBCL versus DLBCL distinction. Naive Bayes (NB) and weighted voting (WV) classifiers that included 10 to 1000

genes were built and evaluated with regard to prediction errors using leave-one-out cross-validation (LOO-CV) (see Supplemental Document). The classifier with the lowest balanced error rate (proportion of samples wrongly classified, averaged within classes) was chosen for further analysis (see Supplemental Document). A more rigorous estimate of the error rate was also computed based on a 2-level CV procedure in which the choice of how many genes to use is automatically made within the CV loop (see Supplemental Document).

Analysis of coregulated genes

Genes that were coregulated (ie, genes whose expression values follow a similar pattern) with the top-ranked genes in MLBCL were identified using a 1 minus the Pearson correlation coefficient as the distance metric (see Supplemental Document).

Enrichment test for cHL genes in MLBCL

An enrichment test was used to evaluate the significance of the observed similarity between the MLBCL and cHL signatures. The cHL signature was defined using a set of genes independently identified by Schwering et al as differentially expressed in Hodgkin Reed-Sternberg (HRS) cell lines and normal purified B cells (centroblasts, centrocytes, and naive and memory B cells) using Affymetrix U95 oligonucleotide arrays.¹⁷ The similarity between the MLBCL and HRS signatures was assessed using the following recently described procedure.¹⁸ A total of 15 000 genes were selected from the U133A/B chips according to a MAD-based variation filter and ranked according to their SNR with respect to the "MLBCL versus DLBCL" class membership. The 294 genes with reduced expression in HRS cell lines¹⁷ were then located within our ranked list of 15 000 genes, and their proximity to the genes with lower levels of expression in MLBCL was measured by a Kolmogorov-Smirnoff (KS) score (with a higher score corresponding to a higher proximity). A similar procedure was used to place the 195 genes with increased expression in HRS cell lines^{17,19} within the 15 000 genes ranked in the opposite direction. Thereafter, permutation of the "MLBCL versus DLBCL" sample labels, associated reranking of the 15 000 genes, and computation of the corresponding KS scores were performed multiple times ($n = 1000$) so as to compare the observed KS scores with those that could be expected by chance under a random class labeling. Empirical P values were then computed to quantify the significance of the observed similarities between the genes with increased and decreased expression in the HRS cell lines and MLBCLs (p_{max}). An alternative, less stringent, method was also used to compute an empirical P value (p_{min}). This method was based on the computation of the KS score for multiple ($n = 1000$) random sets of 294 or 195 genes. This approach determines how likely it would be to obtain the observed KS score if a random set of genes were selected (see Supplemental Document).

Immunohistochemistry

Immunohistochemistry with anti-signal transducer and activator of transcription-1 (anti-STAT1) (9H2, Cell Signaling Technology, Beverly, MA) or anti-tumor necrosis factor (TNF) receptor-associated factor-1 (anti-TRAF1) (H3; Santa Cruz Biotechnology, Santa Cruz, CA) murine monoclonal antibodies was performed using 3 to 5 μ -thick formalin-fixed, paraffin-embedded specimens by standard immunohistochemical methods. Slides were deparaffinized and pretreated with 10 mM citrate, pH 6.0, (Zymed, South San Francisco, CA) in a steam pressure cooker (Decloaking Chamber, BioCare Medical, Walnut Creek, CA). Slides were then treated with Peroxidase Block (DAKO USA, Carpinteria, CA) for 5 minutes to quench endogenous peroxidase activity and incubated with a 1:5 dilution of goat serum in 50 mM Tris (tris(hydroxymethyl)aminomethane)-HCl, pH 7.4, for 20 minutes to block nonspecific binding sites. The anti-STAT1 antibody and anti-TRAF1 antibodies were diluted in 50 mM Tris-HCl, pH 7.4, with 3% goat serum (anti-STAT1 1:1000, anti-TRAF1 1:250) and applied to slides at room temperature for 1 hour. Goat antiumouse horseradish peroxidase-conjugated antibody (Envision detection kit, DAKO USA) was applied for 30 minutes and developed using a diaminobenzidine (DAB) chromogen kit (DAKO USA).

c-REL immunohistochemistry was performed as outlined above with the following modifications. Formalin-fixed, paraffin-embedded sections were dewaxed and antigen retrieved in 1 mM EDTA (ethylenediaminetetraacetic acid) as published.²⁰ Following blocking of nonspecific binding with 5% nonfat dried milk in Tris-buffered saline, pH 7.5, slides were incubated overnight with either affinity-purified rabbit anti-c-REL antibody at 1 μ g/mL (PC139, Oncogene Research Products, Darmstadt, Germany) or pooled nonspecific rabbit Ig (Sigma, St Louis, MO). The slides were then washed and incubated with biotin-conjugated goat antirabbit secondary antibody (Vector, Burlingame, CA), followed by fluorescein isothiocyanate (FITC)-avidin (Molecular Probes, Eugene, OR). The sections were mounted and counterstained with propidium iodide (Molecular Probes). In separately reported studies, the association between nuclear c-REL immunostaining and nuclear factor- κ B (NF- κ B) activation was confirmed using molecular techniques and similarly prepared untreated and CD40 ligand-activated B cells (G.C. et al, manuscript in preparation).

Fluorescence in situ hybridization (FISH)

Air-dried touch preparations were prepared on Superfrost Plus slides from fresh frozen tumor specimens and stored in a desiccator at room temperature until use. Nuclei were hybridized to commercially available probes flanking or spanning the *IgH*, *BCL2*, and *BCL6* loci (Vysis, Downer's Grove, IL) using conditions recommended by the manufacturer. After counterstaining with 4,6 diamidino-2-phenylindole (DAPI), interphase nuclei were scored for various chromosomal aberrations by fluorescence microscopy.

Results

Clinical and pathologic features of MLBCL

The clinical characteristics of the MLBCL and DLBCL patients included in this analysis are outlined in Table 1. The 34 MLBCL patients were predominantly young females (median age, 32 years) with localized and bulky disease (78% stage I/II). Those MLBCL patients with advanced-stage disease had contiguous involvement

of the pleura and/or pericardium. In contrast, DLBCL patients were older (median age, 64 years) with an equal sex distribution and a higher incidence of advanced-stage disease (65% stage III/IV, Table 1). Pathologic review of the diagnostic MLBCL specimens confirmed the presence of sclerosis in 94% of cases; 77% of cases with available Ig immunohistochemistry had undetectable Ig, and 12% had equivocal Ig expression (Table 1). In the subset of MLBCLs with available genetic data, there was only one tumor with a *BCL6* rearrangement (n = 19) and none with *BCL2* translocations (n = 17) (Table 1), consistent with previous reports.¹²

MLBCLs have a unique transcriptional profile

The distinctive clinical and pathologic characteristics of MLBCL—predominantly young women with Ig-negative, localized, and sclerotic tumors—suggested that this entity might also have a unique molecular signature. To address this possibility, diagnostic tumor specimens from the MBLCL and DLBCL cohorts were subjected to transcriptional profiling. The genes were sorted by their degree of correlation with the MLBCL versus DLBCL distinction according to the signal-to-noise metric (see “Patients, materials, and methods”). Permutation of the sample labels indicated that MLBCL had significantly lower expression of more than 1000 genes and significantly higher expression of more than 1000 additional genes, compared with DLBCLs ($P < .01$) (Figure 1A).

The top 50 genes with significantly higher or lower expression in MLBCL (as compared with DLBCL) are visually displayed in Figure 1B. The MLBCL transcriptional signature also included 3 genes previously reported to be expressed at high levels in this disease: the cell surface protein and lipid raft component, *MAL*; the recently described interleukin-4 (IL-4)-induced gene, *FIG1*; and the adhesion molecule, *CD58/LFA3* (Table 2).^{8,21,22} These tumors also had lower levels of *IgM*, consistent with the previously noted absence of Ig in most cases of MLBCL (Figure 1B and Table 3).⁴⁰

Class prediction

The discriminatory power of the gene expression signature was evaluated by building naive Bayes (NB) and weighted voting classifiers for the MLBCL versus DLBCL distinction. These classifiers were tested using a leave-one-out cross-validation strategy. A 100-gene NB model achieved the lowest balanced error rate (11%) (Figure 1B and Supplemental Document). Of interest, 6 patients whose DLBCL was classified by the NB model as MLBCL had predominant, although not exclusive, mediastinal disease (Figure 1B), and 2 patients whose tumors did not involve the mediastinum exhibited *c-REL* amplification (data not shown). In addition, 45 patients whose disease involved regional mediastinal lymph nodes as well as other nodal and extranodal sites were identified by the NB model as having DLBCL rather than primary MLBCL (Figure 1B). These data highlight the potential value of a molecular classifier in entities that are currently defined with only clinical and pathologic criteria.

MLBCL transcriptional profile resembles that of cHL

Inspection of the MLBCL transcriptional profile revealed striking similarities to that of cHL (Figure 1B; Tables 2-3). Like Hodgkin Reed-Sternberg (HRS) cells, MLBCLs had low levels of expression of multiple B-cell signaling components and coreceptors (Table 3 and Figure 2).^{17,42-44} MLBCLs also had high levels of expression of cytokine pathway components, TNF family members, and extracellular matrix elements previously identified in cHL (Table 2).⁴⁵ These observations are of particular interest because

Table 1. Clinical and pathological features of MLBCL and DLBCL patients and tumors

Clinical feature	MLBCL, n = 34	DLBCL, n = 176
Median age, y (range)	32 (14-59)	64 (20-92)
Sex (%)		
Female	24 (71)	84 (48)
Male	10 (29)	92 (52)
Stage (%)		
I/II	25 (74)	53 (30)
III/IV	9 (26)	115 (65)
Unknown	0 (0)	8 (5)
Pathology (%)		N/A
Sclerosis		
Present	32 (94)	
Absent	2 (6)	
Immunoglobulin by IHC*		
Present	3 (12)	
Absent	20 (77)	
Equivocal	3 (12)	
Not available	8	
FISH†		N/A‡
<i>BCL2</i> rearrangements	0	
<i>BCL6</i> rearrangements	1	

N/A indicates not available.

*MLBCL immunoglobulin status by immunohistochemistry (IHC) available for 26 patients (% positive calculated on available cases).

†MLBCL interphase FISH available: *BCL2*, 17 patients; *BCL6*, 19 patients.

‡To be described in a subsequent analysis.




Figure 1. DLBCL and MLBCL genes. (A) Permutation analyses of differentially expressed genes in MLBCL and DLBCL. The observed signal-to-noise ratios (SNRs, x-axis) of ranked genes with lower levels of expression (left) and higher levels of expression (right) (black) in MLBCL versus DLBCL are compared with those expected by chance in 99% (red), 95% (yellow), and 50% (blue) of permutations. The y-axis indicates the number of genes that are differentially expressed in each direction. (B) Comparative gene expression profiles of DLBCL and MLBCL. At the top, the actual clinical/pathologic diagnosis of DLBCL versus MLBCL (green versus red), presence or absence of mediastinal disease (pink versus light green), and molecular prediction of DLBCL versus MLBCL (green versus red) are compared. The top 50 genes that were expressed at higher levels in MLBCL are shown in the upper half of the figure and the top 50 that were more abundant in DLBCL are shown in the lower half of the figure. Red indicates high relative expression; blue, low expression. Color scale at the bottom indicates relative expression in standard deviations from the mean. Each column is a sample and each row is a gene. Genes are clustered using hierarchical clustering. Expression profiles of 176 DLBCLs are on the left; profiles of the 34 MLBCLs are on the right.

MLBCL and the most common subtype of cHL (nodular sclerosis) have similar clinical presentations—in younger patients with local/mediastinal tumors characterized by reactive fibrosis.

To determine the statistical significance of a potential MLBCL-cHL connection, we performed an enrichment test using an independently identified series of differentially expressed genes in HRS cell lines (Figure 3).^{17,19} When compared with random class labelings, the Kolmogorov-Smirnov (KS) statistic indicated that the observed MLBCL-cHL similarity was highly significant. Interestingly, this similarity was primarily attributable to genes with *low levels of expression* in HRS cell lines (versus normal B cells) and primary MLBCL (versus DLBCL) ($p_{\max} = .012$, $p_{\min} < .001$) (Figure 3). In contrast, the similarity between genes with *high levels of expression* in HRS cell lines and primary MLBCL was less significant ($p_{\max} = .213$, $p_{\min} = .007$). This may reflect the contribution of the tumor microenvironment to the gene signatures of primary MLBCL and/or the absence of this contribution from profiles of HRS cell lines maintained *in vitro*.^{19,45}

Decreased expression of BCR signaling cascade components in MLBCL

Like cHL, MLBCLs had low levels of expression of multiple components of the BCR signaling cascade including the cell surface immunoglobulin receptor, IgM; the Ig α / β -associated SRC-family protein tyrosine kinase (PTK), BLK; the SYK PTK-regulated scaffolding protein, BLNK; and the Bruton tyrosine kinase (BTK) binding protein, SAB (Table 3 and Figure 2).⁴⁶⁻⁴⁸ MLBCLs also had low levels of phospholipase C γ 2 (PLC γ 2)-regulated molecules including the critical lymphoid transcription

factor, nuclear factor of activated T cells (NFATc), and protein kinase C β 1 (PKC β), also reported to be decreased in HRS cells (Table 3 and Figure 2).⁴⁶

The phosphatidylinositol-3 kinase (PI-3K)-regulated downstream kinase, AKT, and a major AKT target, FORKHEADP1 (FOXP1), were also expressed at low levels in MLBCL^{46,49} (Table 3 and Figure 2). Lymphoid transcription factors including Spi-B and Ikaros were also less abundant in MLBCL (Table 3 and Figure 2).⁵⁰

In addition to the above-mentioned B-cell transcription factors and components of the BCR signaling cascade, MLBCLs, like HRS cells, had low levels of expression of the CD22 coreceptor and several major histocompatibility complex (MHC) class II molecules (Table 3 and Figure 2).^{46,51} MLBCLs also expressed low levels of the germinal center (GC) metalloendopeptidase, CD10 (Table 3).

Increased expression of cytokine pathway components, TNF family members, and extracellular matrix elements in MLBCL

Like primary cHL, MLBCLs had increased expression of specific cytokine pathway members. For example, MLBCLs expressed high levels of the interleukin-13 receptor α 1 (IL-13R α 1) and additional downstream effectors of IL-13 signaling previously identified in primary cHL (Table 2).^{23,24,45} In normal B cells, IL-13R signaling stimulates cellular proliferation and triggers Ig class switching.⁴⁵ IL-13R α 1 associates with IL-4R α to form a high-affinity IL-13R complex; signaling through this receptor results in phosphorylation of Janus kinase-2 (JAK2) and STAT1.⁵² Of note, MLBCLs expressed high levels of JAK2 and STAT1 and 2

Table 2. Genes with higher levels of expression in MLBCL*

Gene class and specific genes	Fold change†
Cytokines and cytokine pathways	
IL-13R α 1‡	1.9
JAK2‡	2.0
STAT1‡	1.7
Interleukin-4–induced gene 1 (FIG1)§	1.7
NF-IL-3‡	2.0
γ IFN-inducible protein 10 (IP10/CXCR3 ligand)‡	2.0
γ IFN-inducible protein 30 (IFI30/GILT)	1.6
IFN-induced protein 35	1.5
RANTES (CCL5)‡	1.9
Chemokine receptor 1 (CCR1)	1.5
RON (macrophage-stimulating 1 receptor/c-met–related TK)	1.7
Interleukin-15	1.7
TNF family members and induced proteins	
OX40 ligand (TNFSF4)‡	2.0
FAS ligand receptor/CD95 (TNFRSF6)‡§	1.7
Herpes virus entry mediator (HVEM, TNFRSF14)	1.5
TRAF1‡	1.8
TNF- α –induced protein 3 (A20)‡	1.6
TNF- α –induced protein 6	2.3
Costimulatory molecules and inflammatory mediators	
SLAM	1.7
CD80 (B7.1)‡	1.4
CD86 (B7.2)‡	1.4
Cyclooxygenase 1	1.9
Prostaglandin receptor (IP)	2.6
Cell adhesion molecules/extracellular matrix components	
Fibroblast growth factor receptor-1	1.4
CD58 (lymphocyte-function associated antigen 3)‡§	1.5
Integrin α M (CD11b)	1.8
Fibronectin	3.0
Fibroblast activation protein α (seprase)	2.0
Decorin	2.0
Collagen type III α 1	2.1
Collagen type IV α 1	2.3
Collagen type IV α 2	1.6
Matrix metalloproteinase 14	2.0
Tissue inhibitor of metallo-proteinase 1‡	1.9
Other	
MAL§	2.8
SNFT (Jun dimerization protein)	2.8
Adenosine deaminase	1.9

*Gene list derived from the top 400 genes with higher levels of expression in MLBCL. All genes significant at $P < .01$.

†Fold change compared with DLBCL. ‡Previously identified in Hodgkin Reed-Sternberg cells: IL-13R,^{23,24} JAK2,^{25,26} STAT1,²⁷ NF-IL-3,²⁸ IP10,²⁹ RANTES,²⁹ OX40L,³⁰ CD95,³⁰ TRAF1,^{31,32} A20,³³ CD80,³⁴ CD86,³⁵ CD58,³⁶ and TIMP1.³⁷

§Previously identified in MLBCL: FIG1,²¹ FAS/Apo1/CD95,³⁸ MAL,^{8,39} and CD58.^{22,36}

IL-4–induced genes, *FIG1* and *NF-IL-3* (Table 2).^{21,53} Given the composition of the IL-13R α 1/IL-4 α high-affinity receptor, the identification of 2 IL-4–induced genes (*FIG1* and *NF-IL-3*) in MLBCL suggests that the pathway may be active in this disease (Table 2). Consistent with this possibility, *NF-IL-3* was the gene most closely coregulated with the *IL-13R* in MLBCL (data not shown).

MLBCLs also had high levels of expression of γ interferon (IFN)–induced proteins including the chemokine and CXCR3 ligand, IP10⁵⁴; IFI30/GILT⁵⁵; and IFI35 (Table 2).⁵⁶ HRS cell lines and primary cHLs also express γ IFN–induced proteins such as IP10, a chemoattractant for T and natural killer (NK) cells²⁹ (Table 2). Like cHL, MLBCLs also had high levels of expression of the T-helper cell chemokine, RANTES (regulated on activation normal

Table 3. Genes with lower levels of expression in MLBCL*

Gene class and specific genes	Fold change†
B-cell receptor components and signaling cascade	
IgM‡§	5.1
B-cell lymphocyte kinase (BLK)‡	2.7
SH3 domain-binding protein (SAB) (BTK-associated binding protein)	2.0
B-cell linker protein (BLNK)‡	1.5
PKC β I‡	2.0
NFATc‡	2.0
AKT-1	1.5
CD22‡	2.6
Transcription factors	
Spi-B‡	1.6
Ikaros	1.5
FOXP1	2.0
Class II MHC components‡§	
DO β ‡	1.9
DQ β	2.0
DM α	1.5
Germinal center and other B-cell markers	
CD10	3.1
SPAP1 (SH2 domain-containing phosphatase anchor protein, IRTA 4)	2.4
Protein tyrosine phosphatase receptor type C (PTPRCAP)	1.4
TNF family members	
Transmembrane activator and CAML interactor (TAC1) (TNFRSF13B)	1.5
CD27 (TNFRSF7)	1.7
TRAF5‡	1.5
Cytokines	
Interleukin-16‡	1.4

*Gene list derived from the top 400 genes with lower levels of expression in MLBCL. All genes significant at $P < .01$.

†Fold change compared with DLBCL.

‡Previously identified as having reduced expression in Hodgkin Reed-Sternberg cells: IgM, CD22, BLK, PKC β I/II, BLNK, NFATc, Spi-B, class II DO β , TRAF5, and IL-16.¹⁷

§Previously identified as having reduced expression in MLBCL: Ig² and MHC class II.⁴¹

T cells expressed and secreted) (Table 2)²⁹ and the costimulatory molecules CD80 (B7.1) and CD86 (B7.2).^{34,35} MLBCLs also had increased expression of signaling lymphocytic activation molecule (SLAM), a B- and T-cell surface molecule that potentiates lymphocyte expansion in a CD28-independent manner and serves as a




Figure 2. BCR signaling cascade components in MLBCL. Components of the BCR signaling pathway that are expressed at significantly lower levels in MLBCL are indicated in blue, and components that are reported to be expressed at reduced levels in HRS cells are circled in black. Genes are derived from the top 400 with low levels of expression in MLBCL (Table 3).




Figure 3. Enrichment test for differentially expressed HRS genes in MBLCL. A graphic rendition of the computation of the KS scores for HRS genes that are expressed at low (A) and at high (B) levels in MBLCL is shown. The bottom panel for each signature indicates with vertical bars the location of the HRS genes within the ranked set of 15 000 differentially expressed genes in MLBCL and DLBCL (with the MLBCL genes to the left and the DLBCL genes to the right). The top panel shows the assignment of "rewards" and "penalties" to the overall KS score as the list of 15 000 ranked genes is scanned from the MLBCL end (left) to the DLBCL end (right). Every "hit" (ie, the encounter of an HRS gene during the scan) increases the KS score, and every "miss" (the encounter of a non-HRS gene) decreases the score, resulting in the indicated stepwise curve. The final score corresponds to the highest value (on the y-axis) in the plot. High enrichment would correspond to a steep climb upward to the left, whereas lack of enrichment would correspond to a lack of clear upward trend in the curve. The KS statistic indicates that genes reported to have low levels of expression in HRS cell lines are similarly decreased in MLBCL ($p_{\max} = .012$, $p_{\min} < .001$). In contrast, the similarity between genes with higher levels of expression in HRS cell lines and primary MLBCL was less significant ($p_{\max} = .213$, $p_{\min} = .007$), likely reflecting the importance of tumor microenvironment to primary MLBCL and cHL signatures⁴⁵ and the absence of these features in the signatures of isolated HRS cell lines maintained in vitro.¹⁹

measles virus receptor.⁵⁷ In addition, components of the prostaglandin pathway, cyclooxygenase 1 and prostaglandin IP receptor, were also expressed at high levels in MLBCLs.⁵⁸

Equally striking in MLBCL was the increased expression of TNF superfamily (SF) members previously implicated in the pathogenesis of cHL, including the OX40 ligand (TNFSF4) and the FAS ligand receptor (TNFRSF6) previously identified in MLBCL³⁸ (Table 2). The herpes virus entry mediator (HVEM, TNFRSF14) and TRAF1 were also expressed at high levels in MLBCL. These results are of interest because HVEM associates with TRAF1 to stimulate NF- κ B activation⁵⁹ and TRAF-1 is expressed in HRS cells (Table 2) and up-regulated following NF- κ B activation (Table 2).^{31,32,60}

Signaling through TNF receptors triggers distinct signaling pathways leading to either activation of NF- κ B transcription factors or apoptosis.⁶¹ In this regard, TRAF1 plays a unique regulatory role in lymphoid cells, interacting with cellular inhibitor-of-apoptosis proteins (cIAPs) to suppress TNF-induced apoptosis.⁶² In addition to TRAF1, MLBCLs also expressed higher levels of A20, another TNF- α -induced protein (Table 2) that protects against TNF-induced apoptosis.⁶³ Of note, HRS cell lines with high basal expression of A20 and TRAF1 are reported to be resistant to apoptosis.³³ Taken together, these data suggest that in MLBCL, like

cHL, signaling through TNF receptors and associated factors favors NF- κ B activation and resistance to apoptosis.

Primary MLBCL and cHL often exhibit sclerosis, underscoring the potential importance of extracellular matrix components in both diseases. Consistent with these observations, MLBCLs had higher levels of expression of a TNF- α -induced hyaluronan-binding protein (TN-FAIP6)⁶⁴; adhesion molecules including CD58/LFA3, integrin α M/CD11b, and fibronectin; fibroblast growth factor receptors and activation proteins; and additional extracellular matrix components including several types of collagen and decorin (Table 2).^{65,66}

Expression of STAT1 and TRAF1 proteins in MLBCL

The critical roles of IL-13 signaling and NF- κ B activation in cHL and the observed similarities between cHL and MLBCL prompted us to obtain protein confirmation for specific pathway members in MLBCL. Given the known functions of STAT1 in IL-13 signaling and TRAF1 in downstream activation of NF- κ B,^{52,67} we performed STAT1 and TRAF1 immunohistochemistry in a series of MLBCLs and DLBCLs from our cohort with available formalin-fixed paraffin-embedded tissue as well as cases of nodular sclerosis cHL. As summarized in Table 4 and illustrated in Figure 4A, all MLBCLs exhibited tumor cell expression of STAT1, whereas none of the DLBCLs had definite STAT1 staining. The cHLs also expressed STAT1 in the HRS cells and surrounding inflammatory cells (likely macrophages) (Table 4).

There was also uniform TRAF1 staining in the malignant cells of all MLBCLs, whereas the DLBCLs lacked TRAF1 expression (Figure 4A and Table 4). Consistent with prior reports of TRAF1 transcripts and protein in HRS cells of cHL,^{31,32} all of the cHL cases exhibited prominent TRAF1 staining (Table 4).

MLBCLs have immunohistochemical evidence of NF- κ B activation

MLBCLs that lack Ig and associated BCR survival signals have likely developed alternative mechanisms for escaping cell death, similar to

Table 4. Immunohistochemical analysis of STAT1 and TRAF1 expression in DLBCL, MLBCL, and nodular sclerosis HL

Diagnosis (no.)	STAT1 (no.)	TRAF1 (no.)
DLBCL (5)	Negative* (5)	Negative (5)
MLBCL (6)	Positive† (6)	Positive (5) Weakly positive‡ (1)
Nodular sclerosis HL (12)	Positive (6) Weakly positive (5) Negative (1)	Positive (12)

*Staining observed in less than 5% of tumor cells.

†Strong staining observed in more than 90% of tumor cells.

‡Weak staining observed in more than 90% of tumor cells.




Figure 4. STAT1, TRAF1, and c-REL immunohistochemistry in MLBCL. (A) STAT1 and TRAF1 proteins in MLBCL. A control DLBCL (i, iii) and representative MLBCL (ii, iv) were stained for STAT1 (i, ii) and TRAF1 (iii, iv). Original magnification, $\times 1000$. (B) c-REL immunohistochemistry in MLBCL. A control DLBCL (i,iii,v) and representative MLBCL (ii,iv,vi) are doubly stained for c-REL (iii,iv; green) and propidium iodide (i,ii; nuclear, red). The 2 channels of each case are superimposed in subpanels v and vi. Note predominant nuclear c-REL staining of MLBCL at right (vi) compared with the predominantly cytoplasmic c-REL staining of the control DLBCL on the left (v). Original magnification, $\times 400$.

those described in cHL.⁴⁵ In cHL, activation of the NF- κ B pathway enhances the survival of surface immunoglobulin-negative (sIg⁻) Reed-Sternberg cells.⁴⁴ Given the TRAF1 expression in MLBCL and components of the MLBCL transcriptional profiles suggestive of NF- κ B activation (Tables 2 and 4; Figure 4A), we determined the subcellular location of the c-REL subunit of the NF- κ B heterodimer in primary MLBCLs.⁶⁸ In 5 of 6 MLBCLs assessed by c-REL immunohistochemistry, the protein was localized to the nucleus, consistent with activation of the NF- κ B pathway (Figure 4B). These observations highlight a shared survival pathway in MLBCL and cHL.

Discussion

The unique profile of primary MLBCL—low levels of expression of BCR signaling pathway components, a distinctive cytokine pathway signature, and activation of NF- κ B—is strikingly similar to that of a clinically related disorder, cHL. Both MLBCL and cHL of the nodular sclerosis subtype commonly present in young patients as mediastinal tumors with absent surface Ig and prominent fibrosis. MLBCL and classical HRS cells also exhibit common genetic abnormalities including gains of chromosome 2p and 9p. Found in about 20% of MLBCL and up to 50% of cHL, gains in chromosome 2p are associated with amplification of the *REL* locus, one potential mechanism for increased NF- κ B activity and tumor cell resistance to apoptosis.^{11,25,68,69} Gains in chromosome 9p and the *JAK2* locus are observed in about 75% of MLBCLs and 25% of cHLs and are unique to these diseases.^{10,11,26} In addition to these clinical, immunologic, and molecular similarities, there are also rare reported cases of composite cHL and MLBCL, further supporting a pathogenetic relationship between these tumors.⁷⁰ The clonal relationship between HRS cells and non-Hodgkin lymphoma (NHL) cells in composite lymphomas was one of the strongest initial pieces of evidence that cHL was a B-cell lymphoma.⁷¹ In our own series, one patient with MLBCL subsequently relapsed with cHL (nodular sclerosis subtype).

The extent of similarity between the nodular sclerosis subtype of cHL and MLBCL is perhaps surprising given the degree to which the histopathologies and outcome of these 2 entities differ. Cells resembling HRS cells are usually not seen in MLBCL, and MLBCL cells retain the expression of multiple B-cell markers, including CD20, CD79a, BOB1, and OCT2,^{6,72} that are lost in HRS cells.^{17,73} In addition, nodular sclerosis cHL is characteristically associated with a polymorphous inflammatory infiltrate rich in plasma cells, neutrophils, and eosinophils, reactive cell types that are usually absent in MLBCL.⁴⁵ The reactive infiltrate in cHL is recruited and maintained by additional chemokines and cytokines, such as IL-5, IL-6, and IL-10,⁴⁵ that are not part of the MLBCL signature. Based on the existence of unusual patients with both nodular sclerosis cHL and MLBCL, we suggest these 2 entities arise from a common precursor cell with a growth and survival advantage stemming from genetic lesions that result in constitutive activation of NF- κ B. Acquisition of distinct sets of secondary genetic lesions then results in either MLBCL or nodular sclerosis cHL, potentially explaining the differences in the clinical course of these diseases.

The reason for the lower levels of expression of Ig and BCR signaling components in MLBCL is not yet known. In a subset of cHLs, the lack of a functional BCR has been attributed to “crippling” somatic mutations of the rearranged immunoglobulin gene.⁷⁴ In cHLs with functional gene rearrangements, the absence of Ig has been ascribed to deficiencies in transcription factors (OCT2, BOB1, PU1) necessary for Ig synthesis.^{73,75} In a recent study, 12 of 13 MLBCLs exhibited functional *IgVH* gene rearrangements, suggesting that destructive somatic mutations were not the primary cause of reduced surface Ig in this disease.⁹ More recently, abundant OCT2 and BOB1 transcription factors and Ig transcripts with a switched isotype were detected in MLBCL, suggesting that the transcriptional machinery for Ig synthesis was intact in these tumors.⁷² However, our MLBCL transcriptional profile reveals reduced levels of other critical B-cell transcription factors such as

NFATc and Spi-B, which have known roles in regulating the expression of specific Ig isotypes.^{76,77}

MLBCL has a prominent cytokine pathway signature, which likely reflects dynamic interactions between the tumor cells, infiltrating inflammatory cells, and the surrounding tumor stroma. Like cHLs, MLBCLs have increased abundance of high-affinity IL-13 receptor subunits and downstream effectors (JAK2 and STAT1, NF-IL-3), implicating this cytokine pathway in disease pathogenesis.⁴⁵ Our studies also indicate that MLBCLs overexpress specific TNF family members that are known to interact with TRAF1 and subsequently activate NF- κ B. TRAF1 transcripts and protein were also more abundant in MLBCL. Consistent with a

likely role for NF- κ B activation in MLBCL, MLBCLs exhibited c-REL nuclear staining.

The nuclear localization of c-REL and likely activation of NF- κ B in MLBCL provide new clues regarding more rational targeted therapy in this disease. Given the limitations of current combined modality therapy, including treatment failures and long-term radiation toxicity, the NF- κ B pathway and supporting cytokine network may represent more specific MLBCL treatment targets. In this regard, the current study both defines molecular differences in morphologically similar tumors (MLBCL versus DLBCL) and identifies shared features of morphologically distinct neoplasms (MLBCL and cHL).

References

- Savage KJ, Shipp MA. Initial evaluation: staging and prognostic factors. In: Mauch PM, Armitage JO, Harris NL, Dalla-Favera R, Coiffier B, eds. *Non-Hodgkin's Lymphomas*. Philadelphia, PA: Lippincott Williams and Wilkins. In press.
- Barth TF, Leithauser F, Joos S, Bentz M, Moller P. Mediastinal (thymic) large B-cell lymphoma: where do we stand? *Lancet Oncol*. 2002;3:229-234.
- Banks PM, Warnke RA. Mediastinal (thymic) large B-cell lymphoma. In: Jaffe ES, Harris NL, Stein H, Vardiman JW, eds. *World Health Organization Classification of Tumours: Pathology and Genetics of Tumours of Hematopoietic and Lymphoid Tissues*. Lyon, France: IARC; 2001:175-178.
- Lazzarino M, Orlandi E, Paulli M, et al. Primary mediastinal B-cell lymphoma with sclerosis: an aggressive tumor with distinctive clinical and pathologic features. *J Clin Oncol*. 1993;11:2306-2313.
- Lamarre L, Jacobson JO, Aisenberg AC, Harris NL. Primary large cell lymphoma of the mediastinum. *Am J Surg Pathol*. 1989;13:730-739.
- Kanavaros P, Gaulard P, Charlotte F, et al. Discordant expression of immunoglobulin and its associated molecule mb-1/CD79a is frequently found in mediastinal large B cell lymphomas. *Am J Pathol*. 1995;146:735-741.
- Isaacson P, Norton A, Addis B. The human thymus contains a novel population of B-lymphocytes. *Lancet*. 1987;2:1488-1490.
- Copie-Bergman C, Plonquet A, Alonso MA, et al. MAL expression in lymphoid cells: further evidence for MAL as a distinct molecular marker of primary mediastinal large B-cell lymphomas. *Mod Pathol*. 2002;15:1172-1180.
- Leithauser F, Bauerle M, Huynh MQ, Moller P. Isotype-switched immunoglobulin genes with a high load of somatic hypermutation and lack of ongoing mutational activity are prevalent in mediastinal B-cell lymphoma. *Blood*. 2001;98:2762-2770.
- Bentz M, Barth TF, Bruderlein S, et al. Gain of chromosome arm 9p is characteristic of primary mediastinal B-cell lymphoma (MLB): comprehensive molecular cytogenetic analysis and presentation of a novel MLB cell line. *Genes Chromosomes Cancer*. 2001;30:393-401.
- Joos S, Otano-Joos MI, Ziegler S, et al. Primary mediastinal (thymic) B-cell lymphoma is characterized by gains of chromosomal material including 9p and amplification of the REL gene. *Blood*. 1996;87:1571-1578.
- Tsang P, Cesarman E, Chadburn A, Liu YF, Knowles DM. Molecular characterization of primary mediastinal B cell lymphoma. *Am J Pathol*. 1996;148:2017-2025.
- van Besien K, Kelta M, Bahaguna P. Primary mediastinal B-cell lymphoma: a review of pathology and management. *J Clin Oncol*. 2001;19:1855-1864.
- Chronowski GM, Wilder RB, Tucker SL, et al. Analysis of in-field control and late toxicity for adults with early-stage Hodgkin's disease treated with chemotherapy followed by radiotherapy. *Int J Radiat Oncol Biol Phys*. 2003;55:36-43.
- Golub TR, Slonim DK, Tamayo P, et al. Molecular classification of cancer: class discovery and class prediction by gene expression monitoring. *Science*. 1999;286:531-537.
- Shipp MA, Ross KN, Tamayo P, et al. Diffuse large B-cell lymphoma outcome prediction by gene-expression profiling and supervised machine learning. *Nat Med*. 2002;8:68-74.
- Schwering I, Brauninger A, Klein U, et al. Loss of the B-lineage-specific gene expression program in Hodgkin and Reed-Sternberg cells of Hodgkin lymphoma. *Blood*. 2003;101:1505-1512.
- Mootha VK, Lingren CM, Eriksson KF, et al. PDG-1alpha responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nat Genet*. 2003;34:267-273.
- Kuppers R, Klein U, Schwering I, Distler V, et al. Identification of Hodgkin and Reed-Sternberg cell-specific genes by gene expression profiling. *J Clin Invest*. 2003;111:529-537.
- Cattoretti G, Pileri S, Parravicini C, et al. Antigen unmasking on formalin-fixed, paraffin-embedded tissue sections. *J Pathol*. 1993;171:83-98.
- Copie-Bergman C, Boulland ML, Dehouille C, et al. Interleukin 4-induced gene 1 is activated in primary mediastinal large B-cell lymphoma. *Blood*. 2003;101:2756-2761.
- Eichelmann A, Koretz K, Mechtersheimer G, Moller P. Adhesion receptor profile of thymic B-cell lymphoma. *Am J Pathol*. 1992;141:729-741.
- Skinider BF, Elia AJ, Gascoyne RD, et al. Interleukin 13 and interleukin 13 receptor are frequently expressed by Hodgkin and Reed-Sternberg cells of Hodgkin lymphoma. *Blood*. 2001;97:250-255.
- Ohshima K, Akaiwa M, Umeshita R, Suzumiya J, Izuhara K, Kikuchi M. Interleukin-13 and interleukin-13 receptor in Hodgkin's disease: possible autocrine mechanism and involvement in fibrosis. *Histopathology*. 2001;38:368-375.
- Joos S, Granzow M, Holtgreve-Grez H, et al. Hodgkin's lymphoma cell lines are characterized by frequent aberrations on chromosomes 2p and 9p including REL and JAK2. *Int J Cancer*. 2003;103:489-495.
- Joos S, Kupper M, Ohl S, et al. Genomic imbalances including amplification of the tyrosine kinase gene JAK2 in CD30⁺ Hodgkin cells. *Cancer Res*. 2000;60:549-552.
- Garcia JF, Camacho FI, Morente M, et al. Hodgkin and Reed-Sternberg cells harbor alterations in the major tumor suppressor pathways and cell-cycle checkpoints: analyses using tissue microarrays. *Blood*. 2003;101:681-689.
- Kapp U, Yeh WC, Patterson B, et al. Interleukin 13 is secreted by and stimulates the growth of Hodgkin and Reed-Sternberg cells. *J Exp Med*. 1999;189:1939-1946.
- Maggio E, van den Berg A, Diepstra A, Kluiver J, Visser L, Poppema S. Chemokines, cytokines and their receptors in Hodgkin's lymphoma cell lines and tissues. *Ann Oncol*. 2002;13(suppl 1):52-56.
- Gruss HJ, Duyster J, Herrmann F. Structural and biological features of the TNF receptor and TNF ligand superfamilies: interactive signals in the pathobiology of Hodgkin's disease. *Ann Oncol*. 1996;7(suppl 4):19-26.
- Durkop H, Foss HD, Demel G, Klotzbach H, Hahn C, Stein H. Tumor necrosis factor receptor-associated factor 1 is overexpressed in Reed-Sternberg cells of Hodgkin's disease and Epstein-Barr virus-transformed lymphoid cells. *Blood*. 1999;93:617-623.
- Izban KF, Ergin M, Martinez RL, Alkan S. Expression of the tumor necrosis factor receptor-associated factors (TRAFs) 1 and 2 is a characteristic feature of Hodgkin and Reed-Sternberg cells. *Mod Pathol*. 2000;13:1324-1331.
- Durkop H, Hirsch B, Hahn C, Foss HD, Stein H. Differential expression and function of A20 and TRAF1 in Hodgkin lymphoma and anaplastic large cell lymphoma and their induction by CD30 stimulation. *J Pathol*. 2003;200:214-221.
- Delabie J, Ceuppens JL, Vandenberghe P, de Boer M, Coorevits L, De Wolf-Peeters C. The B7/BB1 antigen is expressed by Reed-Sternberg cells of Hodgkin's disease and contributes to the stimulating capacity of Hodgkin's disease-derived cell lines. *Blood*. 1993;82:2845-2852.
- Van Gool SW, Delabie J, Vandenberghe P, Coorevits L, De Wolf-Peeters C, Ceuppens JL. Expression of B7-2 (CD86) molecules by Reed-Sternberg cells of Hodgkin's disease. *Leukemia*. 1997;11:846-851.
- Delabie J, Chan WC, Weisenburger DD, De Wolf-Peeters C. The antigen-presenting cell function of Reed-Sternberg cells. *Leuk Lymphoma*. 1995;18:35-40.
- Oelmann E, Herbst H, Zuhlsdorf M, et al. Tissue inhibitor of metalloproteinases 1 is an autocrine and paracrine survival factor, with additional immune-regulatory functions, expressed by Hodgkin/Reed-Sternberg cells. *Blood*. 2002;99:258-267.
- Moller P, Henne C, Leithauser F, et al. Coregulation of the APO-1 antigen with intercellular adhesion molecule-1 (CD54) in tonsillar B cells and coordinate expression in follicular center B cells and in follicle center and mediastinal B-cell lymphomas. *Blood*. 1993;81:2067-2075.
- Copie-Bergman C, Gaulard P, Maouche-Chretien L, et al. The MAL gene is expressed in primary mediastinal large B-cell lymphoma. *Blood*. 1999;94:3567-3575.

40. Moller P, Moldenhauer G, Momburg F, et al. Mediastinal lymphoma of clear cell type is a tumor corresponding to terminal steps of B cell differentiation. *Blood*. 1987;69:1087-1095.
41. Moller P, Lammler B, Herrmann B, Otto HF, Moldenhauer G, Momburg F. The primary mediastinal clear cell lymphoma of B-cell type has variable defects in MHC antigen expression. *Immunology*. 1986;59:411-417.
42. Hertel CB, Zhou XG, Hamilton-Dutoit SJ, Junker S. Loss of B cell identity correlates with loss of B cell-specific transcription factors in Hodgkin/Reed-Sternberg cells of classical Hodgkin lymphoma. *Oncogene*. 2002;21:4908-4920.
43. Marafioti T, Hummel M, Foss HD, et al. Hodgkin and Reed-Sternberg cells represent an expansion of a single clone originating from a germinal center B-cell with functional immunoglobulin gene rearrangements but defective immunoglobulin transcription. *Blood*. 2000;95:1443-1450.
44. Kuppers R, Scherwing I, Brauning A, Rajewsky K, Hansmann ML. Biology of Hodgkin's lymphoma. *Ann Oncol*. 2002;13(suppl 1):11-18.
45. Skinnider BF, Mak TW. The role of cytokines in classical Hodgkin lymphoma. *Blood*. 2002;99:4283-4297.
46. Niirio H, Clark EA. Regulation of B-cell fate by antigen-receptor signals. *Nat Rev Immunol*. 2002;2:945-956.
47. Yamadori T, Baba Y, Matsushita M, et al. Bruton's tyrosine kinase activity is negatively regulated by Sab, the Btk-SH3 domain-binding protein. *Proc Natl Acad Sci U S A*. 1999;96:6341-6346.
48. Fu C, Turck CW, Kurosaki T, Chan AC. BLNK: a central linker protein in B cell activation. *Immunity*. 1998;9:93-103.
49. Banham AH, Beasley N, Campo E, et al. The FOXP1 winged helix transcription factor is a novel candidate tumor suppressor gene on chromosome 3p1. *Cancer Res*. 2001;61:8820-8829.
50. Kirstetter P, Thomas M, Dierich A, Kastner P, Chan S. Ikaros is critical for B cell differentiation and function. *Eur J Immunol*. 2002;32:720-730.
51. Otipoby KL, Andersson KB, Draves KE, et al. CD22 regulates thymus-independent responses and the lifespan of B cells. *Nature*. 1996;384:634-637.
52. Roy B, Bhattacharjee A, Xu B, Ford D, Maizel AL, Cathcart MK. IL-13 signal transduction in human monocytes: phosphorylation of receptor components, association with Jaks, and phosphorylation/activation of Stats. *J Leukoc Biol*. 2002;72:580-589.
53. Chu CC, Paul WE. Expressed genes in interleukin-4 treated B cells identified by cDNA representational difference analysis. *Mol Immunol*. 1998;35:487-502.
54. Neville LF, Mathiak G, Bagasra O. The immunobiology of interferon-gamma inducible protein 10 kD (IP-10): a novel, pleiotropic member of the C-X-C chemokine superfamily. *Cytokine Growth Factor Rev*. 1997;8:207-219.
55. Arunachalam B, Phan UT, Geuze HJ, Cresswell P. Enzymatic reduction of disulfide bonds in lysosomes: characterization of a gamma-interferon-inducible lysosomal thiol reductase (GILT). *Proc Natl Acad Sci U S A*. 2000;97:745-750.
56. Bange FC, Vogel U, Flohr T, Kiekenbeck M, Denecke B, Bottger EC. IFP 35 is an interferon-induced leucine zipper protein that undergoes interferon-regulated cellular redistribution. *J Biol Chem*. 1994;269:1091-1098.
57. Hahm B, Arbour N, Naniche D, Homann D, Manchester M, Oldstone MB. Measles virus infects and suppresses proliferation of T lymphocytes from transgenic mice bearing human signaling lymphocytic activation molecule. *J Virol*. 2003;77:3505-3515.
58. Tilley SL, Coffman TM, Koller BH. Mixed messages: modulation of inflammation and immune responses by prostaglandins and thromboxanes. *J Clin Invest*. 2001;108:15-23.
59. Marsters SA, Ayres TM, Skubatch M, Gray CL, Rothe M, Ashkenazi A. Herpesvirus entry mediator, a member of the tumor necrosis factor receptor (TNFR) family, interacts with members of the TNFR-associated factor family and activates the transcription factors NF-kappaB and AP-1. *J Biol Chem*. 1997;272:14029-14032.
60. Hinz M, Loser P, Mathas S, Krappmann D, Dorken B, Scheiderei C. Constitutive NF-kappaB maintains high expression of a characteristic gene network, including CD40, CD86, and a set of antiapoptotic genes in Hodgkin/Reed-Sternberg cells. *Blood*. 2001;97:2798-2807.
61. Locksley RM, Killeen N, Lenardo MJ. The TNF and TNF receptor superfamilies: integrating mammalian biology. *Cell*. 2001;104:487-501.
62. Bradley JR, Pober JS. Tumor necrosis factor receptor-associated factors (TRAFs). *Oncogene*. 2001;20:6482-6491.
63. He KL, Ting AT. A20 inhibits tumor necrosis factor (TNF) alpha-induced apoptosis by disrupting recruitment of TRADD and RIP to the TNF receptor 1 complex in Jurkat T cells. *Mol Cell Biol*. 2002;22:6034-6045.
64. Lee TH, Wisniewski HG, Vilcek J. A novel secretory tumor necrosis factor-inducible protein (TSG-6) is a member of the family of hyaluronate binding proteins, closely related to the adhesion receptor CD44. *J Cell Biol*. 1992;116:545-557.
65. Goldstein LA, Ghersi G, Pineiro-Sanchez ML, et al. Molecular cloning of seprase: a serine integral membrane protease from human melanoma. *Biochim Biophys Acta*. 1997;1361:11-19.
66. Timar J, Lapis K, Dudas J, Sebestyén A, Kopper L, Kovalszky I. Proteoglycans and tumor progression: Janus-faced molecules with contradictory functions in cancer. *Semin Cancer Biol*. 2002;12:173-186.
67. Chung JY, Park YC, Ye H, Wu H. All TRAFs are not created equal: common and distinct molecular mechanisms of TRAF-mediated signal transduction. *J Cell Sci*. 2002;115:679-688.
68. Barth TF, Martin-Subero JI, Joos S, et al. Gains of 2p involving the REL locus correlate with nuclear c-Rel protein accumulation in neoplastic cells of classical Hodgkin's lymphoma. *Blood*. 2003;101:3681-3686.
69. Martin-Subero JI, Gesk S, Harder L, et al. Recurrent involvement of the REL and BCL11A loci in classical Hodgkin lymphoma. *Blood*. 2002;99:1474-1477.
70. Rudiger T, Jaffe ES, Delsol G, et al. Workshop report on Hodgkin's disease and related diseases ('grey zone' lymphoma). *Ann Oncol*. 1998;9(suppl 5):S31-S38.
71. Brauning A, Hansmann ML, Strickler JG, et al. Identification of common germinal-center B-cell precursors in two patients with both Hodgkin's disease and non-Hodgkin's lymphoma. *N Engl J Med*. 1999;340:1239-1247.
72. Pileri SA, Gaidano G, Zinzani PL, et al. Primary mediastinal B-cell lymphoma: high frequency of BCL-6 mutations and consistent expression of the transcription factors OCT-2, BOB.1, and PU.1 in the absence of immunoglobulins. *Am J Pathol*. 2003;162:243-253.
73. Re D, Muschen M, Ahmadi T, et al. Oct-2 and Bob-1 deficiency in Hodgkin and Reed Sternberg cells. *Cancer Res*. 2001;61:2080-2084.
74. Kanzler H, Kuppers R, Hansmann ML, Rajewsky K. Hodgkin and Reed-Sternberg cells in Hodgkin's disease represent the outgrowth of a dominant tumor clone derived from (crippled) germinal center B cells. *J Exp Med*. 1996;184:1495-1505.
75. Jundt F, Kley K, Anagnostopoulos I, et al. Loss of PU.1 expression is associated with defective immunoglobulin transcription in Hodgkin and Reed-Sternberg cells of classical Hodgkin disease. *Blood*. 2002;99:3060-3062.
76. Ranger AM, Hodge MR, Gravalles EM, et al. Delayed lymphoid repopulation with defects in IL-4-driven responses produced by inactivation of NF-ATc. *Immunity*. 1998;8:125-134.
77. Su GH, Ip HS, Cobb BS, Lu MM, Chen HM, Simon MC. The Ets protein Spi-B is expressed exclusively in B cells and T cells during development. *J Exp Med*. 1996;184:203-214.