

STRATIGRAPHIC NOTE

Jurassic-Cretaceous Arabian orbital stratigraphy: The AROS-JK Chart

Moujahed Al-Husseini and Robley K. Matthews

An orbital-forcing, glacio-eustatic model predicts that the depositional durations of transgressive-regressive sequences manifest a periodic hierarchy. Model fourth-, third- and second-order sequences have discrete depositional periods that range from 405 thousand years (Ky) to 14.58 million years (My) (Figure 1; Matthews and Frohlich, 1998, 2002; Immenhauser and Matthews, 2004; Al-Husseini and Matthews, 2005a). This note is part of the ongoing Arabian Orbital Stratigraphy (AROS) program aimed at correlating the model to the sequence stratigraphy of the Arabian Plate.

This note accompanies the enclosed Jurassic-Cretaceous Chart (AROS-JK Chart) and focuses on recently documented and dated Jurassic and Cretaceous Arabian sequences. The framework is based on tuning the LA04 orbital-forcing computation of Laskar et al. (2004) and glacio-eustatic parametric forward modeling (Matthews and Frohlich, 2002, p. 515) by R.K. Matthews. Second-order sequence boundaries are here predicted to have occurred every 14.58 My, starting at either 11.3 ± 0.3 or 16.1 ± 0.3 Ma, depending on the choice among geologically reasonable model parameters. The correlation to Jurassic and Cretaceous regional unconformities-hiatuses, with ages estimated from GTS 2004 (Gradstein et al., 2004), supported a correlation to the 16.1 Ma second-order model (Figure 2).

In this note we position representative second-, third- and possible fourth-order Arabian depositional sequences, maximum flooding surfaces and time-marker units in the model framework (AROS JK-Chart). With the exception of the Arabian Plate maximum flooding surfaces (Sharland et al., 2001) most of these sequences, surfaces and time-marker units are only defined locally to semi-regionally (e.g. outcrop, concession, petroleum field, etc.); their extent (or lack of) across Arabia remains to be evaluated. The bibliographic references used to date and correlate the sequences and time-rock units are listed in descending stratigraphic age below.

RECENT STRATIGRAPHIC SOURCES

Geological Time Scale GTS 2004 (Gradstein et al., 2004).

Arabian Plate Maximum Flooding Surface, MFS (Sharland et al., 2001; Davies et al., 2003; Simmons et al., 2007).

Aruma Group (Turonian-Paleogene)

Saudi Arabia, Aruma Formation (Vaslet et al., 1983, 1991; Le Nindre et al., 1990; Philip et al., 2002; Al-Husseini and Matthews 2005b; Y.-M. Le Nindre et al., *in preparation*): Paleogene Lina Member; Late Maastrichtian Hajajah Member, *Aruma Sequence 3* (sub-sequences Ar 3a to 3f) and *Aruma Sequence 2* (sub-sequences Ar2a to 2d); and Late Campanian-Maastrichtian Khanasir Member, *Aruma Sequence 1* (sub-sequences Ar1a to Ar1c).

Iraq, Aruma Group (van Bellen et al., 1959; Jassim and Goff, 2006; Haddad and Ameen, 2007): Campanian-Maastrichtian Shiranish Formation; Campanian Lower Hartha Formation, *Sequence C3*; Santonian?-Campanian Sa'adi Formation, *Sequences C1 and C2*; Santonian-Campanian? Tanuma Formation, *Sequences B2 and B3*; and Turonian?-Coniacian Khasib Formation, *Sequence B1*.

Turonian Unconformity-Hiatus (Searle, 2007)

Wasia Group (Late Aptian-Turonian)

Kuwait, Wasia Group (Al-Fares et al., 1998): Cenomanian Mishrif Formation, Cenomanian Rumaila Formation, Cenomanian-Albian Ahmadi Formation, Albian Wara Formation, Albian Mauddud Formation, Albian Burgan Formation and Late Aptian "Unnamed Clastics Formation".




Figure 1: The order of transgressive-regressive depositional sequences (DS) and sequence boundaries (SB) is indicated by superscripts 2 to 4. A model second-order sequence (DS² N) has a depositional period of 14.58 million years (My) and is bounded by regional second-order sequence boundaries (SB² N and SB² N-1, unconformities in proximal settings). A complete DS² consists of six third-order sequences (DS³ N.1 to DS³ N.6) with depositional periods of 2.025 My (short), 2.43 My (nominal), 2.835 My (long) that are bounded by third-order sequence boundaries (SB³). Even-odd DS³ pairs (e.g. DS³ N.3 + N.4) always form double-third-order sequences with a period of 4.86 My, but may not always clearly separate as two DS³s. The sequential arrangement of DS³s (e.g. short-long or nominal-nominal, etc.) is shown schematically. Fourth-order sequences have a period of 405,000 years (405 Ky) and are uniquely identified as DS⁴ N.M.1 to N.M.5, or to N.M.6 or to N.M.7.




Figure 2: Cretaceous and Jurassic rock units (groups, formations and key members) are shown in terms of the second-order AROS framework. Hiatuses are indicated in grey between unconformity below, and transgressive surface above. Detailed representative successions in the AROS-JK Chart are highlighted in green.

Kuwait, Burgan Formation (Kirby and Al-Humoudi, 1996; Kirby et al., 1998): Albian Upper Third Sand (Mauddud Formation and mfs 3.U.D to 3.U.A); Albian Middle Third Sand (mfs 3.M.F to 3.M.A); Albian Lower Third Sand (mfs 3.L.E to 3.L.A); Albian Dair Limestone; Albian Upper Fourth Sand (mfs 4.U.D to 4.U.A); Late Aptian?-Albian Middle Fourth Sand (Main Sand); and Late Aptian Lower Fourth Sand ("Unnamed Clastics Formation").

Thamama Group (Late Tithonian-Aptian)

Oman and UAE, Kharaiib and Shu'aiba Formations (Boichard et al., 1995; Taher, 1997; van Buchem et al., 2002; Boote and Mou, 2003; Immenhauser et al., 2004; van Buchem, *in preparation*): Late Aptian *Bab Sequence*; Early Aptian *Lower Shu'aiba Sequence* (including Hawar sub-sequence); Late Barremian *Upper Kharaiib Sequence* (Thamama Zone B and Dense B); and Early Barremian *Lower Kharaiib Sequence* (Thamama Zone C and Dense C).

UAE, Abu Dhabi, Lekhwair Formation, Nasr Member (Taher, 1997): Hauterivian?-Barremian Thamama Zone D to Dense E.

UAE, Abu Dhabi (Aziz and El Sattar, 1997): Early Hauterivian *Zakum Sequence IV* (Thamama Zone F to Dense H); Late Valanginian *Habshan Sequence III*; Early Valanginian *Habshan Sequence II*; and Berriasian *Habshan Sequence I*.

UAE, Abu Dhabi, Tithonian Manifa Formation (Grotsch et al., 2003).

Shaqra Group (Jurassic)

Saudi Arabia, Shaqra Group (Powers, 1968; Wilson, 1985; Manivit et al., 1985a, b, 1986, 1990; Enay, 1987; Vaslet et al., 1983, 1991; Le Nindre et al., 1990; Meyer et al., 1996; Hughes, 1996, 2004, 2006; Fischer et al., 2001; Haq and Al-Qahtani, 2005; G. Grabowski, written communication, 2006; Al-Husseini et al., 2006; Hughes et al., 2006; Enay et al., *in preparation*): Tithonian Hith Anhydrite: Manifa Member, Transition Member and Main Hith Anhydrite; Tithonian?-Kimmeridgian Arab A, B and C members; Kimmeridgian Arab-D Member, Arab-D Anhydrite, Arab-D Reservoir, Arab-D Reservoir Zones 1 to 4, Arab-D Layers 1 to 18, Biozones D1 to D3; Kimmeridgian Jubaila Formation, units J1 and J2; Oxfordian Hanifa Formation, *Ulayyah and Hawtah Sequences*; Callovian Tuwaiq Mountain Limestone, units T3 to T1, Daddiya, Maysiyah and Bladiyah members; Dhurma Formation, Lower Member (Dhibi Limestone and Dhurma Shale), Middle Member, and Upper Member (Hisyan and Atash); Dhurma Formation: Bajocian units D1 to D3, Bathonian units D4 to D6 and Callovian unit D7.

Kuwait, Late Sinemurian? to Early Aalenian? Marrat Formation (Yousif and Nauman, 1997; Al-Sahlan, 2005; G. Grabowski, written communication, 2006).

REFERENCES

- Al-Fares, A.A., M. Bouman and P. Jeans 1998. A new look at the Middle to Lower Cretaceous stratigraphy, offshore Kuwait. *GeoArabia*, v. 3, no. 4, p. 543-560.
- Al-Husseini, M.I. and R.K Matthews, 2005a. Arabian Orbital Stratigraphy: Periodic second-order sequence boundaries? *GeoArabia*, v. 10, no. 2, p 165-184.
- Al-Husseini, M.I. and R.K. Matthews 2005b. Stratigraphic note: Orbital-forcing calibration of the Late Cretaceous and Paleocene Aruma Formation, Saudi Arabia. *GeoArabia*, v. 10, no. 3, p 173-176.
- Al-Husseini, M.I. and R.K Matthews, 2006. Stratigraphic Note: Orbital calibration of the Arabian Jurassic second-order sequence stratigraphy. *GeoArabia*, v. 11, no. 3, p. 161-170.
- Al-Husseini, M.I., R.K. Matthews and J. Mattner 2006. Stratigraphic Note: Orbital-forcing calibration of the Late Jurassic (Oxfordian-early Kimmeridgian) Hanifa Formation, Saudi Arabia. *GeoArabia*, v. 11, no. 1, p. 145-149.
- Al-Sahlan, G. 2005. Letter to the Editor. *GeoArabia*, v. 10, no. 3, p. 193-194.
- Aziz, S.K. and M.M.A. El-Sattar 1997. Sequence stratigraphic modeling of the lower Thamama Group, east onshore Abu Dhabi, United Arab Emirates. *GeoArabia*, v. 2, no. 2, p. 179-202.

- Boichard, R.A.P., A.S. Al-Suwaidi and H. Karakhanian 1995. Sequence boundary types and related porosity evolutions: example of the upper Thamama Group in field "A" offshore Abu Dhabi, United Arab Emirates. In, M.I. Al-Husseini (Ed.), Middle East Petroleum Geosciences Conference, GEO'94. Gulf PetroLink, Bahrain, v. 1, p. 191-201.
- Boote, D.R.D. and D. Mou 2003. Safah field, Oman: retrospective of a new-concept exploration play, 1980 to 2000. *GeoArabia*, v. 8, no. 3, p. 367-430.
- Davies, Roger B., David M. Casey, Andrew D. Horbury, Peter R. Sharland and Michael D. Simmons 2002. Early to mid-Cretaceous mixed carbonate-clastic shelfal systems: examples, issues, and models from the Arabian Plate. *GeoArabia*, v. 7, no. 3, p. 541-598.
- Enay, R. 1987. Le Jurassique d'Arabie Saoudite Centrale. *Geobios*, Lyons, Memoir Special, no. 9, 314 p.
- Fischer, J.-C., Y.-M. Le Nindre, J. Manivit and D. Vaslet 2001. Jurassic gastropod faunas of central Saudi Arabia. *GeoArabia*, v. 6, no. 1, p. 63-100.
- Gradstein, F.M., J.G. Ogg, A.G. Smith, et al. 2004. *A Geologic Time Scale 2004*. Cambridge University Press, 589 p.
- Grotsch, J., O. Suwaina, G. Ajlani, A. Taher, R. Khassawneh, S. Lokier, G. Coy, E. van der Weerd, S. Masalmeh and J. van Dorp 2003. The Arab Formation in central Abu Dhabi: 3-D reservoir architecture and static and dynamic modeling, *GeoArabia*, v. 8, no. 1, p. 47-86.
- Haq, B.U. and A.M. Al-Qahtani 2005. Phanerozoic cycles of sea-level change on the Arabian Platform. *GeoArabia*, v. 10, no. 2, p. 127-160.
- Hughes, G.W. 1996. A new bioevent stratigraphy of Late Jurassic Arab-D carbonates of Saudi Arabia. *GeoArabia*, v. 1, no. 3, p. 417-434.
- Hughes, G.W. 2004. Middle to Upper Jurassic Saudi Arabian carbonate petroleum reservoirs: biostratigraphy, micropalaeontology and palaeoenvironments. *GeoArabia*, v. 9, no. 3, p. 79-114.
- Hughes, G.W. 2006 (in press). Biofacies and Palaeoenvironments of the Jurassic Shaqra Group of Saudi Arabia.
- Hughes, G.W., M. Al-Khaled and O. Varol 2006 (in press). Oxfordian biofacies and palaeoenvironments of Saudi Arabia.
- Immenhauser, A. and R.K. Matthews 2004. Albian sea-level cycles in Oman: the 'Rosetta Stone' approach. *GeoArabia*, v. 9, no. 3, p. 11-46.
- Immenhauser, A., H. Hillgärtner, U. Sattler, G. Bertotti, P. Schoepfer, P. Homewood, V. Vahrenkamp, T. Steuber, J.-P. Masse, H. Droste, J. Taal-van Koppen, B. van der Kooij, E. van Bentum, K. Verwer, E.H. Strating, W. Swinkels, J. Peters, I. Immenhauser-Pothast and S. Al Maskery 2004. Barremian-lower Aptian Qishn Formation, Haushi-Huqf area, Oman: a new outcrop analogue for the Kharaiib/Shu'aiba reservoirs. *GeoArabia*, v. 9, no. 1, p. 153-194.
- Jassim, S.Z. and J.C. Goff 2006. *Geology of Iraq*. Published by Dolin, Prague and Moravian Museure, Brno, Czech Republic, 341 p.
- Kirby, R.H. and J.A. Al-Humoud 1996. Genetic Stratigraphy, Depositional Systems and Reservoir Performance, Burgan Formation, Greater Burgan Field, Kuwait. Paper presented at GEO 1994, Bahrain.
- Kirby, R.H., B.S. Carr, J.A. Al-Humoud, A.I. Safar, D. A. Al-Matar and W. Naser 1998. Characterization of a vertically compartmentalized reservoir in a supergiant field, Burgan Formation, Greater Burgan field, Part 1, Stratigraphy and water encroachment. Society of Petroleum Engineers SPE Paper 49214, p. 509-520.
- Laskar, J., P. Robutel, F. Joutel, M. Gastineau, A.C.M. Correia and B. Levrard 2004. A long term numerical solution for the insolation quantities of the Earth. *Astronomy and Astrophysics*, v. 428, p. 261-285. DOI: 10.1051/0004-6361:20041335.
- Le Nindre, Y.-M., J. Manivit and D. Vaslet 1990. Stratigraphie séquentielle du Jurassique et du Crétacé en Arabie Saoudite. *Bulletin, Société géologique de France*, Paris, sér. 8, v. VI, p. 1025-1034.
- Manivit, J., C. Pellaton, D. Vaslet, Y.-M. Le Nindre, J.-M. Brosse and J. Fourniguet 1985a. Explanatory notes to the geologic map of the Wadi Al Mulayh Quadrangle, Kingdom of Saudi Arabia. Geoscience Map GM-92C, scale 1:250,000, sheet 22H. Deputy Ministry for Mineral Resources, Ministry of Petroleum and Mineral Resources, Kingdom of Saudi Arabia. 32 p.
- Manivit, J., C. Pellaton, D. Vaslet, Y.-M. Le Nindre, J.-M. Brosse, J.-P. Breton, J. Fourniguet and J.-C. Prevot 1985b. Explanatory notes to the geologic map of the Darma' Quadrangle, Kingdom of Saudi Arabia. Geoscience map GM-101C, scale 1:250,000, sheet 24H. Deputy Ministry for Mineral Resources, Ministry of Petroleum and Mineral Resources, Kingdom of Saudi Arabia. 33 p.
- Manivit, J., D. Vaslet, A. Berthiaux, P. Le Strat and J. Fourniguet 1986. Explanatory notes to the geologic map of the Buraydah Quadrangle, Kingdom of Saudi Arabia. Geoscience Map GM-114 C, scale 1:250,000, sheet 26G. Deputy Ministry for Mineral Resources, Ministry of Petroleum and Mineral Resources, Kingdom of Saudi Arabia. 32 p.

- Manivit, J., Y.-M. Le Nindre and D. Vaslet 1990. Histoire géologique de la bordure occidentale de plate-forme Arabe, v. 4: Le Jurassique d'Arabie centrale. Bureau de Recherches Géologiques et Minières, Document no. 194, 560 p., 74 figures, 1 table, 12 plates, 3 appendices.
- Matthews, R.K. and C. Frohlich 1998. Forward modelling of sequence stratigraphy and diagenesis: application to rapid, cost-effective carbonate reservoir characterization. *GeoArabia*, v. 3, no. 3, p. 359-384.
- Matthews, R.K. and C. Frohlich 2002. Maximum flooding surface and sequence boundaries: comparisons between observation and orbital forcing in the Cretaceous and Jurassic (65-190 Ma). *GeoArabia*, v. 7, no. 3, p. 503-538.
- Meyer, F.O., R.C. Price, I.A. Al-Ghamdi, I.M. Al-Goba, S.M. Al-Raimi and J.C. Cole 1996. Sequential stratigraphy of outcropping strata equivalent to Arab-D reservoir, wadi Nisah, Saudi Arabia. *GeoArabia*, v. 1, no. 3, p. 435-456.
- Philip, J.-P., J. Roger, D. Vaslet, F. Cecca, S. Gardin and A.M.S. Memesh 2002. Sequence stratigraphy, biostratigraphy and paleontology of the (Maastrichtian-Paleocene) Aruma Formation in outcrop in Saudi Arabia. *GeoArabia*, v. 7, no. 4, p. 699-718.
- Powers, R.W. 1968. Lexique stratigraphique international. Volume III, Asie, Fas. 10 b1, Arabia Saoudite. Centre Nationale de la Recherche Scientifique, Paris, 177 p.
- Haddad, S.N.S., and M.A. Ameen, 2007. Mid-Turonian-early Campanian sequence stratigraphy of northeast Iraq. *GeoArabia*, v. 12, no. 2, p. 135-176.
- Searle, M.P. 2007. Structural geometry, style and timing of deformation in the Hawasina Window, Al Jabal al Akhdar and Saih Hatat culminations, Oman Mountains. *GeoArabia*, v. 12, no. 2, p. 99-130.
- Sharland, P.R., R. Archer, D.M. Casey, R.B. Davies, S.H. Hall, A.P. Heward, A.D. Horbury and M.D. Simmons 2001. Arabian Plate Sequence Stratigraphy. *GeoArabia Special Publication 2*, Gulf PetroLink, Bahrain, 371 p.
- Simmons, M.D., P.R. Sharland, D.M. Casey, R.B. Davies and O.E. Sutcliffe 2007. Arabian Plate sequence stratigraphy: Potential implications for global chronostratigraphy. *GeoArabia*, v. 12, no. 4, p. 101-130.
- Taher, A.A. 1997. Delineation of organic richness and thermal history of the lower Cretaceous Thamama Group, east Abu Dhabi: a modelling approach for oil exploration. *GeoArabia*, v. 2, no. 1, p. 65-88.
- Vaslet, D., J. Manivit, Y.-M. Le Nindre, J.-M. Brosse, J. Fourniguet and J. Delfour 1983. Explanatory notes to the geologic map of the Wadi Ar Rayn Quadrangle, Kingdom of Saudi Arabia. Geoscience Map GM-63C, scale 1:250,000, sheet 23H. Deputy Ministry for Mineral Resources, Ministry of Petroleum and Mineral Resources, Kingdom of Saudi Arabia. 46 p.
- Vaslet, D., M.S. Al-Muallem, S.S. Maddah, J.-M. Brosse, J. Fourniguet, J.-P. Breton and Y.-M. Le Nindre 1991. Explanatory notes to the geologic map of the Ar Riyad Quadrangle, Kingdom of Saudi Arabia. Geoscience map GM-121, scale 1:250,000, sheet 24I. Deputy Ministry for Mineral Resources, Ministry of Petroleum and Mineral Resources, Kingdom of Saudi Arabia. 54 p.
- van Bellen, R.C., H.V. Dunnington, R. Wetzel and D.M. Morton 1959. Iraq. Lexique Stratigraphique International Centre National de la Recherche Scientifique, III, Asie, Fasc. 10a, Paris, 333 p. Reprinted by *GeoArabia* 2005.
- van Buchem, Frans S.P., Bernard Pittet, Heiko Hillgärten, Jurgen Grötsch, A. Al-Mansouri, Ian M. Billing, Henk H.J. Droste, Heiko Oterdoom and Mia van Steenwinkel 2002. High-resolution sequence stratigraphic architecture of the Barremian-Aptian carbonate systems in northern Oman and the United Arab Emirates Kharaiib and Shu'aiba Formations. *GeoArabia*, v. 7, no. 3, p. 461-500.
- Wilson, A.O. 1985. Depositional and diagenetic facies in the Jurassic Arab-C and -D reservoirs, Qatif field, Saudi Arabia. In, P.O. Roehl and P.W. Choquette (Eds.), *Carbonate Petroleum Reservoirs: A Casebook*. Springer-Verlag, New York, p. 319-340.
- Yousif, S. and G. Nouman 1997. Jurassic geology of Kuwait. *GeoArabia*, v. 2, no. 1, p. 91-110.

The authors welcome comments on the application of the orbital-forcing glacio-eustatic model. For further information about modeling concepts, write to Dr. Robley K. Matthews (RKM@brown.edu). To comment on the regional geology depicted in Figure 2 and the AROS-JK Chart, write to Dr. Moujahed Al-Husseini (geoarabi@batelco.com.bh). For further information about participating in the Arabian Orbital Stratigraphy (AROS) Consortium write to both authors. AROS proposal available upon request. The design and drafting by Heather Paul is appreciated.