

Surface uplift of Tibet and Cenozoic global cooling

Carmala N. Garziona

Department of Earth and Environmental Sciences, University of Rochester, Rochester, New York 14620, USA

Continental weathering on a global scale influences ocean chemistry and imposes a net drawdown of atmospheric CO_2 that modulates global climate (Walker et al., 1981; Berner et al., 1983). This observation, in addition to seawater Sr records that suggest an increase in continental weathering after ca. 40 Ma, led researchers to suggest that the uplift and erosion of the Himalayan-Tibetan orogen over the past 40 m.y. has drawn down atmospheric CO_2 and cooled the globe, leading to the glacial climate that persists today (e.g., Raymo and Ruddiman, 1992; Edmond, 1992). Modeling of tectonic-climate feedbacks associated with increased weathering of the Himalaya and Tibet suggests that two primary biogeochemical processes, silicate weathering and organic carbon burial, can account for the lowering of atmospheric CO_2 necessary to force global cooling (Raymo et al., 1988; Zachos and Kump, 2005). At the time of these inferences, accurate records of both atmospheric $p\text{CO}_2$ and the surface uplift history of the Himalaya and Tibet were lacking. Now, in light of a growing body of atmospheric $p\text{CO}_2$ reconstructions and paleoelevation records, we can test the theory that increased weathering associated with growing mountain belts leads to the drawdown of atmospheric CO_2 and global cooling.

The Eocene-Oligocene transition (EOT) at ca. 34 Ma marks the first major decline in Cenozoic global temperatures and the initiation of Antarctic glaciation. Critical to the argument that continental weathering has caused the drawdown of atmospheric CO_2 is documentation of the temporal relationship between the surface uplift of the Himalayan-Tibetan orogen relative to major cooling events, such as the EOT. Dupont-Nivet et al. (2008, p. 987 in this issue) and other researchers focused on the early elevation history of Tibet (Cyr et al., 2005; Graham et al., 2005; Rowley and Currie, 2006; DeCelles et al., 2007) to provide constraints on the timing and magnitude of the early surface uplift of Tibet (Fig. 1). Rowley and Currie (2006) and DeCelles et al. (2007) use stable isotope paleoaltimetry to show that central Tibet has been at an elevation similar to its modern elevation since ca. 40–26 Ma. Precise age constraints reported by DeCelles et al. (2007) indicate a minimum age of 26 Ma at which high elevations were obtained. However, neither of these studies documents a change in surface elevation that pinpoints the timing of surface uplift in central Tibet. Another paleoelevation study based on stable isotopes in the Hoh Xil basin of north-central Tibet (Cyr et al., 2005) suggests ≤ 2 km paleoelevation at a loosely constrained age of ca. 35–40 Ma. However, the ability to determine the precise timing of surface uplift of the Hoh Xil basin is limited by the poorly constrained ages of Hoh Xil lake deposits, large uncertainties in applying conventional stable isotope paleoaltimetry in northern Tibet (e.g., Quade et al., 2007), and the lack of a recorded increase in paleoelevation in the Hoh Xil basin. Yet another stable isotope study by Graham et al. (2005) documents a regional trend toward more positive $\delta^{18}\text{O}$ values in Eocene to Oligocene time in the southern Tarim basin and western Qaidam basin, north of Tibet, which they attribute to aridification associated with the initial topographic growth of the Tibetan plateau. The lack of good age constraints on these Tarim and Qaidam basin deposits precludes the ability to discern whether this positive shift in $\delta^{18}\text{O}$ values preceded the EOT or simply reflects climate change associated with the EOT.

Dupont-Nivet et al. (2008) present pollen data from well-dated sedimentary deposits in the Xining basin in northeast Tibet to evaluate the regional tectonic versus global climatic influences on the flora of northeast Tibet. The value of the Dupont-Nivet et al. study is that it




Figure 1. Shaded relief map of the Tibetan plateau and surrounding regions, showing localities of Eocene to Oligocene age basins that have yielded quantitative estimates of paleoelevation.

records a transition to cooler-climate plant taxa within a precisely dated succession of lacustrine deposits. The sudden appearance and increasing abundance of conifer pollen (in particular, *Picea*) between 38.3 Ma and 37.3 Ma cannot be accounted for by global cooling because this appearance predates the major cooling trend observed at the EOT. Under warm climate conditions, prior to the EOT, the surface uplift of northeast Tibet provides the simplest explanation for the appearance of cool-climate plant taxa. With the knowledge that distal parts of Tibet, far removed from the collision zone between India and Asia, were significantly elevated (>2 – 3 km) by ca. 38 Ma, it is clear that the surface uplift of vast tracts of Tibet preceded major Cenozoic global cooling. Dupont-Nivet et al. conclude that the surface uplift of central and northern Tibet, prior to the EOT, is therefore consistent with the idea that erosion and silicate weathering associated with large mountain belts lowers atmospheric CO_2 , thereby forcing global cooling.

The relation of surface uplift histories to records of continental weathering and $p\text{CO}_2$ is important in understanding the role of mountain belts in the drawdown of atmospheric CO_2 . Specifically, does the new constraint on the timing of surface uplift of northeast Tibet from Dupont-Nivet et al. (2008) correspond with an increase in continental weathering and a concomitant decrease in $p\text{CO}_2$? Over the Cenozoic, seawater $^{87}\text{Sr}/^{86}\text{Sr}$ shows a marked increase, beginning after ca. 40 Ma (Fig. 2), that has been commonly attributed to erosion of the Himalayan-Tibetan orogen (Edmond, 1992; Richter et al., 1992). Interestingly, the greatest change in the rate of $^{87}\text{Sr}/^{86}\text{Sr}$ increase occurs at ca. 38 Ma at the precise time that northeast Tibet experienced surface uplift based on the appearance of conifer pollen (Fig. 2). Over the entire Cenozoic, multiple proxies for atmospheric $p\text{CO}_2$ indicate a decrease over the middle Cenozoic, from values of several thousand ppmV in the early Cenozoic to values of <500 ppmV by ca. 25 Ma (Fig. 2). During the time period of 40–25 Ma, the only proxy for atmospheric $p\text{CO}_2$ comes from alkenones, and these data suggest the most dramatic decline in $p\text{CO}_2$ occurred between ca. 37 and 25 Ma (Pagani et al., 2005). The close




Figure 2. Cenozoic seawater Sr and paleoatmospheric CO₂ in ppmV (parts per million by volume). Sr curve is shown as black line (McArthur et al., 2001) inverted for comparison to the pCO₂ curve. Compilation of Cenozoic atmospheric pCO₂ proxy records (modified from Tiple and Pagani, 2007). Red circles with error estimates are from boron isotope records (Pearson and Palmer, 2000), yellow circles with error bars are from pedogenic carbonate records (Ekart et al., 1999; Royer et al., 2004), and the blue field shows the upper and lower estimates from alkenone records (Pagani et al., 1999, 2005). Timing of the Eocene-Oligocene transition (EOT) is labeled, and timing of the first appearance of conifers (inferred to reflect surface uplift) in northeast Tibet is labeled with a green bar.

correspondence between the seawater Sr curve and pCO₂ supports the inference that continental weathering imparts a significant effect on the drawdown of atmospheric CO₂. The observation that at least northeast Tibet had attained moderate to high elevations by ca. 38 Ma implicates the Himalayan-Tibetan orogen as the source of continental erosion.

What additional information can help to resolve the question “What is the role of the surface uplift and erosion of Tibet on the drawdown of atmospheric CO₂ over the Cenozoic?” Additional paleoelevation studies aimed at understanding both the timing and spatial extent of early surface uplift of the Himalaya-Tibetan orogen are critical to evaluating the role of Tibet in Cenozoic global cooling. Better resolution of the sedimentation records in marginal marine basins will provide more accurate reconstructions of the sedimentary flux from the Himalaya and Tibet (e.g., Clift, 2006) to evaluate whether long-term erosion rates were sufficiently high to impart a global effect on atmospheric CO₂. Recent studies of riverine Sr flux (Bickle et al., 2005, and references therein) have established a solid understanding of the relative contribution of silicate weathering versus carbonate weathering to the riverine flux of Sr to the oceans, and the conclusion for the Himalaya is that approximately half of dissolved Sr in Himalayan rivers is derived from silicates (Bickle et al., 2005). A remaining challenge is to understand the relative contribution of silicate weathering in the Himalayan-Tibetan orogen within the global silicate weathering budget.

REFERENCES CITED

Berner, R.A., Lasaga, A.C., and Garrels, R.M., 1983, The carbonate-silicate geochemical cycles and its effect on atmospheric carbon dioxide over the past 100 million years: *American Journal of Science*, v. 283, p. 641–683.
 Bickle, M.J., Chapman, H.J., Bunbury, J., Harris, N.B.W., Fairchild, I.J., Ahmad, T., and Pomés, C., 2005, Relative contributions of silicate and carbonate

rocks to riverine Sr fluxes in the headwaters of the Ganges: *Geochimica et Cosmochimica Acta*, v. 69, p. 2221–2240, doi: 10.1016/j.gca.2004.11.019.
 Clift, P.D., 2006, Controls on the erosion of Cenozoic Asia and the flux of clastic sediment to the ocean: *Earth and Planetary Science Letters*, v. 241, p. 571–580, doi: 10.1016/j.epsl.2005.11.028.
 Cyr, A., Currie, B.S., and Rowley, D.B., 2005, Geochemical and stable isotopic evaluation of Fenghuoshan Group lacustrine carbonates, north-central Tibet: Implications for the paleoaltimetry of Late Eocene Tibetan Plateau: *The Journal of Geology*, v. 113, p. 517–533, doi: 10.1086/431907.
 DeCelles, P.G., Quade, J., Kapp, P., Fan, M., Dettman, D.L., and Lin, D., 2007, High and dry in central Tibet during the Late Oligocene: *Earth and Planetary Science Letters*, v. 253, p. 389–401, doi: 10.1016/j.epsl.2006.11.001.
 Dupont-Nivet, G., Hoorn, C., and Konert, M., 2008, Tibetan uplift prior to the Eocene-Oligocene climate transition: Evidence from pollen analysis of the Xining Basin: *Geology*, v. 36, p. 987–990, doi: 10.1130/G25063A.1.
 Edmond, J.M., 1992, Himalayan tectonics, weathering processes and the strontium isotope record in marine limestones: *Science*, v. 258, p. 1594–1597, doi: 10.1126/science.258.5088.1594.
 Ekart, D.D., Cerling, T.E., Montañez, I.P., and Tabor, N.J., 1999, A 400 year carbon isotope record of pedogenic carbonate: Implications for paleoatmospheric carbon dioxide: *American Journal of Science*, v. 299, p. 805–827, doi: 10.2475/ajs.299.10.805.
 Graham, S.A., Chamberlain, C.P., Yue, Y.J., Ritts, B.D., Hanson, A.D., Horton, T.W., Waldbauer, J.R., Poage, M.A., and Feng, X., 2005, Stable isotope records of Cenozoic climate and topography, Tibetan plateau and Tarim basin: *American Journal of Science*, v. 305, p. 101–118, doi: 10.2475/ajs.305.2.101.
 McArthur, J.M., Howarth, R.J., and Bailey, T.R., 2001, Strontium isotope stratigraphy: LOWESS version 3: Best fit to the marine Sr-isotope curve for 0–509 Ma and accompanying look-up table for deriving numerical age: *The Journal of Geology*, v. 109, p. 155–170, doi: 10.1086/319243.
 Pagani, M., Arthur, M.A., and Freeman, K.H., 1999, Late Miocene atmospheric CO₂ concentrations and the expansion of C₄ grass: *Science*, v. 285, p. 876–878, doi: 10.1126/science.285.5429.876.
 Pagani, M., Zachos, J., Freeman, K.H., Tiple, B.J., and Bohaty, S., 2005, Marked change in atmospheric carbon dioxide concentrations during the Oligocene: *Science*, v. 309, p. 600–603, doi: 10.1126/science.1110063.
 Pearson, P.N., and Palmer, M.R., 2000, Atmospheric carbon dioxide concentrations over the last 60 million years: *Nature*, v. 406, p. 695–699, doi: 10.1038/35021000.
 Quade, J., Garzzone, C., and Eiler, J., 2007, Paleoelevation reconstructions using pedogenic carbonates: *Reviews in Mineralogy and Geochemistry*, v. 66, p. 53–87, doi: 10.2138/rmg.2007.66.3.
 Raymo, M.E., Ruddiman, W.F., and Froelich, P.N., 1988, Influence of late Cenozoic mountain building on ocean geochemical cycles: *Geology*, v. 16, p. 649–653, doi: 10.1130/0091-7613(1988)016<0649:IOLCMB>2.3.CO;2.
 Raymo, M.E., and Ruddiman, W.F., 1992, Tectonic forcing of late Cenozoic climate: *Nature*, v. 359, p. 117–122, doi: 10.1038/359117a0.
 Richter, F.M., Rowley, D.B., and DePaolo, D.J., 1992, Sr isotope evolution of seawater: The role of tectonics: *Earth and Planetary Science Letters*, v. 109, p. 11–23, doi: 10.1016/0012-821X(92)90070-C.
 Rowley, D.B., and Currie, B.S., 2006, Palaeo-altimetry of the late Eocene to Miocene Lunpola basin, central Tibet: *Nature*, v. 439, p. 677–681, doi: 10.1038/nature04506.
 Royer, D.L., Berner, R.A., Montañez, I.P., Tabor, N.J., and Beerling, D.J., 2004, CO₂ as a primary driver of Phanerozoic climate change: *GSA Today*, v. 14, p. 4–10, doi: 10.1130/1052-5173(2004)014<4:CAAPDO>2.0.CO;2.
 Tiple, B.J., and Pagani, M., 2007, The early origins of C₄ photosynthesis: *Annual Review of Earth and Planetary Sciences*, v. 35, p. 435–461, doi: 10.1146/annurev.earth.35.031306.140150.
 Walker, J.C.G., Hays, P.B., and Kasting, J.F., 1981, A negative feedback mechanism for the long-term stabilization of Earth's surface temperature: *Journal of Geophysical Research*, v. 86, p. 9776–9782, doi: 10.1029/JC086iC10p09776.
 Zachos, J.C., and Kump, L.R., 2005, Carbon cycle feedbacks and the initiation of Antarctic glaciation in the earliest Oligocene: *Global and Planetary Change*, v. 47, p. 51–66, doi: 10.1016/j.gloplacha.2005.01.001

Printed in USA