

Network Analysis of Sport-Related Concussion Research During the Past Decade (2010–2019)

Shawn R. Eagle, PhD, ATC*; Anthony P. Kontos, PhD*; Micky W. Collins, PhD*; Chris Connaboy, PhD†; Shawn D. Flanagan, PhD†

*Department of Orthopaedic Surgery and †Neuromuscular Research Laboratory, University of Pittsburgh, PA

Context: Research into sport-related concussion (SRC) has grown substantially over the past decade, yet no authors to date have synthesized developments over this critical time period.

Objective: To apply a network-analysis approach in evaluating trends in the SRC literature using a comprehensive search of original, peer-reviewed research articles involving human participants published between January 1, 2010, and December 15, 2019.

Design: Narrative review.

Main Outcome Measure(s): Bibliometric maps were derived from a comprehensive search of all published, peer-reviewed SRC articles in the Web of Science database. A clustering algorithm was used to evaluate associations among journals, organizations or institutions, authors, and key words. The online search yielded 6130 articles, 528 journals, 7598 authors, 1966 organizations, and 3293 key words.

Results: The analysis supported 5 thematic clusters of journals: (1) biomechanics/sports medicine (n = 15), (2) pediatrics/rehabilitation (n = 15), (3) neurotrauma/neurology/neurosurgery (n = 11), (4) general sports medicine (n = 11), and

(5) neuropsychology (n = 7). The analysis identified 4 organizational clusters of hub institutions: (1) University of North Carolina (n = 19), (2) University of Toronto (n = 19), (3) University of Michigan (n = 11), and (4) University of Pittsburgh (n = 10). Network analysis revealed 8 clusters for SRC key words, each with a central topic area: (1) epidemiology (n = 14), (2) rehabilitation (n = 12), (3) biomechanics (n = 11), (4) imaging (n = 10), (5) assessment (n = 9), (6) mental health/chronic traumatic encephalopathy (n = 9), (7) neurocognition (n = 8), and (8) symptoms/impairments (n = 5).

Conclusions: The findings suggest that during the past decade, SRC research has (1) been published primarily in sports medicine, pediatric, and neuro-focused journals, (2) involved a select group of researchers from several key institutions, and (3) concentrated on new topical areas, including treatment or rehabilitation and mental health.

Key Words: mild traumatic brain injury, assessment, rehabilitation

Key Points

- Our findings represent the first network analysis of sport-related concussion (SRC) research and suggest that during the past decade, this literature has been published primarily in sports medicine, pediatric, and neuro-focused journals.
- Research on SRCs has emphasized epidemiology, assessment, and return to play, as well as emerging areas including rehabilitation and mental health.
- This network analysis of SRC research highlighted its benefits for identifying appropriate journals for article submissions, potential authors and institutions with whom to collaborate, and topics reflecting the direction in which the field is headed.

Sport-related concussion (SRC) continues to be a significant health concern affecting millions of athletes of all ages and levels, resulting in an estimated US economic burden up to \$17 billion annually. This heterogeneous injury involves myriad signs, symptoms, and impairments, and although many athletes recover within a few weeks,¹ some take months or longer.² Given the large number of affected athletes and burden of this injury, the field of SRC research has experienced tremendous growth during the past decade. In fact, a search of the term *sport-related concussion* in PubMed on January 24, 2020, yielded 1822 peer-reviewed articles published during the previous 10 years. This work has enhanced our empirical and clinical knowledge of SRC in areas such as epidemiology, identification, assessment, recovery, and rehabilitation.³ However, investigators have yet to empir-

ically synthesize developments and trends in SRC research during this period of rapid growth.

Research and clinical practice in concussion have evolved substantially since the late 1990s and early 2000s, when the literature emphasized SRC identification by focusing on grading scales, symptoms, and the role of loss of consciousness.⁴ As the field evolved from 2005 to 2010, the focus shifted to biomechanics, assessments of balance and cognitive performance, age and sex differences, recovery time, rest-based intervention, and consensus statements designed to inform clinical care.⁵ More recently, SRC research has progressed to include ocular and vestibular assessments, advanced neuroimaging, fluid biomarkers, clinical subtypes or profiles, and rehabilitation.^{3,6,7} During the past decade, the paradigm of prescribed rest as the primary or sole management strategy for SRC

was challenged by empirical evidence that supported more-active approaches.⁸ The field has also begun to shift toward earlier, more-active interventions that target specific symptoms and impairments.³ Empirical evidence has provided the basis for these theoretical and clinical paradigm shifts and will continue to do so as the work moves forward. However, the field of SRC research has lacked introspective analyses of past research trends to help guide future paradigms and approaches to clinical care. An empirical analysis of trends in SRC research during the past decade could inform future lines of inquiry and areas for clinical advancement.

Network analysis provides a useful tool for objectively assessing the bibliographic trends of a scientific field.⁹ On the basis of the outcome of interest (eg, journals, institutions, and authors) network analysis produces an interconnected bibliometric map from associations among outcomes.^{9,10} Given the rapid growth in SRC research over the previous 10 years, this method could offer a greater understanding of recent trends that may otherwise go unnoticed, including journals, key research institutions, and authors. This information could inform a foundational basis for future studies and collaborations.

The purpose of our study was to evaluate trends in the SRC literature using network analysis based on a comprehensive search of original, peer-reviewed research articles involving human participants published between January 1, 2010, and December 15, 2019. Specifically, we used network analysis to evaluate associations in the following areas: (1) peer-reviewed journals, (2) organizations and institutions, (3) authors, and (4) relevant key words.

METHODS

We identified articles in the Web of Science Core Collection using key terms to encompass all forms of SRC (eg, *concussion*, *sport-related concussion*, *cerebral concussion*) or mild traumatic brain injury (mTBI; eg, *mild traumatic brain injury*, *mild brain injury*, *mild TBI*) research from 2010 to 2019 that involved human participants. The search was conducted on December 15, 2019. The search was delimited to the previous decade to focus on recent trends in SRC research. The search was further refined to only original articles, reducing the total number to 6130 from 10086. All bibliometric data, including title, authors, organizations, abstract, key words, and bibliography, were imported into network-analysis software (version 1.6.13; VOSviewer, Leiden University, Leiden, The Netherlands).⁹ A detailed mathematical description of the software and how it applies a variant of a multidimensional scaling algorithm can be found in a report by van Eck et al.⁹ Search terms and their variants were removed from the analysis. Clusters (ie, groups of interconnected nodes) were derived for each map using a resolution of 1.0, attraction of 2, repulsion of 1, minimum cluster size of 5, and resolution of 2.¹¹ Clusters were subjectively named on the basis of the overarching themes of the individual cluster groups. With regard to interpreting the maps, connecting lines indicate an association between nodes. More lines indicate more dense connections between the given node and its connecting nodes.⁹ Similarly, closer proximity

between 2 nodes reflects a stronger association than 2 nodes at a farther distance.⁹ Central location of a node indicates more connections, whereas more distal locations indicate fewer connections.

The online search yielded 6130 articles, 528 journals, 1966 organizations, 7598 authors, and 3293 key words. *Citation analyses*, or assessments of the number of times journals, organizations, or authors cite one another, were conducted to analyze journals, organizations, and authors using specific minimum inclusion criteria: 5 publications for journals, 5 publications for organizations, and 10 publications for authors. Outcomes for each analysis were the number of articles, citations, total link strength, and clusters. For organization- and author-cluster reporting, the *hub*, or the most highly interconnected organization or individual author, is reported as the cluster name. To analyze the top key words in concussion research, a *co-occurrence analysis* (ie, assessment of the number of articles in which multiple key words occur together) was performed with a co-occurrence minimum of 10 for inclusion. Variants of the search terms and populations were removed from the analysis in order to investigate the key topics that have been evaluated. Outcomes for key-word analysis are occurrences, total link strength, and clusters; the top 10 (ordered by most SRC publications within the study period) were also analyzed for journals, organizations, authors, and key words using full counting.

RESULTS

The number of publications, citations, and total link strength for the top 10 journals, organizations, and authors as well as the number of occurrences, citations, and total link strength for the top key words from 2010 to 2019 are presented in Table 1. The top 10 journals, organizations, and authors accounted for 48.1%, 34.1%, and 20.3% of the total SRC publications, respectively (journals = 59, organizations = 50, authors = 148). The top 10 key words from 2010 to 2019 accounted for 32.3% of the total occurrences ($n = 78$). The top 5 journals per cluster are shown in Table 2.

Network analysis revealed 5 journal clusters (categories) for SRC research (Figure): (1) biomechanics/sports medicine ($n = 15$), (2) pediatrics/rehabilitation ($n = 15$), (3) “neuro focused” (eg, neurology, neurotrauma, neurosurgery; $n = 11$), (4) general sports medicine ($n = 11$), and (5) neuropsychology ($n = 7$). Among institutions that published SRC research, 4 clusters were identified (Figure B): (1) University of North Carolina ($n = 19$), (2) University of Toronto ($n = 19$), (3) University of Michigan ($n = 11$), and (4) University of Pittsburgh ($n = 10$). Among authors who published SRC research, 7 clusters were identified (Figure C): (1) Kontos ($n = 32$), (2) Iverson ($n = 27$), (3) McCrea ($n = 27$), (4) Broglio ($n = 25$), (5) Kerr ($n = 16$), (6) Kroshus ($n = 14$), and (7) Guskiewicz ($n = 7$). Network analysis revealed 8 key-word clusters for SRC research published from 2010 to 2019 (Figure D): (1) epidemiology ($n = 14$), (2) rehabilitation ($n = 12$), (3) biomechanics ($n = 11$), (4) imaging ($n = 10$), (5) assessment ($n = 9$), (6) mental health/chronic traumatic encephalopathy ($n = 9$), (7) neurocognition ($n = 8$), and (8) symptoms/impairments ($n = 5$).

Table 1. Overall Top 10 Journals, Organizations, Authors, and Key Words in Sport-Related Concussion Publications, 2010–2019

Variable	Articles	Citations	Link Strength
Journal			
<i>American Journal of Sports Medicine</i>	115	4763	2516
<i>Archives of Clinical Neuropsychology</i>	43	692	594
<i>Brain Injury</i>	121	1397	1125
<i>British Journal of Sports Medicine</i>	87	3426	1472
<i>Clinical Journal of Sport Medicine</i>	112	2250	1428
<i>Frontiers in Neurology</i>	44	204	474
<i>Journal of Athletic Training</i>	132	2530	1701
<i>Journal of Head Trauma Rehabilitation</i>	46	743	582
<i>Journal of Neurotrauma</i>	145	2906	1445
<i>Journal of Science and Medicine in Sport</i>	60	591	455
Organization			
Boston Children's Hospital	74	1269	2935
Harvard Medical School	83	441	1961
Harvard University	81	2699	3068
Medical College of Wisconsin	72	1426	2478
University of Calgary	95	1587	2446
University of Michigan	101	2190	2646
University of North Carolina	155	3503	4181
University of Pittsburgh	108	2975	3254
University of Toronto	135	1790	2049
University of Washington	88	1268	2232
Author			
Collins	46	2030	2460
Comstock	51	1761	1795
Covassin	56	1629	2291
Guskiewicz	76	2997	3460
Kerr	68	1245	2336
Kontos	57	1805	2569
Marshall	52	1437	1992
McCrea	55	1398	2562
Meehan/Broglio (tie)	61/61	1779/1786	2420/2804
Mihalik	53	1020	1859
Key Words			
	Occurrences	Citations (Average)	Link Strength
Balance	51	12.8	60
Biomechanics	55	17.9	25
Chronic traumatic encephalopathy	53	17.7	19
Epidemiology	82	18.9	25
Impact	73	19.3	53
Postconcussion syndrome	49	26.8	37
Recovery	41	16.7	39
Rehabilitation	46	13.6	42
Return to play	70	20.5	46
Symptoms	50	33.4	52

DISCUSSION

We are the first to use network analysis to evaluate trends in SRC literature among peer-reviewed journals, institutions, authors, and key words. The results demonstrate that SRC research has been published primarily in sports medicine and secondarily in pediatric and neuro-focused (ie, combined neurology, neurosurgery, neurotrauma) journals. Four key, highly interconnected institutions—the University of North Carolina, University of Toronto, University of Michigan, and University of Pittsburgh—disseminated a disproportionate amount of SRC research (Figure B). Seven author clusters were also identified, with

many of the author hubs located at the key institutions (Figure C). Eight key-word clusters were identified (Figure D), suggesting that epidemiology, assessment evaluation, and return to play remain central themes, with rehabilitation and mental health concerns emerging as newer focus areas.

Journals

Given that the overarching topic of the current analysis was concussion in sport, it was not surprising that sports medicine journals such as the *Journal of Athletic Training*, *American Journal of Sports Medicine*, *Clinical Journal of*

Downloaded from <http://meridian.allenpress.com/jat/article-pdf/56/4/398/3019823/1062-6050-56-4-398.pdf> by guest on 16 July 2024

Table 2. Top 5 Journals Within Each Cluster of Sport-Related Concussion Publications, 2010–2019

Cluster	Journals	Publications	Citations	Total Link Strength
Biomechanics/sports medicine	<i>British Journal of Sports Medicine</i>	87	3426	1472
	<i>Journal of Science and Medicine in Sport</i>	60	591	455
	<i>Medicine & Science in Sports & Exercise</i>	43	1163	761
	<i>Annals of Biomedical Engineering</i>	29	756	347
	<i>Journal of Sport Rehabilitation</i>	23	82	121
Pediatrics/rehabilitation	<i>Clinical Journal of Sport Medicine</i>	112	2250	1428
	<i>Journal of Head Trauma Rehabilitation</i>	46	743	582
	<i>Orthopedic Journal of Sports Medicine</i>	42	276	329
	<i>Pediatrics</i>	42	2091	1017
Neuro focused	<i>Physician and Sportsmedicine</i>	39	256	356
	<i>Journal of Neurotrauma</i>	145	2906	1445
	<i>Frontiers in Neurology</i>	44	204	474
	<i>Journal of Neurosurgery: Pediatrics</i>	33	602	542
	<i>PLoS One</i>	31	552	276
General sports medicine	<i>Neurosurgical Focus</i>	24	505	339
	<i>Journal of Athletic Training</i>	132	2530	1701
	<i>American Journal of Sports Medicine</i>	115	4763	2516
	<i>Sports Health</i>	36	246	359
	<i>International Journal of Athletic Therapy and Training</i>	23	22	143
Neuropsychology	<i>Journal of Neurological Sciences</i>	14	501	221
	<i>Brain Injury</i>	121	1397	1125
	<i>Archives of Clinical Neuropsychology</i>	43	692	594
	<i>Journal of the International Neuropsychology Society</i>	28	605	417
	<i>Applied Neuropsychology: Child</i>	26	149	311
	<i>Journal of Clinical and Experimental Neuropsychology</i>	25	291	199

Sport Medicine, and *British Journal of Sports Medicine* were primary outlets for SRC research (Tables 1 and 2). The largest journal cluster ($n = 15$) was biomechanics/sports medicine. As motion-analysis and engineering technology has progressed over the past decade, this technology has been applied to SRC research to investigate topics such as the role of subconcussive impacts,¹² a force threshold for sustaining an SRC,¹³ sex differences in neck and head control,^{14,15} and optimizing helmets for SRC prevention and identification.¹⁶ Pediatrics/rehabilitation was tied for the largest cluster ($n = 15$), with the most peer-reviewed publications during the study period (478, 19% more than the next-highest cluster). This body of research consisted of risk-factor identification for prolonged recovery or persistent post-SRC symptoms, such as initial symptom burden, preinjury mood disorders, personal or family history of migraine, or vision or vestibular dysfunction preinjury or postinjury.^{3,17,18} This work provided important contributions to our clinical understanding of SRC because clinician knowledge about preexisting risk factors can help guide management and intervention strategies. Additional management strategies and safe return-to-play or return-to-school guidelines were also a primary focus of this research cluster, reflecting the emphasis on pediatrics.^{19–21}

The third largest cluster was neuro focused, which addressed an enhanced clinical understanding of SRC heterogeneity, treatment efficacy, and assessment of objective biomarkers (eg, cerebral blood flow, magnetic resonance imaging, diffusion tensor imaging, and blood biomarkers; Tables 1 and 2).^{22,23} Specifically, the classification of different SRC presentations into “profiles” or “subtypes” and the development of tools to identify these different SRC types were in this cluster.⁸ The purpose of identifying SRC profiles is to provide a framework for interventions that target individual responses to SRC.²⁴

General sports medicine was the fourth largest cluster and focused on validation and implementation of assessments such as balance and postural stability, gait, vestibular and ocular impairment, and neurocognitive tests for identification and prognostic utility. Sideline assessments, such as the Sideline Concussion Assessment Tool and symptom-reporting scales, were also emphasized in this cluster. The final journal cluster was neuropsychology ($n = 7$). A focus of this cluster was neurocognitive and neuropsychological testing because these measures are commonly used to inform the clinical diagnosis.^{25,26} This cluster also highlighted symptom-reporting measures and their diagnostic and prognostic value to the clinician. Although the topical areas in this cluster were broad, the core was research centered on improving the neuropsychologist’s assessment and management of patients with SRC.

Institutions and Authors

A select group of key institutions and authors acted as the primary SRC research hubs over the past decade (Table 1 and Figure B–D). The University of North Carolina has published SRC research in topical areas such as knowledge of and attitudes toward SRC among players and health care professionals, biomechanics of head impacts, and balance assessments over the past decade.^{13,27–29} Epidemiology has also been a focus of this group.^{30,31} The University of Toronto has addressed the potential role of imaging and other diagnostic tools for identifying concussion and persistent symptoms, as well as pediatric concussion.^{32,33} The University of Pittsburgh was another hub ($n = 10$) identified in this analysis. Its authors emphasized a clinical profile approach to SRC management, risk factors, and vestibular and ocular assessment and impairment after the injury.^{6,7,34} This group has also focused on sex differences in presentation and recovery, clinical assessments, and




Figure. A, Network map of journals that published sport-related concussion research from 2010–2019. Five clusters were identified: (1) biomechanics/sports medicine (red), (2) pediatrics/rehabilitation (green), (3) neurotrauma/neurology/neurosurgery (blue), (4) general sports medicine (yellow), and (5) neuropsychology (purple). B, Network map of organizations that published sport-related concussion research from 2010–2019. Four clusters were identified (named for the primary hub): (1) University of North Carolina (red), (2) University of Toronto (green), (3) University of Michigan (blue), and (4) University of Pittsburgh (yellow). C, Network map of authors who published sport-related concussion research from 2010–2019. Seven clusters were identified (named for the primary hub): (1) Kontos (red), (2) Iverson (green), (3) McCrea (blue), (4) Broglio (yellow), (5) Kroshus (light blue), (6) Kerr (purple), and (7) Guskiewicz (orange). D, Network map of key words used in sport-related concussion publications in peer-reviewed journals from 2010–2019. Eight clusters were identified 1. epidemiology (red), 2. rehabilitation (green), 3. biomechanics (navy blue), 4. imaging (yellow), 5. assessment (purple), 6. mental health/chronic traumatic encephalopathy (light blue), 7. neurocognition (orange), 8. symptoms/impairments (brown).

predictors of prolonged recovery from SRC.^{35,36} The University of Michigan, the final hub (n = 11) identified in this analysis, published research on the utility and efficacy of clinical assessments³⁷ and the role of biomechanics/head impacts.³⁸ Together, these institutions represent the most prolific hubs of SRC research over the previous decade.

Key Words

As the largest cluster and top co-occurring key word, *epidemiology* was a primary area of SRC research over the decade, with close associations to knowledge, education, and prevention (Tables 1 and 2; Figure D). Rehabilitation

was the second-largest cluster, with 3 of the top 10 most co-occurring key words (eg, *balance*, *rehabilitation*, *recovery*; Table 1). The rehabilitation cluster was associated with key words from research on balance assessments, gait, and physical activity, among other topics. Similarly, the biomechanics cluster included research on head impacts and acceleration. Imaging and assessment were the only 2 clusters not represented by a key word in the top 10 most co-occurring key words of the decade, which may be related to the broad nature of their associations (Figure D). The imaging cluster encompassed research using imaging modalities, such as magnetic resonance imaging, as well as investigations of neuropsychology and cognition. The inclusion of neuropsychology and cognition in this cluster

was likely related to neurocognitive metrics being the clinical assessment of choice during imaging. The assessment cluster was linked with research that used key words such as *video* and *vision analysis*, among others.

During the last decade, research on SRC has also emphasized the long-term effects of the injury, as evidenced by the mental health/chronic traumatic encephalopathy cluster's inclusion of 3 of the top 10 key words: *return to play*, *postconcussion syndrome*, and *chronic traumatic encephalopathy* (Table 1). This cluster also included *anxiety* and *depression*, 2 mental health topics that are increasingly viewed as important areas of focus in SRC research. The co-occurrence of these key words is possibly related to increased public concern over the association of SRC with the development of neurodegenerative diseases or mood disorders later in life. The neurocognition cluster included *impact* as a top 10 key word, which may reflect the commonly used computerized neurocognitive test (ie, Immediate Post-concussion Assessment and Cognitive Testing [ImPACT Applications, Inc]) or head impacts or impact exposure. Finally, the symptoms/impairments cluster contained typical impairments associated with the injury, such as vestibular, oculomotor, and neurocognitive impairments, as well as general symptoms.

Strengths and Limitations

Our findings contribute to the broader concussion literature by describing the relationships among topical areas (ie, journal clusters) and individual journals that often publish research on those topics. This study provides a useful resource for clinicians and researchers alike to identify the top peer-reviewed information sources on their topic of interest (Table 2). The key-word analysis (Table 1 and Figure D) supplies our major contribution to the literature in revealing the co-occurrences, which can elucidate trends during this period of enormous growth in SRC publications. For example, *rehabilitation* being the 9th-ranked key word of the decade is a critical finding in our opinion (Table 1). This result is important, given that approximately 15 to 20 years ago, SRC was still predominantly viewed as a homogeneous injury that required a “one-size-fits-all” treatment approach.³⁹ Thus, rehabilitation would likely not have been a central focus of research during that time because everyone with concussion was usually treated with the same approach. Researchers can build on the key findings of the past decade to initiate more-informed therapeutic trials to enhance our understanding of the efficacy of certain rehabilitation strategies. To maximize the potential of the next decade, SRC investigators can use the foundational research we presented to identify the most pertinent areas for future inquiry.

Another critical aspect of informing treatment practice is the mitigation of potential long-term effects of SRC, such as postconcussion syndrome and chronic traumatic encephalopathy. Our key-word analysis of the past decade indicated substantial interest in these long-term effects (Table 1). *Recovery* was also a top key word. The significant co-occurrence of these key words was possibly related to the increased public concern over an SRC history and the development of neurodegenerative diseases later in life.⁴⁰ Potential long-term effects of concussion are another area of SRC research that can benefit from intervention

studies to obtain high-quality evidence. Our results can be used by clinicians and investigators to encourage collaboration and inform future lines of research in this area, among others.

The limitations to this type of analysis are worth noting. The study was delimited to SRC research published during 2010–2019, so the findings presented here do not reflect concussion research before the inclusion date or in other populations (eg, military, accidents, assaults). In addition, each cluster and key word could not be covered exhaustively due to space constraints. Therefore, the summary of journal clusters and key words is informed by our subjective decision making based on the results of the analysis. The software used in the network analysis is limited to word detection and cannot detect capitalizations of words. As such, the key word *impact* could be referring to a physical contact that may have resulted in a concussion or a popular computerized neurocognitive test (ie, ImPACT). Given the close proximity of *impact* and *neurocognitive testing* in the network map (Figure 1D), we suspect it was the latter, but this assumption cannot be determined definitively. Another limitation involved the space constraints in the network maps that resulted in certain nodes not having labels.

CONCLUSIONS

Our understanding of SRC identification, assessment, and rehabilitation has grown considerably over the past decade. The current findings represent the first network analysis of SRC research and suggest that, during the past decade, SRC research has (1) been published primarily in sports medicine, pediatric, and neuro-focused journals, (2) involved several key institutions within a broad field, and (3) centered on epidemiology, assessment, and return to play, as well as emerging areas including rehabilitation and mental health. This study also highlights the benefits of using a network analysis of SRC research to identify appropriate journals for article submissions, potential authors and institutions with whom to collaborate, and topics reflecting the directions in which the field is headed.

REFERENCES

1. Harmon KG, Clugston JR, Dec K, et al. American Medical Society for Sports Medicine position statement on concussion in sport. *Br J Sports Med.* 2019;53(4):213–225. doi:10.1136/bjsports-2018-100338
2. Thomas DJ, Coxe K, Li H, et al. Length of recovery from sports-related concussions in pediatric patients treated at concussion clinics. *Clin J Sport Med.* 2018;28(1):56–63. doi:10.1097/JSM.0000000000000413
3. Harmon KG, Clugston JR, Dec K, et al. American Medical Society for Sports Medicine position statement on concussion in sport. *Br J Sports Med.* 2019;53(4):213–225.
4. McCrory P, Johnston K, Meeuwisse W, et al. Summary and agreement statement of the 2nd International Conference on Concussion in Sport, Prague 2004. *Br J Sports Med.* 2005;39(4):196–204. doi:10.1136/bjism.2005.018614
5. McCrory P, Meeuwisse W, Johnston K, et al. Consensus Statement on Concussion in Sport: the 3rd International Conference on Concussion in Sport held in Zurich, November 2008. *Br J Sports Med.* 2009;43(suppl 1):i76–i90. doi:10.1136/bjism.2009.058248
6. Mucha A, Collins MW, Elbin R, et al. A brief vestibular/ocular motor screening (VOMS) assessment to evaluate concussions:

- preliminary findings. *Am J Sports Med.* 2014;42(10):2479–2486. doi:10.1177/0363546514543775
7. Kontos AP, Sufrinko A, Sandel N, Emami K, Collins MW. Sport-related concussion clinical profiles: clinical characteristics, targeted treatments, and preliminary evidence. *Curr Sports Med Rep.* 2019;18(3):82–92. doi:10.1249/JSR.0000000000000573
 8. Collins MW, Kontos AP, Reynolds E, Murawski CD, Fu FH. A comprehensive, targeted approach to the clinical care of athletes following sport-related concussion. *Knee Surg Sports Traumatol Arthrosc.* 2014;22(2):235–246. doi:10.1007/s00167-013-2791-6
 9. van Eck NJ, Waltman L, Dekker R, van den Berg J. A comparison of two techniques for bibliometric mapping: multidimensional scaling and VOS. *J Am Soc Inf Sci Technol.* 2010;61(12):2405–2416. doi:10.1002/asi.21421
 10. van Eck NJ, Waltman L, van Raan AF, Klautz RJ, Peul WC. Citation analysis may severely underestimate the impact of clinical research as compared to basic research. *PLoS One.* 2013;8(4):e62395. doi:10.1371/journal.pone.0062395
 11. Bullmore E, Sporns O. Complex brain networks: graph theoretical analysis of structural and functional systems. *Nat Rev Neurosci.* 2009;10(3):186–198. doi:10.1038/nrn2575
 12. Beckwith JG, Greenwald RM, Chu JJ, et al. Head impact exposure sustained by football players on days of diagnosed concussion. *Med Sci Sports Exerc.* 2013;45(4):737–746. doi:10.1249/MSS.0b013e3182792ed7
 13. Guskiewicz KM, Mihalik JP. Biomechanics of sport concussion: quest for the elusive injury threshold. *Exerc Sport Sci Rev.* 2011;39(1):4–11. doi:10.1097/JES.0b013e318201f53e
 14. Lynall RC, Clark MD, Grand EE, et al. Head impact biomechanics in women's college soccer. *Med Sci Sports Exerc.* 2016;48(9):1772–1778. doi:10.1249/MSS.0000000000000951
 15. McCuen E, Svaldi D, Breedlove K, et al. Collegiate women's soccer players suffer greater cumulative head impacts than their high school counterparts. *J Biomech.* 2015;48(13):3720–3723. doi:10.1016/j.jbiomech.2015.08.003
 16. McGuine TA, Hetzel S, McCrea M, Brooks MA. Protective equipment and player characteristics associated with the incidence of sport-related concussion in high school football players: a multifactorial prospective study. *Am J Sports Med.* 2014;42(10):2470–2478. doi:10.1177/0363546514541926
 17. Kontos AP, Elbin RJ, Sufrinko A, Marchetti G, Holland CL, Collins MW. Recovery following sport-related concussion: integrating pre- and postinjury factors into multidisciplinary care. *J Head Trauma Rehabil.* 2019;34(6):394–401. doi:10.1097/HTR.0000000000000536
 18. Womble MN, McAllister-Deitrick J, Marchetti GF, et al. Risk factors for vestibular and oculomotor outcomes after sport-related concussion [published online June 11, 2019]. *Clin J Sport Med.* doi:10.1097/JSM.0000000000000761
 19. Halstead ME, McAvoy K, Devore CD, Carl R, Lee M, Logan K; Council on Sports Medicine and Fitness; Council on School Health. Returning to learning following a concussion. *Pediatrics.* 2013;132(5):948–957. doi:10.1542/peds.2013-2867
 20. DeMatteo CA, Lin CA, Foster G, et al. Evaluating adherence to return to school and activity protocols in children after concussion [published online December 24, 2019]. *Clin J Sport Med.* doi:10.1097/JSM.0000000000000800
 21. D'Lauro C, Johnson BR, McGinty G, Allred CD, Campbell DE, Jackson JC. Reconsidering return-to-play times: a broader perspective on concussion recovery. *Orthop J Sports Med.* 2018;6(3):2325967118760854. doi:10.1177/2325967118760854
 22. Churchill N, Hutchison M, Richards D, Leung G, Graham S, Schweizer TA. Brain structure and function associated with a history of sport concussion: a multi-modal magnetic resonance imaging study. *J Neurotrauma.* 2017;34(4):765–771. doi:10.1089/neu.2016.4531
 23. Gardner A, Iverson GL, Stanwell P. A systematic review of proton magnetic resonance spectroscopy findings in sport-related concussion. *J Neurotrauma.* 2014;31(1):1–18. doi:10.1089/neu.2013.3079
 24. Lumba-Brown A, Teramoto M, Bloom OJ, et al. Concussion guidelines step 2: evidence for subtype classification. *Neurosurgery.* 2020;86(1):2–13. doi:10.1093/neuros/nyz332
 25. Littleton AC, Schmidt JD, Register-Mihalik JK, et al. Effects of attention deficit hyperactivity disorder and stimulant medication on concussion symptom reporting and computerized neurocognitive test performance. *Arch Clin Neuropsychol.* 2015;30(7):683–693. doi:10.1093/arclin/acv043
 26. Echemendia RJ, Iverson GL, McCrea M, et al. Role of neuropsychologists in the evaluation and management of sport-related concussion: an inter-organization position statement. *Arch Clin Neuropsychol.* 2012;27(1):119–122. doi:10.1093/arclin/acr077
 27. Register-Mihalik JK, Kay MC, Kerr ZY, et al. Influence of concussion education exposure on concussion-related educational targets and self-reported concussion disclosure among first-year service academy cadets. *Mil Med.* 2020;185(3–4):e403–e409. doi:10.1093/milmed/usz414
 28. Kerr ZY, Collins CL, Mihalik JP, Marshall SW, Guskiewicz KM, Comstock RD. Impact locations and concussion outcomes in high school football player-to-player collisions. *Pediatrics.* 2014;134(3):489–496. doi:10.1542/peds.2014-0770
 29. Mihalik JP, Guskiewicz KM, Marshall SW, Blackburn JT, Cantu RC, Greenwald RM. Head impact biomechanics in youth hockey: comparisons across playing position, event types, and impact locations. *Ann Biomed Eng.* 2012;40(1):141–149. doi:10.1007/s10439-011-0405-3
 30. Haarbauer-Krupa JK, Comstock RD, Lionbarger M, Hirsch S, Kavee A, Lowe B. Healthcare professional involvement and RTP compliance in high school athletes with concussion. *Brain Inj.* 2018;32(11):1337–1344. doi:10.1080/02699052.2018.1542000
 31. Khodae M, Currie DW, Asif IM, Comstock RD. Nine-year study of US high school soccer injuries: data from a national sports injury surveillance programme. *Br J Sports Med.* 2017;51(3):185–193. doi:10.1136/bjsports-2015-095946
 32. Ledoux AA, Barrowman NJ, Boutis K, et al; Pediatric Emergency Research Canada PedCARE team. Multicentre, randomised clinical trial of paediatric concussion assessment of rest and exertion (PedCARE): a study to determine when to resume physical activities following concussion in children. *Br J Sports Med.* 2019;53(3):195. doi:10.1136/bjsports-2017-097981
 33. Panwar J, Hsu CC, Tator CH, Mikulis D. Magnetic resonance imaging criteria for post-concussion syndrome: a study of 127 post-concussion syndrome patients. *J Neurotrauma.* 2020;37(10):1190–1196. doi:10.1089/neu.2019.6809
 34. Eagle SR, Nindl BC, Johnson CD, Kontos AP, Connaboy C. Does concussion affect perception-action coupling behavior? Action boundary perception as a biomarker for concussion [published online February 27, 2019]. *Clin J Sport Med.* doi:10.1097/JSM.0000000000000731
 35. Sufrinko AM, Mucha A, Covassin T, et al. Sex differences in vestibular/ocular and neurocognitive outcomes after sport-related concussion. *Clin J Sport Med.* 2017;27(2):133–138. doi:10.1097/JSM.0000000000000324
 36. Sufrinko A, Pearce K, Elbin RJ, et al. The effect of preinjury sleep difficulties on neurocognitive impairment and symptoms after sport-related concussion. *Am J Sports Med.* 2015;43(4):830–838. doi:10.1177/0363546514566193
 37. Broglio SP, Guskiewicz KM, Norwig J. If you're not measuring, you're guessing: the advent of objective concussion assessments. *J Athl Train.* 2017;52(3):160–166. doi:10.4085/1062-6050-51.9.05

38. Eckner JT, Sabin M, Kutcher JS, Broglio SP. No evidence for a cumulative impact effect on concussion injury threshold. *J Neurotrauma*. 2011;28(10):2079–2090. doi:10.1089/neu.2011.1910
39. Collins MW, Kontos AP, Okonkwo DO, et al. Statements of agreement from the Targeted Evaluation and Active Management (TEAM) Approaches to Treating Concussion meeting held in Pittsburgh, October 15–16, 2015. *Neurosurgery*. 2016;79(6):912–929. doi:10.1227/NEU.0000000000001447
40. McAllister T, McCrea M. Long-term cognitive and neuropsychiatric consequences of repetitive concussion and head-impact exposure. *J Athl Train*. 2017;52(3):309–317. doi:10.4085/1062-6050-52.1.14

Address correspondence to Shawn R. Eagle, PhD, ATC, Department of Orthopaedic Surgery, University of Pittsburgh, 3850 South Water Street, Pittsburgh, PA 15203. Address email to seagle@pitt.edu.