

Evidence Review for Preventing Osteoarthritis After an Anterior Cruciate Ligament Injury: An Osteoarthritis Action Alliance Consensus Statement

Osteoarthritis Action Alliance, Secondary Prevention Task Group

Jeffrey B. Driban, PhD, ATC, CSCS*; Heather K. Vincent, PhD†; Thomas H. Trojian, MD‡; Kirsten R. Ambrose, MS§; Shelby Baez, PhD, ATC||; Nicholas Beresic, EdD, CSCS§; David J. Berkoff, MD¶; Leigh F. Callahan, PhD§; Bruce Cohen, PhD, CSCS#; Madison Franek, PT, DPT**; Yvonne M. Golightly, PT, MS, PhD††; Matthew Harkey, PhD, ATC||; Christopher M. Kuenze, PhD, ATC||; Mary Catherine Minnig, MS‡‡; Ali Mobasher, BSc, MSc, DPhil (Oxon)§§; Adam Naylor, EdD|||; Connie B. Newman, MD, MACP¶¶; Darin A. Padua, PhD, ATC###; Brian Pietrosimone, PhD, ATC***¶##; Daniel Pinto, PT, PhD†††; Hayley Root, PhD, MPH, ATC‡‡‡; Matthew Salzler, MD§§§; Laura Schmitt, PT, MPT, PhD|||||; Lynn Snyder-Mackler, PT, ATC, ScD¶¶¶; Jeffrey B. Taylor, PT, DPT, PhD, CSCS####; Louise M. Thoma, PT, DPT, PhD****; Kevin R. Vincent, MD, PhD†; Elizabeth Wellsandt, PT, DPT, PhD††††; Monette Williams, BS‡‡‡‡

*Division of Rheumatology, Allergy, and Immunology, Tufts Medical Center, Boston, MA; †UF Health Sports Performance Center, Department of Physical Medicine and Rehabilitation, University of Florida, Gainesville; ‡Drexel University College of Medicine, Philadelphia, PA; §Osteoarthritis Action Alliance, Thurston Arthritis Research Center, University of North Carolina at Chapel Hill; ||Department of Kinesiology, Michigan State University, East Lansing; ¶Department of Orthopedics, University of North Carolina, Chapel Hill; #Excusercise, LLC, Waltham, MA; **University of North Carolina Therapy Services, UNC Wellness Center at Meadowmont, Chapel Hill; ††Department of Epidemiology, Thurston Arthritis Research Center, Injury Prevention Research Center, Osteoarthritis Action Alliance, University of North Carolina at Chapel Hill; ‡‡Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina at Chapel Hill; §§Research Unit of Medical Imaging, Physics and Technology, University of Oulu, Finland; Department of Regenerative Medicine, State Research Institute Centre for Innovative Medicine, Vilnius, Lithuania; Department of Joint Surgery, First Affiliated Hospital of Sun Yat-sen University, Guangzhou, China; World Health Organization Collaborating Centre for Public Health Aspects of Musculoskeletal Health and Aging, Liege, Belgium; |||Telos SPC, Boston, MA; ¶¶Department of Medicine, Division of Endocrinology, Diabetes & Metabolism, NYU Grossman School of Medicine, New York, NY; ###Department of Exercise and Sport Science, University of North Carolina at Chapel Hill; ***MOTION Science Institute, Department of Exercise and Sport Science, University of North Carolina at Chapel Hill; †††Department of Physical Therapy, Marquette University, Milwaukee, WI; ‡‡‡Department of Physical Therapy and Athletic Training, Northern Arizona University, Flagstaff; §§§Division of Sports Medicine, Department of Orthopaedics and Division of Sports Medicine, Orthopaedic Research, Tufts Medical Center, Boston, MA; |||||Division of Physical Therapy, School of Health and Rehabilitation Sciences, Ohio State University, Columbus; ¶¶¶Department of Physical Therapy, University of Delaware, Newark; ####Department of Physical Therapy, Congdon School of Health Sciences, High Point University, NC; ****Division of Physical Therapy, Department of Allied Health Sciences, University of North Carolina at Chapel Hill; ††††Division of Physical Therapy Education, University of Nebraska Medical Center, Omaha; ‡‡‡‡Triangle Multisport, Durham, NC

Context: The Osteoarthritis Action Alliance formed a secondary prevention task group to develop a consensus on secondary prevention recommendations to reduce the risk of osteoarthritis after a knee injury.

Objective: Our goal was to provide clinicians with secondary prevention recommendations that are intended to reduce the risk of osteoarthritis after a person has sustained an anterior cruciate ligament injury. Specifically, this manuscript describes our methods, literature reviews, and dissenting opinions to elaborate on the rationale for our recommendations and to identify critical gaps.

Design: Consensus process.

Setting: Virtual video conference calls and online voting.

Patients or Other Participants: The Secondary Prevention Task Group consisted of 29 members from various clinical backgrounds.

Main Outcome Measure(s): The group initially convened online in August 2020 to discuss the target population, goals, and key topics. After a second call, the task group divided into 9 subgroups to draft the recommendations and supportive text for crucial content areas. Twenty-one members completed 2 rounds

of voting and revising the recommendations and supportive text between February and April 2021. A virtual meeting was held to review the wording of the recommendations and obtain final votes. We defined *consensus* as >80% of voting members supporting a proposed recommendation.

Results: The group achieved consensus on 15 of 16 recommendations. The recommendations address patient education, exercise and rehabilitation, psychological skills training, graded-exposure therapy, cognitive-behavioral counseling (lacked consensus), outcomes to monitor, secondary injury prevention, system-level social support, leveraging technology, and coordinated care models.

Conclusions: This consensus statement reflects information synthesized from an interdisciplinary group of experts based on the best available evidence from the literature or personal experience. We hope this document raises awareness among clinicians and researchers to take steps to mitigate the risk of osteoarthritis after an anterior cruciate ligament injury.

Key Words: patient education, rehabilitation, physical activity, injury prevention, psychological stress

Key Points

- Besides optimizing short-term outcomes (eg, return to activity), a focus on secondary prevention of osteoarthritis can help preserve lifelong quality of life and wellness.
- With an intent to reduce the risk of osteoarthritis after a person sustains an anterior cruciate ligament injury, we offered recommendations to comprehensively address physical and psychosocial impairments.

Many knee injuries occur in young, physically active people. More than 1 in 3 anterior cruciate ligament (ACL) reconstructions are performed in high school or college athletes in the United States.¹⁻³ Reconstruction of the ACL often leads to positive outcomes, such as a return to physical activity (eg, sport, occupational, recreational). Unfortunately, for at least 1 in 3 young patients, a knee injury is a catalyst to living with knee osteoarthritis for most of their lives.⁴⁻⁶ Many younger adults with knee osteoarthritis experience poor quality of life, diminished physical activity levels, and decreased physical function, all of which may lead to long-term psychosocial or psychological (eg, depression) and economic (eg, high medical costs) concerns and comorbidities (eg, cardiovascular disease).⁷⁻¹⁰ In the United States, the economic burden of individuals with a history of ACL reconstruction is greater than \$7.6 billion per year.⁷

Prevention of the long-term burden of knee osteoarthritis among people after an ACL injury is urgently needed. Unfortunately, the current literature on secondary prevention for people with an ACL injury lacks high-quality evidence. This deficiency was highlighted by the lack of secondary prevention strategies in the “National Public Health Agenda for Osteoarthritis: 2020 Update.”¹¹ In the absence of consistent evidence-based conclusions from the literature, consensus guidelines can be established by leveraging expert opinion and clinician experience. Hence, the Osteoarthritis Action Alliance formed the interdisciplinary Secondary Prevention Task Group (Supplemental Table) to develop consensus-based secondary prevention recommendations for clinicians intended to reduce the risk of osteoarthritis after a person has an ACL injury and to help define directions needed for future research. The task group proposed a comprehensive list of 16 recommendations, conducted extensive literature reviews, and complet-

ed 3 rounds of voting and revisions. Consensus was reached on 15 recommendations that encompass a broad approach to addressing the mental and physical well-being of patients to mitigate their risk of osteoarthritis from the time of injury until the diagnosis of osteoarthritis, when osteoarthritis treatment guidelines should be consulted.¹²⁻¹⁴ The following sections describe the consensus methods, the literature reviews to justify each recommendation, and all dissenting opinions by members of our task force to facilitate future discussion. We included the 16th recommendation, which failed to achieve consensus, because it may inspire further discussions and research. Overall, our expert panel offered this consensus statement based on their interpretation of the literature and clinical experience to fill a unique gap in an emerging field with insufficient evidence to inform clinical practice guidelines. These recommendations require high-quality research to justify and refine them and ultimately to help inform clinical practice guidelines. We envision that this document is a starting point and will need to be revised in 5 to 10 years to reassess the recommendations based on emerging evidence.

METHODS

The Osteoarthritis Action Alliance is a national coalition of concerned organizations committed to elevating osteoarthritis (OA) as a national health priority and promoting effective policy solutions that address the individual and national tolls of OA. The Alliance was mobilized by the Arthritis Foundation and the Centers for Disease Control and Prevention, with the Thurston Arthritis Research Center at the University of North Carolina at Chapel Hill as the lead agency. The Osteoarthritis Action Alliance formed the Secondary Prevention Task Group to develop a consensus on secondary prevention recommendations in order to

reduce the risk of OA after a knee injury. The associate director of the Osteoarthritis Action Alliance sent an electronic invitation to members of the Steering Committee and Primary Prevention Task Group. The Steering Committee includes representatives from lead partner organizations: the American Medical Society for Sports Medicine, Arthritis Foundation, Association of Rheumatology Professionals, Osteoarthritis Research Society International, and US Bone and Joint Initiative. Other organizations are also represented on the steering committee, including the National Athletic Trainers' Association, American Chronic Pain Association, Movement is Life, Alliance for Balanced Pain Management, National Recreation and Park Association, and American Medical Women's Association. The invitation encouraged recipients to share it with other contacts who might be interested. The task group consisted of 29 members, including 4 staff members from the Osteoarthritis Action Alliance (Supplemental Table). The remainder of the task group were volunteers from the constituent organizations with various backgrounds, including athletic trainers (ATs), orthopaedic surgeons, physical therapists, psychologists, and sports medicine physicians.

On August 28, 2020, 25 task force members met online to discuss the target population, goals, and key topics. The group agreed that the target population would be people ≤ 40 years of age with an ACL injury from the time of injury until a diagnosis of knee OA. We focused on individuals ≤ 40 years old because this population would include most people who experience an ACL injury and would be unlikely to have OA at the time of injury. The short- to mid-term goals were to

1. Promote a patient's role in shared and informed decision-making (foster informed consumers);
2. Educate people about their injury and recovery (eg, what to expect, long-term risks, risk modifiers), coping strategies, maintaining or improving self-efficacy, the importance of adherence, stress management, and goal setting;
3. Optimize physical activity (eg, a total return to previous activities or activity modification);
4. Prevent or reduce chronic symptoms (eg, knee pain, poor quality of life, functional limitations, fear);
5. Prevent subsequent injuries; and
6. Reduce psychological stress (eg, related to return to activity, communication with a clinician).

The recommendations were designed with the overarching long-term goal of providing clinicians with evidence and treatment strategies intended to prevent or delay the onset of symptomatic OA, prevent OA-related disability, and improve quality of life. The group also agreed to the need to recommend key outcomes to monitor and interventions to consider. Specifically, the group agreed to consider education, exercise, exposure therapy, mindfulness therapies, secondary injury prevention, use of technology, and coordinated care strategies. One task force member drafted a summary of key topics from the first call and shared it with the task force on September 2, 2020. These topics were further refined during an online meeting on October 14, 2020.

On November 3, 2020, the task group was partitioned into 9 subgroups to draft recommendations and supportive text for crucial content areas based on published evidence and clinical experience (Supplemental Table). The task

group received a draft of these recommendations and supportive text, and the first round of voting occurred from January 25 to February 15, 2021. We defined *consensus* as $>80\%$ of voting members supporting a proposed recommendation. A voter who could not support the current wording was asked to provide recommended changes to the text. Twenty-two members (76%) voted in round 1. Based on feedback from round 1 voting, the subgroups revised the recommendations and supportive text for each content area (February 16–April 13, 2021). In addition, some subgroups reached out to other content experts within the task group to assist with the revisions and ensure that the revisions were responsive to feedback. A second round of voting occurred online from April 16 to 30, 2021. Twenty-one of the 22 members (96%) from round 1 voted in round 2.

After the second round of voting, the task group convened a virtual meeting to review the wording of the recommendations and provide final votes. During this meeting, attendees agreed that the text for the psychological interventions should be reevaluated and updated. Then over the next 10 days, online voting for the new psychological recommendations occurred. Furthermore, task force members who could not attend the May 27 meeting voted on all the recommendations. Everyone who voted in round 2 voted in round 3.

After the final round of voting, 3 task group members (J.B.D., H.K.V., T.H.T.) harmonized the text into a single document. Next, they forwarded this document to individuals who voted against a specific recommendation to ask for dissenting opinions. Finally, a complete draft was forwarded to the 21 people who voted in rounds 2 and 3. After feedback was received, the document was shared with the task group members who participated in the initial development of the consensus statement but did not write any of the recommendations or vote. This subset served as a final review team. The lead author (J.B.D.) revised the document based on this feedback and shared the document with the task group and the Osteoarthritis Action Alliance Steering Committee for final approval.

We ordered the recommendations in part on the timing of when they might be implemented for each patient. However, the last 4 overarching recommendations (No. 13–16; eg, social support, use of technology, and coordinated care programs) could help implement the preceding recommendations.

RECOMMENDATIONS

1. Provide accessible resources that health care providers can distribute to a patient

Patients who sustain traumatic musculoskeletal injuries need to better understand their risk of developing OA. Only 27% of patients with an ACL injury remembered discussing their OA-related risk with a health care provider.¹⁵ Discussions about OA should center on what patients can do to mitigate the development and progression of the condition. In addition, health care providers can implement communication methods to reduce the potential concerns associated with limited patient health literacy. These methods include avoiding medical jargon, engaging in patient questions, explaining unfamiliar terms, and using “teach-back” to ensure understanding.¹⁶ Finally, health care providers, especially those overseeing rehabilitation after an

ACL injury, need to become more aware of their patients' risk for OA and how to educate patients about this risk.^{17,18}

Health care providers should prompt questions, as patients often use outside sources to obtain information.^{19,20} It is pertinent for the provider to engage with patients about their access to available information.²⁰ Providers' notes accessible through electronic health records are not good information sources for most patients.²¹ AskShareKnow²² is a proven method of engaging patients to ask more questions. It is composed of 3 generic questions: (1) What are my options? (2) What are the possible benefits and harms of those options? (3) How likely are each of these benefits and harms to happen to me? These questions can be a springboard for further discussion with patients about reducing the risk of OA after an injury.

2. Provide educational opportunities to health care professionals regarding how to best educate patients about OA prevention

Incorporating information about effective methods of educating patients about OA after ACL injury, and especially about prevention, into entry-level education for all health care providers is essential. Furthermore, we need more continuing education opportunities for health care providers regarding how to best educate patients after ACL injury about OA. Future researchers should examine the most effective training methods for providers and identify which methods optimize patient literacy about the OA risk.

3. Develop a tool kit for the caregiver to identify a patient's willingness for rehabilitation and the patient's preferred mode of obtaining self-management resources

We recommend that a tool-kit questionnaire consisting of 2 to 4 simple questions be developed to incorporate ways of delivering self-management strategies. The tool kit would identify the patient's willingness to manage the injury and identify limitations to obtaining, learning, and retaining these findings. If the patient is originally unwilling to manage the situation, then health care professionals should provide the patient or a willing or trusted family member with more information about the rationale for managing the injury and some management strategies. The tool kit would also help the health care provider determine the best delivery mode (eg, discussion, doctor's notes, electronic medical records, handouts, multimedia educational tools, telephone coaching). Delivery of care during the COVID-19 pandemic has taught us much about the value of technology-enabled interactions with patients. Patients have favorably endorsed virtual or phone sessions to enhance their education after an injury.²³ Health care providers should explore newer and upcoming technology to deliver health information, such as podcasts or social media, about OA.²⁴ Patient education about the potential for OA needs to begin at the time of the initial injury and recur at regular intervals after injury or reconstruction. Future investigation is needed to establish the best frequency for this education. Still, initial efforts could coincide with traditional follow-up visits (eg, initial clinic visit, acute postoperative visit, 6-month follow-up, return-to-participation assessment) to minimize the initial burden on the health care system.

Dissenting Opinion. Not all people who experience an ACL injury have a *caregiver* (eg, a family member or paid person who regularly looks after the patient) who will be involved in the ACL injury recovery. The term *caregiver* may lead readers to assume the goal is to develop a tool kit for parents, especially for patients <18 years of age. This recommendation encourages the development of a tool kit for patients, parents or caregivers, and health care professionals.

4. After an ACL injury or reconstruction, individuals should undergo a supervised, comprehensive, and progressive rehabilitation program to address impairments and neuromuscular deficits, specifically those related to quality of movement, knee range of motion, quadriceps muscle strength and performance, and functional performance before return to activity

Impairments in body structure and function that persist after an ACL injury or reconstruction are associated with subsequent radiographic abnormalities (eg, joint space narrowing), making it critical to resolve these impairments during postoperative rehabilitation early after the injury or ACL reconstruction. Nearly 50% of young and active individuals after ACL reconstruction are cleared for return-to-sport participation with altered movement patterns.^{25–28} Furthermore, many patients return to sport with impaired muscle function, asymmetries in vertical ground reaction force and loading rate, and biomechanical deficits associated with quadriceps weakness.^{25–28} Early optimization of knee range of motion, quadriceps strength, and movement patterns could help prevent or delay the onset of OA in this high-risk population.

Restore Range of Motion of the Involved Knee to Match the Uninvolved Knee, Emphasizing Extension. After an ACL reconstruction, more than 25% of individuals have side-to-side differences in knee extension $>5^\circ$ and this can persist up to a month.²⁹ Early loss of knee range of motion (as small as 3° to 5°) after an ACL reconstruction, especially with extension, may make the patient more than twice as likely to develop radiographic OA, particularly when coupled with concomitant meniscectomy or cartilage damage at the time of ACL reconstruction.^{30–32} Therefore, early joint motion after ACL reconstruction is an important component of comprehensive rehabilitation. Immediate passive and active motion (≤ 1 week) after ACL reconstruction should be implemented to increase joint range of motion (from full extension to $\geq 90^\circ$ of flexion), reduce joint pain, and lessen the risk of adverse responses of surrounding soft tissues (eg, arthrofibrosis).³⁰ In weeks 2 through 8 after surgery, a gradual but consistent increase in motion to full flexion can help restore knee mobility.³⁰

Restore Quadriceps Femoris Strength and Function to Improve Physical Function and Protect Joint Structures. The quadriceps muscles are the primary stabilizers of the knee and play a critical role in joint dynamic stability and loading, particularly in ground reaction force and knee kinematic and kinetic patterns that may protect cartilage.³³ Eccentric action of the quadriceps muscles during joint-loading activities facilitates knee flexion to absorb shock. Quadriceps weakness is a significant risk factor for the development and progression of knee OA.³⁴ Unfortunately, approximately 50% of young and active individuals are

cleared for high-level sports activities with significant quadriceps femoris strength deficits in the involved limb compared with the uninvolved limb.^{26,35,36} These deficits can range from 15% to 40%.³⁰ The deficit adversely affects mechanical loading³⁷ and contributes to unfavorable changes to cartilage (eg, lower proteoglycan density, more severe patellofemoral cartilage damage).^{38–42} In addition to deficits in muscle peak torque or force output, the rate at which torque develops independently contributes to the individual's ability to dissipate lower limb loading and generate the rapid, forceful movements that are common in sports participation.^{43,44} Women have bilateral quadriceps femoris weakness and a slower rate of torque development than men during the first year after ACL reconstruction.⁴⁵

Reversing strength loss after ACL injury or reconstruction can change the trajectories of physical function and clinical markers of OA. Although quadriceps femoris strength deficits relate to impaired knee-related function, functional performance, altered movement patterns, and cartilage changes,^{26,35,36,40} resolving these deficits produces functional performance similar to that observed in healthy individuals.³⁵ Authors⁴⁶ of a 5-year prospective study showed that more symmetrical quadriceps femoris strength was associated with lower odds of clinical knee OA. For every 1% increase in limb quadriceps femoris strength symmetry, the odds of clinical knee OA were reduced by 4%. Restoring quadriceps femoris strength and performance are key criteria for progressing rehabilitation and preparing for the return to unrestricted sports participation.^{47,48} Implementing functional closed chain strengthening exercises, such as limited-range-of-motion wall slides and step-ups, early in the recovery process (>2 weeks) and slowly progressing weight bearing over 10 weeks³⁰ can help promote strength return. Electrical stimulation for 6 to 8 weeks after an ACL reconstruction can augment muscle strengthening and improve neural drive to increase quadriceps muscle strength and enhance short-term functional outcomes.⁴⁹ Weight-bearing and non-weight-bearing concentric and eccentric exercises should be implemented early after ACL reconstruction, 2 to 3 times/wk for 6 to 10 months, to increase thigh muscle strength and functional performance. Recommendations for a safe return to activity should include achieving quadriceps strength symmetry >90%.³⁰

Resolve Altered Movement Patterns Early After ACL Reconstruction. Although quadriceps femoris strength is an important component of return to activity, persistent aberrations in gait or side-to-side loading mechanics need to be corrected with strengthening.⁴⁶ An ACL injury or reconstruction may produce variable effects on knee motion. After an ACL reconstruction, many people have increased sagittal-plane knee range of motion during the first year after surgery and an acute reduction and then gradual increase in knee-extensor moment.⁵⁰ Furthermore, they have lingering signs of anteroposterior instability and tibial rotation during the stance phase of walking.⁵⁰ The variability of kinematic and kinetic measures during gait suggests that patients have unique responses to their specific injury or surgery. Some individuals after ACL reconstruction demonstrate elevated knee-adduction-moment impulses, which are associated with altered cartilage composition.⁵¹ Others show reductions in peak knee-flexion moment and flexion angle during walking gait from 6 months to year 3.⁵² Biomechanical patterns early after ACL

reconstruction may predict persistent abnormalities in the cartilage and meniscus. During drop-jump landings, vertical ground reaction forces, knee-flexion angles, and knee moments are 4% to 46% lower at 6 months post-ACL reconstruction. These patterns are associated with persistent meniscal and cartilage abnormalities 3 years later, despite the resolution of biomechanical differences.^{52,53}

During a comprehensive and progressive rehabilitation program, side-to-side lower limb asymmetries in movement coordination and performance should be monitored, and neuromuscular reeducation training should be incorporated with muscle-strengthening exercises for the quadriceps femoris and other muscles that influence the lower extremity kinetic chain (eg, hip abductors, hip rotators). In addition, persistent deficits in strength and neuromuscular control or poor quality of movement should continue to be addressed in the clinic or through sport-specific rehabilitation programs before return to activity. Future researchers should follow people long after an ACL injury to clarify the effects of strength and range-of-motion changes on long-term joint health.

5. Before full reintegration into a sport, individuals should gradually resume sport-specific training to restore metabolic conditioning, build tolerance to chronic training loads, and adopt desired movement strategies

Premature return to sport may be associated with early OA⁵⁴ and ACL reinjury.⁵⁵ In addition, the ACL injury incidence rate is 15 times higher among people with a prior ACL reconstruction than in people without that injury history.⁵⁶ These adverse outcomes may result from insufficient physical conditioning and strength,^{57–60} inadequate tolerance of real training loads,⁵⁵ and lingering biomechanical asymmetries^{61–63} at the time of return to sport. Thus, a layered approach to preparing athletes for return to sport that includes progressive and overlapping conditioning activities may help develop tolerance to training loads and improve contextual movement strategies to prevent poor outcomes.

Physical Conditioning and Building Chronic Training Tolerance. Indirect evidence suggests that fatigue affects the ACL injury risk factors of neuromotor function, muscle rate and amount of force development, and movement strategies.⁵⁵ Furthermore, more ACL injuries occur in the second halves of competitions in soccer and a variety of high school sports.^{64,65} Monitoring fatigue in both the involved and uninvolved limbs during recovery while the patient performs functional hop tests may add insight on the readiness to return to sport. The uninvolved limb should be monitored because a fatiguing hop-test battery affected this limb more than the ACL reconstruction limb among athletes >6 months post-ACL reconstruction.⁶⁶ Greater fatigue in the uninvolved limb is likely a function of detraining and should be considered when deciding on a person's readiness to reengage in activity.⁶⁶ Developing aerobic and anaerobic conditioning with muscle strength in both limbs during functional recovery enables athletes to work at higher intensities for longer durations and to return to activity without the risks associated with fatigue.⁵⁵ A progressive increase in chronic training loads (total volume and frequency of training sessions, increased intensity of

activity over time toward peak workload) induces tissue adaptation. It also prevents potential underloading before introducing real sport demands.^{55,59} Readiness to progress the workload and return to sport is complicated and depends on indices of tissue health, stresses on the tissue, and modifiers of risk tolerance.⁶⁷

A gradual, multifaceted sport reintegration plan with balanced benefits and risks⁶⁸ can restore fitness and neuromotor performance. For example, in a 10-month intervention study,⁶⁹ 50 soccer players with ACL reconstruction participated in on-field training and gradually introduced ball-handling activities, higher running intensities, and an increased technical level of ball activities. This program resulted in 23% to 86% gains in the anaerobic threshold knee–flexion–extension strength at the end of rehabilitation. Gradual reintroduction of controlled activities in a sport-specific environment can occur as performance and confidence improve. Then the complexity and involvement with teammates can increase. Emerging evidence and expert opinion indicate that rehabilitation practice should support an athlete’s return to a specific sport environment. Athletes who compete on the field, court, track, or rink benefit from exposure to activities on-site while conditioning is being restored.^{55,70,71} Early reintroduction to sport-specific demands can include graded exposure to different directions of motion (forward, backward, lateral direction changes, or varying patterns). Intermediate on-site training can include low-intensity agility and neuromuscular drills and power development during controlled changes in direction. These early stages emphasize functional strengthening and neuromuscular enhancement.^{71,72} In later rehabilitation phases, an athlete can focus on responding to unanticipated demands (eg, changes in direction and speed with greater perceptive and neurocognitive demands)⁶⁸ and developing more endurance. If return-to-sport criteria are achieved, the athlete can participate in progressively more intensive noncontact team drills, practices, and scrimmages over time.⁷⁰

Adopt Desired Movement Strategies to Facilitate Return to Sport. Nine months post–ACL reconstruction, several motion features associated with ACL injury occur during unplanned 90° cutting maneuvers,⁶² single-legged jumps and hop tasks, and double-legged drop jumps.⁶³ These features include interlimb asymmetries in internal knee–valgus moment, internal knee–rotation angle, and external ankle–rotation moment and a shift of center of mass toward the involved side. Thus, beyond the restoration of greater than 90% limb symmetry in strength, additional advanced training may be needed to restore the explosive performance⁵⁵ needed when transitioning back to sport. Experts^{8,73} describe the importance of developing contextual sport movement through a phased progression from foundational movement training with loading (eg, back squats, unilateral tasks, bilateral landing control) to high-load sport-specific movement retraining (eg, plyometrics, speed, and acceleration) and finally to on-field sport motion. In addition, training interventions that improve performance of risky movements may help prevent recurrent ACL injury. Consensus groups^{74,75} agree that individuals should be able to pass a standard battery of objective movement tests after interventions and before full return to sport (eg, movement quality, hop tests, strength, range of motion), but the specific tests vary. For example,

athletes who fail to complete their rehabilitation and fail all hopping, running speed, agility, and strength tests are 4 times more likely to rerupture the ACL than athletes who pass these tests.⁷⁶

Future researchers should determine clear definitions of return to sport for each sport activity and which metrics should comprise the ideal test battery for progression in rehabilitation and return to sport.^{67,74}

6. Individuals after an ACL injury or ACL reconstruction should be encouraged to meet the “Physical Activity Guidelines for Americans” (2nd edition)

Sport is commonly the main component of physical activity for young people.⁷⁷ When individuals with an ACL injury or ACL reconstruction stop sport participation, often physical activity is reduced,^{77,78} which elevates the risk of chronic morbidity associated with an inactive lifestyle.⁷⁹ Historically, resuming physical activity after ACL reconstruction for overall health benefit has been underemphasized in clinical practice or research. Lower physical activity levels are prognostic of future knee pain and symptoms as well as a lower quality of life.⁸⁰ Clinicians have an opportunity to shape a positive long-term health trajectory, as physical activity can help counter comorbid disease onset and risk factors associated with OA onset and progression.

Physical Activity Patterns and Related Health Considerations. The current “Physical Activity Guidelines for Americans,”⁸¹ issued in 2018 by the US Department of Health and Human Services, recommend that adults participate in at least 150 minutes of moderate-to-vigorous aerobic physical activity and at least 2 days of strengthening activity each week. Unfortunately, young people with a history of an ACL injury are 2 to 4 times less likely to meet these guidelines, even after return to sport.⁷⁹ Furthermore, up to 67 months after an ACL reconstruction, young adults spent 15% to 41% less time in moderate-to-vigorous aerobic physical activity. They were also less likely to meet national physical activity guidelines than those without a history of injury.^{45,77,82–84} Specifically, individuals with a history of ACL reconstruction

- accumulated an average of 16% fewer steps per day,^{82,84}
- met a daily 10 000 step count target at a proportionally lower rate,⁸²
- accrued fewer steps at moderate-to-vigorous cadence,⁸⁴ and
- failed to meet physical activity guidelines as frequently as healthy individuals without a history of ACL injury.^{45,82}

Consequently, people with a history of ACL reconstruction have elevated risks for other lifestyle-related chronic diseases, including cardiovascular disease, hypertension,^{10,85} obesity,⁷⁸ and depression.⁸⁶ Significant weight gain can occur in adolescents during the first year after an ACL reconstruction.⁷⁸ Adults who incur a second ACL injury are 3.8 times more likely to be on a weight gain trajectory than those who are not reinjured.⁸⁷ Concerning the cardiovascular system, the incidence of myocardial infarction increases by 50% with a history of ACL injury.¹⁰ Furthermore, the depression incidence may be as high as

42% among people after an ACL reconstruction.^{86,88} Thus, physical activity may promote overall health and well-being via weight maintenance and systemic health.

Physical Activity for Long-Term Health. There is a clear and substantial benefit of regular physical activity (land and aquatic) on knee pain symptoms and physical function, with moderate effects on quality of life, regardless of variations in dosing.^{89,90} Resistance exercises also significantly affect knee-related quality of life.⁹⁰ Cardiovascular disease risk decreases 5% to 21% for every 1000 daily steps taken,⁹¹ and a dose-response relationship exists between physical activity levels and incident hypertension.⁹² Also, strong evidence from the Physical Activity Guidelines Committee exists to show that prevention of weight gain is most effective with participation in more than 150 minutes of moderate-to-vigorous physical activities a week.⁹³ Furthermore, meeting physical activity guidelines can protect against depression.⁹⁴

Health care professionals should engage patients in discussions on how to safely meet physical activity guidelines throughout the recovery process. These discussions will require health care professionals to respect precautions and a person's ability early in recovery. Furthermore, discussions should address how to progress to allowable activities throughout rehabilitation and emphasize maintaining recommended physical activity levels throughout life. Sport activities are a safe and effective strategy to meet physical activity guidelines. Participation in various sport or recreational activities may even protect against the development and progression of knee symptoms and OA.^{89,95–99} Irrespective of sport or activity type, moderate daily recreational activities (performed as recommended for health benefit) are not a consistent risk factor for clinical or radiographic knee OA.^{34,100} Regular engagement in select sports with high load or high injury risk (eg, soccer or elite-level long-distance running, weight lifting, or wrestling) may, however, elevate the risk for knee OA¹⁰¹ or medial-compartment cartilage damage.¹⁰² People with greater underlying joint pathology demonstrate more knee-joint structural change with step counts greater than 10 000 than with step counts less than 10 000.¹⁰³ Therefore, among persons with a history of an ACL injury or ACL reconstruction, it is prudent to achieve physical activity guidelines using a variety of physical activities with moderate knee loading over the long term, especially among subgroups of people at higher injury risk.

Authors of future studies need to provide more specific recommendations for prescription and dosing and promotion of physical activity, including aerobic and strengthening activities,⁹⁰ after ACL reconstruction and determine which patient subgroups are most protected against OA by various activity types.

7. Psychological skills training should be considered as part of the short- and long-term care plans after ACL injury and reconstruction to prevent reinjury, improve overall health and wellness, and encourage engagement in and adherence to physical activity

Despite successful ACL reconstruction, only 44% of these athletes return to competitive levels of physical activity.¹⁰⁴ Primary barriers to return to physical activity

are elevated injury-related fear (ie, *kinesiophobia*), decreases in *self-efficacy* (ie, the belief in one's ability to complete a task), and pain. Psychological skills training includes mindfulness meditation, relaxation skills, self-knowledge mastery, mental imagery, goal setting, and self-talk.¹⁰⁵ Psychological skills training fosters focus, acceptance, attentional control, decision-making, and a mindful state.¹⁰⁶ It may improve overall well-being, stress coping, development of the most productive mindset, and confidence building before surgery and during rehabilitation or return to activity.¹⁰⁷ Skills are often coupled into a multimodal intervention (eg, relaxation and imagery,¹⁰⁸ goal setting, and mindfulness¹⁰⁹). Various trained clinicians (eg, physicians, nurses, physical therapists, ATs) and other sport science-trained mental health professionals (eg, mental performance consultants, psychologists, mental health counselors) can teach these skills.

Mindfulness, Mental Imagery, and Other Psychological Skills for Early Recovery and Improved Long-Term Outcomes After ACL Injury or Reconstruction. *Mindfulness* is defined as paying attention in a particular way, on purpose, in the present moment, and nonjudgmentally.¹¹⁰ In group settings, the teaching of mindfulness involves developing an awareness of thoughts, feelings, or bodily sensations; bringing attention to breathing; and expanding attention to the body. Other techniques include developing balance in logical and emotional thought and the ability to focus on “how” skills (eg, nonjudgmental, one-mindful [being present in the moment], and effective) and “what” skills (eg, observing, describing, and participating). A person can practice mindfulness during short pauses in everyday life and incorporate mindfulness into daily movements, such as walking and quiet sitting. Mindfulness can also be merged with yoga, tai chi, or other sports. Among individuals with a history of an ACL injury, mindfulness could bring balance to logical and emotional thought, improve situational awareness for subsequent injury, reduce stress and kinesiophobia, and prevent new injuries in the short to mid term. Mindfulness can also improve acute or postsurgical pain.¹¹¹ From studies of people with knee OA, we can infer that if chronic pain occurs after an ACL injury, a mindfulness intervention (eg, tai chi, yoga, meditation) may improve the pain, pain coping, anxiety, stress, depression, kinesiophobia, quality of life, physical function, postural control, and responsiveness to other therapies.^{112–119}

Mental imagery is the self-directed imagining or visualizing of specific events, actions, or outcomes, including associated feelings and responses, to increase motivation toward a target action or task.¹²⁰ Commonly used imagery practices after an injury include healing imagery (eg, imagining tissue healing), soothing imagery (eg, imagining lying on a beach), and performance imagery (eg, imagining completing a task, such as knee extension). Mental imagery techniques can reduce pain before surgery, injury-related fear, knee symptoms, and the number of days people use crutches after an ACL injury or reconstruction.^{121–123} Among athletes with an ACL reconstruction, guided imagery and relaxation initiated within 1 week of surgery lowered pain by 85% and reinjury anxiety by 87% and was associated with greater gains in knee isokinetic strength.¹⁰⁸ Furthermore, a guided imagery-physical ther-

apy intervention (ie, goal setting, motivational healing, exercise rehearsal, relaxation) improves self-efficacy.

Mental Flexibility for Prevention of Reinjury. The musculoskeletal reinjury risk among high school or collegiate athletes may be reduced by increasing mental flexibility. *Mental flexibility* consists of reducing the perception of stress, supporting nonjudgmental awareness, and increasing levels of a positive emotional state.^{124,125} Thus, adding techniques such as mindfulness meditation or mental imagery into the daily routine may improve safety during activity after ACL injury.

Authors of future high-quality investigations should systematically study the differential efficacy of psychological skills training on knee health trajectories, progression of OA, and quality of life years after an ACL injury or reconstruction. Moreover, researchers need to identify the best responders to different psychological skills training programs in order to adopt a precision approach to preventing disability and optimizing quality of life. These studies will strengthen the evidence for psychological skills training after an ACL injury or reconstruction.

Dissenting Opinion. The language of the recommendation is too strong for this emerging field, which draws on evidence from other sports injuries or OA. Many patients experience injury-related fear after an ACL reconstruction. Elevated injury-related fear, or kinesiophobia, relates to a greater risk of secondary ACL injury. Although psychological skills training may be beneficial for patients after an ACL reconstruction, we lack sufficient evidence to integrate psychological skills training into the rehabilitative care of patients after an ACL reconstruction. Specifically, it remains unclear if psychological skills training has a clinically meaningful effect on injury-related fear, kinesiophobia, or, most importantly, the risk of secondary injury.

8. Graded-exposure therapy can be added to therapy programs after ACL injury and reconstruction to overcome fear and prevent reinjury

In addition to fear of reinjury, kinesiophobia is also related to asymmetries in ground reaction force and rectus femoris and biceps femoris activation at 6 months after an ACL reconstruction.¹²⁶ Patients who exhibit heightened levels of injury-related fear after traumatic knee injury are 13 times more likely to sustain a secondary knee injury within 24 months of clearance to return to sport.¹²⁷ Up to 10 years post-ACL reconstruction, kinesiophobia relates to knee symptoms and function.¹²⁸ Therefore, overcoming psychosocial barriers to therapy participation and preventing reductions in activity are important after ACL injury or ACL reconstruction to promote better long-term knee health.

Graded-Exposure Therapy and Early Recovery After ACL Injury or ACL Reconstruction. Graded activity and graded exposure address pain-related avoidance behaviors and pain-related fears, respectively, by providing positive reinforcement when a patient successfully completes the fearful task without experiencing pain or injury.¹⁰⁹ *Graded activity* addresses pain-related avoidance behaviors in a paced, collaborative, and goal-driven manner that requires establishing a time- or intensity-based quota to complete challenging tasks patients avoid because of pain.

The Photographic Series of Sports Activities for Anterior Cruciate Ligament Reconstruction¹²⁹ can be used to identify specific activities that may be considered fearful after ACL reconstruction. Clinicians can monitor the effectiveness of the graded-exposure therapy on injury-related fear using instruments such as the Tampa Scale of Kinesiophobia,¹³⁰ the Tampa Scale of Kinesiophobia-11,¹³¹ or the Anterior Cruciate Ligament Return to Sport After Injury Scale.¹³² Athletes with an ACL reconstruction indicated that graded exposure was part of a “preparation of body and mind” framework for returning to sport.¹⁰⁷

The efficacy of graded activity or graded exposure on reducing injury-related fear has been well studied in patients with musculoskeletal injuries, such as low back pain or total knee arthroplasty.^{109,133–140} In a pilot study¹⁴¹ among women after an ACL reconstruction, graded-exposure therapy led to decreases in fear about specific functional tasks (eg, jumping, hopping, pivoting) but not global changes in injury-related fear (ie, decreases on the Tampa Scale of Kinesiophobia-11). Unfortunately, no authors since 2012 have demonstrated the effectiveness of graded activity in reducing pain-related avoidance behaviors compared with therapeutic exercise as the standard of care.¹⁰⁹ Prospective studies are needed to systematically determine the effects of graded-exposure therapy on risk behaviors, psychological stress related to activities that load the knee, and adaptive behaviors against reinjury.

Dissenting Opinion. The language of the recommendation is too strong. Many patients experience injury-related fear after an ACL reconstruction, which increases the risk of secondary ACL injury. Although graded-exposure therapy may be beneficial for patients after an ACL reconstruction, we lack sufficient evidence to integrate it into the rehabilitative care of these patients. Specifically, it remains unclear if graded-exposure therapy offers a clinically meaningful effect on injury-related fear, kinesiophobia, or, most importantly, the risk of secondary injury.

9. Cognitive-behavioral counseling should be considered to promote patient engagement with therapies and exercise, reinjury prevention, and management and improvement of overall health and wellness after ACL injury or reconstruction

Cognitive-behavioral therapy is a treatment approach that focuses on efforts to change thinking patterns to overcome (1) flawed or nonhelpful ways of thinking, (2) learned patterns of unhelpful behaviors, and (3) barriers to coping with psychological problems. A variety of strategies are used to

- Recognize one’s distortions in thinking that lead to problems,
- Improve understanding of motivation and behavior of others,
- Problem solve to better cope with stress, and
- Increase confidence in personal abilities.

These strategies include learning to calm oneself and relax the body. In addition, people may use role play to prepare for difficult interactions with others and face fears instead of avoiding them. Cognitive-behavioral therapy

helps individuals to problem solve and positively change behaviors in real-life situations. This comprehensive therapy is administered by trained, certified mental health specialists (ie, psychologists, psychiatrists, clinical social workers, professional counselors).¹⁴² Investigators¹⁴³ have also employed other therapists trained by clinical psychologists to perform therapies at larger scales.

Cognitive-behavioral therapy may be helpful for some patients early after ACL injury or ACL reconstruction, but the evidence in this area is minimal. One aspect of cognitive-behavioral therapy, acceptance and commitment therapy, was beneficial in a case series of 4 athletes recovering from an ACL reconstruction (eg, improved acceptance of frustration, anxiety about or fear of reinjury, and motivation; better focused on the present; paying attention to the present during rehabilitation; and balanced values with committed action to recovery). Furthermore, in a pilot study¹⁰⁹ of 8 people with ACL reconstruction, a telephone-based program with various elements of therapy (eg, grounding and activity plan, monitoring self-talk, present mindedness, guided imagery) led to variable responses in kinesiophobia, knee-specific efficacy, and pain catastrophizing. These participants did not adopt these skills after the research intervention was completed. Therefore, this approach may work for some but not all individuals.

Cognitive-Behavioral Therapy Applications to Knee Symptom Management. The effects of long-term management of individuals with ACL injury or ACL reconstruction using cognitive-behavioral therapy are not yet known. Cognitive-behavioral therapy for people with knee OA improves physical performance, pain, the ability to cope with depression, anxiety, and pain catastrophizing when combined with interventions specific to the participants, such as pain psychology and weight management.^{144,145} Cognitive-behavioral therapy can be effectively administered via the internet to reduce depression and improve self-efficacy, joint symptoms, and physical function among people with knee OA.¹⁴⁶ Unfortunately, it is unclear whether in-person or other methods of cognitive-behavioral therapy delivery influence the effectiveness of the therapy on outcomes. Additional research is needed to determine the optimal cognitive-behavioral therapy to improve function and quality of life and promote healthy engagement in regular physical activity after an ACL injury. Moreover, additional research is needed to determine which patients may respond best to cognitive-behavioral therapy, whether cognitive-behavioral therapy is more effective after injury or ACL reconstruction, and during which window of time the greatest effects on outcomes are produced with this therapy.

Dissenting Opinion. The language of the recommendation is too strong. There is a theoretical basis for this recommendation and evidence that cognitive-behavioral counseling has been effective for patients with acute and chronic musculoskeletal conditions. Cognitive-behavioral counseling may be beneficial for many aspects of care for a patient after an ACL injury. However, we lack sufficient evidence that cognitive-behavioral counseling will benefit this patient population, especially if a goal is reinjury prevention.

10. Patients should be monitored regularly after an ACL injury using a comprehensive approach, including patient-reported outcomes, performance-based outcomes, and measures of disease progression

No standard list of recommended patient-reported outcomes has been compiled for individuals after an ACL injury to monitor recovery, limiting evaluation of treatment effects and informed modifications to treatment plans. Incorporating outcomes from key experiential domains relevant to people with a history of ACL injury could help guide (1) return-to-sport decision-making and (2) treatments and approaches focused on slowing or halting the progression of OA. Domains that affect the quality of life include pain and other symptoms, function, fatigue, sleep, psychological factors, fear of movement, stress, depression, anxiety, and physical activity levels. Therefore, monitoring self-reported physical and mental health is an essential element of care to achieve treatment goals. Relevant measures commonly used in orthopaedic or rheumatology research for people with an ACL injury are presented in Table 1. Measures previously tested and used in populations with ACL injuries are listed as “recommended.” Given the existing link between ACL injury and the future development of OA, measures that have been tested in populations with OA may also add value to the clinical assessment.

Objective measures from 6 domains provide complementary insight into patient-reported outcomes as part of the overall understanding of a patient’s risk for knee OA after ACL injury: (1) clinical impairment related, (2) modifiable lifestyle related, (3) functional performance, (4) gait biomechanics, (5) imaging, and (6) biomarkers. These outcomes range from clinically accessible techniques that can easily be incorporated into the clinic to outcomes that provide insight into disease pathogenesis that are currently limited to a research setting (Table 2).

11. Use a multifaceted return-to-sport test battery to inform a shared decision involving all stakeholders (eg, patient, parents, health care team, and coach) when determining readiness to return to play

One of 4 adolescents and young adults who return to cutting and pivoting sports after an ACL injury will sustain a second ACL injury.¹⁸³ The risk for a second ACL injury in young athletes who return to sport after an ACL injury is 30 to 40 times greater than that for their uninjured peers.¹⁸³ A second ACL injury leads to far worse outcomes than the first ACL injury, including twice the risk of having below-normal knee function or developing knee OA.¹⁸⁴

Current evidence-based return-to-sport criteria include benchmarks of quadriceps muscle strength, performance-based metrics such as single-legged hop tests, and patient-reported outcomes of subjective knee function.^{36,185–187} Passing a return-to-sport test battery reduces the risk of an ACL or meniscal injury by more than 84%.^{48,188} In contrast, failing to meet return-to-sport criteria is associated with a 4 times greater risk for sustaining an ACL graft rupture.⁷⁶ Quadriceps strength may be the most important criterion for predicting the reinjury risk.⁴⁸ Despite evidence that passing return-to-sport criteria reduces the reinjury risk, only 14% of patients met the recommended criteria for strength tests,

Table 1. Recommendations for Patient-Reported Outcomes Domains and Example Measures

Domain	Example Measure	No. of Items	Tested in Patients With ACL Injury?	Recommended or Suggested ^a
Pain/symptoms	NRS: pain ^{147,148} (ask separately at tibiofemoral and patellofemoral joints)	2	Yes	Recommended
	IKDC: Symptoms ¹⁴⁹	7	Yes	Recommended
	KOOS: Pain subscale ^{150–152}	9	Yes	Recommended
	KOOS: Symptoms subscale ^{150–152}	7	Yes	Recommended
Function	KOOS: Physical Function Short Form ^{152,154}	7	Yes	Recommended
	IKDC: Sports Activities ¹⁴⁹	10	Yes	Recommended
	IKDC: Function ¹⁴⁹	2	Yes	Recommended
	KOOS: Activities of Daily Living subscale ^{150–152}	17	Yes	Recommended
Fatigue	KOOS: Sports and Recreation subscale ^{150–152}	5	Yes	Recommended
	Stanford NRS fatigue ¹⁵⁵	1	No, used in OA	Suggested
Sleep	Fatigue: Perceived scale of exertion ¹⁵⁶	1	Yes	Recommended
	PROMIS: Fatigue ¹⁵⁷	7	No, used in OA	Suggested
Psychological	PROMIS-Sleep Disturbance Short Form ¹⁵⁸	8	No, used in OA	Suggested
Fear of movement	Anterior Cruciate Ligament Return to Sport After Injury ¹³²	12	Yes	Recommended
	Brief Fear of Movement Scale for osteoarthritis ¹⁵⁸	6	No, used in OA	Suggested
Stress	Tampa Scale of Kinesiophobia-11 ¹⁵⁹	11	Yes	Recommended
	PROMIS Perceived Stress ¹⁵⁷	10	No, used in OA	Suggested
Depression	PROMIS Emotional Distress: Depression Short Form ¹⁵⁷	8	No, used in OA	Suggested
Anxiety	PROMIS Emotional Distress: Anxiety Short Form ¹⁵⁷	7	No, used in OA	Suggested
Physical activity	International Physical Activity Questionnaire Short Form ¹⁶⁰	7	Yes	Recommended
	Tegner Activity Scale ¹⁶¹	1	Yes	Recommended
	Marx Activity Scale ¹⁶²	4	Yes	Recommended
Quality of life	KOOS: QOL subscale ^{150–152}	4	Yes	Recommended
Global rating of health	Health Status Questionnaire: self-rated general health ¹⁶³	1	No, used in OA	Recommended

Abbreviations: ACL, anterior cruciate ligament; IKDC, International Knee Documentation Committee; KOOS, Knee injury and Osteoarthritis Outcome Score; NRS, numeric rating system; OA, osteoarthritis, PROMIS, Patient-Reported Outcomes Measurement Information System; QOL, quality of life.

^a Measures previously tested and used in ACL-injured populations are listed as recommended.

Table 2. Recommendations for Objective Outcome Domains and Example Measures Assessed in Clinical or Research Settings

Domain	Example Measures	Research Only	Tested in Patients with ACL Injury?	Recommended or Suggested ^a
Clinical examination features	Joint line tenderness ¹⁶⁴		No, used in OA	Suggested
	Crepitus ¹⁶⁵		No, used in OA	Suggested
	Sweep/bulge test for joint effusion ¹⁶⁶		Yes	Recommended
	Joint alignment		Yes	Recommended
Modifiable lifestyle related	Body mass index ¹⁶⁷		Yes	Recommended
	Waist circumference		Yes	Recommended
	Fat mass (eg, dual-energy x-ray absorptiometry)		Yes	Recommended
	Physical activity (eg, step counts, % time moderate-vigorous activity) ¹⁶⁸	Yes	Yes	Recommended
Functional performance	Single-legged hop test ¹⁶⁹		Yes	Recommended
	Star Excursion Balance Test ¹⁷⁰		Yes	Recommended
	Chair stand test ¹⁷¹		Yes	Recommended
	Walking speed test ¹⁷²		Yes	Recommended
Gait biomechanics	Muscle function (eg, isometric strength) ³⁹		Yes	Recommended
	Vertical ground reaction force ¹⁷³	Yes	Yes	Recommended
	Knee abduction moment ¹⁷⁴	Yes	Yes	Recommended
	Spatiotemporal measures ¹⁷⁵	Yes	Yes	Recommended
Imaging	Radiograph (eg, Kellgren-Lawrence, joint space width) ^{176,177}		Yes	Recommended
	Semiquantitative MRI (eg, whole-organ MRI score, Anterior Cruciate Ligament Osteoarthritis Score) ¹⁷⁸	Yes	Yes	Recommended
	Quantitative morphological MRI (eg, cartilage thickness, effusion-synovitis volume, bone marrow lesion volume) ¹⁷⁹	Yes	Yes	Recommended
Biomarkers	Quantitative Compositional MRI (eg, T1 ρ , T2, dGEMRIC) ¹⁸⁰	Yes	Yes	Recommended
	Cartilage metabolism (eg, COMP, C2C, CPII) ¹⁸¹	Yes	Yes	Recommended
	Inflammation (eg, IL-6, TNF- α) ¹⁸²	Yes	Yes	Recommended

Abbreviations: ACL, anterior cruciate ligament; COMP, cartilage oligomeric matrix protein; CPII, type 2 collagen synthesis C-propeptide; C2C, cleavage of type II collagen; dGEMRIC, delayed gadolinium-enhanced MRI of cartilage; MRI, magnetic resonance imaging; OA, osteoarthritis.

^a Measures previously tested and used in ACL-injured populations are listed as recommended.

hop tests, and patient-reported outcomes after medical clearance to return to activity at approximately 8 months postoperatively.³⁶ Hence, some individuals may not meet the criteria at a pivotal moment in their return to activity.

Time since ACL reconstruction should not be the sole determinant of return-to-sport readiness.¹⁸⁹ It is, however, a relevant factor, as early return to sport is associated with a greater risk of reinjury.^{48,190} For example, returning to sport before 9 months after an ACL reconstruction increased the risk for secondary injury.^{48,190} Thus, return-to-sport testing should be coupled with assessing the time since ACL reconstruction to ensure appropriate knee function and biological healing. Other factors such as patient age, graft type, and concomitant injury may affect the timing of readiness for return to sport.^{191,192} Furthermore, preliminary evidence^{127,193,194} suggests that psychological readiness, kinesiophobia, and asymmetric biomechanical movement patterns may also be linked to the reinjury risk after returning to sport and thereby warrant assessment. As a result, clinicians should share decisions regarding return-to-sport readiness with the patient, parents of adolescents, medical team, and coaches based on functional return-to-sport testing, time since ACL reconstruction, and individual patient factors to minimize the secondary knee injury risk.

12. Implement a multifaceted preventive training program that includes strategies to improve agility, balance, flexibility, strength, and movement quality to reduce the risk of secondary injury

Several consensus statements and clinical practice guidelines, involving athletic training,¹⁹⁵ physical therapy,¹⁹⁶ the International Olympic Committee,¹⁹⁷ and the Osteoarthritis Action Alliance,¹⁹⁸ recommend primary injury-prevention programs. Although the bulk of the existing injury-prevention literature focuses on primary injury prevention, the continued increase in youth and adolescent ACL injuries creates a growing population needing secondary prevention strategies to prevent reinjury and OA.^{183,199}

Primary preventive training programs that optimize prophylactic benefits are multicomponent exercise programs incorporating agility, balance, flexibility, plyometrics, and strength exercises.²⁰⁰ Which of the exercise subtypes is most influential in reducing injury risk and rate is unknown, yet the combination of various exercises appears important. Also, participants should receive continual feedback regarding proper technique, such as reminders to bend at the knees and hips, land softly, keep the knees over the toes, and avoid dynamic knee valgus (ie, knees caving inward).^{201–204} These programs aim to improve lower extremity biomechanics and elements of neuromuscular control and are strongly recommended to reduce lower extremity injury risks and rates.^{196–198}

To optimize widespread adoption, ACL injury-prevention programs are typically 10 to 20 minutes long and used as a preactivity warm-up.²⁰⁵ When possible, advocates for preventive training programs should work with coaches, team captains, or other stakeholders. They should review the stakeholder's existing warm-up program to determine if minor adjustments could be made to ensure that it incorporates key exercise types. Existing resources, such

as the “Remain in the Game: A Joint Effort” tool kit (remaininthegame.org), are available to help health professionals and coaches develop tailored programs for their athletes. Stakeholders should consider factors such as the type of sport, level of competition, and physical ability of the athletes to tailor the program and appropriately challenge them if needed. Preventive training programs can reduce the risk of primary ACL injuries by 53%.²⁰⁶ Based on the success of primary ACL injury-prevention programs, similar prevention programs to maintain and enhance recovery during postoperative rehabilitation are also likely to reduce the incidence of second ACL injury.

Secondary ACL injury-prevention programs are effective in altering lower extremity biomechanics in those with an ACL reconstruction.^{207–209} Similar to primary ACL injury-prevention programs, the successful alteration of lower extremity biomechanics may lead to a reduced incidence of second ACL injury. During a 2-year intervention with 1435 athletes, those with a self-reported history of ACL injury in the control group had an ACL reinjury rate 5 times greater than that of athletes with a self-reported history of ACL injury who completed a preventive training program. Furthermore, no athlete with a prior ACL injury who completed the preventive training program sustained a noncontact reinjury.²¹⁰ In 2 other studies,^{211,212} the incidence of second ACL injury was only 2.5% among male and 23% among female athletes who performed a secondary prevention program, compared with an incidence rate of 30% for a second ACL injury based on earlier publications.^{48,56,183,213,214} Hence, secondary injury-prevention programs may be effective in male athletes but less so in female athletes. However, female athletes in this cohort had only a 14% incidence of contralateral second ACL injury, which is a promising finding. Across all studies^{210–212} examining preventive training exercises in those with an ACL reconstruction, the incidence of second ACL injury was 8%. These data provide initial evidence to support the recommendation of implementing preventive training programs to reduce the risk of secondary ACL injury; still, future research is needed in this area.

Dissenting Opinion. In theory, this is a great recommendation. A multicomponent training program is important. Nonetheless, the authors of 2 systematic reviews^{215,216} concluded that multicomponent programs were effective but raised concerns that balance may not be a beneficial component and that time previously dedicated to balance activities may be more beneficial if used for technique training, strengthening, or another component.

13. Organizations should optimize socially supportive environments for athletes, service members, employees, providers, families, and caregivers. This can be achieved by recognizing and providing support for psychosocial stressors during the rehabilitation process, including injury-related stigma, uncertainty about return to activity, and threats to personal identity

People with a knee injury who return to activity sooner than they are psychologically ready may experience reinjury, injury to different body parts, anxiety, fear, depression, demotivation, and a decrease in sport performance.^{217–219} Extrinsic motivations such as “feeling

pressured by their coach to return” or feeling “guilty for letting others down if they do not return” were associated with negative psychological outcomes including increased worry and concern.²²⁰ Threats to successful rehabilitation after an injury involve fear and lack of confidence in returning to activity (eg, sport or active military duty) and unrealistic expectations of one’s capabilities, support, and motivation (eg, to regain previous performance standards, to get back in the fight).^{221,222} Similarly, injured soldiers who return to duty too quickly because of a strong sense of personal commitment and expectations to do so may endanger war fighter proficiency, the safety and effectiveness of the unit, and their mission.^{223,224} Hence, we need to educate patients about normal injury recovery times and standardize communication from surgeon to therapist to other key stakeholders to ensure that everyone “remains on the same page” regarding prognosis expectations.²²⁵

Social support and perceived social support influence injury rehabilitation through a buffering effect that “protects” an individual from the detrimental effects of injury-related stressors involving rehabilitation.²²⁶ In addition, social support has a positive effect on an athlete’s adherence to an injury rehabilitation program.^{227–230} Suitable support structures, such as trust in a health care provider, trust in a rehabilitation program, and a facilitative environment,^{227,229,230} are crucial in optimizing adherence during injury rehabilitation. Although emotional support is most effective when provided by family and friends, task support may be most effective when provided by a rehabilitation clinician, such as a physical therapist or AT.²²⁸ Rehabilitation clinicians can serve a patient’s psychosocial needs using system-level task support and comprehensive strategies, including structured goal setting, action planning, and preoperative modeling.^{231,232}

Organizational leadership should recognize the personal social stigma associated with injury, including the sense of burden placed on the team.^{233,234} It may be beneficial for organizations to better normalize the rehabilitation experience by including teammates in the postinjury and rehabilitation experience and increase structural support via scheduling services, because executive function and mental processes are negatively affected during injury and rehabilitation.^{235–237} Further exploration of the role of organizational support in reducing the risk of OA after an injury would also be helpful.

14. Use technology to enable an individual with a knee injury to monitor physical and psychological well-being over time, access evidence-based educational materials developed specifically for their needs, and engage in health promotion

In recent years, technology-driven tools have been developed for primary musculoskeletal injury prevention, promotion of health behaviors, and assessment of health-related outcomes among a broad spectrum of demographic groups.^{238–240} For example, the “Walk With Ease” program for individuals with OA^{241–243} leverages a web-based assessment tool to deliver a hybrid self-organized and community-based walking intervention to reduce OA-related symptoms and improve overall health. Similarly, the Osteoarthritis Action Alliance developed and disseminated the “Remain in the Game” web-based platform to

provide educational materials and evidence-based knee-injury risk-mitigation strategies to active individuals. Unfortunately, these interventions do not address the needs of patients who have sustained a knee injury. Importantly, they are not delivered using methods preferred by the young and active individuals who are most likely to experience an ACL injury (eg, mobile applications and commercially available wearable technology).^{244,245}

Mobile and web-based platforms can measure physical activity and biomechanical outcomes, particularly when linked to wearable technology. Furthermore, these platforms may inform individuals about their progress toward achieving short- and long-term goals.^{244–246} Regrettably, we lack a comprehensive technological solution that integrates all the aforementioned recommendations about the support and educational needs of individuals who have sustained a knee injury. Mobile applications, wearable technology, and web-based education that address areas relevant to this population (eg, see the physical, psychological, and social recommendations) may assist in monitoring patient progress or provide education, monitoring, activity promotion, social support, and alerts. A mobile application and potentially commercially available wearable sensors (eg, smart watches, activity trackers) should be used to collect information regarding the characteristics (eg, frequency, intensity, and duration) of exercise with which an individual participates, knee-related function or other patient-reported outcomes of interest, and symptoms during activities of daily living and exercise. A mobile application and potentially commercially available wearable sensors should also be leveraged to deliver individualized and self-driven exercise interventions to maintain knee and overall health after an individual’s completion of formalized rehabilitation.^{239,240} Moreover, mobile applications can provide a portal for chat or other forms of engagement with a community of individuals experiencing a similar recovery process. Finally, mobile notifications should alert an individual when the patient-reported outcome measures indicate that engagement with a health care provider is indicated. This mobile application could also provide links to health care providers in the local area to reduce user friction with the health care system.

15. Practitioners ought to consider both the ethics and efficacy of how and what technology is adopted for monitoring health outcomes and OA prevention after joint injury

Considering the pace of technological health and wellness solutions entering the marketplace, a “buyer-beware” mentality is wise for both patients and practitioners. In addition, the development and implementation of technology-based solutions need to occur in a manner that allows a user to maintain control of the data and the interaction with the new technology. For example, the development of mobile applications to deliver rehabilitation information and increase adherence to exercise should be engaged wisely. Much of our current technology has been designed to use cognitive-behavioral prompts (ie, “nudges”) to users toward certain behaviors and adherence to intervention goals.^{247,248} Prompts can create adherence, but when misused, they can lead to the unhealthy use of technology and fail to benefit the patient. The potential for

technology to provide great access to care and support healthy lifestyles is ideal. As with in-person treatment, it is often prudent to seek the services of competent professionals.

Dissenting Opinion. The recommendation may lack the specificity needed to inform practitioners. Monitoring of health outcomes is different from using technology to promote adherence to or participation in injury prevention. The safety of personal health information on any system should be protected per standard protocols. Existing protocols are not fully secure from spyware and phishing attacks. The use of electronic health records has put all patient information at risk.

Educative nudges²⁴⁹—similar to those used in technology for ACL injury prevention—act to affect people’s choice making by engaging their insightful abilities. Recipients are likely to receive informational materials with this expectation, and they would not expect to be affected in other ways. The ethical concern of reminding patients with educative nudges has not been substantiated.²⁴⁹ The potential for benefit from this technology and virtual visits is enormous, with minimal risk or hazard.^{247,248}

16. An interdisciplinary, coordinated, patient-centered care strategy is advised to comprehensively address the needs of patients with a history of knee injury

Clinicians need to recognize the multifaceted nature of a patient’s state. An ideal strategy to implement the aforementioned recommendations to address a patient’s needs may be to deploy a personalized, interdisciplinary comprehensive approach that uses technology-driven solutions. Clinicians who manage patients with chronic symptoms (eg, OA) have increasingly adopted coordinated interdisciplinary care models to optimize outcomes.^{250–253} For example, the Osteoarthritis Chronic Care Program includes exercise and occupational therapy, psychological support, and medical management for Australians with hip or knee OA.^{252,254} Many people who complete the program become unwilling to undergo knee surgery, partially because of improved knee pain and function. These unperformed arthroplasties may contribute to saving greater than \$1 million per year at some clinical sites.^{252,254} Despite established coordinated care models for certain chronic conditions, testing of coordinated care strategies tailored to younger people with chronic symptoms after an ACL reconstruction is needed. For example, these programs could offer this population

- a. Anti-inflammatory medications (as needed),
- b. Adaptive supervised rehabilitation with a home exercise program and injury-prevention exercises,
- c. Education to maintain or improve self-efficacy and promote adherence and stress management, and
- d. Social support.

Furthermore, coordinated interdisciplinary care models could also be adapted for when a person first engages with the health care team after an ACL injury to optimize outcomes. Members of the coordinated care team may include (but are not limited to) primary care physicians, orthopaedic surgeons, physical therapists, ATs, psychologists or mental health consultants, strength and conditioning specialists, and nutritionists. Despite the need for more

evidence to support these types of programs in this population, the low risk and intuitive advantage of these strategies make them appealing.

DISCUSSION

These consensus-based recommendations provide clinicians with secondary prevention strategies intended to reduce the risk of OA among patients after an ACL injury. We acknowledge the dearth of high-quality evidence to inform some of these recommendations and offer dissenting opinions to facilitate discussion. These consensus recommendations are not a formal clinical practice guideline based on a synthesis of high-quality evidence. Instead, this document reflects information synthesized from an interdisciplinary group of experts based on the best available evidence from the literature or personal experience. Task group members often weighed the level of evidence with the perceived risk-to-benefit ratio of advocating (or not advocating) for these recommendations. Members were predominately based in the United States, which may influence how clinicians outside the United States interpret the applicability of these recommendations to their clinical practice. We hope these recommendations foster more discussion on how we can help ensure the long-term wellness of people after joint injuries in general, not just ACL injuries. We trust that these recommendations will spark discussion about what patients need to know to be informed consumers in a health care system. We acknowledge the need for more high-quality research studies to shape future recommendations, and we hope this document raises awareness among clinicians and researchers so that they will take steps to mitigate the risk of OA after an ACL injury.

ACKNOWLEDGMENTS

This publication was supported by the Centers for Disease Control and Prevention (CDC) of the US Department of Health and Human Services (HHS) as part of a financial assistance award (1 NU58 DP006980-01) totaling \$461 914 with 65% funded by CDC/HHS and \$250 000 and 35% funded by nongovernment source(s). The contents are those of the authors and do not necessarily represent the official views of, nor an endorsement by, CDC/HHS or the US government.

REFERENCES

1. Mall NA, Chalmers PN, Moric M, et al. Incidence and trends of anterior cruciate ligament reconstruction in the United States. *Am J Sports Med.* 2014;42(10):2363–2370. doi:10.1177/0363546514542796
2. Hootman JM, Dick R, Agel J. Epidemiology of collegiate injuries for 15 sports: summary and recommendations for injury prevention initiatives. *J Athl Train.* 2007;42(2):311–319.
3. Joseph AM, Collins CL, Henke NM, Yard EE, Fields SK, Comstock RD. A multisport epidemiologic comparison of anterior cruciate ligament injuries in high school athletics. *J Athl Train.* 2013;48(6):810–817. doi:10.4085/1062-6050-48.6.03
4. Frobell RB, Roos HP, Roos EM, Roemer FW, Ranstam J, Lohmander LS. Treatment for acute anterior cruciate ligament tear: five year outcome of randomised trial. *BMJ.* 2013;346(7895):f232. doi:10.1136/bmj.f232
5. Harris KP, Driban JB, Sittler MR, Cattano NM, Balasubramanian E, Hootman JM. Tibiofemoral osteoarthritis after surgical or nonsurgical treatment of anterior cruciate ligament rupture: a

- systematic review. *J Athl Train*. 2017;52(6):507–517. doi:10.4085/1062-6050-49.3.89
6. Luc B, Gribble PA, Pietrosimone BG. Osteoarthritis prevalence following anterior cruciate ligament reconstruction: a systematic review and numbers-needed-to-treat analysis. *J Athl Train*. 2014;49(6):806–819. doi:10.4085/1062-6050-49.3.35
 7. Mather RC III, Koenig L, Kocher MS, et al; MOON Knee Group. Societal and economic impact of anterior cruciate ligament tears. *J Bone Joint Surg Am*. 2013;95(19):1751–1759. doi:10.2106/JBJS.L.01705
 8. Driban JB, Harkey MS, Liu SH, Salzler M, McAlindon TE. Osteoarthritis and aging: young adults with osteoarthritis. *Curr Epidemiol Rep*. 2020;7(1):9–15. doi:10.1007/s40471-020-00224-7
 9. Snoeker B, Turkiewicz A, Magnusson K, et al. Risk of knee osteoarthritis after different types of knee injuries in young adults: a population-based cohort study. *Br J Sports Med*. 2020;54(12):725–730. doi:10.1136/bjsports-2019-100959
 10. Meehan WP, Weisskopf MG, Krishnan S, et al. Relation of anterior cruciate ligament tears to potential chronic cardiovascular diseases. *Am J Cardiol*. 2018;122(11):1879–1884. doi:10.1016/j.amjcard.2018.08.030
 11. National public health agenda for osteoarthritis: 2020 update. Osteoarthritis Action Alliance. Accessed May 4, 2022. <https://oaaaction.unc.edu/policy/oa-agenda-2020-update>
 12. Bannuru RR, Osani MC, Vaysbrot EE, et al. OARSI guidelines for the non-surgical management of knee, hip, and polyarticular osteoarthritis. *Osteoarthritis Cartilage*. 2019;27(11):1578–1589. doi:10.1016/j.joca.2019.06.011
 13. Kolasinski SL, Neogi T, Hochberg MC, et al. 2019 American College of Rheumatology/Arthritis Foundation guideline for the management of osteoarthritis of the hand, hip, and knee. *Arthritis Rheumatol*. 2020;72(2):220–233. doi:10.1002/art.41142
 14. Nelson AE, Allen KD, Golightly YM, Goode AP, Jordan JM. A systematic review of recommendations and guidelines for the management of osteoarthritis: the chronic osteoarthritis management initiative of the US Bone and Joint Initiative. *Semin Arthritis Rheum*. 2014;43(6):701–712. doi:10.1016/j.semarthrit.2013.11.012
 15. Bennell KL, van Ginckel A, Kean CO, et al. Patient knowledge and beliefs about knee osteoarthritis after anterior cruciate ligament injury and reconstruction. *Arthritis Care Res (Hoboken)*. 2016;68(8):1180–1185. doi:10.1002/acr.22794
 16. Paterick TE, Patel N, Tajik AJ, Chandrasekaran K. Improving health outcomes through patient education and partnerships with patients. *Proc (Bayl Univ Med Cent)*. 2017;30(1):112–113. doi:10.1080/08998280.2017.11929552
 17. Pietrosimone B, Blackburn JT, Golightly YM, et al. Certified athletic trainers' knowledge and perceptions of posttraumatic osteoarthritis after knee injury. *J Athl Train*. 2017;52(6):541–559. doi:10.4085/1062-6050-51.2.13
 18. Palmieri-Smith RM, Cameron KL, DiStefano LJ, et al. The role of athletic trainers in preventing and managing posttraumatic osteoarthritis in physically active populations: a consensus statement of the Athletic Trainers' Osteoarthritis Consortium. *J Athl Train*. 2017;52(6):610–623. doi:10.4085/1062-6050-52.2.04
 19. Dalhaug EM, Haakstad LAH. What the health? information sources and maternal lifestyle behaviors. *Interact J Med Res*. 2019;8(3):e10355. doi:10.2196/10355
 20. Lee JL, Rawl SM, Dickinson S, et al. Communication about health information technology use between patients and providers. *J Gen Intern Med*. 2020;35(9):2614–2620. doi:10.1007/s11606-020-05903-1
 21. Belyeu BM, Klein JW, Reisch LM, et al. Patients' perceptions of their doctors' notes and after-visit summaries: a mixed methods study of patients at safety-net clinics. *Health Expect*. 2018;21(2):485–493. doi:10.1111/hex.12641
 22. Mariano DJ, Liu A, Eppler SL, et al. Does a question prompt list improve perceived involvement in care in orthopaedic surgery compared with the AskShareKnow questions? A pragmatic randomized controlled trial. *Clin Orthop Relat Res*. 2021;479(2):225–232. doi:10.1097/cor.0000000000001582
 23. Hinman RS, Delany CM, Campbell PK, Gale J, Bennell KL. Physical therapists, telephone coaches, and patients with knee osteoarthritis: qualitative study about working together to promote exercise adherence. *Phys Ther*. 2016;96(4):479–493. doi:10.2522/ptj.20150260
 24. Mobasheri A, Costello K. Podcasting: an innovative tool for enhanced osteoarthritis education and research dissemination. *Osteoarthr Cartil Open*. 2020;3(1):100130. doi:10.1016/j.ocarto.2020.100130
 25. Beischer S, Hamrin Senorski E, Thomeé C, Samuelsson K, Thomeé R. Young athletes return too early to knee-strenuous sport, without acceptable knee function after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc*. 2018;26(7):1966–1974. doi:10.1007/s00167-017-4747-8
 26. Palmieri-Smith RM, Lepley LK. Quadriceps strength asymmetry after anterior cruciate ligament reconstruction alters knee joint biomechanics and functional performance at time of return to activity. *Am J Sports Med*. 2015;43(7):1662–1669. doi:10.1177/0363546515578252
 27. Ithurburn MP, Paterno MV, Ford KR, Hewett TE, Schmitt LC. Young athletes with quadriceps femoris strength asymmetry at return to sport after anterior cruciate ligament reconstruction demonstrate asymmetric single-leg drop-landing mechanics. *Am J Sports Med*. 2015;43(11):2727–2737. doi:10.1177/0363546515602016
 28. Schmitt LC, Paterno MV, Ford KR, Myer GD, Hewett TE. Strength asymmetry and landing mechanics at return to sport after anterior cruciate ligament reconstruction. *Med Sci Sports Exerc*. 2015;47(7):1426–1434. doi:10.1249/mss.0000000000000560
 29. Sugimoto D, Heyworth BE, Yates BA, Kramer DE, Kocher MS, Micheli LJ. Effect of graft type on thigh circumference, knee range of motion, and lower-extremity strength in pediatric and adolescent males following anterior cruciate ligament reconstruction. *J Sport Rehabil*. 2020;29(5):555–562. doi:10.1123/jsr.2018-0272
 30. Adams D, Logerstedt DS, Hunter-Giordano A, Axe MJ, Snyder-Mackler L. Current concepts for anterior cruciate ligament reconstruction: a criterion-based rehabilitation progression. *J Orthop Sports Phys Ther*. 2012;42(7):601–614. doi:10.2519/jospt.2012.3871
 31. Shelbourne KD, Gray T. Minimum 10-year results after anterior cruciate ligament reconstruction: how the loss of normal knee motion compounds other factors related to the development of osteoarthritis after surgery. *Am J Sports Med*. 2009;37(3):471–480. doi:10.1177/0363546508326709
 32. Shelbourne KD, Urch SE, Gray T, Freeman H. Loss of normal knee motion after anterior cruciate ligament reconstruction is associated with radiographic arthritic changes after surgery. *Am J Sports Med*. 2012;40(1):108–113. doi:10.1177/0363546511423639
 33. Palmieri-Smith RM, Thomas AC, Wojtys EM. Maximizing quadriceps strength after ACL reconstruction. *Clin Sports Med*. 2008;27(3):405–424, vii–ix. doi:10.1016/j.csm.2008.02.001
 34. Øiestad BE, Juhl CB, Eitzen I, Thorlund JB. Knee extensor muscle weakness is a risk factor for development of knee osteoarthritis: a systematic review and meta-analysis. *Osteoarthritis Cartilage*. 2015;23(2):171–177. doi:10.1016/j.joca.2014.10.008
 35. Schmitt LC, Paterno MV, Hewett TE. The impact of quadriceps femoris strength asymmetry on functional performance at return to sport following anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther*. 2012;42(9):750–759. doi:10.2519/jospt.2012.4194

36. Toole AR, Ithurburn MP, Rauh MJ, Hewett TE, Paterno MV, Schmitt LC. Young athletes cleared for sports participation after anterior cruciate ligament reconstruction: how many actually meet recommended return-to-sport criterion cutoffs? *J Orthop Sports Phys Ther.* 2017;47(11):825–833. doi:10.2519/jospt.2017.7227
37. Kuenze CM, Hertel J, Hart JM. Quadriceps muscle function after exercise in men and women with a history of anterior cruciate ligament reconstruction. *J Athl Train.* 2014;49(6):740–746. doi:10.4085/1062-6050-49.3.46
38. Macias-Hernández SI, Miranda-Duarte A, Ramírez-Mora I, et al. Knee muscle strength correlates with joint cartilage T2 relaxation time in young participants with risk factors for osteoarthritis. *Clin Rheumatol.* 2016;35(8):2087–2092. doi:10.1007/s10067-016-3333-7
39. Pietrosimone B, Pfeiffer SJ, Harkey MS, et al. Quadriceps weakness associates with greater T1ρ relaxation time in the medial femoral articular cartilage 6 months following anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2019;27(8):2632–2642. doi:10.1007/s00167-018-5290-y
40. Brunst C, Ithurburn MP, Zbojniewicz AM, Paterno MV, Schmitt LC. Return-to-sport quadriceps strength symmetry impacts 5-year cartilage integrity after anterior cruciate ligament reconstruction: a preliminary analysis. *J Orthop Res.* 2022;40(1):285–294. doi:10.1002/jor.25029
41. Tourville TW, Jarrell KM, Naud S, Slauterbeck JR, Johnson RJ, Beynon BD. Relationship between isokinetic strength and tibiofemoral joint space width changes after anterior cruciate ligament reconstruction. *Am J Sports Med.* 2014;42(2):302–311. doi:10.1177/0363546513510672
42. Wang HJ, Ao YF, Jiang D, et al. Relationship between quadriceps strength and patellofemoral joint chondral lesions after anterior cruciate ligament reconstruction. *Am J Sports Med.* 2015;43(9):2286–2292. doi:10.1177/0363546515588316
43. Angelozzi M, Madama M, Corsica C, et al. Rate of force development as an adjunctive outcome measure for return-to-sport decisions after anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther.* 2012;42(9):772–780. doi:10.2519/jospt.2012.3780
44. Blackburn JT, Pietrosimone B, Harkey MS, Luc BA, Pamukoff DN. Quadriceps function and gait kinetics after anterior cruciate ligament reconstruction. *Med Sci Sports Exerc.* 2016;48(9):1664–1670. doi:10.1249/mss.0000000000000963
45. Kuenze C, Lisee C, Birchmeier T, et al. Sex differences in quadriceps rate of torque development within 1 year of ACL reconstruction. *Phys Ther Sport.* 2019;38:36–43. doi:10.1016/j.ptsp.2019.04.008
46. Arhos EK, Thoma LM, Grindem H, Logerstedt D, Risberg MA, Snyder-Mackler L. Association of quadriceps strength symmetry and surgical status with clinical osteoarthritis five years after anterior cruciate ligament rupture. *Arthritis Care Res (Hoboken).* 2022;74(3):386–391. doi:10.1002/acr.24479
47. Gokeler A, Welling W, Zaffagnini S, Seil R, Padua D. Development of a test battery to enhance safe return to sports after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2017;25(1):192–199. doi:10.1007/s00167-016-4246-3
48. Grindem H, Snyder-Mackler L, Moksnes H, Engebretsen L, Risberg MA. Simple decision rules can reduce reinjury risk by 84% after ACL reconstruction: the Delaware-Oslo ACL Cohort Study. *Br J Sports Med.* 2016;50(13):804–808. doi:10.1136/bjsports-2016-096031
49. Hauger AV, Reiman MP, Bjordal JM, Sheets C, Ledbetter L, Goode AP. Neuromuscular electrical stimulation is effective in strengthening the quadriceps muscle after anterior cruciate ligament surgery. *Knee Surg Sports Traumatol Arthrosc.* 2018;26(2):399–410. doi:10.1007/s00167-017-4669-5
50. Moore JM, Cessford K, Willmott AP, et al. Lower limb biomechanics before and after anterior cruciate ligament reconstruction: a systematic review. *J Biomech.* 2020;106:109828. doi:10.1016/j.jbiomech.2020.109828
51. Kumar D, Su F, Wu D, et al. Frontal plane knee mechanics and early cartilage degeneration in people with anterior cruciate ligament reconstruction: a longitudinal study. *Am J Sports Med.* 2018;46(2):378–387. doi:10.1177/0363546517739605
52. Shimizu T, Markes AR, Samaan MA, et al. Patients with abnormal limb kinetics at 6 months after anterior cruciate ligament reconstruction have an increased risk of persistent medial meniscal abnormality at 3 years. *Orthop J Sports Med.* 2020;8(1):2325967119895248. doi:10.1177/2325967119895248
53. Shimizu T, Samaan MA, Tanaka MS, et al. Abnormal biomechanics at 6 months are associated with cartilage degeneration at 3 years after anterior cruciate ligament reconstruction. *Arthroscopy.* 2019;35(2):511–520. doi:10.1016/j.arthro.2018.07.033
54. Culvenor AG, Patterson BE, Guermazi A, Morris HG, Whitehead TS, Crossley KM. Accelerated return to sport after anterior cruciate ligament reconstruction and early knee osteoarthritis features at 1 year: an exploratory study. *PM R.* 2018;10(4):349–356. doi:10.1016/j.pmrj.2017.09.005
55. Buckthorpe M. Optimising the late-stage rehabilitation and return-to-sport training and testing process after ACL reconstruction. *Sports Med.* 2019;49(7):1043–1058. doi:10.1007/s40279-019-01102-z
56. Paterno MV, Rauh MJ, Schmitt LC, Ford KR, Hewett TE. Incidence of second ACL injuries 2 years after primary ACL reconstruction and return to sport. *Am J Sports Med.* 2014;42(7):1567–1573. doi:10.1177/0363546514530088
57. Almeida AM, Santos Silva PR, Pedrinelli A, Hernandez AJ. Aerobic fitness in professional soccer players after anterior cruciate ligament reconstruction. *PLoS One.* 2018;13(3):e0194432. doi:10.1371/journal.pone.0194432
58. de Mille P, Osmak J. Performance: bridging the gap after ACL surgery. *Curr Rev Musculoskelet Med.* 2017;10(3):297–306. doi:10.1007/s12178-017-9419-2
59. Taylor JB, Owen E, Ford KR. Incorporating workload measures into rehabilitation after anterior cruciate ligament reconstruction: a case report. *Int J Sports Phys Ther.* 2020;15(5):823–831. doi:10.26603/ijsp.20200823
60. Welling W, Benjaminse A, Seil R, Lemmink K, Zaffagnini S, Gokeler A. Low rates of patients meeting return to sport criteria 9 months after anterior cruciate ligament reconstruction: a prospective longitudinal study. *Knee Surg Sports Traumatol Arthrosc.* 2018;26(12):3636–3644. doi:10.1007/s00167-018-4916-4
61. King E, Richter C, Daniels KAJ, et al. Biomechanical but not strength or performance measures differentiate male athletes who experience ACL reinjury on return to level 1 sports. *Am J Sports Med.* 2021;49(4):918–927. doi:10.1177/0363546520988018
62. King E, Richter C, Franklyn-Miller A, et al. Biomechanical but not timed performance asymmetries persist between limbs 9 months after ACL reconstruction during planned and unplanned change of direction. *J Biomech.* 2018;81:93–103. doi:10.1016/j.jbiomech.2018.09.021
63. King E, Richter C, Franklyn-Miller A, et al. Whole-body biomechanical differences between limbs exist 9 months after ACL reconstruction across jump/landing tasks. *Scand J Med Sci Sports.* 2018;28(12):2567–2578. doi:10.1111/sms.13259
64. Hawkins RD, Hulse MA, Wilkinson C, Hodson A, Gibson M. The association football medical research programme: an audit of injuries in professional football. *Br J Sports Med.* 2001;35(1):43–47. doi:10.1136/bjbm.35.1.43
65. Nagle K, Johnson B, Brou L, Landman T, Sochanska A, Comstock RD. Timing of lower extremity injuries in competition and practice

- in high school sports. *Sports Health*. 2017;9(3):238–246. doi:10.1177/1941738116685704
66. Tallard JC, Hedt C, Lambert BS, McCulloch PC. The role of fatigue in return to sport testing following anterior cruciate ligament reconstruction. *Int J Sports Phys Ther*. 2021;16(4):1043–1051. doi:10.26603/001c.25687
 67. Ardern CL, Glasgow P, Schneiders A, et al. 2016 consensus statement on return to sport from the First World Congress in Sports Physical Therapy, Bern. *Br J Sports Med*. 2016;50(14):853–864. doi:10.1136/bjsports-2016-096278
 68. Taberner M, Allen T, Cohen DD. Progressing rehabilitation after injury: consider the “control-chaos continuum.” *Br J Sports Med*. 2019;53(18):1132–1136. doi:10.1136/bjsports-2018-100157
 69. Della Villa S, Boldrini L, Ricci M, et al. Clinical outcomes and return-to-sports participation of 50 soccer players after anterior cruciate ligament reconstruction through a sport-specific rehabilitation protocol. *Sports Health*. 2012;4(1):17–24. doi:10.1177/1941738111417564
 70. Capin JJ, Behrns W, Thatcher K, Arundale A, Smith AH, Snyder-Mackler L. On-ice return-to-hockey progression after anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther*. 2017;47(5):324–333. doi:10.2519/jospt.2017.7245
 71. Waters E. Suggestions from the field for return to sports participation following anterior cruciate ligament reconstruction: basketball. *J Orthop Sports Phys Ther*. 2012;42(4):326–336. doi:10.2519/jospt.2012.4030
 72. Bizzini M, Hancock D, Impellizzeri F. Suggestions from the field for return to sports participation following anterior cruciate ligament reconstruction: soccer. *J Orthop Sports Phys Ther*. 2012;42(4):304–312. doi:10.2519/jospt.2012.4005
 73. Buckthorpe M. Recommendations for movement re-training after ACL reconstruction. *Sports Med*. 2021;51(8):1601–1618. doi:10.1007/s40279-021-01454-5
 74. Meredith SJ, Rauer T, Chmielewski TL, et al; Panther Symposium ACL Injury Return to Sport Consensus Group. Return to sport after anterior cruciate ligament injury: Panther Symposium ACL Injury Return to Sport Consensus Group. *Orthop J Sports Med*. 2020;8(6):2325967120930829. doi:10.1177/2325967120930829
 75. van Melick N, van Cingel RE, Brooijmans F, et al. Evidence-based clinical practice update: practice guidelines for anterior cruciate ligament rehabilitation based on a systematic review and multidisciplinary consensus. *Br J Sports Med*. 2016;50(24):1506–1515. doi:10.1136/bjsports-2015-095898
 76. Kyritsis P, Bahr R, Landreau P, Miladi R, Witvrouw E. Likelihood of ACL graft rupture: not meeting six clinical discharge criteria before return to sport is associated with a five times greater risk of rupture. *Br J Sports Med*. 2016;50(15):946–951. doi:10.1136/bjsports-2015-095908
 77. Triplett AN, Kuenze CM. Characterizing body composition, cardiorespiratory fitness, and physical activity in women with anterior cruciate ligament reconstruction. *Phys Ther Sport*. 2021;48:54–59. doi:10.1016/j.ptsp.2020.12.014
 78. MacAlpine EM, Talwar D, Storey EP, Doroshov SM, Lawrence JTR. Weight gain after ACL reconstruction in pediatric and adolescent patients. *Sports Health*. 2020;12(1):29–35. doi:10.1177/1941738119870192
 79. Kuenze C, Collins K, Pfeiffer KA, Lisee C. Assessing physical activity after ACL injury: moving beyond return to sport. *Sports Health*. 2022;14(2):197–204. doi:10.1177/19417381211025307
 80. Filbay S, Andersson C, Gauffin H, Kvist J. Prognostic factors for patient-reported outcomes at 32 to 37 years after surgical or nonsurgical management of anterior cruciate ligament injury. *Orthop J Sports Med*. 2021;9(8):23259671211021592. doi:10.1177/23259671211021592
 81. Piercy KL, Troiano RP, Ballard RM, et al. The Physical Activity Guidelines for Americans. *JAMA*. 2018;320(19):2020–2028. doi:10.1001/jama.2018.14854
 82. Bell DR, Pfeiffer KA, Cadmus-Bertram LA, et al. Objectively measured physical activity in patients after anterior cruciate ligament reconstruction. *Am J Sports Med*. 2017;45(8):1893–1900. doi:10.1177/0363546517698940
 83. Kuenze C, Cadmus-Bertram L, Pfeiffer K, et al. Relationship between physical activity and clinical outcomes after ACL reconstruction. *J Sport Rehabil*. 2019;28(2):180–187. doi:10.1123/jsr.2017-0186
 84. Lisee CM, Montoye AHK, Lewallen NF, Hernandez M, Bell DR, Kuenze CM. Assessment of free-living cadence using ActiGraph accelerometers between individuals with and without anterior cruciate ligament reconstruction. *J Athl Train*. 2020;55(9):994–1000. doi:10.4085/1062-6050-425-19
 85. Butler CR, Allen K, DiStefano LJ, Lopley LK. Protracted cardiovascular impairments after anterior cruciate ligament injury: a critically appraised topic. *J Sport Rehabil*. 2020;29(5):680–683. doi:10.1123/jsr.2019-0175
 86. Wu HH, Liu M, Dines JS, Kelly JD, Garcia GH. Depression and psychiatric disease associated with outcomes after anterior cruciate ligament reconstruction. *World J Orthop*. 2016;7(11):709–717. doi:10.5312/wjo.v7.i11.709
 87. Thoma LM, Snyder-Mackler L, Risberg M, White DK. Trajectories of weight gain in young adults following anterior cruciate ligament rupture: the Delaware-Oslo ACL cohort study. *Osteoarthritis Cartilage*. 2019;27(suppl 1):S274–S275. doi:10.1016/j.joca.2019.02.653
 88. Garcia GH, Wu HH, Park MJ, et al. Depression symptomatology and anterior cruciate ligament injury: incidence and effect on functional outcome—a prospective cohort study. *Am J Sports Med*. 2016;44(3):572–579. doi:10.1177/0363546515612466
 89. Kraus VB, Sprow K, Powell KE, et al; 2018 Physical Activity Guidelines Advisory Committee. Effects of physical activity in knee and hip osteoarthritis: a systematic umbrella review. *Med Sci Sports Exerc*. 2019;51(6):1324–1339. doi:10.1249/mss.0000000000001944
 90. Turner MN, Hernandez DO, Cade W, Emerson CP, Reynolds JM, Best TM. The role of resistance training dosing on pain and physical function in individuals with knee osteoarthritis: a systematic review. *Sports Health*. 2020;12(2):200–206. doi:10.1177/1941738119887183
 91. Hall KS, Hyde ET, Bassett DR, et al. Systematic review of the prospective association of daily step counts with risk of mortality, cardiovascular disease, and dysglycemia. *Int J Behav Nutr Phys Act*. 2020;17(1):78. doi:10.1186/s12966-020-00978-9
 92. Pescatello LS, Buchner DM, Jakicic JM, et al; 2018 Physical Activity Guidelines Advisory Committee. Physical activity to prevent and treat hypertension: a systematic review. *Med Sci Sports Exerc*. 2019;51(6):1314–1323. doi:10.1249/mss.0000000000001943
 93. Jakicic JM, Powell KE, Campbell WW, et al; 2018 Physical Activity Guidelines Advisory Committee. Physical activity and the prevention of weight gain in adults: a systematic review. *Med Sci Sports Exerc*. 2019;51(6):1262–1269. doi:10.1249/mss.0000000000001938
 94. Dishman RK, McDowell CP, Herring MP. Customary physical activity and odds of depression: a systematic review and meta-analysis of 111 prospective cohort studies. *Br J Sports Med*. 2021;55(16):926–934. doi:10.1136/bjsports-2020-103140
 95. Alentorn-Geli E, Samuelsson K, Musahl V, Green CL, Bhandari M, Karlsson J. The association of recreational and competitive running with hip and knee osteoarthritis: a systematic review and meta-analysis. *J Orthop Sports Phys Ther*. 2017;47(6):373–390. doi:10.2519/jospt.2017.7137
 96. Chakravarty EF, Hubert HB, Lingala VB, Zatarain E, Fries JF. Long distance running and knee osteoarthritis: a prospective study.

- Am J Prev Med.* 2008;35(2):133–138. doi:10.1016/j.amepre.2008.03.032
97. Øiestad BE, Holm I, Risberg MA. Return to pivoting sport after ACL reconstruction: association with osteoarthritis and knee function at the 15-year follow-up. *Br J Sports Med.* 2018;52(18):1199–1204. doi:10.1136/bjsports-2017-097718
 98. Øiestad BE, Quinn E, White D, et al. No association between daily walking and knee structural changes in people at risk of or with mild knee osteoarthritis: prospective data from the Multicenter Osteoarthritis Study. *J Rheumatol.* 2015;42(9):1685–1693. doi:10.3899/jrheum.150071
 99. Timmins KA, Leech RD, Batt ME, Edwards KL. Running and knee osteoarthritis: a systematic review and meta-analysis. *Am J Sports Med.* 2017;45(6):1447–1457. doi:10.1177/0363546516657531
 100. Lefèvre-Colau MM, Nguyen C, Haddad R, et al. Is physical activity, practiced as recommended for health benefit, a risk factor for osteoarthritis? *Ann Phys Rehabil Med.* 2016;59(3):196–206. doi:10.1016/j.rehab.2016.02.007
 101. Driban JB, Hootman JM, Sitler MR, Harris KP, Cattano NM. Is participation in certain sports associated with knee osteoarthritis? a systematic review. *J Athl Train.* 2017;52(6):497–506. doi:10.4085/1062-6050-50.2.08
 102. Friedman JM, Su F, Zhang AL, et al. Patient-reported activity levels correlate with early cartilage degeneration after anterior cruciate ligament reconstruction. *Am J Sports Med.* 2021;49(2):442–449. doi:10.1177/0363546520980431
 103. Doré DA, Winzenberg TM, Ding C, et al. The association between objectively measured physical activity and knee structural change using MRI. *Ann Rheum Dis.* 2013;72(7):1170–1175. doi:10.1136/annrheumdis-2012-201691
 104. Medvecky MJ, Nelson S. Kinesiophobia and return to sports after anterior cruciate ligament reconstruction. *Conn Med.* 2015;79(3):155–157.
 105. Heil J. *Psychology of Sport Injury.* Human Kinetics Publishers; 1993.
 106. Schuman-Olivier Z, Trombka M, Lovas DA, et al. Mindfulness and behavior change. *Harv Rev Psychiatry.* 2020;28(6):371–394. doi:10.1097/hrp.0000000000000277
 107. Mahood C, Perry M, Gallagher P, Sole G. Chaos and confusion with confidence: managing fear of re-injury after anterior cruciate ligament reconstruction. *Phys Ther Sport.* 2020;45:145–154. doi:10.1016/j.ptsp.2020.07.002
 108. Cupal DD, Brewer BW. Effects of relaxation and guided imagery on knee strength, reinjury anxiety, and pain following anterior cruciate ligament reconstruction. *Rehabil Psychol.* 2001;46(1):28–43. doi:10.1037/0090-5550.46.1.28
 109. Coronado RA, Sterling EK, Fenster DE, et al. Cognitive-behavioral-based physical therapy to enhance return to sport after anterior cruciate ligament reconstruction: an open pilot study. *Phys Ther Sport.* 2020;42:82–90. doi:10.1016/j.ptsp.2020.01.004
 110. Kabat-Zinn J. Mindfulness-based interventions in context: past, present, and future. *Clin Psychol Sci Pract.* 2003;10(2):144–156. doi:10.1093/clipsy.bpg016
 111. Shires A, Sharpe L, Davies JN, Newton-John TRO. The efficacy of mindfulness-based interventions in acute pain: a systematic review and meta-analysis. *Pain.* 2020;161(8):1698–1707. doi:10.1097/j.pain.0000000000001877
 112. Lee AC, Harvey WF, Price LL, et al. Mindfulness is associated with treatment response from nonpharmacologic exercise interventions in knee osteoarthritis. *Arch Phys Med Rehabil.* 2017;98(11):2265–2273.e1. doi:10.1016/j.apmr.2017.04.014
 113. Lee AC, Harvey WF, Price LL, Morgan LPK, Morgan NL, Wang C. Mindfulness is associated with psychological health and moderates pain in knee osteoarthritis. *Osteoarthritis Cartilage.* 2017;25(6):824–831. doi:10.1016/j.joca.2016.06.017
 114. Innes KE, Selfe TK, Kandati S, Wen S, Huysmans Z. Effects of mantra meditation versus music listening on knee pain, function, and related outcomes in older adults with knee osteoarthritis: an exploratory randomized clinical trial (RCT). *Evid Based Complement Alternat Med.* 2018;2018:7683897. doi:10.1155/2018/7683897
 115. Wang C, Schmid CH, Iversen MD, et al. Comparative effectiveness of tai chi versus physical therapy for knee osteoarthritis: a randomized trial. *Ann Intern Med.* 2016;165(2):77–86. doi:10.7326/M15-2143
 116. Ghandali E, Moghadam ST, Hadian MR, Olyaei G, Jalaie S, Sajjadi E. The effect of tai chi exercises on postural stability and control in older patients with knee osteoarthritis. *J Bodyw Mov Ther.* 2017;21(3):594–598. doi:10.1016/j.jbmt.2016.09.001
 117. Ye J, Zheng Q, Zou L, et al. Mindful exercise (Baduanjin) as an adjuvant treatment for older adults (60 years old and over) of knee osteoarthritis: a randomized controlled trial. *Evid Based Complement Alternat Med.* 2020;2020:9869161. doi:10.1155/2020/9869161
 118. Ahn H, Zhong C, Miao H, et al. Efficacy of combining home-based transcranial direct current stimulation with mindfulness-based meditation for pain in older adults with knee osteoarthritis: a randomized controlled pilot study. *J Clin Neurosci.* 2019;70:140–145. doi:10.1016/j.jocn.2019.08.047
 119. Ebnezar J, Nagarathna R, Yogitha B, Nagendra HR. Effect of integrated yoga therapy on pain, morning stiffness and anxiety in osteoarthritis of the knee joint: a randomized control study. *Int J Yoga.* 2012;5(1):28–36. doi:10.4103/0973-6131.91708
 120. Conroy D, Hagger MS. Imagery interventions in health behavior: a meta-analysis. *Health Psychol.* 2018;37(7):668–679. doi:10.1037/hea0000625
 121. Rodriguez RM, Marroquin A, Cosby N. Reducing fear of reinjury and pain perception in athletes with first-time anterior cruciate ligament reconstructions by implementing imagery training. *J Sport Rehabil.* 2019;28(4):385–389. doi:10.1123/jsr.2017-0056
 122. Maddison R, Prapavessis H, Clatworthy M. Modeling and rehabilitation following anterior cruciate ligament reconstruction. *Ann Behav Med.* 2006;31(1):89–98. doi:10.1207/s15324796abm3101_13
 123. Zaffagnini S, Russo RL, Marcheggiani Muccioli GM, Marcacci M. The Videosight® method: improving rehabilitation following anterior cruciate ligament reconstruction—a preliminary study. *Knee Surg Sports Traumatol Arthrosc.* 2013;21(4):851–858. doi:10.1007/s00167-013-2392-4
 124. Gledhill A, Forsdyke D, Murray E. Psychological interventions used to reduce sports injuries: a systematic review of real-world effectiveness. *Br J Sports Med.* 2018;52(15):967–971. doi:10.1136/bjsports-2017-097694
 125. Petterson H, Olson BL. Effects of mindfulness-based interventions in high school and college athletes for reducing stress and injury, and improving quality of life. *J Sport Rehabil.* 2017;26(6):578–587. doi:10.1123/jsr.2016-0047
 126. Tajdini H, Letafatkar A, Brewer BW, Hosseinzadeh M. Association between kinesiophobia and gait asymmetry after ACL reconstruction: implications for prevention of reinjury. *Int J Environ Res Public Health.* 2021;18(6):3264. doi:10.3390/ijerph18063264
 127. Paterno MV, Flynn K, Thomas S, Schmitt LC. Self-reported fear predicts functional performance and second ACL injury after ACL reconstruction and return to sport: a pilot study. *Sports Health.* 2018;10(3):228–233. doi:10.1177/1941738117745806
 128. Van Wyngaarden JJ, Jacobs C, Thompson K, et al. Quadriceps strength and kinesiophobia predict long-term function after ACL reconstruction: a cross-sectional pilot study. *Sports Health.* 2021;13(3):251–257. doi:10.1177/1941738120946323

129. van Lankveld W, van Melick N, Habets B, Roelofsens E, Staal JB, van Cingel R. Measuring individual hierarchy of anxiety invoking sports related activities: development and validation of the Photographic Series of Sports Activities for Anterior Cruciate Ligament Reconstruction (PHOSA-ACL). *BMC Musculoskeletal Disord.* 2017;18(1):287. doi:10.1186/s12891-017-1643-9
130. Miller RP, Kori SH, Todd DD. The Tampa Scale: a measure of kinesiophobia. *Clin J Pain.* 1991;7(1):51.
131. Woby SR, Roach NK, Urmston M, Watson PJ. Psychometric properties of the TSK-11: a shortened version of the Tampa Scale for Kinesiophobia. *Pain.* 2005;117(1-2):137-144. doi:10.1016/j.pain.2005.05.029
132. Webster KE, Feller JA, Lambros C. Development and preliminary validation of a scale to measure the psychological impact of returning to sport following anterior cruciate ligament reconstruction surgery. *Phys Ther Sport.* 2008;9(1):9-15. doi:10.1016/j.ptsp.2007.09.003
133. Ariza-Mateos MJ, Cabrera-Martos I, Ortiz-Rubio A, Torres-Sánchez I, Rodríguez-Torres J, Valenza MC. Effects of a patient-centered graded exposure intervention added to manual therapy for women with chronic pelvic pain: a randomized controlled trial. *Arch Phys Med Rehabil.* 2019;100(1):9-16. doi:10.1016/j.apmr.2018.08.188
134. Cai L, Gao H, Xu H, Wang Y, Lyu P, Liu Y. Does a program based on cognitive behavioral therapy affect kinesiophobia in patients following total knee arthroplasty? a randomized, controlled trial with a 6-month follow-up. *J Arthroplasty.* 2018;33(3):704-710. doi:10.1016/j.arth.2017.10.035
135. George SZ, Wittmer VT, Fillingim RB, Robinson ME. Comparison of graded exercise and graded exposure clinical outcomes for patients with chronic low back pain. *J Orthop Sports Phys Ther.* 2010;40(11):694-704. doi:10.2519/jospt.2010.3396
136. Macedo LG, Smeets RJ, Maher CG, Latimer J, McAuley JH. Graded activity and graded exposure for persistent nonspecific low back pain: a systematic review. *Phys Ther.* 2010;90(6):860-879. doi:10.2522/ptj.20090303
137. Vlaeyen JW, de Jong J, Geilen M, Heuts PH, van Breukelen G. Graded exposure in vivo in the treatment of pain-related fear: a replicated single-case experimental design in four patients with chronic low back pain. *Behav Res Ther.* 2001;39(2):151-166. doi:10.1016/s0005-7967(99)00174-6
138. Vlaeyen JW, de Jong J, Geilen M, Heuts PH, van Breukelen G. The treatment of fear of movement/(re)injury in chronic low back pain: further evidence on the effectiveness of exposure in vivo. *Clin J Pain.* 2002;18(4):251-261. doi:10.1097/00002508-200207000-00006
139. Woods MP, Asmundson GJG. Evaluating the efficacy of graded in vivo exposure for the treatment of fear in patients with chronic back pain: a randomized controlled clinical trial. *Pain.* 2008;136(3):271-280. doi:10.1016/j.pain.2007.06.037
140. Leeuw M, Goossens MEJB, van Breukelen GJP, et al. Exposure in vivo versus operant graded activity in chronic low back pain patients: results of a randomized controlled trial. *Pain.* 2008;138(1):192-207. doi:10.1016/j.pain.2007.12.009
141. Baez S, Cormier M, Andreaatta R, Gribble P, Hoch JM. Implementation of in vivo exposure therapy to decrease injury-related fear in females with a history of ACL-reconstruction: a pilot study. *Phys Ther Sport.* 2021;52:217-223. doi:10.1016/j.ptsp.2021.09.009
142. Welcome to the Association for Behavioral and Cognitive Therapies (ABCT). Association for Cognitive and Behavioral Therapies. Accessed March 22, 2021. <https://www.abct.org/Home/>
143. Foo CN, Arumugam M, Lekhray R, Lye MS, Mohd-Sidik S, Jamil Osman Z. Effectiveness of health-led cognitive behavioral-based group therapy on pain, functional disability and psychological outcomes among knee osteoarthritis patients in Malaysia. *Int J Environ Res Public Health.* 2020;17(17):6179. doi:10.3390/ijerph17176179
144. Bennell KL, Ahamed Y, Jull G, et al. Physical therapist-delivered pain coping skills training and exercise for knee osteoarthritis: randomized controlled trial. *Arthritis Care Res (Hoboken).* 2016;68(5):590-602. doi:10.1002/acr.22744
145. Ismail A, Moore C, Alshishani N, Yaseen K, Alshehri MA. Cognitive behavioural therapy and pain coping skills training for osteoarthritis knee pain management: a systematic review. *J Phys Ther Sci.* 2017;29(12):2228-2235. doi:10.1589/jpts.29.2228
146. O'Moore KA, Newby JM, Andrews G, et al. Internet cognitive-behavioral therapy for depression in older adults with knee osteoarthritis: a randomized controlled trial. *Arthritis Care Res (Hoboken).* 2018;70(1):61-70. doi:10.1002/acr.23257
147. Jensen MP, McFarland CA. Increasing the reliability and validity of pain intensity measurement in chronic pain patients. *Pain.* 1993;55(2):195-203.
148. Farrar JT, Young JPI, LaMoreaux L, Werth JL, Poole RM. Clinical importance of changes in chronic pain intensity measured on an 11-point numerical pain rating scale. *Pain.* 2001;94(2):149-158.
149. Irrgang JJ, Anderson AF, Boland AL, et al. Development and validation of the International Knee Documentation Committee Subjective Knee Form. *Am J Sports Med.* 2001;29(5):600-613. doi:10.1177/03635465010290051301
150. Roos EM, Roos HP, Lohmander LS, Ekdahl C, Beynon BD. Knee Injury and Osteoarthritis Outcome Score (KOOS)—development of a self-administered outcome measure. *J Orthop Sports Phys Ther.* 1998;28(2):88-96.
151. Roos EM, Lohmander LS. Knee Injury and Osteoarthritis Outcome Score (KOOS): from joint injury to osteoarthritis. *Health Qual Life Outcomes.* 2003;1:64.
152. Collins NJ, Misra D, Felson DT, Crossley KM, Roos EM. Measures of knee function: International Knee Documentation Committee (IKDC) Subjective Knee Evaluation Form, Knee Injury and Osteoarthritis Outcome Score (KOOS), Knee Injury and Osteoarthritis Outcome Score Physical Function Short Form (KOOS-PS), Knee Outcome Survey Activities of Daily Living Scale (KOS-ADL), Lysholm Knee Scoring Scale, Oxford Knee Score (OKS), Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC), Activity Rating Scale (ARS), and Tegner Activity Score (TAS). *Arthritis Care Res.* 2011;63(suppl 11):S208-S228.
153. Davis AM, Perruccio AV, Canizares M, et al. Comparative, validity and responsiveness of the HOOS-PS and KOOS-PS to the womac physical function subscale in total joint replacement for osteoarthritis. *Osteoarthritis Cartilage.* 2009;17(7):843-847. doi:10.1016/j.joca.2009.01.005
154. Perruccio AV, Stefan Lohmander L, Canizares M, et al. The development of a short measure of physical function for knee OA KOOS-Physical Function Shortform (KOOS-PS)—an OARSI/OMERACT initiative. *Osteoarthritis Cartilage.* 2008;16(5):542-550. doi:10.1016/j.joca.2007.12.014
155. Evaluation tools. Stanford Patient Education Research Center. Accessed January 20, 2023. <https://selfmanagementresource.com/resources/evaluation-tools/english-evaluation-tools/>
156. Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc.* 1982;14(5):377-381.
157. Cella D, Riley W, Stone A, et al. The Patient-Reported Outcomes Measurement Information System (PROMIS) developed and tested its first wave of adult self-reported health outcome item banks: 2005-2008. *J Clin Epidemiol.* 2010;63(11):1179-1194. doi:10.1016/j.jclinepi.2010.04.011
158. Shelby RA, Somers TJ, Keefe FJ, et al. Brief fear of movement scale for osteoarthritis. *Arthritis Care Res (Hoboken).* 2012;64(6):862-871. doi:10.1002/acr.21626

159. Tkachuk GA, Harris CA. Psychometric properties of the Tampa Scale for Kinesiophobia-11 (TSK-11). *J Pain*. 2012;13(10):970–977. doi:10.1016/j.jpain.2012.07.001
160. Naal FD, Impellizzeri FM, Leunig M. Which is the best activity rating scale for patients undergoing total joint arthroplasty? *Clin Orthop Relat Res*. 2009;467(4):958–965.
161. Tegner Y, Lysholm J. Rating systems in the evaluation of knee ligament injuries. *Clin Orthop Relat Res*. 1985;198:43–49.
162. Marx RG, Stump TJ, Jones EC, Wickiewicz TL, Warren RF. Development and evaluation of an activity rating scale for disorders of the knee. *Am J Sports Med*. 2001;29(2):213–218. doi:10.1177/03635465010290021601
163. Idler EL, Russell LB, Davis D. Survival, functional limitations, and self-rated health in the NHANES I epidemiologic follow-up study, 1992: first National Health and Nutrition Examination Survey. *Am J Epidemiol*. 2000;152(9):874–883. doi:10.1093/aje/152.9.874
164. Wang Y, Teichtahl AJ, Wluka AE, et al. Associations of joint line tenderness and patellofemoral grind with long-term knee joint outcomes: data from the Osteoarthritis Initiative. *Arthritis Care Res (Hoboken)*. 2020;72(6):778–786. doi:10.1002/acr.23906
165. Lo GH, Strayhorn MT, Driban JB, Price LL, Eaton CB, McAlindon TE. Subjective crepitus as a risk factor for incident symptomatic knee osteoarthritis: data from the Osteoarthritis Initiative. *Arthritis Care Res (Hoboken)*. 2018;70(1):53–60. doi:10.1002/acr.23246
166. Wang Y, Martel-Pelletier J, Teichtahl AJ, et al. The bulge sign—a simple physical examination for identifying progressive knee osteoarthritis: data from the Osteoarthritis Initiative. *Rheumatology (Oxford)*. 2020;59(6):1288–1295. doi:10.1093/rheumatology/kez443
167. Lane AR, Harkey MS, Davis HC, et al. Body mass index and type 2 collagen turnover in individuals after anterior cruciate ligament reconstruction. *J Athl Train*. 2019;54(3):270–275. doi:10.4085/1062-6050-525-17
168. Kuenze C, Lisee C, Pfeiffer KA, et al. Sex differences in physical activity engagement after ACL reconstruction. *Phys Ther Sport*. 2019;35:12–17. doi:10.1016/j.ptsp.2018.10.016
169. Moksnes H, Risberg MA. Performance-based functional evaluation of non-operative and operative treatment after anterior cruciate ligament injury. *Scand J Med Sci Sports*. 2009;19(3):345–355. doi:10.1111/j.1600-0838.2008.00816.x
170. Kanko LE, Birmingham TB, Bryant DM, et al. The star excursion balance test is a reliable and valid outcome measure for patients with knee osteoarthritis. *Osteoarthritis Cartilage*. 2019;27(4):580–585. doi:10.1016/j.joca.2018.11.012
171. Dobson F, Hinman RS, Roos EM, et al. OARSI recommended performance-based tests to assess physical function in people diagnosed with hip or knee osteoarthritis. *Osteoarthritis Cartilage*. 2013;21(8):1042–1052. doi:10.1016/j.joca.2013.05.002
172. Pietrosimone B, Blackburn JT, Harkey MS, et al. Walking speed as a potential indicator of cartilage breakdown following anterior cruciate ligament reconstruction. *Arthritis Care Res (Hoboken)*. 2016;68(6):793–800. doi:10.1002/acr.22773
173. Pietrosimone B, Blackburn JT, Harkey MS, et al. Greater mechanical loading during walking is associated with less collagen turnover in individuals with anterior cruciate ligament reconstruction. *Am J Sports Med*. 2016;44(2):425–432. doi:10.1177/0363546515618380
174. Pfeiffer SJ, Spang J, Nissman D, et al. Gait mechanics and T1ρ MRI of tibiofemoral cartilage 6 months after acl reconstruction. *Med Sci Sports Exerc*. 2019;51(4):630–639. doi:10.1249/MSS.0000000000001834
175. Majewska J, Szczepanik M, Szymczyk D, et al. Evaluation of selected gait parameters in patients prior to and at 6 months following early anterior cruciate ligament reconstruction. *Ortop Traumatol Rehabil*. 2017;19(3):273–283.
176. Kellgren JH, Lawrence JS. Radiological assessment of osteoarthritis. *Ann Rheum Dis*. 1957;16(4):494–502. doi:10.1136/ard.16.4.494
177. Kellgren JH, Lawrence JS. *Atlas of Standard Radiographs*. Oxford; 1963.
178. Roemer FW, Frobell R, Lohmander LS, Niu J, Guermazi A. Anterior Cruciate Ligament OsteoArthritis Score (ACLOAS): longitudinal MRI-based whole joint assessment of anterior cruciate ligament injury. *Osteoarthritis Cartilage*. 2014;22(5):668–682. doi:10.1016/j.joca.2014.03.006
179. Frobell RB, Le Graverand MP, Buck R, et al. The acutely ACL injured knee assessed by MRI: changes in joint fluid, bone marrow lesions, and cartilage during the first year. *Osteoarthritis Cartilage*. 2009;17(2):161–167. doi:10.1016/j.joca.2008.06.020
180. Li X, Kuo D, Theologis A, et al. Cartilage in anterior cruciate ligament-reconstructed knees: MR imaging T1ρ and T2—initial experience with 1-year follow-up. *Radiology*. 2011;258(2):505–514.
181. Harkey MS, Luc BA, Golightly YM, et al. Osteoarthritis-related biomarkers following anterior cruciate ligament injury and reconstruction: a systematic review. *Osteoarthritis Cartilage*. 2015;23(1):1–12. doi:10.1016/j.joca.2014.09.004
182. Hunt ER, Villasanta-Tezanos AG, Butterfield TA, Lattermann C, Jacobs CA. Upregulation of systemic inflammatory pathways following anterior cruciate ligament injury relates to both cartilage and muscular changes: a pilot study. *J Orthop Res*. 2020;38(2):387–392. doi:10.1002/jor.24467
183. Wiggins AJ, Grandhi RK, Schneider DK, Stanfield D, Webster KE, Myer GD. Risk of secondary injury in younger athletes after anterior cruciate ligament reconstruction: a systematic review and meta-analysis. *Am J Sports Med*. 2016;44(7):1861–1876. doi:10.1177/0363546515621554
184. Grassi A, Ardern CL, Marcheggiani Muccioli GM, Neri MP, Marcacci M, Zaffagnini S. Does revision ACL reconstruction measure up to primary surgery? A meta-analysis comparing patient-reported and clinician-reported outcomes, and radiographic results. *Br J Sports Med*. 2016;50(12):716–724. doi:10.1136/bjsports-2015-094948
185. Burgi CR, Peters S, Ardern CL, et al. Which criteria are used to clear patients to return to sport after primary ACL reconstruction? a scoping review. *Br J Sports Med*. 2019;53(18):1154–1161. doi:10.1136/bjsports-2018-099982
186. Ashgibi EYK, Banzer W, Niederer D. Return to sport tests' prognostic value for reinjury risk after anterior cruciate ligament reconstruction: a systematic review. *Med Sci Sports Exerc*. 2020;52(6):1263–1271. doi:10.1249/mss.0000000000002246
187. Granan LP, Baste V, Engebretsen L, Inacio MC. Associations between inadequate knee function detected by KOOS and prospective graft failure in an anterior cruciate ligament-reconstructed knee. *Knee Surg Sports Traumatol Arthrosc*. 2015;23(4):1135–1140. doi:10.1007/s00167-014-2925-5
188. Grindem H, Engebretsen L, Axe M, Snyder-Mackler L, Risberg MA. Activity and functional readiness, not age, are the critical factors for second anterior cruciate ligament injury—the Delaware-Oslo ACL Cohort Study. *Br J Sports Med*. 2020;54(18):1099–1102. doi:10.1136/bjsports-2019-100623
189. Chu CR. Can we afford to ignore the biology of joint healing and graft incorporation after ACL reconstruction? *J Orthop Res*. 2022;40(1):55–64. doi:10.1002/jor.25145
190. Laboute E, Savalli L, Puig P, et al. Analysis of return to competition and repeat rupture for 298 anterior cruciate ligament reconstructions with patellar or hamstring tendon autograft in sportspeople. *Ann Phys Rehabil Med*. 2010;53(10):598–614. doi:10.1016/j.rehab.2010.10.002
191. Smith AH, Capin JJ, Zarzycki R, Snyder-Mackler L. Athletes with bone-patellar tendon-bone autograft for anterior cruciate ligament

- reconstruction were slower to meet rehabilitation milestones and return-to-sport criteria than athletes with hamstring tendon autograft or soft tissue allograft: secondary analysis from the ACL-SPORTS trial. *J Orthop Sports Phys Ther.* 2020;50(5):259–266. doi:10.2519/jospt.2020.9111
192. Balendra G, Jones M, Borque KA, Willinger L, Pinheiro VH, Williams A. Factors affecting return to play and graft re-rupture after primary ACL reconstruction in professional footballers. *Knee Surg Sports Traumatol Arthrosc.* 2022;30(7):2200–2208. doi:10.1007/s00167-021-06765-8
 193. Paterno MV, Huang B, Thomas S, Hewett TE, Schmitt LC. Clinical factors that predict a second ACL injury after ACL reconstruction and return to sport: preliminary development of a clinical decision algorithm. *Orthop J Sports Med.* 2017;5(12):2325967117745279. doi:10.1177/2325967117745279
 194. Paterno MV, Schmitt LC, Ford KR, et al. Biomechanical measures during landing and postural stability predict second anterior cruciate ligament injury after anterior cruciate ligament reconstruction and return to sport. *Am J Sports Med.* 2010;38(10):1968–1978. doi:10.1177/0363546510376053
 195. Padua DA, DiStefano LJ, Hewett TE, et al. National Athletic Trainers' Association position statement: prevention of anterior cruciate ligament injury. *J Athl Train.* 2018;53(1):5–19. doi:10.4085/1062-6050-99-16
 196. Arundale AJH, Bizzini M, Giordano A, et al. Exercise-based knee and anterior cruciate ligament injury prevention. *J Orthop Sports Phys Ther.* 2018;48(9):A1–A42. doi:10.2519/jospt.2018.0303
 197. Ardern CL, Ekås GR, Grindem H, et al. 2018 International Olympic Committee consensus statement on prevention, diagnosis and management of paediatric anterior cruciate ligament (ACL) injuries. *Br J Sports Med.* 2018;52(7):422–438. doi:10.1136/bjsports-2018-099060
 198. Trojjan T, Driban J, Nuti R, et al. Osteoarthritis action alliance consensus opinion—best practice features of anterior cruciate ligament and lower limb injury prevention programs. *World J Orthop.* 2017;8(9):726–734. doi:10.5312/wjo.v8.i9.726
 199. Gans I, Retzky JS, Jones LC, Tanaka MJ. Epidemiology of recurrent anterior cruciate ligament injuries in National Collegiate Athletic Association sports: the Injury Surveillance Program, 2004–2014. *Orthop J Sports Med.* 2018;6(6):2325967118777823. doi:10.1177/2325967118777823
 200. Sugimoto D, Myer GD, Barber Foss KD, Pepin MJ, Micheli LJ, Hewett TE. Critical components of neuromuscular training to reduce ACL injury risk in female athletes: meta-regression analysis. *Br J Sports Med.* 2016;50(20):1259–1266. doi:10.1136/bjsports-2015-095596
 201. Alentorn-Geli E, Myer GD, Silvers HJ, et al. Prevention of non-contact anterior cruciate ligament injuries in soccer players, II: a review of prevention programs aimed to modify risk factors and to reduce injury rates. *Knee Surg Sports Traumatol Arthrosc.* 2009;17(8):859–879. doi:10.1007/s00167-009-0823-z
 202. Herman K, Barton C, Malliaras P, Morrissey D. The effectiveness of neuromuscular warm-up strategies, that require no additional equipment, for preventing lower limb injuries during sports participation: a systematic review. *BMC Med.* 2012;10:75. doi:10.1186/1741-7015-10-75
 203. Michaelidis M, Koumantakis GA. Effects of knee injury primary prevention programs on anterior cruciate ligament injury rates in female athletes in different sports: a systematic review. *Phys Ther Sport.* 2014;15(3):200–210. doi:10.1016/j.ptsp.2013.12.002
 204. Pappas E, Nightingale EJ, Simic M, Ford KR, Hewett TE, Myer GD. Do exercises used in injury prevention programmes modify cutting task biomechanics? A systematic review with meta-analysis. *Br J Sports Med.* 2015;49(10):673–680. doi:10.1136/bjsports-2014-093796
 205. Rahlh AL, Zech A. Comparison of 10 vs. 20 min neuromuscular training for the prevention of lower extremity injuries in male youth football: a cluster randomised controlled trial. *J Sports Sci.* 2020;38(19):2177–2185. doi:10.1080/02640414.2020.1776459
 206. Huang YL, Jung J, Mulligan CMS, Oh J, Norcross MF. A majority of anterior cruciate ligament injuries can be prevented by injury prevention programs: a systematic review of randomized controlled trials and cluster-randomized controlled trials with meta-analysis. *Am J Sports Med.* 2020;48(6):1505–1515. doi:10.1177/0363546519870175
 207. Nagelli CV, Wordeman SC, Di Stasi S, Hoffman J, Marulli T, Hewett TE. Neuromuscular training improves biomechanical deficits at the knee in anterior cruciate ligament-reconstructed athletes. *Clin J Sport Med.* 2021;31(2):113–119. doi:10.1097/jsm.0000000000000723
 208. Nagelli C, Di Stasi S, Tatarski R, et al. Neuromuscular training improves self-reported function and single-leg landing hip biomechanics in athletes after anterior cruciate ligament reconstruction. *Orthop J Sports Med.* 2020;8(10):2325967120959347. doi:10.1177/2325967120959347
 209. Nagelli C, Wordeman S, Di Stasi S, Hoffman J, Marulli T, Hewett TE. Biomechanical deficits at the hip in athletes with ACL reconstruction are ameliorated with neuromuscular training. *Am J Sports Med.* 2018;46(11):2772–2779. doi:10.1177/0363546518787505
 210. Gilchrist J, Mandelbaum BR, Melancon H, et al. A randomized controlled trial to prevent noncontact anterior cruciate ligament injury in female collegiate soccer players. *Am J Sports Med.* 2008;36(8):1476–1483. doi:10.1177/0363546508318188
 211. Arundale AJH, Capin JJ, Zarzycki R, Smith AH, Snyder-Mackler L. Two year ACL reinjury rate of 2.5%: outcomes report of the men in a secondary ACL injury prevention program (ACL-SPORTS). *Int J Sports Phys Ther.* 2018;13(3):422–431.
 212. Johnson JL, Capin JJ, Arundale AJH, Zarzycki R, Smith AH, Snyder-Mackler L. A secondary injury prevention program may decrease contralateral anterior cruciate ligament injuries in female athletes: 2-year injury rates in the ACL-SPORTS randomized controlled trial. *J Orthop Sports Phys Ther.* 2020;50(9):523–530. doi:10.2519/jospt.2020.9407
 213. Kamath GV, Murphy T, Creighton RA, Viradia N, Taft TN, Spang JT. Anterior cruciate ligament injury, return to play, and reinjury in the elite collegiate athlete: analysis of an NCAA Division I cohort. *Am J Sports Med.* 2014;42(7):1638–1643. doi:10.1177/0363546514524164
 214. Gerber JP, Marcus RL, Dibble LE, Greis PE, Burks RT, LaStayo PC. Effects of early progressive eccentric exercise on muscle structure after anterior cruciate ligament reconstruction. *J Bone Joint Surg Am.* 2007;89(3):559–570. doi:10.2106/jbjs.F.00385
 215. Petushek EJ, Sugimoto D, Stoolmiller M, Smith G, Myer GD. Evidence-based best-practice guidelines for preventing anterior cruciate ligament injuries in young female athletes: a systematic review and meta-analysis. *Am J Sports Med.* 2019;47(7):1744–1753. doi:10.1177/0363546518782460
 216. Taylor JB, Waxman JP, Richter SJ, Shultz SJ. Evaluation of the effectiveness of anterior cruciate ligament injury prevention programme training components: a systematic review and meta-analysis. *Br J Sports Med.* 2015;49(2):79–87. doi:10.1136/bjsports-2013-092358
 217. Glazer DD. Development and preliminary validation of the Injury-Psychological Readiness to Return to Sport (I-PRRS) scale. *J Athl Train.* 2009;44(2):185–189. doi:10.4085/1062-6050-44.2.185
 218. Quinn AM, Fallon BJ. The changes in psychological characteristics and reactions of elite athletes from injury onset until full recovery. *J Appl Sport Psychol.* 1999;11(2):210–229. doi:10.1080/10413209908404201

219. Wiese DM, Weiss MR. Psychological rehabilitation and physical injury: implications for the sportsmedicine team. *Sport Psychol.* 1987;1(4):318–330. doi:10.1123/tsp.1.4.318
220. Podlog L, Eklund RC. Return to sport after serious injury: a retrospective examination of motivation and psychological outcomes. *J Sport Rehabil.* 2005;14(1):20–34. doi:10.1123/jsr.14.1.20
221. Podlog L, Banham SM, Wadey R, Hannon JC. Psychological readiness to return to competitive sport following injury: a qualitative study. *Sport Psychol.* 2015;29(1):1–14. doi:10.1123/tsp.2014-0063
222. Giusti J. *Operation Return to Duty: Developing and Validating an Instrument to Assess Soldiers' Mental Readiness to Resume Their Duties Following Injury Rehabilitation.* Dissertation. University of Idaho; 2018.
223. Radomski MV, Weightman MM, Davidson LF, et al. Development of a measure to inform return-to-duty decision making after mild traumatic brain injury. *Mil Med.* 2013;178(3):246–253. doi:10.7205/milmed-d-12-00144
224. Booth-Kewley S, Larson GE, Highfill-McRoy RM. Psychosocial predictors of return to duty among marine recruits with musculoskeletal injuries. *Mil Med.* 2009;174(2):139–152. doi:10.7205/milmed-d-01-2708
225. Gehring MB, Lerret S, Johnson J, et al. Patient expectations for recovery after elective surgery: a common-sense model approach. *J Behav Med.* 2020;43(2):185–197. doi:10.1007/s10865-019-00097-2
226. Mitchell ID, Neil R, Wadey R, Hanton S. Gender differences in athletes' social support during injury rehabilitation. *J Sport Exerc Psychol.* 2007;29(suppl):S189–S190.
227. Evans L, Mitchell I, Jones S. Psychological responses to sport injury: a review of current research. In: Hanton S, Mellalieu SD, eds. *Literature Reviews in Sport Psychology.* Nova Science Publishers; 2006:289–319.
228. Levy AR, Polman RC, Clough PJ. Adherence to sport injury rehabilitation programs: an integrated psycho-social approach. *Scand J Med Sci Sports.* 2008;18(6):798–809. doi:10.1111/j.1600-0838.2007.00704.x
229. Niven A. Rehabilitation adherence in sport injury: sport physiotherapists' perceptions. *J Sport Rehabil.* 2007;16(2):93–110. doi:10.1123/jsr.16.2.93
230. Pizzari T, McBurney H, Taylor NF, Feller JA. Adherence to anterior cruciate ligament rehabilitation: a qualitative analysis. *J Sport Rehabil.* 2002;11(2):90–102. doi:10.1123/jsr.11.2.90
231. Alexanders J, Chesterton P, Brooks A, Kaye JA. An exploration of UK student physiotherapists' goal setting practices within anterior cruciate ligament rehabilitation. *Musculoskeletal Care.* 2021;19(2):172–179. doi:10.1002/msc.1519
232. Gennarelli SM, Brown SM, Mulcahey MK. Psychosocial interventions help facilitate recovery following musculoskeletal sports injuries: a systematic review. *Phys Sportsmed.* 2020;48(4):370–377. doi:10.1080/00913847.2020.1744486
233. Vassallo AJ, Pappas E, Stamatakis E, Hiller CE. Injury fear, stigma, and reporting in professional dancers. *Saf Health Work.* 2019;10(3):260–264. doi:10.1016/j.shaw.2019.03.001
234. Pescosolido BA, Martin JK. The stigma complex. *Annu Rev Sociol.* 2015;41:87–116. doi:10.1146/annurev-soc-071312-145702
235. Thompson MM, McCreary DR. Enhancing mental readiness in military personnel. In: Adler AB, Castro CA, Britt TW, eds. *Military Life: The Psychology of Serving in Peace and Combat: Operational Stress.* Praeger Security International; 2006:54–79.
236. Christino MA, Fantry AJ, Vopat BG. Psychological aspects of recovery following anterior cruciate ligament reconstruction. *J Am Acad Orthop Surg.* 2015;23(8):501–509. doi:10.5435/jaaos-d-14-00173
237. Walker N, Thatcher J, Lavalley D. Psychological responses to injury in competitive sport: a critical review. *J R Soc Promot Health.* 2007;127(4):174–180. doi:10.1177/1466424007079494
238. Dobkin BH, Dorsch A. The promise of mHealth: daily activity monitoring and outcome assessments by wearable sensors. *Neurorehabil Neural Repair.* 2011;25(9):788–798. doi:10.1177/1545968311425908
239. Gal R, May AM, van Overmeeren EJ, Simons M, Monninkhof EM. The effect of physical activity interventions comprising wearables and smartphone applications on physical activity: a systematic review and meta-analysis. *Sports Med Open.* 2018;4(1):42. doi:10.1186/s40798-018-0157-9
240. Brickwood KJ, Watson G, O'Brien J, Williams AD. Consumer-based wearable activity trackers increase physical activity participation: systematic review and meta-analysis. *JMIR Mhealth Uhealth.* 2019;7(4):e11819. doi:10.2196/11819
241. Wyatt B, Mingo CA, Waterman MB, White P, Cleveland RJ, Callahan LF. Impact of the Arthritis Foundation's Walk with Ease program on arthritis symptoms in African Americans. *Prev Chronic Dis.* 2014;11:E199. doi:10.5888/pcd11.140147
242. Conte KP, Odden MC, Linton NM, Harvey SM. Effectiveness of a scaled-up arthritis self-management program in Oregon: Walk With Ease. *Am J Public Health.* 2016;106(12):2227–2230. doi:10.2105/AJPH.2016.303478
243. Silverstein RP, VanderVos M, Welch H, Long A, Kaboré CD, Hootman JM. Self-directed Walk With Ease workplace wellness program—Montana, 2015–2017. *MMWR Morb Mortal Wkly Rep.* 2018;67(46):1295–1299. doi:10.15585/mmwr.mm6746a3
244. Wang JB, Cataldo JK, Ayala GX, et al. Mobile and wearable device features that matter in promoting physical activity. *J Mob Technol Med.* 2016;5(2):2–11. doi:10.7309/jmtm.5.2.2
245. Omura JD, Carlson SA, Paul P, Watson KB, Fulton JE. National physical activity surveillance: users of wearable activity monitors as a potential data source. *Prev Med Rep.* 2017;5:124–126. doi:10.1016/j.pmedr.2016.10.014
246. Islam MM, Poly TN, Walther BA, Jack Li YC. Use of mobile phone app interventions to promote weight loss: meta-analysis. *JMIR Mhealth Uhealth.* 2020;8(7):e17039. doi:10.2196/17039
247. Mrazek AJ, Mrazek MD, Cherolini CM, et al. The future of mindfulness training is digital, and the future is now. *Curr Opin Psychol.* 2019;28:81–86. doi:10.1016/j.copsyc.2018.11.012
248. Hertwig R, Grune-Yanoff T. Nudging and boosting: steering or empowering good decisions. *Perspect Psychol Sci.* 2017;12(6):973–986. doi:10.1177/1745691617702496
249. Busch J, Madsen EK, Fage-Butler AM, Kjær M, Ledderer L. Dilemmas of nudging in public health: an ethical analysis of a Danish pamphlet. *Health Promot Int.* 2020;36(4):1140–1150. doi:10.1093/heapro/daaa146
250. Eyles JP, Lucas BR, Patterson JA, et al. Does clinical presentation predict response to a nonsurgical chronic disease management program for endstage hip and knee osteoarthritis? 2014;41(11):2223–2231. doi:10.3899/jrheum.131475
251. Musculoskeletal network: osteoarthritis chronic care program model of care. ACI Musculoskeletal Network. Accessed October 21, 2022. https://www.aci.health.nsw.gov.au/_data/assets/pdf_file/0003/165306/Osteoarthritis-Chronic-Care-Program-Mode-of-Care-High-Resolution.pdf
252. Allen KD, Choong PF, Davis AM, et al. Osteoarthritis: models for appropriate care across the disease continuum. *Best Pract Res Clin Rheumatol.* 2016;30(3):503–535. doi:10.1016/j.berh.2016.09.003
253. Hunter DJ, Hinman RS, Bowden JL, et al; PARTNER Study Team. Effectiveness of a new model of primary care management on knee pain and function in patients with knee osteoarthritis:

protocol for the PARTNER study. *BMC Musculoskelet Disord.* 2018;19(1):132. doi:10.1186/s12891-018-2048-0

254. Teoh LSG, Eyles JP, Makovey J, Williams M, Kwok CK, Hunter DJ. Observational study of the impact of an individ-

ualized multidisciplinary chronic care program for hip and knee osteoarthritis treatment on willingness for surgery. *Int J Rheum Dis.* 2017;20(10):1383–1392. doi:10.1111/1756-185X.12950

Address correspondence to Jeffrey B. Driban, PhD, ATC, CSCS, Division of Rheumatology, Allergy, and Immunology, Tufts Medical Center, 800 Washington Street, Box 406, Boston, MA 02111. Address email to jeffrey.driban@tufts.edu.