

Comparing the Primary Concerns of Injured Collegiate Athletes With the Content of Patient-Reported Outcome Measures

Jennifer Tinsley Kubala, MS, LAT, ATC*; Heather L. Pannill, DPT, LAT, ATC†; Kimberly S. Fasczewski, PhD‡; Laurie A. Rivera, MEd, LAT, ATC§; Erin D. Bouldin, MPH, PhD||; Jennifer S. Howard, PhD, LAT, ATC§

*Department of Athletic Training and Clinical Nutrition, University of Kentucky, Lexington; †Rehabilitation Services, Appalachian Regional Healthcare System, Boone, NC; ‡Department of Public Health and Exercise Science and §Department of Rehabilitation Science, Appalachian State University, Boone, NC; || Department of Internal Medicine, University of Utah, Salt Lake City

Context: Patient-reported outcome measures (PROMs) have been endorsed for providing patient-centered care. However, PROMs must represent their target populations.

Objective: To identify the primary concerns of collegiate athletes experiencing injury and compare those with the content of established PROMs.

Design: Cross-sectional study.

Setting: Collegiate athletic training facilities.

Patients or Other Participants: Collegiate athletes experiencing injury (N = 149).

Main Outcome Measure(s): Open-ended responses to the Measure Yourself Medical Outcome Profile were used to identify primary concerns, which were linked to International Classification of Functioning, Disability and Health taxonomy codes. Items of the Patient-Reported Outcomes Measurement Information System; modified Disablement of the Physically Active Scale; Lower Extremity Functional Scale; Knee injury and Osteoarthritis Outcome Score (KOOS); International Knee Documentation Committee Subjective Knee Form (IKDC); Foot and Ankle Ability Measure; Disabilities of the Arm, Shoulder, and Hand; Functional Arm Scale for Throwers; and Kerlan-Jobe Orthopaedic Clinic questionnaire were linked to International Classification of Functioning, Disability and Health codes. We calculated χ^2 single-sample goodness-of-fit tests to determine if 70% of the content was shared between PROMs and participant-generated codes.

Results: Participant-generated concerns were primarily related to sport participation (16%) and pain (23%). Chi-square tests showed that the Lower Extremity Functional Scale and Foot and Ankle Ability Measure presented significant content differences, with common participant-generated lower extremity responses at all levels. The Patient-Reported Outcomes Measurement Information System; modified Disablement of the Physically Active Scale; KOOS; IKDC; Disabilities of the Arm, Shoulder and Hand; Functional Arm Scale for Throwers; and Kerlan-Jobe Orthopaedic Clinic questionnaire did not have significant content differences for level 2 codes; still, significant differences were present for level 3 analyses except for the KOOS and IKDC ($P < .001$). All measures except the IKDC contained significant superfluous content ($P < .05$).

Conclusions: The presence of significant content differences supports clinician-perceived barriers regarding the relevance of established PROMs. However, the IKDC was a relevant and efficient PROM for evaluating the primary concerns of collegiate athletes experiencing lower extremity injury. Clinicians should consider using patient-generated measures to support coverage of patient-specific concerns in care.

Key Words: patient-centered care, disablement, health-related quality of life

Key Points

- Pain and sport participation were the primary concerns of participants, with limited variation based on injury location or classification.
- A large amount of extraneous and unrelated content was present among commonly used and established patient-reported outcome measures.
- Only the International Knee Documentation Committee Subjective Knee Form maintained coverage of participant-generated concerns when analyzed at both International Classification of Functioning, Disability and Health levels 2 and 3 and did not demonstrate a significant amount of superfluous content.

Patient-reported outcome measures (PROMs) are defined as instruments patients complete that provide information about the effect of their health condition or injury on their health status or health-related quality of life (HRQOL) and that highlight the patient

perspective.^{1–3} Their use is intended to give the clinician a greater understanding of the patient's physical and psychosocial response to treatment.^{1,3} This improved clarity is thought to result in a more educated and engaged patient, a more individualized treatment plan, and an improved

patient-clinician relationship and therefore an improved treatment outcome.¹⁻⁵ Despite these constructs supporting PROMs, traditional clinical measures (eg, strength measures) and clinician experience remain primary drivers of clinical decision-making.²⁻⁴

Dependence upon disease- and clinician-oriented measures may be linked to perceptions by athletic trainers (ATs) that many PROMs are not relevant to their patient population.^{2,5,6} This may be the reality, as many PROMs reportedly being used most commonly by ATs (eg, Lower Extremity Functional Scale [LEFS] and Disabilities of the Arm, Shoulder and Hand scale [DASH])⁶⁻⁸ were designed with lower-demand populations and may not be ideal in a high-demand athletic population. Additionally, the time to complete and analyze PROMs has also been documented as a barrier to PROM implementation.^{2,5,6} To optimize the utility of PROMs in traditional athletic training settings, PROMs must be both relevant, by addressing the primary concerns of athletes experiencing injury, and efficient, by having limited content that is not related to those primary concerns.

The International Classification of Functioning, Disability and Health (ICF) framework, developed by the World Health Organization, is intended to provide a standard language to serve as a reference for describing and comparing health states.⁹⁻¹¹ Rather than focusing on a diagnosis or condition, the aim of the ICF is to emphasize experience and changes in functional ability, encouraging recognition of the patient's perception and individualized response to the treatment, condition, or both.^{9,11-13} Common PROMs have been analyzed to identify which ICF domains (health condition, body structure and function, activity, participation, environmental factors, and personal factors)¹⁴ they capture; however, neither PROM content nor self-identified patient concerns have been analyzed and compared using the full ICF coding taxonomy. Therefore, the ICF provides an ideal framework by which to classify patient primary concerns and the content covered in commonly used PROMs.^{10,11,13,15}

The primary aims of our study were to use the ICF framework to identify the primary concerns of collegiate athletes experiencing an injury, determine if these primary concerns varied based on phase of injury or injury region, and establish if these primary concerns were sufficiently and efficiently represented in the following established PROMs: the Patient-Reported Outcomes Measurement Information System Global Health Scale version 1.2 (PROMIS), modified Disablement of the Physically Active Scale (mDPAS), LEFS, Knee injury and Osteoarthritis Outcome Score (KOOS), International Knee Documentation Committee Subjective Knee Form (IKDC), Foot and Ankle Ability Measure (FAAM; including both the Activities of Daily Living and Sports subscales), DASH, Functional Arm Scale for Throwers (FAST), and Kerlan-Jobe Orthopaedic Clinic questionnaire (KJOC). We hypothesized that (1) differences in primary concerns would exist among participants in an acute, subacute, or chronic phase of injury; (2) the PROMIS and those region-specific PROMs reported as being most commonly used by ATs for lower extremity injuries (LEFS) or upper extremity injuries (DASH)^{6-8,16} would fail to contain relevant content and efficiently address the primary concerns of collegiate athletes experiencing injury; and (3) those PROMs

designed for use with highly active populations (mDPAS, KOOS, IKDC, FAST, and KJOC)¹⁷⁻²² would contain relevant content and efficiently address the most commonly occurring participant-generated ICF codes.

METHODS

This observational cross-sectional study used the Measure Yourself Medical Outcome Profile (MYMOP-2)²³ to identify the items most relevant to injured collegiate student-athletes and compare those symptoms and activities with those represented on established PROMs. This study was approved by the Appalachian State University Institutional Review Board.

Participants

Participants consisted of a convenience sample of student-athletes experiencing an injury at 4 collegiate institutions in 2 states in the mid-Atlantic region of the United States, including National Collegiate Athletic Association Division I (Football Bowl Subdivision), Division II (2 schools), and Division III schools. Any student-athletes over the age of 18 receiving care for any injury, fluent in English, who were identified by their treating ATs as having modified or restricted their sport participation in games or practices because of injury and were willing to sign a Health Insurance Portability and Accountability Act (HIPAA) release for their ATs to share information regarding their injuries were eligible for the study. Participants were not enrolled more than once for a given body region.

Procedures

Participants were asked to complete the MYMOP-2 and a demographics form requesting their treating AT, age, year in school, sport, date of injury, and location and description of injury. Data collection occurred 1 time per participant per region of injury. At the time of enrollment, the treating ATs were asked for their assessment of the injury phase of each participant (acute, subacute, or chronic).

Linking to the ICF

Each participant's responses to the MYMOP-2 and each item on the included established PROMs were linked to an ICF code via the established ICF linking rules described by Cieza et al.¹¹ In brief, 3 raters reviewed all MYMOP-2 responses independently and assigned a code in the deepest relevant level of the ICF taxonomy. The raters were 2 licensed ATs and an individual licensed as both an AT and a physical therapist. Before rating responses, reviewers underwent additional training in the ICF model via the ICF e-learning tool; reviewed recommendations from the literature outlining and examining the ICF model, the World Health Organization ICF manuals, and previously conducted studies with similar methods; and completed pilot linking sessions and discussed and compared results.^{9-11,13,15} The ICF model is a flexible, multitiered, hierarchical framework split into 2 parts. Part 1 contains the domains of body functions (b), body structures (s), and activity and participation (d). Part 2 contains contextual factors called environmental factors (e) and personal factors

(pf). Each contains up to 4 levels. The first level is referred to as the chapter, and each chapter contains up to 3 nested levels (second, third, and fourth) increasing in specificity.^{9,10} This hierarchical structure was used to link each patient concern to an ICF code, guided by the ICF browser definitions, inclusions, and exclusions available at the second and third levels.²⁴ For example, if a response was “pitching a baseball,” it would be coded as “d4454 Throwing.” The definition for this code states: “Using fingers, hands and arms to lift something and propel it with some force through the air, such as when tossing a ball.”²⁴ The process to arrive at this code is as follows:

- d Activities and participation (domain)
- d4 Mobility (chapter)
- d430–d449 Lifting and carrying objects (component)
- d445 Hand and arm use (level 2)
- d4454 Throwing (level 3)

Raters were permitted to assign multiple ICF codes to a single MYMOP-2 response in order to fully represent the participant’s concern. For example, if a participant listed “pain when I run,” separate codes were assigned to represent pain (b280) and running (d4552). Similarly, multiple codes could be assigned as needed for PROM items. Once the individual rating was complete, the raters met to review the identified codes. When disagreement existed among raters, they discussed the code(s) until a consensus was reached. Fleiss κ statistics indicated moderate to very good (0.433 to 0.846) interrater agreement for coding of the MYMOP-2 responses at both levels 2 and 3 before the consensus process that was used to arrive at final codes for each participant response and each PROM item.²⁵ The linking process resulted in a final single set of agreed-upon codes for each participant response and PROM item. Throughout the review process, previously agreed-upon codes were reviewed to ensure consistency.

Instrumentation

The MYMOP-2 is a patient-generated outcome measure. It asks the patient to “choose one or two symptoms (physical or mental) that bother you the most” and “choose one activity (physical, social, or mental) that is important to you, and that your problem makes difficult or prevents you from doing.”²³ The MYMOP-2 has been examined in various acute and chronic settings and has been reported to detect changes in several populations, including the military^{23,26,27}; therefore, its application to an athletic population is not unreasonable. As an open-ended tool, it allows for a patient-centered evaluation of symptoms across multiple phases of the injury, disease, or condition and across multiple pathologies.^{23,26,27} Although not completed by participants, all possible items of the investigated PROMs were linked to ICF codes. These instruments were selected because the LEFS and DASH have been identified as the most common upper and lower extremity-specific PROMs used by practicing ATs⁶ and the PROMIS has been widely proposed as a generic HRQOL instrument that can be applied across populations.¹⁶ The mDPAS, KOOS, IKDC, FAAM (including Sports subscale), FAST, and KJOC were included because of their use in athletic populations. The mDPAS is a general HRQOL PROM designed to evaluate both mental and physical components

of disability in physically active populations.¹⁸ The KOOS is a knee-specific PROM that assesses pain, symptoms, activities of daily living, sport and recreation function, and knee-related quality of life.¹⁹ The IKDC is a region-specific measure of symptoms, function, and sports activity for those experiencing knee-related problems.²¹ The FAAM is a region-specific PROM assessing concerns related to activities of daily living and sport-specific concerns for those experiencing injuries of the lower leg, ankle, and foot.²² The FAST is an upper extremity, region-specific PROM targeting overhead-throwing athletes experiencing multiple domains of disablement.²⁰ Finally, the KJOC is a region-specific PROM that evaluates function and performance for overhead athletes experiencing upper extremity injuries.¹⁷

Statistical Analysis

To reduce the data to those codes representing the items most important to the majority of participants, we performed frequency counts to identify those codes cumulatively representing 80% of patient-generated ICF codes overall, by phase of injury, and for the upper and lower extremities. Those codes in the cumulative 80% were considered the primary concerns and were used for our analyses. Primary concerns were examined by phase of injury and region for substantial content differences. To explore various levels of specificity, codes were examined at both levels 2 and 3 of the ICF taxonomy. To address the relevance of the established PROMs, we determined that a minimum of 70% of the primary participant-generated codes must be encompassed in a PROM if it was to be considered representative of patient concerns (ie, 70% of the cumulative 80% of generated codes were addressed by the established PROM). The a priori 70% threshold was based on established logic that indicates if a special test is considered clinically useful.²⁸ Additionally, an approximately 70% threshold has been applied in recent Delphi analyses examining both return-to-sport decisions (70%)²⁹ and PROM content validity among both researchers and practicing clinicians (67%).³⁰ Collectively, this past use of a 70% threshold supports its application as an established and acceptable standard for clinical decision-making and PROM-related research. To evaluate the efficiency of commonly used PROMs and determine the amount of extraneous content present, we examined if 70% of the established PROM codes were among the primary participant-generated codes. Chi-square, 1-sample goodness-of-fit tests were performed to test the a priori selected 70% thresholds. If less than 70% ($P < .05$) of the relevant most common participant-generated codes were represented in the established PROMs, they were considered nonrepresentative of the participant-generated primary concerns. If less than 70% of the established PROM-generated codes were encompassed by the most common participant-generated codes, the PROM was considered to have a significant amount of extraneous content.

RESULTS

Participants were 149 collegiate athletes (74 women, 75 men, age = 19.6 \pm 1.3 years) and represented 150 injuries. Lower extremity injuries accounted for 77% of our sample and upper extremity injuries, 15%. Back, head, and neck

Table 1. Sport Distribution

| Sport | % (No.) | Cumulative % |
|-----------------|------------|--------------|
| Soccer | 30.67 (46) | 30.67 |
| Football | 16 (24) | 46.67 |
| Lacrosse | 11.33 (17) | 58.00 |
| Track and field | 8.67 (13) | 66.67 |
| Volleyball | 6.67 (10) | 73.34 |
| Basketball | 6 (9) | 79.34 |
| Softball | 5.33 (8) | 84.67 |
| Baseball | 3.33 (5) | 88.00 |
| Cheerleading | 3.33 (5) | 91.33 |
| Cross-country | 2 (3) | 93.33 |
| Cycling | 1.33 (2) | 94.66 |
| Field hockey | 1.33 (2) | 95.99 |
| Swimming | 1.33 (2) | 97.33 |
| Tennis | 1.33 (2) | 98.66 |
| Wrestling | 1.33 (2) | 100 |
| Total | 100 (150) | 100 |

injuries constituted 6%, with 2% missing the region of injury. Most participants were classified by their treating ATs as being in the acute phase of injury at the time of collection (50%), followed by chronic (35%), subacute (7%), and missing (7%). Participant sport distributions are shown in Table 1. Forty participants chose to list only a single symptom on the MYMOP-2 in response to the question prompting the listing of 1 or 2 symptoms. The final set of agreed-upon ICF codes for participant responses yielded 594 total codes, containing 90 unique codes. The majority were represented in the body function (56%) and activities and participation (41%) domains. The body structure domain represented 1%, and 2% were not codable. Code frequency distributions for the commonly occurring codes overall are displayed in Tables 2 and 3. Code frequency distributions by extremity and phase of injury overlapped substantially (see Supplemental Tables 1–6, available online at <http://dx.doi.org/10.4085/1062-6050-0516.21.S1>). Frequency counts of participant-generated ICF codes overall, by extremity, and by phase of injury are provided in Table 4. A summary of shared codes between the primary concerns of participants and established PROMs can be seen in Table 5.

Global PROMs

Comparisons between all participant-generated common codes and the PROMIS were mixed, with PROMIS content

relevant at level 2 ($P = .061$) but falling significantly below the 70% threshold for relevance at level 3 ($P < .001$). In contrast, the PROMIS efficiently evaluated primary concerns at level 3 ($P = .081$) but not at level 2 ($P = .011$). When all participant-generated codes were compared with codes generated from the mDPAS, level 2 codes appeared relevant to the primary concerns of participants ($P = .506$). However, this relevance was lost for level 3 codes ($P = .004$). Additionally, based on codes generated from the mDPAS, it was not efficient in evaluating the primary concerns of injured student-athletes at either level ($P < .001$, $P < .001$).

Lower Extremity PROMs

Among participants with lower extremity injuries, analyses of level 2 and level 3 codes revealed that the LEFS was not relevant to the primary concerns of participants ($P < .001$). Similarly, the LEFS was not efficient in evaluating participant concerns at either level 2 or level 3 ($P < .001$). Comparison of participant-generated concerns for lower extremity injuries with the IKDC indicated relevance for both level 2 and level 3 codes ($P = .801$, $P = .143$). Additionally, the IKDC was efficient in evaluating primary concerns at both levels 2 ($P = .159$) and 3 ($P = .078$). Comparison of codes for those experiencing lower extremity injuries with the KOOS demonstrated relevancy at both level 2 ($P = .705$) and 3 ($P = .143$) analyses. However, the KOOS was not efficient at either level ($P = .003$, $P < .001$). Comparison of participant-generated codes with the FAAM displayed a lack of relevance at both level 2 ($P = .001$) and level 3 ($P = .001$) analyses. Similarly, the FAAM was not efficient at either level ($P = .001$, $P < .001$).

Upper Extremity PROMs

Among those with upper extremity injuries, level 2 ICF code analysis indicated that 7 of the 8 primary codes (88%) were represented in the DASH, reflecting PROM relevance ($P = .28$). However, this relevance was not maintained at level 3 ($P < .001$). Furthermore, examination of the ICF codes generated from the DASH revealed that it did not efficiently evaluate the primary concerns of participants at either level 2 or level 3 ($P < .001$). The FAST was relevant in capturing participant concerns at level 2 ($P = .758$) but not at level 3 ($P = .006$). Additionally, the FAST was not

Table 2. Most Common Participant-Generated International Classification of Functioning, Disability and Health Codes Overall (Level 2)

| Code | Description | % (No.) | Cumulative % |
|------|---|-------------|--------------|
| b280 | Sensation of pain | 22.73 (135) | 22.73 |
| d920 | Recreation and leisure | 15.63 (93) | 38.38 |
| d455 | Moving around | 10.23 (61) | 48.65 |
| b780 | Sensations related to muscles and movement functions | 7.57 (45) | 56.23 |
| b152 | Emotional functions | 5.38 (32) | 61.62 |
| d450 | Walking | 4.04 (24) | 65.66 |
| b289 | Sensation of pain, other specified and unspecified | 3.36 (20) | 69.02 |
| b798 | Movement functions, other specified and unspecified | 2.52 (15) | 71.55 |
| b710 | Mobility of joint functions | 2.02 (12) | 73.57 |
| b439 | Functions of the hematological and immunological systems, other specified and unspecified | 1.85 (11) | 75.42 |
| d445 | Hand and arm use | 1.68 (10) | 77.10 |
| ns | Not codable | 1.68 (10) | 78.79 |
| b199 | Mental functions, unspecified | 1.51 (9) | 80.30 |

Table 3. Most Common Participant-Generated International Classification of Functioning, Disability and Health Codes Overall (Level 3)

| Code | Description | % (No.) | Cumulative % |
|-------|---|------------|--------------|
| b2801 | Pain in body part | 15.66 (93) | 15.66 |
| d9201 | Sport | 13.81 (82) | 29.46 |
| b280 | Sensation of pain | 6.91 (41) | 36.36 |
| b7808 | Sensations related to muscles and movement functions, unspecified | 6.23 (37) | 42.59 |
| d4552 | Running | 5.90 (35) | 48.48 |
| b1528 | Emotional functions, other specified | 3.87 (23) | 52.36 |
| d4509 | Walking, unspecified | 3.53 (21) | 55.89 |
| b289 | Sensation of pain, other specified and unspecified | 3.36 (20) | 59.26 |
| b798 | Neuromusculoskeletal and movement-related functions, other specified | 2.52 (15) | 61.78 |
| b439 | Functions of the hematological and immunological systems, other specified and unspecified | 1.85 (11) | 63.64 |
| d4558 | Moving around, other specified | 1.68 (10) | 65.32 |
| Ns | Not codable | 1.68 (10) | 67.00 |
| b1522 | Range of emotion | 1.51 (9) | 68.52 |
| b199 | Mental functions, unspecified | 1.51 (9) | 70.03 |
| d4553 | Jumping | 1.51 (9) | 71.55 |
| b7800 | Sensation of muscle stiffness | 1.34 (8) | 72.90 |
| d4300 | Lifting | 1.34 (8) | 74.24 |
| b298 | Sensory functions and pain, other specified | 1.17 (7) | 75.42 |
| b7100 | Mobility of a single joint | 1.17 (7) | 76.60 |
| b799 | Neuromusculoskeletal and movement-related functions, unspecified | 1.01 (6) | 77.61 |
| b1349 | Sleep functions, unspecified | 0.84 (5) | 78.45 |
| b7109 | Mobility of joint functions, unspecified | 0.84 (5) | 79.30 |
| d4551 | Climbing | 0.84 (5) | 80.13 |

efficient at either level 2 ($P = .027$) or level 3 ($P < .001$). Comparison of participant codes for upper extremity injuries with KJOC-generated codes at level 2 supported KJOC relevance and efficiency ($P = .758$, $P = .076$). However, analysis of the more specific level 3 ICF codes did not support continuation of this relevance or efficiency ($P = .006$, $P < .001$).

DISCUSSION

The objective of our study was to identify the primary concerns held by injured collegiate athletes and compare those concerns, as measured by the MYMOP-2, with the content of established PROMs. Sensations of pain and sport participation were the most commonly reported participant concerns, representing 39% of the total. Codes related to running or moving around, emotional function (stress, confidence, frustration, anxiety, etc), mobility (range of motion), swelling, and strength or muscle power were also consistently among the primary participant concerns overall. It is important to note that regardless of the level of analysis (level 2 or 3 of the ICF taxonomy) or region of injury, “not codable” was consistently present as well. These primary areas of concerns were in keeping with previous works linking participant concerns to ICF codes.¹⁵ Pain, sport, and movement function were among the most

common concerns reported across all phases of injury. Notably fewer participants were in the subacute and chronic phases; however, their primary concerns were consistent with those of participants in the acute phase, regardless of the region of injury. These results may suggest that use of the same PROM may be appropriate throughout the rehabilitation process. However, it is critical that selected PROMs reflect the patient’s concerns and be sensitive enough to detect change at various stages of rehabilitation.

Global PROMs

Both the PROMIS and the mDPAS are global PROMs intended to evaluate HRQOL; nonetheless, the mDPAS was specifically developed for use with physically active populations. For both global measures, the percentage of agreement between PROM content and primary concerns at ICF level 2 was not statistically different from 70%. Yet the mDPAS addressed more level 2 participant concerns (62%) than the PROMIS (46%). This greater representation in the mDPAS likely occurred because it was specifically developed for active populations. Significant content differences were present at level 3 for both measures. Uniquely for the PROMIS, we observed significant superfluous content at level 2 but not at level 3. These results indicate that both measures have potential utility in

Table 4. Number of International Classification of Functioning, Disability and Health Codes Represented or Not Represented Among the Most Common Participant-Generated Responses

| Level of Analysis | All | Extremity | | Phase of Injury | | |
|-------------------------------|-----|-----------|-------|-----------------|----------|---------|
| | | Upper | Lower | Acute | Subacute | Chronic |
| Level 2 unique codes | 58 | 20 | 51 | 48 | 14 | 36 |
| Level 3 unique codes | 90 | 26 | 73 | 75 | 20 | 54 |
| Top 80%, level 2 | 13 | 8 | 12 | 6 | 1 | 3 |
| Top 80%, level 3 | 23 | 12 | 20 | 10 | 1 | 4 |
| Codes not in top 80%, level 2 | 45 | 12 | 39 | 42 | 13 | 33 |
| Codes not in top 80%, level 3 | 67 | 14 | 53 | 65 | 19 | 50 |

Table 5. Summary of International Classification of Functioning, Disability and Health (ICF) Codes Shared by Most Common Participant Concerns and Established Patient-Reported Outcome Measures (PROMs)

| PROM | % of Most Common Concerns Addressed (Codes Addressed/ Total Common Concerns) | <i>P</i> Value of Test for 70% of Common Concerns | % of PROM Content Representing Most Common Concerns (Common Codes/ Total Codes Represented) | <i>P</i> Value of Test for 70% of Content |
|--|--|---|---|---|
| Patient-Reported Outcome Measures Information System Global Health Scale | | | | |
| ICF level 2 | 46 (6/13) ^a | .061 | 40 (6/15) | .011 |
| ICF level 3 | 35 (8/23) | <.001 | 50 (8/16) ^a | .081 |
| Modified Disablement of the Physically Active Scale | | | | |
| ICF level 2 | 62 (8/13) ^a | .506 | 33 (8/23) | <.001 |
| ICF level 3 | 43 (10/23) | .004 | 23 (10/43) | <.001 |
| Lower Extremity Functional Scale | | | | |
| ICF level 2 | 25 (3/12) | <.001 | 23 (3/13) | <.001 |
| ICF level 3 | 25 (5/20) | <.001 | 21 (5/24) | <.001 |
| Knee Injury and Osteoarthritis Outcome Score | | | | |
| ICF level 2 | 75 (9/12) ^a | .705 | 41 (9/22) | .003 |
| ICF level 3 | 55 (11/20) ^a | .143 | 28 (11/39) | <.001 |
| International Knee Documentation Committee Subjective Knee Form | | | | |
| ICF level 2 | 66 (8/12) ^a | .801 | 53 (8/15) ^a | .159 |
| ICF level 3 | 55 (11/20) ^a | .143 | 52 (11/21) ^a | .078 |
| Foot and Ankle Ability Measure | | | | |
| ICF level 2 | 25 (3/12) | .001 | 21 (3/14) | <.001 |
| ICF level 3 | 35 (7/20) | .001 | 29 (7/24) | <.001 |
| Disabilities of the Arm, Shoulder and Hand | | | | |
| ICF level 2 | 88 (7/8) ^a | .28 | 27 (7/26) | <.001 |
| ICF level 3 | 42 (5/12) | <.001 | 15 (5/33) | <.001 |
| Functional Arm Scale for Throwers | | | | |
| ICF level 2 | 75 (6/8) ^a | .758 | 43 (6/14) | .027 |
| ICF level 3 | 42 (5/12) | .006 | 29 (5/17) | <.001 |
| Kerlan-Jobe Orthopaedic Clinic questionnaire | | | | |
| ICF level 2 | 63 (5/8) ^a | .758 | 45 (5/11) ^a | .076 |
| ICF level 3 | 33 (4/12) | .006 | 27 (4/15) | <.001 |

^a Shared content not significantly different from 70% ($P > .05$).

assessing more general level 2 concerns (eg, muscle power functions, recreation and leisure, hand and arm use)²⁴ but not more specific and descriptive level 3 concerns (eg, sports, throwing, catching, running, jumping).²⁴ However, the PROMIS was the more efficient instrument, with 40% to 50% of its content aligning with the primary concerns of patients. These findings are counter to our hypothesis that PROMs designed for highly active individuals would be more relevant and efficient than those designed for use with the general population. A previous analysis¹⁴ of the mDPAS demonstrated a strong focus on the physiological domains of health and the activity domain of the ICF, with less representation in the psychological or participation areas. Conversely, our participants' concerns aligned more with their lived experiences, as exemplified by consistent reports of emotional functions and sport participation concerns. These differences in primary focus may explain the lack of alignment in the specific level 3 analysis.

Lower Extremity PROMs

The LEFS and FAAM content failed to represent the primary concerns of participants as measured by the MYMOP-2. Perhaps most notable was that pain, the most commonly occurring participant concern, was not represented in the LEFS or FAAM content. Our results also indicated that the LEFS and FAAM contain a large amount of extraneous content. It should be noted that we evaluated the full FAAM and the sport-specific subscale, so the findings could differ if only the sport-specific subscale of

the FAAM was examined. Still, these results are consistent with earlier concerns regarding the FAAM's ability to address concerns of an injured athletic population.³¹ Lam et al⁶ previously classified both instruments as being heavily focused on physical health and almost exclusively on the activity domain of the ICF. Given the large number of our patient-generated responses that were related to pain, sport performance, and emotional function, it is not surprising that the LEFS and FAAM appear to have limited relevance to the primary concerns of injured collegiate athletes.

The KOOS and IKDC demonstrated the strongest agreement between content and the primary concerns of collegiate athletes experiencing lower extremity injury as assessed by the MYMOP-2. Both PROMs adequately represented the concerns of participants at both ICF level 2 and level 3. The results endorse the hypothesis that these instruments, designed to address a wide range of activity levels, accurately reflect both broad and specific concerns of participants. Furthermore, the percentage of ICF codes generated from the IKDC at level 2 (53%, 8 of 15) and 3 (52%, 11 of 21) was not different than the 70% criterion value, demonstrating that the IKDC was the most efficient of the investigated PROMs in evaluating the concerns of participants experiencing lower extremity injuries. Conversely, the KOOS provided significant extraneous content at both levels 2 and 3, suggesting that the IKDC is preferable for use with highly active patients with lower extremity conditions. The relevance and efficiency of the IKDC may be products of its development. The IKDC was specifically designed to represent knee impairment-related

symptoms and limitations in function and sports activity.²¹ Moreover, the IKDC was iteratively tested with large samples of patients representing the target population and revised through a formal item-reduction process.²¹ The use of patient engagement in PROM development has been proposed to improve relevance to the patient.³² Creation of the KOOS similarly engaged patients, but the developers purposefully included a subgroup with osteoarthritis.¹⁹ It is possible that items relevant to patients with osteoarthritis may not align with the primary concerns of the college-age participants in our study, resulting in the lack of efficiency for the KOOS. This observation highlights a challenge of PROM selection and development. An instrument that is broadly generalizable may lack specificity, efficiency, or both for a target population. Although generalizability can be beneficial, a lack of perceived relevance by both clinicians and patients can be a barrier to PROM adoption.^{2,5,6} Overall, the IKDC was the most relevant and efficient PROM in this analysis for addressing the primary concerns of student-athletes experiencing injury.

Upper Extremity PROMs

Contrary to our hypothesis, for the upper extremity, relevance to patient concerns was similar among the more general DASH and the more activity-focused FAST and KJOC. In support of our hypothesis, a trend toward greater efficiency with those PROMs designed for highly active individuals, particularly the KJOC, was noted. All investigated upper extremity PROMs were relevant to patient concerns at level 2, but none were relevant at level 3. Therefore, we conclude that the DASH, FAST, and KJOC may be sensitive to the general concerns of participants but are not specific to the precise concerns of participants as assessed by the MYMOP-2. Only the KJOC at level 2 achieved efficiency, with the DASH and FAST containing a significant number of superfluous codes in both the level 2 and level 3 analyses. An earlier analysis¹⁴ of the DASH showed that the physiological, social, physical, and psychological domains of health and the ICF domains of body structure and function, activity, and participation are all encompassed within the DASH. This broad spectrum of health assessment likely contributed to the DASH's addressing the highest percentage of participant concerns (80%) but also having a significant amount of extraneous material (73%–85%). Both the FAST²⁰ and the KJOC¹⁷ were specifically developed with input from baseball and softball players, whereas our sample included those with upper extremity injuries from a variety of sports in addition to baseball and softball. The heterogeneity of our sample with upper extremity conditions may have contributed to the limited agreement between FAST and KJOC content and the primary concerns of our respondents. In particular, this may explain why the broad concerns of level 2 were adequately addressed but not the more specific level 3 concerns.

PROM Content Limitations and Recommendations

The content differences we observed between the evaluated PROMs and patient concerns as assessed by the MYMOP-2 were consistent with the perceived barriers and concerns identified by surveyed clinicians.^{2,5} They were also consistent with the previously reported tendency of

PROMs to focus on clinician-oriented items, such as strength and range of motion, but lacking coverage of items of importance to patients, such as psychological and social factors.³³ Not only did the majority of measures we examined represent the patient-generated concerns only superficially, they also contained a large amount of extraneous content. This extraneous content places additional demands on the patient and adds unnecessary time for clinicians to score PROMs. Furthermore, perceived barriers and concerns regarding PROM implementation may be reinforced by the PROMs being used in practice.⁶ Lam et al⁶ identified the LEFS, FAAM, and DASH as the most common extremity measures being selected by ATs using PROMs for clinical practice. None of these instruments were both relevant and efficient for evaluating the primary concerns of student-athletes experiencing injury. To combat this, we believe that the development and use of PROMs targeting athletic populations, informed by both clinician and patient concerns, are essential to facilitate the clinical utility of PROMs in athletic training.¹ Patient-reported outcome measures must be purposefully developed and selected with their end use in mind, and clinicians and researchers should not assume that a form is useful or appropriate simply because it has been used frequently in the past.

Additionally, we believe these results support the use of a patient-generated outcome measure, such as the MYMOP-2 or the Patient-Specific Functional Scale.³⁴ Such open-ended measures may be beneficial, as patient concerns were highly individualistic (Table 4) and frequently included items beyond standard ICF codes, as evidenced by the consistent presence of “not codable” items in our findings. Patient-generated items may be particularly useful if paired with the commonly used⁶ single-item Numeric Pain Rating Scale,³⁵ given that nearly 30% of the identified codes were related to “sensation of pain.” Similarly, it may be relevant to consider using psychosocial measures if a PROM addresses only physical impairments or limitations. Finally, the limited ability of the investigated PROMs to address concerns at the more specific level 3 of the ICF highlights the importance of using PROMs not only as an endpoint but as another piece of the clinical evaluation that can prompt follow-up questions and lead to deeper and more engaging conversations with patients regarding their health.¹

Limitations of the ICF

For both participant responses and PROM items, many of the final codes were those listed as *other specified or unspecified*²⁴ or “not codable.” Coding language limitations were evident when individuals reported emotional experiences or symptoms such as swelling. Concerns related to a level of function surpassing activities of daily living were also constrained by the existing ICF taxonomy. For example, although codes are available for concerns such as *sport*, *swimming*, and *lifting*, the definitions provided by the ICF do not necessarily encompass the participants' concerns. For example, the ICF code for *lifting* is defined as “raising up an object in order to move it from a lower to a higher level, such as when lifting a glass from the table.”²⁴ This definition can be applied to both lifting weights (frequently listed by our participants) and *lifting or carrying a shopping bag* (as stated on the DASH),⁷ yet

these tasks are not clinically equivalent. Such limitations in the ICF taxonomy may have contributed to some of the overlap in content between the PROM-generated codes and the participant-generated codes.

These concerns regarding the inclusivity of the ICF taxonomy are not unique. Mitra and Shakespeare³⁶ proposed that a revised ICF model would need to consider an individual's ability to participate in components of life deemed personally meaningful. Our data confirm this need, especially related to functional and psychosocial concerns. Specifically, although a small set of codes is related to handling stress, the inclusion criteria do not lend themselves to experiences frequently described by participants, such as anxiety, confidence, or frustration. Therefore, raters linked these concerns to the *emotional functions* component, defined as

Specific mental functions related to the feeling and affective components of the processes of the mind. Inclusions: functions of appropriateness of emotion, regulation and range of emotion; affect, sadness, happiness, love, fear, anger, hate, tension, anxiety, joy, sorrow; lability of emotion; flattening of affect.²⁴

However, this definition is more related to pathologic disorders than to emotional responses to injury or external stimuli. The frequency of emotional experiences among participant concerns highlights the need for improved recognition and evaluation of patient emotional states throughout care.

Study Limitations

Data were collected from 4 collegiate campuses from 2019 to 2020, and the data collection period was shortened because of the COVID-19 pandemic. Although we assumed that the responses provided by participants are generalizable to other collegiate athletes, further investigation is needed to verify this, particularly among individuals with upper extremity injuries, because of our limited sample. For both participant-generated responses and PROM questions, we chose to link every component of a response or question. This generated multiple codes for most responses and questions. If we had linked only the questions themselves, less extraneous content might have been identified; however, common concerns might also have been eliminated. Additionally, we classified injuries by body region and did not separate them by individual joints. We also operationally defined the primary concerns as those cumulatively representing 80% of all patient-generated codes and used 70% content agreement as the threshold by which to evaluate PROMs for relevance and efficiency.

CONCLUSIONS

We aimed to identify the primary concerns of collegiate athletes experiencing injuries and determine if those PROMs commonly used or recommended for use in athletic training adequately and efficiently encompass those primary concerns. These results validate the concerns expressed by clinicians regarding PROM content while also identifying the primary concerns of an athletic population as pain and sport participation ability when

assessed by the MYMOP-2. Most PROMs investigated presented with notable differences in content compared with the primary concerns of participants, particularly at the more descriptive ICF level 3. Similarly, most PROMs analyzed revealed significant amounts of extraneous content. Of the studied PROMs, the IKDC was the most relevant and efficient for collegiate athletes experiencing lower extremity injuries. Additionally, the development or use of measures designed for highly active populations based on the identified primary concerns of participants may be beneficial, as the primary codes were highly consistent (pain, sport, mobility or muscle function of the involved limb, emotional functions or experiences), regardless of the injury phase or location. Finally, improved, patient-centered evaluation of emotional experiences is needed, as this factor was largely inadequately encapsulated by both the evaluated PROMs and the ICF taxonomy. The implementation of patient-generated evidence is imperative to facilitating successful well-rounded practice, as well as the Athletic Training Research Agenda (<https://www.natafoundation.org/research/atresearchagenda>) surrounding PROMs. To continue to further these endeavors, future researchers and clinicians must focus on developing and implementing PROMs targeting athletic populations with a focus on relevance and efficiency.

REFERENCES

- Howard JS, Sciascia A, Hoch JM. Using patient evidence to guide clinical care: consulting the other expert in the room. *Int J Athl Ther Train*. 2018;23(2):53–56. doi:10.1123/ijatt.2018-0020
- Valier AR, Jennings AL, Parsons JT, Vela LI. Benefits of and barriers to using patient-rated outcome measures in athletic training. *J Athl Train*. 2014;49(5):674–683. doi:10.4085/1062-6050-49.3.15
- Sciascia AD. A basic construct for improving outcomes in rehabilitation. *Int J Athl Ther Train*. 2013;18(3):14–19. doi:10.1123/ijatt.18.3.14
- Lopes Sauers AD, Sauers EL, Valier ARS. Quality improvement in athletic health care. *J Athl Train*. 2017;52(11):1070–1078. doi:10.4085/1062-6050-52.10.15
- Coulombe BJ, Games KE, Eberman LE. The use of patient-reported outcome measures: secondary school athletic trainers' perceptions, practices, and barriers. *J Athl Train*. 2019;54(2):142–151. doi:10.4085/1062-6050-86-17
- Lam KC, Harrington KM, Cameron KL, Valier ARS. Use of patient-reported outcome measures in athletic training: common measures, selection considerations, and practical barriers. *J Athl Train*. 2019;54(4):449–458. doi:10.4085/1062-6050-108-17
- Hudak PL, Amadio PC, Bombardier C. Development of an upper extremity outcome measure: the DASH (disabilities of the arm, shoulder, and hand) [corrected]. *Am J Ind Med*. 1996;29(6):602–608. doi:10.1002/(SICI)1097-0274(199606)29:6<602::AID-AJIM4>3.0.CO;2-L
- Binkley JM, Stratford PW, Lott SA, Riddle DL. The Lower Extremity Functional Scale (LEFS): scale development, measurement properties, and clinical application. *Phys Ther*. 1999;79(4):371–383.
- Cieza A, Stucki G. The International Classification of Functioning Disability and Health: its development process and content validity. *Eur J Phys Rehabil Med*. 2008;44(3):303–313.
- Cieza A, Fayed N, Bickenbach J, Prodinger B. Refinements of the ICF Linking Rules to strengthen their potential for establishing comparability of health information. *Disabil Rehabil*. 2019;41(5):574–583. doi:10.3109/09638288.2016.1145258

11. ICF beginner's guide: towards a common language for functioning, disability and health. World Health Organization. Published 2002. Accessed December 7, 2022. <https://www.who.int/publications/m/item/icf-beginner-s-guide-towards-a-common-language-for-functioning-disability-and-health>
12. Silva Drummond A, Ferreira Sampaio R, Cotta Mancini M, Noce Kirkwood R, Stamm TA. Linking the Disabilities of Arm, Shoulder, and Hand to the International Classification of Functioning, Disability, and Health. *J Hand Ther.* 2007;20(4):336–343. doi:10.1197/j.jht.2007.07.008
13. Fairbairn K, May K, Yang Y, Balasundar S, Hefford C, Abbott JH. Mapping Patient-Specific Functional Scale (PSFS) items to the International Classification of Functioning, Disability and Health (ICF). *Phys Ther.* 2012;92(2):310–317. doi:10.2522/ptj.20090382
14. Lam KC, Marshall AN, Snyder Valier AR. Patient-reported outcome measures in sports medicine: a concise resource for clinicians and researchers. *J Athl Train.* 2020;55(4):390–408. doi:10.4085/1062-6050-171-19
15. Smith-Forbes EV, Moore-Reed SD, Westgate PM, Kibler WB, Uhl TL. Descriptive analysis of common functional limitations identified by patients with shoulder pain. *J Sport Rehabil.* 2015;24(2):179–188. doi:10.1123/jsr.2013-0147
16. Hays RD, Bjorner JB, Revicki DA, Spritzer KL, Cella D. Development of physical and mental health summary scores from the Patient-Reported Outcomes Measurement Information System (PROMIS) global items. *Qual Life Res.* 2009;18(7):873–880. doi:10.1007/s11136-009-9496-9
17. Alberta FG, ElAttrache NS, Bissell S, et al. The development and validation of a functional assessment tool for the upper extremity in the overhead athlete. *Am J Sports Med.* 2010;38(5):903–911. doi:10.1177/0363546509355642
18. Vela LI, Denegar C. Transient disablement in the physically active with musculoskeletal injuries, part 1: a descriptive model. *J Athl Train.* 2010;45(6):615–629. doi:10.4085/1062-6050-45.6.615
19. Roos EM, Roos HP, Lohmander LS, Ek Dahl C, Beynnon BD. Knee Injury and Osteoarthritis Outcome Score (KOOS)—development of a self-administered outcome measure. *J Orthop Sports Phys Ther.* 1998;28(2):88–96. doi:10.2519/jospt.1998.28.2.88
20. Sauers EL, Bay RC, Snyder Valier AR, Ellery T, Huxel Bliven KC. The Functional Arm Scale for Throwers (FAST)—part I: the design and development of an upper extremity region-specific and population-specific patient-reported outcome scale for throwing athletes. *Orthop J Sports Med.* 2017;5(3):2325967117698455. doi:10.1177/2325967117698455
21. Irrgang JJ, Anderson AF, Boland AL, et al. Development and validation of the International Knee Documentation Committee Subjective Knee Form. *Am J Sports Med.* 2001;29(5):600–613. doi:10.1177/03635465010290051301
22. Martin RL, Irrgang JJ, Burdett RG, Conti SF, Van Swearingen JM. Evidence of validity for the Foot and Ankle Ability Measure (FAAM). *Foot Ankle Int.* 2005;26(11):968–983. doi:10.1177/107110070502601113
23. Ishaque S, Johnson JA, Vohra S. Individualized health-related quality of life instrument Measure Yourself Medical Outcome Profile (MYMOP) and its adaptations: a critical appraisal. *Qual Res.* 2019;28(4):879–893. doi:10.1007/s11136-018-2046-6
24. ICF browser. World Health Organization. Accessed June 2, 2019. <https://apps.who.int/classifications/icfbrowser/>
25. Portney LG, Watkins MP. *Foundations of Clinical Research: Applications to Practice.* 2nd ed. Prentice Hall Health; 2000.
26. Paterson C. Seeking the patient's perspective: a qualitative assessment of EuroQol, COOP-WONCA charts and MYMOP. *Qual Life Res.* 2004;13(5):871–881. doi:10.1023/B:QURE.0000025586.51955.78
27. Polus BI, Kimpton AJ, Walsh MJ. Use of the Measure Your Medical Outcome Profile (MYMOP2) and W-BQ12 (Well-Being) outcomes measures to evaluate chiropractic treatment: an observational study. *Chiropr Man Therap.* 2011;19:7. doi:10.1186/2045-709X-19-7
28. Starkey C, Brown SD. *Examination of Orthopedic & Athletic Injuries.* 4th ed. FA Davis; 2015.
29. Smith MD, Vicenzino B, Bahr R, et al. Return to sport decisions after an acute lateral ankle sprain injury: introducing the PAASS framework—an international multidisciplinary consensus. *Br J Sports Med.* 2021;55(22):1270–1276. doi:10.1136/bjsports-2021-104087
30. Terwee CB, Prinsen CAC, Chiarotto A, et al. COSMIN methodology for evaluating the content validity of patient-reported outcome measures: a Delphi study. *Qual Life Res.* 2018;27(5):1159–1170. doi:10.1007/s11136-018-1829-0
31. Majewski-Schrage T, Evans TA, Snyder KR. Identifying meaningful patient outcomes after lower extremity injury, part 1: patient experiences during recovery. *J Athl Train.* 2019;54(8):858–868. doi:10.4085/1062-6050-232-18
32. Meadows KA. Patient-reported outcome measures: an overview. *Br J Community Nurs.* 2011;16(3):146–151. doi:10.12968/bjcn.2011.16.3.146
33. Majewski-Schrage T, Evans TA, Snyder KR. Identifying meaningful patient outcomes after lower extremity injury, part 2: linking outcomes to the International Classification of Functioning, Disability and Health. *J Athl Train.* 2019;54(8):869–880. doi:10.4085/1062-6050-233-18
34. Stratford P, Gill C, Westaway M, Binkley J. Assessing disability and change on individual patients: a report of a patient specific measure. *Physiother Can.* 1995;4(47):258–263. doi:10.3138/ptc.47.4.258
35. Downie WW, Leatham PA, Rhind VM, Wright V, Branco JA, Anderson JA. Studies with pain rating scales. *Ann Rheum Dis.* 1978;37(4):378–381. doi:10.1136/ard.37.4.378
36. Mitra S, Shakespeare T. Remodeling the ICF. *Disabil Health J.* 2019;12(3):337–339. doi:10.1016/j.dhjo.2019.01.008

SUPPLEMENTAL MATERIAL

Found at DOI: <http://dx.doi.org/10.4085/1062-6050-0516.21.S1>

Address correspondence to Jennifer Sebert Howard, PhD, LAT, ATC, Appalachian State University, PO Box 32165, Boone, NC 28608. Address email to Howardjs@appstate.edu.