

Common Social and Health Disparities Contribute to Racial Differences in Ambulatory Impairment in Multiple Sclerosis

Farren B.S. Briggs, PhD, ScM; Erika S. Trapl, PhD; Farrah J. Mateen, MD, PhD; Alessandro De Nadai, PhD; Devon S. Conway, MD, MS; Douglas D. Gunzler, PhD

CE INFORMATION

ACTIVITY AVAILABLE ONLINE: To access the article and evaluation online, go to <https://www.highmarksce.com/mscare>.

TARGET AUDIENCE: The target audience for this activity is physicians, advanced practice clinicians, nursing professionals, social workers, and other health care providers involved in the management of patients with multiple sclerosis (MS).

LEARNING OBJECTIVE:

1. Describe the extent to which common social and health disparities contribute to racial differences in ambulatory impairment in MS.
2. Recognize the importance of distinguishing mediators from confounders in multivariable regression models.

ACCREDITATION:



JOINT ACCREDITATION
INTERPROFESSIONAL CONTINUING EDUCATION

In support of improving patient care, this activity has been planned and implemented by the Consortium of Multiple Sclerosis Centers (CMSC) and Intellisphere, LLC. The CMSC is jointly accredited by the Accreditation Council for Continuing Medical Education (ACCME), the Accreditation Council for Pharmacy Education (ACPE), and the American Nurses Credentialing Center (ANCC), to provide continuing education for the health care team.



IPCE CREDIT™

This activity was planned by and for the health care team, and learners will receive .5 Interprofessional Continuing Education (IPCE) credit for learning and change.

PHYSICIANS: The CMSC designates this journal-based activity for a maximum of 0.5 *AMA PRA Category 1 Credit(s)*™. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

NURSES: The CMSC designates this enduring material for .5 contact hour of nursing continuing professional development (NCPD) (none in the area of pharmacology).

PSYCHOLOGISTS: This activity is awarded .5 CE credit.

SOCIAL WORKERS: As a Jointly Accredited Organization, the CMSC is approved to offer social work continuing education by the Association of Social Work Boards (ASWB) Approved Continuing Education (ACE) program. Organizations, not individual courses, are approved under this program. Regulatory boards are the final authority on courses accepted for continuing education credit. Social workers completing this course receive .5 general continuing education credits.

DISCLOSURES: It is the policy of the Consortium of Multiple Sclerosis Centers to mitigate all relevant financial disclosures from planners, faculty, and other persons that can affect the content of this CE activity. For this activity, all relevant disclosures have been mitigated.

Francois Bethoux, MD, editor in chief of the *International Journal of MS Care (IJMSC)*, has served as physician planner for this activity. He has disclosed no relevant financial relationships. **Alissa Mary Willis, MD**, associate editor of *IJMSC*, has disclosed not relevant financial relationships. Authors: **Farren B.S. Briggs, PhD, ScM; Farrah J. Mateen, MD, PhD; Devon Conway, MD, MS; Alessandro de Nadai, PhD; Erika S. Trapl, PhD; and Douglas D. Gunzler, PhD**, have disclosed no relevant financial relationships.

The staff at *IJMSC*, CMSC, and Intellisphere, LLC who are in a position to influence content have disclosed no relevant financial relationships. **Laurie Scudder, DNP, NP**, continuing education director CMSC, has served as a planner and reviewer for this activity. She has disclosed no relevant financial relationships.

METHOD OF PARTICIPATION:

Release Date: January 1, 2024; Valid for Credit through: January 1, 2025

To receive CE credit, participants must:

- (1) Review the continuing education information, including learning objectives and author disclosures.
- (2) Study the educational content.
- (3) Complete the evaluation, which is available at <https://www.highmarksce.com/mscare>.

Statements of Credit are awarded upon successful completion of the evaluation. There is no fee to participate in this activity.

DISCLOSURE OF UNLABELED USE: This educational activity may contain discussion of published and/or investigational uses of agents that are not approved by the FDA. The CMSC and Intellisphere, LLC do not recommend the use of any agent outside of the labeled indications. The opinions expressed in the educational activity are those of the faculty and do not necessarily represent the views of the CMSC or Intellisphere, LLC.

DISCLAIMER: Participants have an implied responsibility to use the newly acquired information to enhance patient outcomes and their own professional development. The information presented in this activity is not meant to serve as a guideline for patient management. Any medications, diagnostic procedures, or treatments discussed in this publication should not be used by clinicians or other health care professionals without first evaluating their patients' conditions, considering possible contraindications or risks, reviewing any applicable manufacturer's product information, and comparing any therapeutic approach with the recommendations of other authorities.

ABSTRACT

BACKGROUND: We previously reported more rapid accrual of ambulatory impairments in Black compared to White individuals with relapsing remitting multiple sclerosis (RRMS) and higher body mass index (BMI). Hypertension and lower neighborhood socioeconomic status (SES) were associated with greater impairment, irrespective of race. We hypothesize that these common social and health inequities may explain a substantial portion of the racial differences in ambulation in American individuals with RRMS.

METHODS: Causal mediation analyses investigated baseline and change-over-time mediators of ambulatory impairment differences between 1795 Black and White individuals with RRMS using a retrospective cohort study comprised of electronic health record data from 8491 clinical encounters between 2008 and 2015 where Timed 25-Foot Walk (T25FW) speeds without assistive devices were recorded. The hypothesis was that BMI, neighborhood SES, and hypertension were possible mediators.

RESULTS: At baseline, Black individuals with RRMS ($n = 175$) had significantly slower T25FW speeds (5.78 vs 5.27 ft/s), higher BMI, a higher prevalence of hypertension, and they were more likely to live in lower-income neighborhoods than White individuals ($n = 1,620$). At baseline, a significant proportion (33.7%; 95% CI, 18.9%-59.4%) of the T25FW difference between Black and White individuals was indirectly due to a higher BMI (12.5%), hypertension burden (9.5%), and living in lower-income neighborhoods (11.2%). Once baseline mediation relationships were accounted for, there were no significant longitudinal mediation relationships.

CONCLUSIONS: The findings implicate social and health disparities as prominent drivers of ambulatory differences between Black and White individuals with RRMS, suggesting that wellness and health promotion are essential components of MS care, particularly for Black individuals.

Int J MS Care. 2024;26(1):36-40. doi: 10.7224/1537-2073.2023-004

conclusions about MS in BAs.² Despite these limitations, we know that the 100,000 BAs with MS are prone to more severe disease and adverse long-term outcomes.^{1,3}

In a retrospective cohort study of electronic health records (EHRs) that comprehensively adjusted for sociodemographic and clinical attributes, we previously reported that BAs with relapsing remitting multiple sclerosis (RRMS) were more prone to rapidly advancing ambulatory impairment than White Americans (WA) with RRMS.⁴ We also observed that higher body mass index (BMI), lower neighborhood socioeconomic status (SES), and hypertension were independently associated with greater ambulatory impairments in individuals with RRMS, and these relationships did not differ between BAs and WAs.^{4,5} Due to structural racism (ie, generational impact of redlining) and social and health inequities in the United States, BAs are prone to live in lower-income neighborhoods and have a greater burden of obesity and hypertension than WAs.⁶ There are similar racial differences in MS, including evidence supporting a higher burden of uncontrolled hypertension in BAs compared with WAs with MS.^{4,7,8} Therefore, we theorize that these common social and health inequities (ie, racial disparities) substantially contribute to the observed racial differences in MS. This is supported, in part, by a 2006 cross-sectional study of North Americans with MS, where Black compared with White participants had greater impairments, and associations attenuated when adjusted for SES.⁹

To date, no study has sought to identify mediators of racial disparities in MS risk or progression. In hindsight, we recognize our prior multivariable longitudinal models of ambulatory impairment that included sociodemographic and clinical attributes assumed “all else being equal”⁴; however, all else is not equal in the United States due to pervasive social and health inequities that disproportionately impact BAs. To characterize the true total magnitude of a racial disparity, it is important to appropriately adjust for confounders. Oftentimes mediators of the racial disparity (ie, descendants of race in a causal diagram that lead to the outcome; mechanisms through which race impacts the outcome) are included as covariates. Thus, the adjusted estimates for race from these models only reflect a portion of the true underlying relationship (ie, the effect of race independent of mediators in the model), and fail to illustrate the true magnitude of the racial difference. Here we conduct mediation analyses, hypothesizing that BMI, neighborhood

There are remarkable racial differences in multiple sclerosis (MS) risk and presentation, and data suggest racial health and social disparities may be key drivers of these observed differences.¹ A decade ago, Black/African Americans (BA) were shown to be the most at-risk American subpopulation for MS, yet contemporary research characterizing racial differences and racial disparities have been notably sparse.¹ This evidence gap is compounded by continued underrepresentation of diverse populations in clinical trials, further impeding efforts to make generalizable

From the Department of Public Health Sciences, Miller School of Medicine, University of Miami, Miami, FL, USA (FBSB); the Department of Population and Quantitative Health Sciences (FBSB, EST, DDG), the Prevention Research Center for Healthy Neighborhoods (EST), and the Center for Health Care Research and Policy (DDG), School of Medicine, Case Western Reserve University, Cleveland, OH, USA; the Department of Neurology, Division of Neuroimmunology and Neuroinfectious Diseases, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA (FJM); McLean Hospital, Harvard Medical School, Belmont, MA, USA (ADN); the Department of Neurology, Neurological Institute, Mellen Center for Multiple Sclerosis Treatment and Research, Cleveland Clinic Foundation, Cleveland, OH, USA (DSC); and the Cleveland Clinic Lerner College of Medicine, Case Western Reserve University, Cleveland, OH, USA (DSC). Correspondence: Farren B.S. Briggs, PhD, ScM, Department of Public Health Sciences, Miller School of Medicine, University of Miami, 1120 NW 14th St, 912 CRC, Miami, FL, 33136; email: fbb31@miami.edu

© 2024 Consortium of Multiple Sclerosis Centers.

SES, and hypertension are prominent mechanisms of the observed racial difference in ambulatory impairment between BAs and WAs with RRMS.

METHODS

We revisited our retrospective cohort study to examine cross-sectional and longitudinal mediators of baseline and change-over-time racial disparities in ambulatory impairment in patients with RRMS. These analyses were approved by the institutional review boards at the Cleveland Clinic Foundation and Case Western Reserve University. The source population consisted of RRMS patients seeking extended care at a tertiary MS referral center in the United States between 2008 and 2015.^{4,5} We excluded patients requiring unilateral or bilateral assistive devices at baseline as their distribution of Timed 25-Foot Walk (T25FW) speed at baseline was right-skewed with wide standard deviations (data not shown), which would introduce heterogeneity in the subsequent models. There were 1795 individuals with RRMS who reported that they did not require an assistive device and T25FW (objective measure of ambulation) was recorded in 8491 clinical encounters, which spanned an average interval of 2.5 years (SD = 0.8). Diagnosis of MS was confirmed by chart review.¹⁰ Age, sex, race (Black or White only), disease duration, disease-modifying therapy (DMT) status (yes/no), smoking status (ever/never), BMI, hypertension (yes/no), and 9-digit ZIP code of residence (mapped to 520 census tracts and linked to 2010 US Census median household income data) were abstracted from EHRs, as previously described.^{4,5} To determine a hypertension diagnosis, 4 EHR domains were inspected (progress notes, problem list, past medical history, medication list) and a patient was classified as having hypertension based on evidence from at least 2 domains (eg, listed in the medical history and having an antihypertensive prescription).⁵

Baseline attributes were compared using Fisher's exact and Wilcoxon rank-sum tests for categorical and continuous variables, respectively. Mediation analyses were informed by path diagrams that illustrate the directionality of the associations of interest (FIGURE S1, available online at IJMSc.org), and direct acyclic graphs were used to identify confounders to be included as covariates (FIGURE S2). Mediators of interest were baseline measures of BMI, hypertension, and median household income of ZIP code of residence. Other abstracted variables were covariates.

Mediation analyses were conducted to investigate mediators of racial disparities in T25FW, informed by directed acyclic graphs (Figure S2). Mediators of interest were baseline measures of BMI, hypertension, and median household income, and covariates included age, sex, disease duration, smoking status, and DMT status. First, a latent growth model evaluated linear growth over time in T25FW using maximum likelihood to estimate

the parameters with robust Huber-White sandwich estimation to calculate standard errors (MLR option in MPlus v8.2).^{11,12} Robust approaches are insensitive to departures from parametric assumptions that may exist in the data, and the model effectively handled ignorable missing data dependent on the data in hand (ie, following a missing at random assumption) via full information maximum likelihood. Therefore, all patients were included in the trajectory analysis for unbiased inference. Second, the latent growth model was extended to include the hypothesized causal pathways, and accounting for covariates, to investigate mediation relationships (Figure S1).^{13,14} That is, the growth of the T25FW (via a latent intercept and linear slope) was regressed on all baseline measures (the mediators, race and covariates), while baseline measures of the multiple mediators were regressed on race and the covariates in a single analysis. There was no significant growth in T25FW after accounting for the baseline mediation relationships; thus, the model was reduced to evaluate cross-sectional mediation alone. Given the cross-sectional setting, we were able to explicitly address nonlinearity that may arise due to hypertension being a binary mediator employing a Bayes estimator and a Markov chain Monte Carlo algorithm based on the Gibbs sampler with a noninformative prior. Direct, specific indirect, total indirect, and total effects were then calculated (IND option in MPlus v8.2) with 95% Bayesian credibility intervals (CI) to test for mediation. A 2-sided alpha of 5% determined statistical significance in our frequentist modeling (ie, latent growth modeling approaches). Data analyses took place in April 2022.

RESULTS

There were 175 BAs and 1620 WAs with RRMS who did not differ by sex or age (TABLE 1). At baseline, BAs had significantly slower T25FW speeds (5.78 vs 5.27 ft/s), shorter disease duration, lower DMT usage, and were less likely to have ever smoked than WAs. Black/African American patients also had significantly higher BMI, a higher prevalence of hypertension, and they were more likely to live in lower-income neighborhoods than WAs seen at the same tertiary institution.

There was significant worsening in T25FW over time (FIGURE S3; intercept = 5.3s, SE = 0.035, $P < .001$; slope = 0.012s/month, SE = 0.001, $P < .001$). However, there was no change-over-time once baseline mediation relationships were accounted for (intercept = 3.4s, SE = 0.205, $P < .001$; slope = -0.011s/month, SE = 0.008, $P = .170$). At baseline, the total direct effect of being a BA was a 0.56s (95% CI, 0.35-0.77) increase in T25FW compared to WAs, which encompasses a direct effect and hypothesized indirect effects adjusted for other covariates (TABLE 2). This total effect is very similar to the unadjusted difference (Table 1), which illustrates that BAs take, on average, about 10% longer to complete the T25FW. A significant proportion (33.7% [95% CI, 18.9%-59.4%]; 0.19s [95% CI, 0.11s-0.27s])

TABLE 1. Study Population Attributes at Baseline

Variable	All RRMS	Black Americans	White Americans	P
Participants	1795	175 (9.8%)	1620 (90.2%)	
T25FW (seconds)	5.32	5.78	5.27	8.1 × 10 ⁻⁸
Slow ambulators (T25FW > 8s)	98 (5.5%)	11 (6.3%)	87 (5.4%)	0.60
BMI (mean; kg/m ²)	28.6	30.7	28.4	3.4 × 10 ⁻⁵
Hypertension	323 (18.0%)	52 (29.7%)	271 (16.7%)	6.9 × 10 ⁻⁴
Median household income by ZIP code (2010)	\$55,300	\$44,100	\$56,500	3.0 × 10 ⁻¹⁹
Female	1329 (74.0%)	136 (77.7%)	1193 (73.6%)	0.28
Age (years)	42.9	41.6	43.0	0.06
Never smoker	730 (40.7%)	94 (53.7%)	636 (39.3%)	0.0013
Disease duration (years)	9.4	7.6	9.6	0.014
On DMT	1265 (70.5%)	108 (61.7%)	1157 (71.4%)	0.0088

BMI, Body Mass Index; DMT disease-modifying therapy; RRMS, relapsing remitting multiple sclerosis; T25FW, Timed 25-Foot Walk.

TABLE 2. Cross-Sectional Mediation Statistics Using Bayes Estimator for Racial Disparity in Timed 25-Foot Walk

Effect of identifying as Black for individuals with RRMS	Coefficient (95% CI)	Percent of total effect (95% CI)
Total effect	0.558s (0.349, 0.768)	100%
Total direct effect	0.369s (0.151, 0.585)	66.3% (40.6, 81.1)
Total indirect effect	0.187s (0.113, 0.269)	33.7% (18.9, 59.4)
Indirect effect	Body mass index	12.5% (5.7, 23.2)
	Hypertension	9.5% (1.1, 23.4)
	Lower income neighborhood	11.2% (3.2, 23.6)

of this racial difference in ambulation speeds was due to racial health disparities as a result of a higher burden of social and health inequities in BAs with RRMS compared to WAs with RRMS, specifically of a higher BMI, a greater burden of hypertension, and a greater likelihood of living in a lower-income neighborhood (Table 2). Higher BMI mediated 12.5% (95% CI, 5.7-23.2) of the racial difference, followed by 11.2% (95% CI, 3.2-23.6) for living in a lower-income neighborhood, and 9.5% (95% CI, 1.1-23.4) for hypertension. The direct effect of being a BA on ambulation, after accounting for the mediated relationships and adjusting for covariates, was 0.37s (95% CI, 0.15-0.58); thus, 66.1% (95% CI, 40.6-81.1) of the racial difference in ambulation between Black and White individuals with RRMS remains unexplained and it is likely that other common drivers of racial disparities may further explain these differences (ie, differential prescribing patterns, allostatic load, earning potential).

DISCUSSION

There are substantial racial differences in MS, and inequities that drive racial health disparities play a prominent role in these observed patterns. Here, we demonstrate that, on average, Black individuals with RRMS take about 10.5% (0.558s/5.32s) longer to complete a timed ambulatory task compared to their WA counterparts. A third of this difference is due to racial health and social disparities that can be attributed to higher BMI (12.5%), a higher burden of hypertension (9.5%), and living in lower-income

neighborhoods (11.2%). Had we treated these 3 mediators as confounders (covariates in a multivariable model), we would have reported that BAs with RRMS were, at most, 6.9% slower (0.369s/5.32s)—an underestimate of 33%. This work highlights the need to judiciously consider which potential covariates may be mediators (downstream of the exposure of interest [being BA in this study] and leading to the outcome) and how it may impact the interpretation of findings. In the case of our prior study of longitudinal change in T25FW, we should have said that the reported difference between BAs and WAs was *only* the direct effect and likely an underestimate of the true difference, which would mean that the accrual of ambulatory impairment in BAs is even greater than we previously reported.

The presented work is not comprehensive, but it is a start to the conversations that must take place because contemporary race-specific or race-comparative studies of MS outcomes continue to be sparse and often rely on multivariable models that may treat mediators as confounders. Our prior study, like most other findings, likely failed to quantify the true underlying racial difference (via direct and indirect effects) in MS outcomes, much less the extent to which disparities drive these relationships. As a result, the prior findings were likely dampened, diminishing the urgency to address them.

While this work is meant to start the discourse, there are several limitations to acknowledge. First, we were only able to focus on 3 common social and health inequities and were unable to examine other individual-level social



As common social and health disparities in the United States contribute to racial differences in ambulatory impairment, clinicians should view wellness and health promotion strategies as essential components of multiple sclerosis care, particularly for Black individuals. ■

determinants of health (eg, financial insecurity, changes in insurance status, housing instability) and other systemic health care biases (ie, that a greater proportion of BAs are not on a DMT). We only examined these relationships in a study population that originated from a single tertiary care setting; although the ratio of BA to WA in this study sample is reflective of the ratio observed within the MS population in the United States, it is not reflective of the racially diverse metropolitan setting (Cleveland, 47% BA)¹⁵ nor the relatively moderately diverse region (Northeast Ohio, 17% BA)¹⁶ of the tertiary facility. It is plausible that selection bias may have factored into the study sample, such that the included patients only represent a subset of the source population, for example, those with specific health-care-seeking behaviors (ie, having health insurance, which facilitates referral to a tertiary care facility) and more active/aggressive disease (therefore, more likely to seek specialty care). We also excluded those participants who required an assistive device at baseline, limiting our generalizability to individuals with RRMS who ambulate without assistance. A notable strength of the study is that we were able to account for key correlates of disease activity/progression and health-care-seeking behaviors (ie, age, sex, smoking, disease duration, and treatment status), as well as examine whether there were cross-sectional and longitudinal mediation effects.

CONCLUSIONS

In summary, racial disparities matter in MS, as illustrated in this first-of-its-kind mediation analysis. We demonstrate that a higher BMI, a higher hypertension burden, and living in a lower-income neighborhood explained a third of the ambulatory impairment difference between BA and WA individuals with RRMS. How neighborhood SES and hypertension may contribute to ambulatory impairments are not directly clear and merit further investigation, particularly

in BA individuals with RRMS. It is critical that future work should further illuminate the extent to which other inequities and racial disparities impact observed clinical relationships in MS, which together can be leveraged to advocate for investing more resources in health promotion and preventive care as integral components of MS care. Such efforts will have the most benefit for the most vulnerable individuals with MS. ■

CONFLICTS OF INTEREST: On behalf of all authors, the corresponding author states that there is no conflict of interest.

PREVIOUS PRESENTATION: This work was presented at the 2021 annual meeting of the Consortium of Multiple Sclerosis Centers; October 25-28; Orlando, Florida. This work was supported by funding from the National Institutes of Health/National Institute of Nursing Research (R56NR019306).

REFERENCES

1. Amezcua L, Rivera VM, Vazquez TC, Baezconde-Garbanati L, Langer-Gould A. Health disparities, inequities, and social determinants of health in multiple sclerosis and related disorders in the US: a review. *JAMA Neurol.* 2021;78(12):1515-1524. doi:10.1001/jamaneurol.2021.3416
2. Mateen FJ. Is it time for quotas to achieve racial and ethnic representation in multiple sclerosis trials? *Front Neurol.* 2021;12:680912. doi:10.3389/fneur.2021.680912
3. Briggs FB, Hill E. Estimating the prevalence of multiple sclerosis using 56.6 million electronic health records from the United States. *Mult Scler.* 2020;26(14):1948-1952. doi:10.1177/1352458519864681
4. Briggs FBS, Thompson NR, Conway DS. Prognostic factors of disability in relapsing remitting multiple sclerosis. *Mult Scler Relat Disord.* 2019;30:9-16. doi:10.1016/j.msard.2019.01.045
5. Conway DS, Thompson NR, Cohen JA. Influence of hypertension, diabetes, hyperlipidemia, and obstructive lung disease on multiple sclerosis disease course. *Mult Scler.* 2017;23(2):277-285. doi:10.1177/1352458516650512
6. Bell CN, Thorpe RJ Jr, Bowie JV, LaVeist TA. Race disparities in cardiovascular disease risk factors within socioeconomic status strata. *Ann Epidemiol.* 2018;28(3):147-152. doi:10.1016/j.annepidem.2017.12.007
7. Briggs FBS, Hill E, Abboud H. The prevalence of hypertension in multiple sclerosis based on 37 million electronic health records from the United States. *Eur J Neurol.* 2021;28(2):558-566. doi:10.1111/ene.14557
8. Conway DS, Briggs FB, Mowry EM, Fitzgerald KC, Hersh CM. Racial disparities in hypertension management among multiple sclerosis patients. *Mult Scler Relat Disord.* 2022;64:103972. doi:10.1016/j.msard.2022.103972
9. Marrie RA, Cutter G, Tyry T, Vollmer T, Campagnolo D. Does multiple sclerosis-associated disability differ between races? *Neurology.* 2006;66(8):1235-1240. doi:10.1212/01.wnl.0000208505.81912.82
10. Polman CH, Reingold SC, Banwell B, et al. Diagnostic criteria for multiple sclerosis: 2010 revisions to the McDonald criteria. *Ann Neurol.* 2011;69(2):292-302. doi:10.1002/ana.22366
11. Muthén LK, Muthén BO. *Mplus: the comprehensive modelling program for applied researchers: user's guide.* Muthén and Muthén; 2012;5.
12. Huber PJ. The behavior of maximum likelihood estimates under nonstandard conditions. In: Le Cam LM, Neyman J, eds. *Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and Probability.* University of California Press; 1967:221-233.
13. Gunzler DD, Perzynski AT, Carle AC. *Structural Equation Modeling for Health and Medicine.* Chapman & Hall; 2021.
14. MacKinnon D. *Introduction to Statistical Mediation Analysis.* Routledge; 2012.
15. United States Census Bureau. QuickFacts Cleveland city, Ohio. Accessed November 10, 2023. <https://www.census.gov/quickfacts/fact/table/clevelandcityohio/PSTo45222>
16. Healthy Northeast Ohio. 2023 demographics. Accessed November 10, 2023. <https://www.healthynortheastohio.org/demographicdata?id=285563§ionId=940>