

The Use of Traditional and Complementary Medicine Among Patients With Multiple Sclerosis in Morocco

Rachid Lotfi, PhD; Mourad Chikhaoui, PhD; Abdessamad Elmourid, PhD; and Fatiha Chigr, PhD

ABSTRACT

BACKGROUND: Multiple sclerosis (MS) is an acquired chronic, autoimmune, and neurodegenerative disease of the central nervous system. In addition to conventional MS therapy, patients are interested in traditional and complementary medicine (T&CM). Our study aims to describe the use of T&CM in a cohort of Moroccan patients with MS.

METHODS: A quantitative descriptive study was adopted to study this subject. For data collection, we opted for an anonymous questionnaire for 98 patients with MS. We gathered data via an electronic survey, using multivariable analysis to examine the effect of specific factors on T&CM use. Data collection took place from March 2022 to June 2022.

RESULTS: The results show that 52% of patients use T&CM. Of those, 29.6% use cupping, 23.5% recite the Holy Quran, 15.3% use phytotherapy, 13.2% use apitherapy, and 10.2% use acupuncture. In addition, 66.3% of the surveyed respondents stated that alternative medicine positively affects their health. Finally, 49.1% of surveyed patients who use alternative medicine are between 30 and 40 years old. The results also show that the rate of T&CM use is higher in patients with progressive MS (OR = 2.540) and patients without financial access to disease-modifying therapy (OR = 2.100).

CONCLUSIONS: This study invites us to consider societal, cultural, and economic factors when studying the use of T&CM among people with MS. Further research is needed to understand the motivations for using T&CM.

Int J MS Care. 2024;26:140-143. doi:10.7224/1537-2073.2022-116

Multiple sclerosis (MS) is the most common chronic inflammatory, degenerative, and demyelinating disease of the central nervous system. Medications are used for MS treatment in 3 main ways: disease-modifying therapies (DMTs), symptom management, and relapse treatment. In addition to drug treatment, patients are interested in traditional and complementary medicine (T&CM).

Traditional medicine is “the total of the knowledge, skill, and practices based on the theories, beliefs, and experiences indigenous to different cultures, whether explicable or not, used to treat physical and mental illness.”¹ Complementary medicine refers to a broad set of health care practices that are not part of that country’s traditional or conventional medicine and are not fully integrated into the dominant health care system.¹ Among the T&CM techniques used in Morocco are cupping (cups placed over the skin to create negative pressure through suction²), reciting the Holy Quran (to encourage the production of endorphins by activating alpha brain waves³), phytotherapy (plants or plant extracts for medicinal uses), apitherapy (bee product therapy), and massage therapy (therapeutic manipulation by the use of hands or mechanical devices for maintaining the suppleness of the body⁴). Other common T&CM techniques that are used to treat MS include acupuncture (insertion of a thin metal needle into a patient’s skin at an acupoint⁵), aromatherapy (promotes health and well-being by utilizing pure plant oils⁶), and yoga (combination of meditation and movement⁷). Studies show that T&CM may improve MS symptoms such as fatigue, motor inflammation, and pain.⁸⁻¹⁰

In addition to dissatisfaction with currently available treatments that do not offer a cure, patients use T&CM for several other reasons, especially the clinical and evolutionary characteristics of MS itself, as it is marked by an intermittent evolution with exacerbation and remission. Sociocultural variables, such as anecdotal reports of the advantages of T&CM, also promote its use.¹¹ As T&CM modalities and the reasons

From the Biological Engineering Laboratory, Faculty of Sciences and Techniques, University Sultan Moulay Slimane, Bèni Mellal, Morocco (RL, FC); Laboratory of Ecology and Environment, Faculty of Sciences Ben M’Sik, Hassan II University of Casablanca, Casablanca, Morocco (MC); and Polyvalent Team in Research and Development, Department of Biology and Geology, Polydisciplinary Faculty, Sultan Moulay Slimane University, Bèni Mellal, Morocco (AE). *Correspondence:* Rachid Lotfi, PhD, Biological Engineering Laboratory, Faculty of Sciences and Techniques, University Sultan Moulay Slimane, Bèni Mellal, 23030, Morocco; *email:* r.lotfi@usms.ma.

© 2024 Consortium of Multiple Sclerosis Centers.



In a study of individuals with multiple sclerosis in Morocco, more than half of the participants use traditional and complementary medicine, and two-thirds were satisfied with this use.

The rate of use of traditional and complementary medicine is higher in patients with progressive multiple sclerosis and those without access to disease-modifying therapies because of cost. ■

for their use have yet to be formally studied in Morocco, our study aims to describe T&CM use and identify some of the reasons patients with MS use it.

METHODS

Study Design

We conducted a descriptive study with a sample of 98 persons representing the following 6 Moroccan regions: Fès-Meknès, Rabat-Salè-Kènitra, Bèni Mellal-Khenifra, Casablanca-Settat, Marrakesh-Safi, and Oriental. To recruit participants, we used the snowball survey method: we identified a few patients with MS and asked them to suggest other people for the survey. In addition, we also contacted associations active in this field, which helped us collect data.

We collected data via an online survey using Sphinx Declic v.4.29 (Sphinx), a platform that specializes in remote scientific surveys. We shared the survey link in WhatsApp groups for individuals with MS.

To ensure the validity of responses, we integrated the platform and WhatsApp groups to explain the objectives and variables of the study to participants. In addition, participants could contact us directly if they needed help understanding the survey questions. We discussed any problems encountered in completing the questionnaire via the WhatsApp groups. Prior to distribution to the complete cohort, we conducted a pilot survey with 5 individuals with MS to adjust and finalize the questionnaire.

We obtained written approval for this study from Sultan Moulay Slimane University and the Health Ministry of Morocco on April 30, 2022 (reference no. 8366-3/3/2021) as well as the university's ethics committee (approval number FST/LGB/2016/28-FEB./2017-DEC.2019). All participants provided informed written consent after the purpose of the study, the importance of their contribution, and their right to

refuse to participate was explained. The data are anonymized and free of personally identifiable information.

Measure

The questionnaire asked for information on sociodemographic characteristics (age, sex, employment status, marital status, education level, place of residence), clinical aspects, and treatments used since MS diagnosis. We also asked participants about the types of T&CM they used, the results of this use (satisfaction with, decrease in number of relapses and physical pain), and factors in their decision to use T&CM (clinical, sociodemographic, and economic characteristics).

Statistical Analysis

We used IBM's SPSS Statistics, version 20, for the statistical analysis and the χ^2 test to identify the link between the use of T&CM and other qualitative variables, namely sociodemographic characteristics, clinical phenotype, and treatment cost. For the link between a quantitative variable and a qualitative variable, we used the Student *t* test and Fisher exact test and presented the data as means plus or minus SD. We used multivariable logistic regression models (binary and multinomial) to examine the effect of specific factors on the use of T&CM.

RESULTS

The survey group was female (68%), generally young (mean age 34.49; SD = 10.49 years), and likely to be employed (42%; as juxtaposed to a 39.7 % employment rate for the entire Moroccan population¹²). More than 53% had no governmental or private medical coverage. As shown in **TABLE 1**, 51 of the respondents used T&CM and more than 9 types of T&CM were mentioned. Types of T&CM used most often were cupping (56.9%), reciting the Holy Quran (45%), phytotherapy (29.4%), apitherapy (25.5%), massage therapy (19.6%), and acupuncture (19.6%). Of the patients who used T&CM, 49.1% were between 30 and 39 years old.

The statistical analysis showed no statistically significant relationship between the use of T&CM and the following factors: age ($P = .122$), sex ($P = .741$), employment status ($P = .878$), financial ability to access DMT ($P = .557$), marital status ($P = .235$), or number of relapses ($P = .714$). However, the rate of T&CM use was higher in patients with progressive MS (OR = 2.540, 95% CI, 0.851-7.575) and patients without financial access to DMT (OR = 2.100, 95% CI, 0.176-25.010) (**TABLE 2**).

DISCUSSION

Studies on T&CM use in the MS population report that, in general, between 27% and 100% of persons with MS reported using, per the language in some of the cited studies, complementary and alternative medicine (CAM).¹³⁻¹⁸ Skovgaard and colleagues showed that the total prevalence of CAM use by people with MS over the previous year varied from 46.0% in Sweden to 58.9% in Iceland.¹⁹ A study by Tekin and colleagues found that 29.9% of individuals with MS used T&CM practices at least once in the hopes of being cured or to feel better physically or psychologically.¹³

TABLE 1. Participants' Survey Responses (N = 98)

	n	%
Use of T&CM		
Yes	51	52
No	47	48
Age (n = 51)		
< 30	12	23.5
30-39	25	49
40-49	8	15.7
50 and over	6	11.8
Type of T&CM (n = 51)		
Cupping	29	56.9
Reciting the Holy Quran	23	45
Phytotherapy	15	29.4
Apitherapy	13	25.5
Massage therapy	10	19.6
Acupuncture	10	19.6
Aromatherapy	5	9.8
Yoga	3	5.9
Other (eg, ozone therapy, diet, dietary supplement)	6	11.8

T&CM, traditional and complementary medicine.

TABLE 2. Multivariable Logistic Regression Model

	OR (CI)	P value
Sex (female)	1.158 (0.484-2.770)	.741
Age	1.032 (0.484-2.770)	.122
Residence (rural)	0.913 (0.247-3.378)	.892
Marital status (single)	1.623 (0.730-3.611)	.235
Employment status (no)	0.939 (0.420-2.087)	.878
Multiple sclerosis type (progressive)	2.540 (0.851-7.575)	.095
Number of relapses	0.963 (0.789-1.176)	.714
Physical pain	1.000 (0.244-4.097)	1.000
Treatment cost (expensive)	2.100 (0.176-25.010)	.557

Note: Significance tests were OR test, Student t test, and Fisher exact test.

We found that the rate of T&CM use is higher in patients with progressive MS and patients without financial access to DMT; in Morocco, 70.4% of patients do not have access to DMT.²⁰ However, analysis does not show a statistically significant relationship between the use of T&CM and the sociodemographic characteristics of age, sex, employment, place of residence, and marital status.

Our study's findings about which types of T&CM were most often used are supported by the findings of other studies. In our study, cupping was the most popular T&CM practice. A study from Saudi Arabia also found that 20.7% of individuals

with MS used cupping.¹⁶ In another study,²¹ wet cupping significantly improved the quality of life for a adult patients with chronic medical issues, particularly in the physical dimension, and athletes who received cupping therapy had lower levels of exercise-induced inflammatory markers. In Turkey, cupping was found among the T&CM practices most used by MS patients, followed by phytotherapy, osteopathy, ozone therapy, and music therapy.¹³

Our study found that the second most common T&CM used by individuals with MS was prayer, such as reciting the Holy Quran. This method was also employed in the Kingdom of Saudi Arabia and Iran.²² Ghavi and colleagues also showed that individuals with MS found spiritual therapy effective for treating stress, depression, and anxiety.²³

The third most common practice was phytotherapy; for some individuals with MS, medicinal plants and plant compounds can reduce neurologic deficits due to spasticity, muscle spasms, neuropathic pain, and urinary tract complications.²⁴ One study showed that phytochemicals present in the butanolic subextract of *Capparis ovata* extract could be beneficial in preventing/treating MS, in which neuroinflammation is part of the pathophysiology.²⁵ The study by Xie and colleagues found that curcumin (a spice derived from turmeric) was used by many people with MS for treating the symptoms of the disease.²⁶

Fourth was apitherapy, a practice that is used for autoimmune disorders such as systemic lupus erythematosus and MS.^{27,28} By modifying the release of cytokines, chemokines, and matrix-degrading enzymes that control inflammatory and immunological responses, Manuka honey has an anti-inflammatory impact.²⁹

Acupuncture and massage therapy were used by the same number of respondents. Other studies have reported that the use of acupuncture therapy ranges from 7.2% to 21% of people with MS,³⁰⁻³² and this has been suggested as a nonpharmacologic method of disease management. Additional studies show that acupuncture may reduce fatigue and decrease muscle spasms and pain, delaying the disease's evolution and reducing relapses.^{31,33} Various massage techniques, particularly Swedish methods, have been shown to improve MS-related impairments and reduce both motor and nonmotor MS symptoms such as fatigue, depression, anxiety, spasticity, and pain.³⁴⁻³⁶

The statistical analysis of the survey data shows no statistically significant relationship between the use of T&CM and age, sex, employment status, marital status, and number of relapses. Our data show that patients with progressive MS who are between 30 and 40 years old and who do not have financial access to DMT most commonly use T&CM. A study by Kim and colleagues showed that T&CM use by individuals with MS was related to sex, education level, DMT therapy, and MS disease progression.³⁷ Another study showed that people with MS in Denmark who were younger, female, educated at a bachelor's degree level or above, and had a high income tended to use T&CM more frequently. Respondents in another study said that the unreliability of conventional therapy, anecdotal evidence from other T&CM users, and medical advice all impacted their use of T&CM.¹⁵

Of the respondents, 66.7% were satisfied with T&CM. This is similar to the results of another study that showed that 65.6% of the participants were satisfied with T&CM.³⁸

Some limitations on the findings of this study are its small sample size that cannot be inclusive of the entirety of the people with MS in Morocco. Our use of the snowball method to recruit respondents may have predisposed our cohort toward the use of T&CM. The online survey also precluded us from having direct contact with the participants. Finally, Morocco has neither a national MS register nor regional registers for easy access to potential study participants.

CONCLUSIONS

This study shows that T&CM occupies an essential place in the treatment options used by individuals with MS in Morocco. These results invite us to consider societal, cultural, and economic factors when studying the use of T&CM among people with MS. Further research is needed to understand the motivations for using T&CM. ■

Funding/Support: None.

Prior Presentation: The abstract of this study was presented to the congress of the Middle East North Africa Committee for Research and Treatment in Multiple Sclerosis; November 11-12, 2022; Cairo, Egypt.

REFERENCES

1. WHO. *Traditional, Complementary and Integrative Medicine*. Accessed January 17, 2024. https://www.who.int/health-topics/traditional-complementary-and-integrative-medicine#tab=tab_1
2. Furhad S, Sina RE, Bokhari AA. Cupping therapy. *StatPearls*; 2023. Accessed May 9, 2023. <http://www.ncbi.nlm.nih.gov/books/NBK538253/>
3. Ghiasi A, Keramat A. The effect of listening to Holy Quran recitation on anxiety: a systematic review. *Iran J Nurs Midwifery Res*. 2018;23(6):411-420. doi:10.4103/ijnmr.IJNMR_173_17
4. Vickers A, Zollman C. ABC of complementary medicine. *Massage therapies*. *BMJ*. 1999;319(7219):1254-1257. doi:10.1136/bmj.319.7219.1254
5. Zhu J, Li J, Yang L, Liu S. Acupuncture, from the ancient to the current. *Anat Rec (Hoboken)*. 2021;304(11):2365-2371. doi:10.1002/ar.24625
6. Babar A, Al-Wabel NA, Shams S, Ahamad A, Khan SA, Anwar F. Essential oils used in aromatherapy: a systemic review. *Asian Pac J Trop Biomed*. 2015;5(8):601-611. doi:10.1016/j.apjtb.2015.05.007
7. Feuerstein G. *The Yoga Tradition: Its History, Literature, Philosophy and Practice*. SCB Distributors; 2012.
8. Hasanpour Dehkordi A. Influence of yoga and aerobics exercise on fatigue, pain and psychosocial status in patients with multiple sclerosis: a randomized trial. *J Sports Med Phys Fitness*. 2016;56(11):1417-1422.
9. Irish AK, Erickson CM, Wahls TL, Snetselaar LG, Darling WG. Randomized control trial evaluation of a modified paleolithic dietary intervention in the treatment of relapsing-remitting multiple sclerosis: a pilot study. *Degener Neurol Neuromuscul Dis*. 2017;7:1-18. doi:10.2147/DNND.S116949
10. Siev-Ner I, Gamus D, Lerner-Geva L, Achiron A. Reflexology treatment relieves symptoms of multiple sclerosis: a randomized controlled study. *Mult Scler*. 2003;9(4):356-361. doi:10.1191/1352458503ms9250a
11. Sirois FM, Gick ML. An investigation of the health beliefs and motivations of complementary medicine clients. *Soc Sci Med* 1982. 2002;55(6):1025-1037. doi:10.1016/S0277-9536(01)00229-5
12. Haut-Commissariat au Plan du Maroc. Note d'information du Haut-Commissariat au Plan relative aux principales caractéristiques de la population active occupée en 2021. January 24, 2022. Updated February 13, 2023. https://www.hcp.ma/Note-D-information-du-haut-commissariat-au-plan-relative-aux-principales-caracteristiques-de-la-population-active_a2825.html
13. Tekin C, Tecelioglu M, Sekerci G, Ozcan A. Traditional and complementary medicine use in multiple sclerosis: a cross-sectional study. *Ann Med Res*. 2022;29(9):1043-1048. <https://annalsmedres.org/index.php/aomr/article/view/4281>
14. Huybregts E, Betz W, Devroey D. The use of traditional and complementary medicine among patients with multiple sclerosis in Belgium. *J Med Life*. 2018;11(2):128-136.
15. Olsen SA. A review of complementary and alternative medicine (CAM) by people with multiple sclerosis. *Occup Ther Int*. 2009;16(1):57-70. doi:10.1002/oti.266
16. Shariff EM, Al-Shamrani FJ, Nazish S, et al. Is non-traditional therapy for multiple sclerosis overwhelming in Saudi Arabia. *Neurosciences (Riyadh)*. 2019;24(3):192-198. doi:10.17712/nsj.2019.3.20180010
17. Masullo L, Papas MA, Cotugna N, Baker S, Mahoney L, Trabulsi J. Complementary and alternative medicine use and nutrient intake among individuals with multiple sclerosis in the United States. *J Community Health*. 2015;40(1):153-160. doi:10.1007/s10900-014-9913-z
18. Campbell DG, Turner AP, Williams RM, et al. Complementary and alternative medicine use in veterans with multiple sclerosis: prevalence and demographic associations. *J Rehabil Res Dev*. 2006;43(1):99-110. doi:10.1682/jrrd.2004.08.0110
19. Skovgaard L, Nicolajsen PH, Pedersen E, et al. Use of complementary and alternative medicine among people with multiple sclerosis in the Nordic countries. *Autoimmune Dis*. 2012;2012:841085. doi:10.1155/2012/841085
20. Lotfi R, Chigr F, Najimi M. The impact of sociodemographic characteristics on diagnosis and access to disease-modifying treatment for patients with multiple sclerosis in Morocco. *Mult Scler Relat Disord*. 2022;67:104193. doi:10.1016/j.msard.2022.104193
21. Ekrami N, Ahmadian M, Nourshahi M, Shakouri GH. Wet-cupping induces anti-inflammatory action in response to vigorous exercise among martial arts athletes: a pilot study. *Complement Ther Med*. 2021;56:102611. doi:10.1016/j.ctim.2020.102611
22. Shayannejad V, Mohamadirizi S. Spiritual health in women with multiple sclerosis and its association with self-esteem. *J Educ Health Promot*. 2020;9(1):64. doi:10.4103/jehp.jehp_144_19
23. Ghavi F, Jahromi MK, Poorgholami F, Rezaee S. A study on the impact of spiritual therapy training on self-esteem in multiple sclerosis patients. *Pak J Med Health Sci*. 2020;14(1):483-486. https://www.researchgate.net/publication/342915128_A_Study_on_the_Impact_of_Spiritual_Therapy_Training_on_Self-Esteem_in_Multiple_Sclerosis_patients
24. Rabiee Z. Phytotherapy as a complementary medicine for multiple sclerosis. *Turk J Pharm Sci*. 2019;16(2):246-251. doi:10.4274/tjps.galenos.2018.90522
25. Sen A, Topcu G, Ozgun O, et al. Anti-neuroinflammatory effect of butanolic subextract of *Capparis ovata* water extract used as an alternative and complementary treatment for multiple sclerosis. *J Neuroimmunol*. 2014;275(1-2):172-173. doi:10.1016/j.jneuroim.2014.08.464
26. Xie L, Li XK, Takahara S. Curcumin has bright prospects for the treatment of multiple sclerosis. *Int Immunopharmacol*. 2011;11(3):323-330. doi:10.1016/j.intimp.2010.08.013
27. Fratellone PM, Tsimis F, Fratellone G. Apitherapy products for medicinal use. *J Altern Complement Med*. 2016;22(12):1020-1022. doi:10.1089/acm.2015.0346
28. Weis WA, Ripari N, Conte FL, et al. An overview about apitherapy and its clinical applications. *Phytomedicine Plus*. 2022;2(2):100239. doi:10.1016/j.phyplu.2022.100239
29. Minden-Birkenmaier BA, Meadows MB, Cherukuri K, et al. The effect of Manuka honey on dHL-60 cytokine, chemokine, and matrix-degrading enzyme release under inflammatory conditions. *Med One*. 2019;9(2):e190005. doi:10.20900/mo.20190005
30. Esmonde L, Long AF. Complementary therapy use by persons with multiple sclerosis: benefits and research priorities. *Complement Ther Clin Pract*. 2008;14(3):176-184. doi:10.1016/j.ctcp.2008.03.001
31. Karpatkin HI, Napolione D, Siminovich-Blok B. Acupuncture and multiple sclerosis: a review of the evidence. *Evid Based Complement Altern Med*. 2014;2014:972935. doi:10.1155/2014/972935
32. Stoll SS, Nieves C, Tabby DS, Schwartzman R. Use of therapies other than disease-modifying agents, including complementary and alternative medicine, by patients with multiple sclerosis: a survey study. *J Am Osteopath Assoc*. 2012;112(1):22-28.
33. Khodaie F, Abbasi N, Kazemi Mottagh AH, Zhao B, Naser Moghadasi A. Acupuncture for multiple sclerosis: a literature review. *Mult Scler Relat Disord*. 2022;60:103715. doi:10.1016/j.msard.2022.103715
34. Schroeder B, Doig J, Premkumar K. The effects of massage therapy on multiple sclerosis patients' quality of life and leg function. *Evid Based Complement Alternat Med*. 2014;2014. doi:10.1155/2014/640916
35. Backus D, Manella C, Bender A, Sweatman M. Impact of massage therapy on fatigue, pain, and spasticity in people with multiple sclerosis: a pilot study. *Int J Ther Massage Bodyw*. 2016;9(4):4. doi:10.3822/ijtm.b.914.327
36. Heidari Z, Shahrbanian S, Chiu C. Massage therapy as a complementary and alternative approach for people with multiple sclerosis: a systematic review. *Disabil Rehabil*. 2022;44(20):5758-5769. doi:10.1080/09638288.2021.1949051
37. Kim S, Chang L, Weinstock-Guttman B, et al. Complementary and alternative medicine usage by multiple sclerosis patients: results from a prospective clinical study. *J Altern Complement Med*. 2018;24(6):596-602. doi:10.1089/acm.2017.0268
38. Skovgaard L, Nicolajsen PH, Pedersen E, et al. Differences between users and non-users of complementary and alternative medicine among people with multiple sclerosis in Denmark: a comparison of descriptive characteristics. *Scand J Public Health*. 2013;41(5):492-499. doi:10.1177/140394813481646