

A Global Systematic Review of Lead (Pb) Exposure and its Health Effects in Wild Mammals

Helle B. Hydeskov,^{1,5} Jon M. Arnemo,^{2,3} Chris Lloyd Mills,⁴ Louise K. Gentle,¹ and Antonio Uzal¹

¹ School of Animal, Rural and Environmental Sciences, Nottingham Trent University, Brackenhurst Campus, Brackenhurst Lane, Southwell, NG25 0QF, UK

² Department of Forestry and Wildlife Management, Faculty of Applied Ecology, Agricultural Sciences and Biotechnology, Inland Norway University of Applied Sciences, Campus Evenstad, Anne Evenstads Vei 80, 2480 Koppang, Norway

³ Department of Wildlife, Fish, and Environmental Studies, Faculty of Forest Sciences, Swedish University of Agricultural Sciences, Skogsmarksgränd, 90736 Umeå, Sweden

⁴ School of Science and Technology, Nottingham Trent University, Clifton Campus, Clifton Lane, Nottingham, NG11 8NS, UK

⁵ Corresponding author (email: hbhvet@gmail.com)

ABSTRACT: Lead (Pb) is a toxic nonessential metal, known mainly for causing poisoning of humans and wild birds. However, little is known about Pb exposure and its associated health effects in wild mammals. We conducted a global systematic literature review to identify peer-reviewed studies published on Pb exposure in wild mammalian species and the health effects they identified. In total, 183 studies, conducted in 35 countries and published over 62 yr (1961–2022), were included in the review. Only 6% (11/183) of the studies were conducted in developing countries. Although 153 mammalian species were studied, most studies focused on species that are easy to access (i.e., hunted species and small mammals that are easy to trap). Therefore, carnivores and scavengers were less frequently studied than herbivores and omnivores. Despite all studies reporting Pb concentrations, only 45 (25%) studies investigated health effects and, of these 45 studies, only 28 (62%) found any health effect in 57 species. All health effects were negative and ranged from subclinical effects to fatality. Methodologies of Pb sampling and quantification and reporting of results varied widely across the studies, making both Pb concentrations and health effects difficult to compare and evaluate. Thus, there is a need for more research on Pb exposure and its health effects on wild mammals, especially as carnivores and scavengers could be used as sentinels for ecosystem health.

Key words: Bioaccumulation, Mammalia, One Health, sentinel species, toxicology, wildlife population health.

INTRODUCTION

Lead (Pb) is a toxic, nonessential metal that has no known biologic function for living organisms, contaminates the environment, and causes negative health effects in humans and other animals (Ma 2011; European Chemicals Agency 2018; Lanphear et al. 2018; Pain et al. 2019). It occurs naturally in the Earth's crust and has historically been used in many products, including water pipes, paints, gasoline, ammunition, aviation fuel, motor vehicle batteries, glassware, and cosmetics (Bergdahl and Skerfving 2022). These anthropogenic activities have increased the amount of environmental Pb to around 1,000 times the natural levels (Renberg et al. 2001), thereby amplifying the potential environmental exposure of wildlife to Pb. Despite legislation banning or reducing the use of Pb, it is still present in many products. For example, very few

countries, states, or regions have banned all Pb-based ammunition (Sonne et al. 2022), and according to the World Health Organization (WHO) as of June 2022, only 45% of countries had made legally binding controls on the production, import, sale, and use of Pb-based paints (WHO 2022a).

Children are generally more at risk of health effects from Pb exposure than adults, as children absorb more ingested Pb from the gastrointestinal system (WHO 2022b). Multiple organ systems, including the nervous, renal, cardiovascular, hematologic, immune, and reproductive systems, are negatively affected by Pb. The health effects of Pb exposure range from chronic and subclinical to acute and fatal (Agency for Toxic Substances and Disease Registry 2020; Bergdahl and Skerfving 2022). Furthermore, there is no known safe level of Pb exposure in humans (WHO 2022b). Despite this, concentrations of concern

vary between health authorities. For example, the Centers for Disease Control and Prevention (CDC) in the US currently have a “blood lead reference value (BLRV)” of 35 $\mu\text{g/L}$, which is used to identify children with high blood Pb concentrations (the 2.5% with the highest concentration) that require additional actions to be taken by health care professionals (CDC 2022). The European Food Safety Authority (EFSA) uses a different system and has set lower confidence limits of 12, 15, and 36 $\mu\text{g/L}$ as benchmark doses for developmental neurotoxicity, effects on the prevalence of chronic kidney disease, and effects on systolic blood pressure, respectively (EFSA Panel on Contaminants in the Food Chain 2013). Furthermore, concentrations of concern set by different authorities may vary over time. Between 2012 and 2021, the CDC’s BLRV was 50 $\mu\text{g/L}$; prior to 2012, it was 100 $\mu\text{g/L}$ and called the “level of concern in children” (CDC 2022). With the continued lowering of the CDC’s BLRV, it is clear that both health authorities agree that even very small concentrations of Pb can negatively affect human health, in alignment with the WHO’s understanding of Pb exposure (WHO 2022b).

Research on Pb exposure and health effects in wildlife has often been undertaken in avian species, where the first poisonings were recognized nearly 150 yr ago (Calvert 1876). In birds, Pb exposure often leads to multisystemic clinical disease or even death, especially in waterfowl, scavenging birds, and predatory birds (e.g., Pain et al. 2019). Consequently, Pb exposure has caused population declines in the scavenging California condor (*Gymnogyps californianus*), a scavenging species (Green et al. 2008). Although toxicity thresholds in different avian species were previously proposed (Franson and Pain 2011), scientists have recently suggested stopping this practice, as it gives the impression that some levels of Pb are acceptable (Pain et al. 2019).

Previous reviews on Pb in mammals have focused primarily on animals in laboratory settings or on domestic species. For example, Ma (2011) showed that humans and laboratory mammals have similar susceptibility to

Pb exposure (i.e., the same Pb dose–health effects relationship). In domestic mammals, young cattle appear most prone to Pb poisoning, probably due to the tendency to lick different items (Constable et al. 2017). However, young cattle may simply absorb Pb more efficiently than adults, as seen in humans (WHO 2022b).

There is a lack of a global systematic literature review focusing on Pb exposure and health effects in wild mammalian species. Synthesizing this scientific evidence is important to understand the breadth of wild mammalian species affected by Pb across the world, as there is no reason to expect wild mammals to be less at risk from Pb exposure than any other vertebrates, including humans.

This article was designed to provide a global systematic review of the literature on environmental Pb exposure and health effects in wild mammals, including assessments of gaps and biases in the research and suggestions on how to advance knowledge of the effects of Pb, particularly in identifying the types of species that may best indicate environmental Pb exposure. The outcomes of this review are important because they can be used to inform future studies, influence policymakers, and contribute toward relevant environmental legislation.

MATERIALS AND METHODS

Systematically identifying peer-reviewed studies to include

This systematic review followed the workflow proposed by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses 2020 statement (Page et al. 2021). A literature search for peer-reviewed scientific articles on Pb exposure in wild mammals was conducted using the database Scopus (Elsevier B.V, Amsterdam, Netherlands) on 21 June 2022. Scopus was chosen over other well-known databases and online search engines, such as Web of Science, PubMed, or Google Scholar due to the ability of Scopus to allow the chemical lead (Pb) as a search term, an option not provided by other databases or online search engines. In addition, Scopus permitted the exclusion of articles containing the word “lead” when used as a verb in the absence of any

occurrence of lead as a chemical in the source. Furthermore, Scopus only indexed peer-reviewed articles and did not have restrictions on the length of search queries.

In Scopus, different query combinations of the following searches were used: CASREGNUMBER (7439-92-1)/CHEMNAME (lead); wild/wildlife/free-living/“free living”/free-ranging/“free ranging”/semi-wild/“semi wild”/free-roaming/“free roaming”/street/feral; captive/zoo/domestic/domesticated/semi-captive/“semi captive”/semi-domestic/“semi domestic”/semi-domesticated/“semi domesticated”; exposure/expose/exposed/exposing/poisoning/poison/poisoned; keywords for mammals, and scientific nomenclature of all mammalian orders, families and genera taken from the International Union for the Conservation of Nature (IUCN) red list of threatened species (IUCN 2021; see Supplementary Material).

The first author conducted the assessment of all records and screening of studies. After identifying records via Scopus, records were checked for duplicates. Unique records were then screened by reading the title and abstract. The selected studies were thoroughly screened by reading the entire article. Articles on Pb exposure in wild mammals that were cited in the reference sections of the Scopus articles passing the full screening were also screened to identify additional publications not found by the original literature searches (i.e., snowball search).

Inclusion criteria included the requirement that studies must be peer-reviewed original research studies reporting Pb concentrations in tissues or body fluids in a nondomestic mammalian (called “wild mammals” in this review) species. Studies written in any language were included, and language was not used as an exclusion criterion. Studies reporting Pb concentrations in captive wild mammals outside laboratory settings were included to illustrate the variety of wild mammalian species documented to be susceptible to Pb exposure from their environment. Exclusion criteria included articles not reporting Pb concentrations in wild mammalian tissues or blood, studies using purely experimental settings, and studies reporting Pb concentrations in free-ranging domestic mammalian species. Data from studies were excluded if the study reported Pb concentration in tissues or body fluids knowingly contaminated with Pb-based ammunition (i.e., from killing the animal). Some studies reported nonstandard units (i.e., liquid concentrations for tissues or units unknown to the authors of this study).

In these cases or when there was a mismatch between the data reported in text and tables, the corresponding author was contacted via the email provided in the publication and given a maximum of 30 d to respond. Without a response, either the data in question or the entire study was excluded.

Data extracted from the studies

For every study included, the following information was extracted, where available: Taxonomy (order, family, common species name, and scientific species name); continent and country where the study was conducted; description of known pollution sources nearby the study site; type of living condition (free-ranging, captive, or semicaptive); year of sample collection; season; age; sex; tissue; sample size, Pb analysis method; reporting in wet weight or dry weight concentrations (when both were reported, the dry weight concentration was preferred); and Pb concentration arithmetic mean (geometric means and weighted arithmetic means were excluded), SD, SE, median, and minimum and maximum values. The Pb concentrations were converted to micrograms per kilogram or micrograms per liter, where applicable, for comparison. Whether the study had investigated and identified any health effects of Pb was also recorded. To investigate in which part of the mammalian food web Pb exposure is most frequently reported, the feeding type of the mammalian species (i.e., herbivore, omnivore, or carnivore) and whether or not the species scavenges on animal material was determined by following species descriptions available in a recently published mammalian textbook (Wilson and Mittermeier 2009–19). Articles from Russia were allocated to the European or Asia continent based on the geographic location of the sample collection. To investigate potential economic resource biases in the publications, the Organisation for Economic Co-operation and Development’s designation of developed and developing countries was used (Organisation for Economic Co-operation and Development 2022).

Different parts of a tissue or body fluid (e.g., whole kidney versus kidney cortex or whole blood versus plasma) were considered separate tissues for the purpose of the review. In addition, cortical bone and antler were differentiated, as cortical bone is permanent bone, whereas antler is deciduous bone (representing only short-term or annual exposure).




FIGURE 1. Modified Preferred Reporting Items for Systematic Reviews and Meta-Analyses 2020 flow diagram (Page et al. 2021) for systematic reviews, indicating the number of studies that were included and excluded in the screening process for studies on lead (Pb) exposure in wild mammals.

RESULTS

Characteristics of the studies

A total of 254 records were retrieved from Scopus (Fig. 1), with no duplicates identified. After the initial screening of the title and abstract, 125 studies remained for full screening, resulting in 69 studies from the Scopus search queries meeting the inclusion criteria. In addition, 114 studies were identified for inclusion through a snowball search, giving a total of 183 studies for systematic review (see Supplementary Material).

The studies were published over 62 yr (1961 to June 2022), with most of the studies (158/183, 86%) published in the second half

of this period (1992 to 2022; Fig. 2). The majority of studies (162/183, 89%) were conducted in Europe (108 studies, 59%) and North America (54 studies, 30%); some studies were conducted in multiple countries (but always within the same continent). The US was the country in which most studies were conducted (38 studies), followed by Canada (17 studies), Poland (16 studies), Germany (12 studies), and the UK (11 studies). Only 6% (11/183) of the studies were conducted in developing countries (Fig. 3).

Pb exposure

Pb was present in all mammalian species studied, with concentrations reported in a total




FIGURE 2. The number of studies published per year on lead (Pb) exposure in wild mammals included in this systematic review (n=183). Studies were published in the period 1961 to June 2022.




FIGURE 3. The number of studies ($n=183$) conducted on lead (Pb) exposure in wild mammals between 1961 and June 2022 per country. Studies were conducted in 35 countries across six continents. Darker color indicates a higher number of studies conducted. Created using Datawrapper (Datawrapper GmbH 2022).

of 153 wild mammalian species belonging to 45 families and 11 orders. The most studied order, family, and species were Cetartiodactyla, Cervidae, and red deer (*Cervus elaphus*), with 67, 46, and 20 studies, respectively (Table 1). In terms of feeding type, herbivores, omnivores,

TABLE 1. Top 5 most studied mammalian orders, families and species by the number of studies published on lead (Pb) exposure in wild mammals between 1961 and June 2022.

Order	No. of studies
Cetartiodactyla	67
Carnivora	55
Rodentia	42
Lagomorpha	15
Eulipotyphla	14
Family	No. of studies
Cervidae	46
Muridae	23
Ursidae	19
Suidae	17
Cricetidae	16
Species	No. of studies
Red deer (<i>Cervus elaphus</i>)	20
European roe deer (<i>Capreolus capreolus</i>)	19
Wild boar (<i>Sus scrofa</i>)	17
Long-tailed field mouse (<i>Apodemus sylvaticus</i>)	12
Brown bear (<i>Ursus arctos</i>)	11

and carnivores were researched in 92, 79, and 49 studies, respectively; 49 studies researched scavenging species.

A total of 33 different tissues and body fluids were tested. The five most studied tissues were the liver (123 studies), kidney (92 studies), skeletal muscle (54 studies), bone (32 studies), and blood (25 studies). At 1,506,000 $\mu\text{g}/\text{kg}$ dry weight, the highest Pb concentration reported across all 11 mammalian orders studied was recorded in the kidney of a Northern short-tailed shrew (*Blarina brevicauda*; Table 2).

Health effects of Pb

A total of 45 of 183 (25%) studies investigated the health effects of Pb. Of these, 28 (62%) studies identified a negative health effect in 57 species. No studies identified a positive health effect. Negative health effects were reported in 32 herbivores, 21 omnivores, and four carnivores and ranged in severity from “interference with developing permanent teeth” to “fatality” (Table 3).

Variation in methodology

The 183 studies used a large variety of sampling approaches and Pb analysis techniques and reported the results in different ways, making it hard to compare findings. For example, some studies only reported mean values, while others reported mean, SD or SE (or both),

median, and range of Pb concentration. Full details on sex and age of the animals tested and on whether the sampling was undertaken in an area with known Pb pollution, were not provided in 158 (86%), 134 (73%), and 101 (55%) studies, respectively.

DISCUSSION

Coverage of Pb exposure in wild mammals in the literature

The number of publications of Pb studies on wild mammalian species has increased over the last three decades. Nevertheless, the frequency of Pb publications on wild avian species still appears to be much higher, indicated by the large number of avian reviews published in recent years (e.g., Pain et al. 2019; Monclús et al. 2020; Ives et al. 2022).

The Pb concentrations were reported in 153 wild mammalian species, highlighting that this class of animals is exposed to Pb on a large scale. Despite this clear evidence of Pb exposure across many taxa, based on IUCN taxonomic classification, only 41% (11/27) of orders, 28% (45/162) of families, and 3% (153/5,968) of species were studied, which evidences a large group of mammals with no information on Pb exposure (IUCN 2021). Understandably, there has been a tendency to study species that are easiest to acquire (i.e., hunted species and small species that are easy to trap without chemical immobilization). Consequently, the studies identified in this review are skewed toward herbivores and omnivores. It is not possible to determine whether tissues were selected due to easy access or because the researcher hypothesized that the selected tissue would provide the most information in the study species. However, liver, kidney, skeletal muscle, bone, and blood are commonly collected and stored for further investigations during wildlife necropsies (McAloose et al. 2018).

Health effects of Pb in wild mammals

Only 28 studies, 62% of those in which health effects were investigated, identified

negative health effects of Pb in wild mammals. As expected, no positive effects were found. From the breadth of health effects identified and organ systems involved, it is clear that Pb affects many parts of the body in wild mammals, as in other species (Pain et al. 2019; Bergdahl and Skerfving 2022). Many of the identified health effects would most likely be subclinical, so it is not possible to determine how exactly these affect wild mammalian populations.

It is expected that due to feeding behaviors, some mammal species are more likely than some bird species to be exposed to smaller amounts of Pb, relative to body mass, but over extended periods of time. For example, waterfowl are known to pick up large numbers of Pb-based shotgun pellets or fishing weights from the bottom of wetland areas when foraging (Pain et al. 2019), yet most wild mammals would not be exposed to such large amounts of Pb at once, as they forage using different methods (e.g., Carnivora species). This difference in feeding behavior would reduce the risk of acute poisonings, seen in many waterfowl and scavenging and predatory bird species, but may cause more chronic and often subclinical health effects (e.g., on the nervous and immune systems). Chronic effects tend to be much less apparent than acute effects in free-ranging wildlife so may be overlooked unless investigated explicitly. Ecke et al. (2017) concluded that sublethal Pb exposure in golden eagles (*Aquila chrysaetos*) might impair flight performance and thus increase the risk of mortality from other causes. Yet, other eagle studies could not find evidence that subclinical Pb concentrations are linked to “increased risk of trauma” (Isomursu et al. 2018) or “other causes of mortality” (Franson and Russell 2014). These examples demonstrate the difficulty in investigating subclinical and chronic health effects in wildlife, especially when studies do not classify effects similarly. Furthermore, ethical permits to conduct Pb toxicological effect studies on wild mammals and other wildlife are, rightly, unlikely to be granted. Therefore, subclinical effects are difficult to confirm in wildlife.

TABLE 2. Highest individual (where $n=1$) or mean lead (Pb) concentration reported in each wild mammalian order studied in studies published between 1961 and June 2022. Solid tissue concentrations are reported in either wet weight (ww) or dry weight (dw).

Order	Concentration	Species	Sample size	Tissue	Location ^a	Country ^b	References
Carnivora	861,900 µg/kg ww	Ringed seal (<i>Pusa hispida</i>)	1	Brain	FR	Canada	(Chan et al. 1995)
Cetartiodactyla	97,300 µg/kg ww	Beluga whale (<i>Delphinapterus leucas</i>)	1	Blubber	FR	Canada	(Chan et al. 1995)
Chiroptera	500,000 µg/kg ww	Grey-headed flying fox (<i>Pteropus poliocephalus</i>)	2	Liver	Captive	US	(Zook et al. 1972)
Dasyuromorphia	62 µg/L	Tasmanian devil (<i>Sarcophilus harrisi</i>)	26	Blood	Captive	Australia	(Hivert et al. 2018)
Didelphimorphia	1,300 µg/kg ww	Virginia opossum (<i>Didelphis virginiana</i>)	4	Liver	FR	US	(Lewis et al. 2001)
Diprotodontia	2,950 µg/kg ^c	Common wombat (<i>Vombatus ursinus</i>)	4	Hair	FR	Australia	(Penrose et al. 2022)
Eulipotyphla	1,506,000 µg/kg dw	Northern short-tailed shrew (<i>Blarina brevicauda</i>)	1	Kidney	FR	US	(Stansley and Roscoe 1996)
Lagomorpha	9,000 µg/kg ww	European hare (<i>Lepus europaeus</i>)	9	Muscle	FR	Austria	(Ertl et al. 2016)
Primates	212,500 µg/kg ww	Stump-tailed macaque (<i>Macaca arctoides</i>)	2	Liver	Captive	US	(Zook et al. 1972)
Rodentia	672,000 µg/kg dw	Long-tailed field mouse (<i>Apodemus sylvaticus</i>)	5	Bone	FR	UK	(Johnson et al. 1978)
Sirenia	100 µg/L	American manatee (<i>Trichechus manatus</i>)	4	Blood	Captive	Brazil	(Anzolim et al. 2012)

^a FR = the wild mammal was from a free-ranging setting; captive: the wild mammal was from a captive setting.

^b Country in which the study was conducted.

^c No indication of whether the concentration is reported in ww or dw.

TABLE 3. Negative health effects associated with lead (Pb) exposure in wild mammals by organ system affected in studies published between 1961 and June 2022.

Organ or system affected	Effects found	References
Whole body	Fatality; DNA damage; oxidative stress	Sauer et al. 1970; Shlosberg et al. 1997; Skerratt et al. 1998; da Silva et al. 2000; Brumbaugh et al. 2010; Burco et al. 2012; North et al. 2015; Lazarus et al. 2020
Nervous system	Neurologic clinical signs; demyelination of spinal cord and subcortical white matter of brain and optic tracts	Sauer et al. 1970; Zook et al. 1973; Diters and Nielsen 1978; Skerratt et al. 1998
Circulatory system	Decreased aminolevulinic acid dehydratase activity; decreased hemoglobin; decreased hematocrit; increased micronucleated erythrocytes; increased total bilirubin; decreased calcium; decreased lactate dehydrogenase	Mouw et al. 1975; Ieradi et al. 1996; Stansley and Roscoe 1996; Rogival et al. 2006; Sánchez-Chardi et al. 2009; Chen et al. 2018
Reproductive system	Testicular structural and functional disruption; abnormal sperm cells; decreased proportion of live sperm	Ieradi et al. 1996; Parelho et al. 2016; Chen et al. 2018
Kidney	Decreased aminolevulinic acid dehydratase activity; necrosis and cell degeneration; eosinophilic acid-fast intranuclear inclusion bodies in proximal tubular epithelium; edema; mitochondrial abnormalities; increased weight of kidneys; hyperplasia of tubules, atrophy of glomeruli, interstitial fibrosis; degenerate cells in the lumen of seminiferous tubules; mineralization and degeneration of proximal tubules; karyocytomegaly; lymphocyte infiltrations; cell debris and change in glomerular aspect and size	Sauer et al. 1970; Zook et al. 1970, 1972; Mouw et al. 1975; Diters and Nielsen 1978; Roberts et al. 1978; Stansley and Roscoe 1996; Skerratt et al. 1998; Ceruti et al. 2002; Damek-Poprawa and Sawicka-Kapusta 2003; Brumbaugh et al. 2010; Tête et al. 2014
Liver	Eosinophilic acid-fast intranuclear inclusion bodies; necrosis; apoptosis; increased number of pyknotic nuclei; interstitial fibrosis, decreased glycogen content; increased liver-body ratio; steatosis; lymphocyte infiltrations	Sauer et al. 1970; Zook et al. 1970, 1972; Diters and Nielsen 1978; Damek-Poprawa and Sawicka-Kapusta 2003; Sánchez-Chardi et al. 2009; Tête et al. 2014
Teeth	Interference with developing permanent teeth	Rappaport et al. 1975; North et al. 2015

Although the impacts of Pb exposure on human, laboratory mammalian, and wild avian health are well-documented, there is still much to learn about the impacts on wild mammal health. In contrast to important infectious diseases in animals, the World Organisation for Animal Health does not have any reference laboratories for any noninfectious diseases such as Pb poisoning, which could guide or

help standardize how the diagnostic steps should be conducted (World Organisation for Animal Health 2022). Potential strategies to research the health effects of Pb on wild mammals include behavioral studies, reproductive success analyses, and immune system function assessments; these are all challenging to conduct in free-ranging animals. Research that would be easier to achieve in

free-ranging settings includes correlations between blood Pb concentrations and hematologic and biochemical parameters, investigations of blood and tissue Pb concentrations to understand the distribution of Pb in the body, and investigations of tissue pathologies in Pb-exposed populations. Ideally, studies should use the most sensitive techniques available for measuring Pb, such as inductively coupled plasma mass spectrometry (Apostoli 2002) and report the results in dry weight for easier comparisons. Regardless of the approach taken by future studies to investigate Pb exposure and health effects in wild mammals, studies should report details of the study area, sex and age of animals, and analytic methods, as well as providing an appropriate range of descriptive statistics (e.g., sample sizes, SD or SE), which would facilitate future evidence-based policies.

Methodologies, kinetics, and thresholds

Our systematic review found a lack of consistency in estimating Pb concentrations, health effects, and how this is reported. The different Pb analytic methods used in the literature reviewed (e.g., atomic absorption spectroscopy and inductively coupled plasma mass spectrometry) vary in sensitivity and differ in susceptibility to matrix effects (Apostoli 2002). Therefore, some studies might have failed to detect Pb, if present in low concentrations. In addition, some studies reported tissue concentrations by dry weight and others by wet weight, impeding direct comparison of such concentrations. Furthermore, Pb uptake and storage distribution potentially may differ between species: As there are no studies available on Pb kinetics in wild mammals, this remains unknown. Variations in methodologies and target species make it difficult to compare Pb concentrations between studies and between species. Consequently, it is not possible to establish toxicity thresholds for wild mammals based on the current body of literature. However, although thresholds may be a valuable tool for some professions (e.g., veterinarians in clinical practice), a pragmatic reason for not attempting to establish threshold values for different wild mammalian

tissues or species is the risk of creating the false impression of acceptable or safe levels of Pb exposure in wild mammals. Therefore, we recommend ceasing the use of threshold values for Pb concentrations in wildlife research, as already suggested by Pain et al. (2019).

Designing future studies on Pb exposure and health effects

From a One Health perspective (FAO et al. 2022), human, animal and ecosystem health are interlinked. Although it would be beneficial to study Pb concentrations in species at differing trophic levels in the wild, the cost of periodic screening programs means that these are likely to be restricted to only a few selected species. Because Pb bioaccumulates in both plants and animals (Clemens 2006; Radomski et al. 2018), testing for Pb in carnivorous and scavenging mammalian species might not only aid monitoring of environmental Pb exposure but also indicate the potential exposure of humans to Pb, as species at these higher trophic levels often have long life spans similar to humans (O'Brien et al. 1993). Using higher trophic-level species as sentinels is already being used in avian Pb research (e.g., Monclús et al. 2020) and in mammalian research on infectious diseases and antimicrobial resistance (e.g., Millán et al. 2014; Sacristán et al. 2020). Another benefit of studying top predatory or scavenging mammals, which are often iconic and charismatic (Albert et al. 2018), is that these mammals might attract more attention from stakeholders and policymakers involved in the regulation of Pb, a phenomenon already seen in conservation, with charismatic species receiving the most attention (Sitas et al. 2009).

To assess or monitor short-term and recent Pb exposure using reduced sampling effort and noninvasive sampling, antlers or hairs might be suitable tissues to consider in the future. Antlers represent Pb exposure of about 6 mo, from spring to autumn, and have previously been used to assess Pb contamination (Ludolph et al. 2022). Antlers do, however, introduce potential biases into a Pb monitoring program. For most cervid species, antlers are only available from

adult and subadult males; thus, detecting differences between age classes and sexes would not be possible using this tissue. Hair is often easily available even from museum specimens (i.e., on pelts), and it can be collected from free-ranging mammals noninvasively using hair traps. The period of Pb exposure can be estimated using laser ablation techniques, provided that potential Pb surface contamination is removed, the particular species' hair growth cycle is known, and the sampling is carried out in a consistent manner (Sela et al. 2007).

To gain information on where Pb is stored in a particular wild mammalian species, researchers would have to analyze multiple tissues and blood, all obtained from the same individual at the same time (Krone 2018; Bergdahl and Skerfving 2022). However, obtaining this quantity of samples would likely only be possible in freshly dead or mildly autolyzed individuals (McAloose et al. 2018). Also, extra expenses from analyzing multiple samples from each individual could be cost prohibitive. Therefore, more controlled studies on the storage and clinical pathology of Pb in captive individuals could help understand the health effects of Pb in wild mammals.

With studies written in any language being included, this review highlights that few studies on Pb exposure in wild mammals have been conducted in developing countries. However, note that detection of studies, in general, always depends on how journals have indexed articles in various databases; therefore, it cannot be ruled out that studies written in other languages than English will be missed. Moreover, it is important to consider that fewer studies published in developing countries does not mean the problem is less prevalent in these countries. Indeed, the lack of studies could be explained by lower access to research funding (Kpokiri et al. 2022). On the contrary, environmental exposure to Pb might be greater in developing countries due to fewer legislative controls on Pb use in products, as seen with Pb-based paints (WHO 2022a). Indeed, children in developing countries are more exposed to Pb than

children in developed countries (Hwang et al. 2019; Bergdahl and Skerfving 2022). Therefore, more research on Pb exposure in wild mammals must be conducted and published in these countries to understand the levels of environmental exposure to Pb, the risk to human and animal health, and to study whether legislation on Pb in products affects the Pb present in the environment. understand the levels of environmental exposure to Pb, the risk to human and animal health, and to study whether legislation on Pb in products affects the Pb present in the environment.

As all organisms are exposed to Pb, studies helping to determine the impacts of Pb on different species will add to current knowledge of this toxic metal. Collectively, this increase in knowledge will create stronger evidence to influence policymakers involved in the regulation of Pb in products that continue to act as sources of environmental contamination and be a major One Health issue. Overall, Pb exposure in wild mammalian species must be recognized as an issue, as this systematic review demonstrates. Control and regulation of Pb need to take a One Health approach and not focus on single taxa groups when developing policies.

ACKNOWLEDGMENT

We thank Victoria Boskett (research librarian at Nottingham Trent University) for help with journal indexing and database search engines.

SUPPLEMENTARY MATERIAL

Supplementary material for this article is online at <http://dx.doi.org/10.7589/JWD-D-23-00055>.

LITERATURE CITED

- Agency for Toxic Substances and Disease Registry. 2020. *Toxicological profile for lead*. <https://www.atsdr.cdc.gov/toxprofiles/tp.asp?id=96&tid=22>. Accessed November 2022.
- Albert C, Luque GM, Courchamp F. 2018. The twenty most charismatic species. *PLoS One* 13:e0199149.
- Anzolin DG, Sarkis JES, Diaz E, Soares DC, Serrano IL, Borges JCG, Souto AS, Taniguchi S, Montone RC,

- et al. 2012. Contaminant concentrations, biochemical and hematological biomarkers in blood of West Indian manatees *Trichechus manatus* from Brazil. *Mar Pollut Bull* 64:1402–1408.
- Apostoli P. 2002. Elements in environmental and occupational medicine. *J Chromatogr B Analyt Technol Biomed Life Sci* 778:63–97.
- Bergdahl IA, Skerfving S. 2022. Lead. In: *Handbook on the toxicology of metals: Specific metals*, 5th Ed., Vol. 2, Nordberg GF, Costa M, editors. Academic Press, London, UK, pp. 427–493.
- Brumbaugh WG, Mora MA, May TW, Phalen DN. 2010. Metal exposure and effects in voles and small birds near a mining haul road in Cape Krusenstern National Monument, Alaska. *Environ Monit Assess* 170:73–86.
- Burco J, Myers AM, Schuler K, Gillin C. 2012. Acute lead toxicosis via ingestion of spent ammunition in a free-ranging cougar (*Puma concolor*). *J Wildl Dis* 48:216–219.
- Calvert HS. 1876. Pheasants poisoned by swallowing shot. *The Field* 47:189.
- CDC (Centers for Disease Control and Prevention). 2022. Blood lead reference value. <https://www.cdc.gov/nceh/lead/data/blood-lead-reference-value.htm>. Accessed November 2022.
- Ceruti R, Ghisleni G, Ferretti E, Cammarata S, Sonzogni O, Scanziani E. 2002. Wild rats as monitors of environmental lead contamination in the urban area of Milan, Italy. *Environ Pollut* 117:255–259.
- Chan HM, Kim C, Khoday K, Receveur O, Kuhnlein HV. 1995. Assessment of dietary exposure to trace metals in Baffin Inuit food. *Environ Health Perspect* 103:740–746.
- Chen YP, Liu Q, Ma QY, Maltby L, Ellison AM, Zhao Y. 2018. Environmental toxicants impair liver and kidney function and sperm quality of captive pandas. *Ecotoxicol Environ Saf* 162:218–224.
- Clemens S. 2006. Toxic metal accumulation, responses to exposure and mechanisms of tolerance in plants. *Biochimie* 88:1707–1719.
- Constable PD, Hinchcliff KW, Done SH, Grünberg W. 2017. Diseases of the nervous system. In: *Veterinary medicine: A textbook of the diseases of cattle, horses, sheep, pigs and goats*, 11th Ed., Constable PD, Hinchcliff KW, Done SH, Grünberg W, editors. Elsevier, St. Louis, Missouri, pp. 1155–1370.
- Damek-Poprawa M, Sawicka-Kapusta K. 2003. Damage to the liver, kidney, and testis with reference to burden of heavy metals in yellow-necked mice from areas around steelworks and zinc smelters in Poland. *Toxicology* 186:1–10.
- da Silva J, de Freitas TRO, Heuser V, Marinho JR, Bittencourt F, Cerski CTS, Kliemann LM, Erdtmann B. 2000. Effects of chronic exposure to coal in wild rodents (*Ctenomys torquatus*) evaluated by multiple methods and tissues. *Mutat Res Genet Toxicol Environ Mutagen* 470:39–51.
- Datwrapperr GmbH. 2022. *Datwrapperr*. <https://www.datwrapperr.de/>. Accessed November 2022.
- Diters RW, Nielsen SW. 1978. Lead poisoning of raccoons in Connecticut. *J Wildl Dis* 14:187–192.
- Ecke F, Singh NJ, Arnemo JM, Bignert A, Helander B, Berglund AMM, Borg H, Bröjer C, Holm K, et al. 2017. Sublethal lead exposure alters movement behavior in free-ranging golden eagles. *Environ Sci Technol* 51:5729–5736.
- EFSA Panel on Contaminants in the Food Chain. 2013. Scientific opinion on lead in food. *EFSA J* 8:1570.
- Ertl K, Kitzer R, Goessler W. 2016. Elemental composition of game meat from Austria. *Food Addit Contam Part B Surveill* 9:120–126.
- European Chemicals Agency. 2018. *Annex XV investigation report: A review of the available information on lead in shot used in terrestrial environments, in ammunition and in fishing tackle*. European Chemicals Agency, Helsinki, Finland, 100 pp.
- FAO, UNEP, WHO, & WOA. 2022. One Health Joint Plan of Action (2022–2026). *Working together for the health of humans, animals, plants and the environment*. FAO; UNEP; WHO; World Organisation for Animal Health (WOAH) (founded as OIE), Rome, Italy, 86 pp. <http://doi.org/10.4060/cc2289en>.
- Franson JC, Pain DJ. 2011. Lead in birds. In: *Environmental contaminants in biota: Interpreting tissue concentrations*, 2nd Ed., Beyer WN, Meador JP, editors. CRC Press, Boca Raton, Florida, pp. 563–593.
- Franson JC, Russell RE. 2014. Lead and eagles: Demographic and pathological characteristics of poisoning, and exposure levels associated with other causes of mortality. *Ecotoxicology* 23:1722–1731.
- Green RE, Hunt WG, Parish CN, Newton I. 2008. Effectiveness of action to reduce exposure of free-ranging California condors in Arizona and Utah to lead from spent ammunition. *PLoS One* 3:e4022.
- Hivert LG, Clarke JR, Peck SJ, Lawrence C, Brown WE, Huxtable SJ, Schaap D, Pemberton D, Grueber CE. 2018. High blood lead concentrations in captive Tasmanian devils (*Sarcophilus harrisi*): A threat to the conservation of the species? *Aust Vet J* 96:442–449.
- Hwang YH, Hsiao CK, Lin PW. 2019. Globally temporal transitions of blood lead levels of preschool children across countries of different categories of Human Development Index. *Sci Total Environ* 659:1395–1402.
- Ieradi LA, Cristaldi M, Mascanzoni D, Cardarelli E, Grossi R, Campanella L. 1996. Genetic damage in urban mice exposed to traffic pollution. *Environ Pollut* 92:323–328.
- Isomursu M, Koivusaari J, Stjernberg T, Hirvelä-Koski V, Venäläinen ER. 2018. Lead poisoning and other human-related factors cause significant mortality in white-tailed eagles. *Ambio* 47:858–868.
- IUCN (International Union for the Conservation of Nature). 2021. *The IUCN red list of threatened species*. Version 2021–3. <https://www.iucnredlist.org>. Accessed June 2022.
- Ives AM, Brenn-White M, Buckley JY, Kendall CJ, Wilton S, Deem SL. 2022. A global review of causes of morbidity and mortality in free-living vultures. *Ecohealth* 19:40–54.
- Johnson MS, Roberts RD, Hutton M, Inskip MJ. 1978. Distribution of lead, zinc and cadmium in small mammals from polluted environments. *Oikos* 30:153–159.
- Kpokiri EE, Sri-Pathmanathan C, Shrestha P, Navaid S, Wivatpanit T, Wijegunawardana A, Umar MA, Jackson D, Alger J, et al. 2022. Crowdfunding for health

- research: A qualitative evidence synthesis and a pilot programme. *BMJ Glob Health* 7:e009110.
- Krone O. 2018. Lead poisoning in birds of prey. In: *Birds of prey: Biology and conservation in the XXI century*, Sarasola JH, Grande JM, Negro JJ, editors. Springer International Publishing, Cham, Switzerland, pp. 251–272.
- Lanphear BP, Rauch S, Auinger P, Allen RW, Hornung RW. 2018. Low-level lead exposure and mortality in US adults: A population-based cohort study. *Lancet Public Health* 3:e177–e184.
- Lazarus M, Orcz T, Sergiel A, Vranković L, Marijić VF, Rašić D, Reljić S, Aladrović J, Zwijacz-Kozica T, et al. 2020. Metal(loid) exposure assessment and biomarker responses in captive and free-ranging European brown bear (*Ursus arctos*). *Environ Res* 183:109166.
- Lewis LA, Poppenga RJ, Davidson WR, Fischer JR, Morgan KA. 2001. Lead toxicosis and trace element levels in wild birds and mammals at a firearms training facility. *Arch Environ Contam Toxicol* 41:208–214.
- Ludolph C, Kierdorf U, Kierdorf H. 2022. Antlers of European roe deer (*Capreolus capreolus*) as monitoring units to assess lead pollution in a floodplain contaminated by historical metal ore mining, processing, and smelting in the Harz Mountains, Germany. *Environ Pollut* 302:119080.
- Ma WC. 2011. Lead in mammals. In: *Environmental contaminants in biota: Interpreting tissue concentrations*, 2nd Ed., Beyer WN, Meador JP, editors. CRC Press, Boca Raton, Florida, pp. 595–608.
- McAloose D, Colegrove KM, Newton AL. 2018. Wildlife necropsy. In: *Pathology of wildlife and zoo animals*, Terio KA, McAloose D, St. Leger J, editors. Academic Press, London, UK, pp. 1–20.
- Millán J, García EJ, Oleaga Á, López-Bao JV, Llana L, Palacios V, Candela MG, Cevidanes A, Rodríguez A, León-Vizcaíno L. 2014. Using a top predator as a sentinel for environmental contamination with pathogenic bacteria: The Iberian wolf and leptospires. *Mem Inst Oswaldo Cruz* 109:1041–1044.
- Monclús L, Shore RF, Krone O. 2020. Lead contamination in raptors in Europe: A systematic review and meta-analysis. *Sci Total Environ* 748:141437.
- Mouw D, Kalitis K, Anver M, Schwartz J, Constan A, Hartung R, Cohen B, Ringler D. 1975. Lead. Possible toxicity in urban vs rural rats. *Arch Environ Health* 30:276–280.
- North MA, Lane EP, Marnewick K, Caldwell P, Carlisle G, Hoffman LC. 2015. Suspected lead poisoning in two captive cheetahs (*Acinonyx jubatus jubatus*) in South Africa, in 2008 and 2013. *J S Afr Vet Assoc* 86:1286.
- O'Brien DJ, Kaneene JB, Poppenga RH. 1993. The use of mammals as sentinels for human exposure to toxic contaminants in the environment. *Environ Health Perspect* 99:351–368.
- Organisation for Economic Co-operation and Development. 2022. *DAC List of ODA recipients. Effective for reporting on 2022 and 2023 flows*. <https://web-archiv.oecd.org/2022-11-07/616607-DAC-List-of-ODA-Recipients-for-reporting-2022-23-flows.pdf>. Accessed November 2022.
- Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, et al. 2021. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *BMJ* 372:n71.
- Pain DJ, Mateo R, Green RE. 2019. Effects of lead from ammunition on birds and other wildlife: A review and update. *Ambio* 48:935–953.
- Parelho C, Bernardo F, Camarinho R, Rodrigues AS, Garcia P. 2016. Testicular damage and farming environments—An integrative ecotoxicological link. *Chemosphere* 155:135–141.
- Penrose B, MacIntosh AE, Parbhakar-Fox A, Smith LBE, Sawyer T, D'Agnesse E, Carver S. 2022. Heavy metal wombats? Metal exposure pathways to bare-nosed wombats (*Vombatus ursinus*) living on remediated tin mine tailings. *Sci Total Environ* 835:155526.
- Radomyski A, Lei K, Giubilato E, Critto A, Lin C, Marcomini A. 2018. Bioaccumulation of trace metals in aquatic food web. A case study, Liaodong Bay, NE China. *Mar Pollut Bull* 137:555–565.
- Rappaport SC, Strebel RF, Garner E, Craston D. 1975. Lead insult to the dentition in the leopard. *Pharmacol Ther Dent* 2:201–207.
- Renberg I, Bindler R, Brännvall ML. 2001. Using the historical atmospheric lead-deposition record as a chronological marker in sediment deposits in Europe. *Holocene* 11:511–516.
- Roberts RD, Johnson MS, Hutton M. 1978. Lead contamination of small mammals from abandoned metalliferous mines. *Environ Pollut* 15:61–69.
- Rogival D, Scheirs J, De Coen W, Verhagen R, Blust R. 2006. Metal blood levels and hematological characteristics in wood mice (*Apodemus sylvaticus* L.) along a metal pollution gradient. *Environ Toxicol Chem* 25:149–157.
- Sacristán I, Esperón F, Acuña F, Aguilar E, García S, López MJ, Cevidanes A, Neves E, Cabello J, et al. 2020. Antibiotic resistance genes as landscape anthropization indicators: Using a wild felid as sentinel in Chile. *Sci Total Environ* 703:134900.
- Sánchez-Chardi A, Oliveira Ribeiro CA, Nadal J. 2009. Metals in liver and kidneys and the effects of chronic exposure to pyrite mine pollution in the shrew *Crocidura russula* inhabiting the protected wetland of Doñana. *Chemosphere* 76:387–394.
- Sauer RM, Zook BC, Garner FM. 1970. Demyelinating encephalomyelopathy associated with lead poisoning in nonhuman primates. *Science* 169:1091–1093.
- Sela H, Karpas Z, Zoriy M, Pickhardt C, Becker JS. 2007. Biomonitoring of hair samples by laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS). *Int J Mass Spectrom* 261:199–207.
- Shlosberg A, Bellaïche M, Regev S, Gal R, Brizzi M, Hanji V, Zaidel L, Nyska A. 1997. Lead toxicosis in a captive bottlenose dolphin (*Tursiops truncatus*) consequent to ingestion of air gun pellets. *J Wildl Dis* 33:135–139.
- Sitas N, Baillie JEM, Isaac NJB. 2009. What are we saving? Developing a standardized approach for conservation action. *Anim Conserv* 12:231–237.
- Skerratt LF, Speare R, Berger L, Winsor H. 1998. Lyssaviral infection and lead poisoning in black flying foxes from Queensland. *J Wildl Dis* 34:355–361.

- Sonne C, Adams DH, Alstrup AKO, Lam SS, Dietz R, Kanstrup N. 2022. Denmark passes total ban of leaded ammunition. *Science* 377:1054–1055.
- Stansley W, Roscoe DE. 1996. The uptake and effects of lead in small mammals and frogs at a trap and skeet range. *Arch Environ Contam Toxicol* 30:220–226.
- Tête N, Durfort M, Rieffel D, Scheifler R, Sánchez-Chardi A. 2014. Histopathology related to cadmium and lead bioaccumulation in chronically exposed wood mice, *Apodemus sylvaticus*, around a former smelter. *Sci Total Environ* 481:167–177.
- Wilson DE, Mittermeier RA, editors. 2009–2019. *Handbook of the mammals of the world*. 9 Vol. Lynx Edicions, Barcelona, Spain, 7693 pp.
- WHO (World Health Organization). 2022a. *Legally-binding controls on lead paint*. <https://www.who.int/data/gho/data/themes/topics/indicator-groups/legally-binding-controls-on-lead-paint>. Accessed November 2022.
- WHO. 2022b. *Lead poisoning*. <https://www.who.int/news-room/fact-sheets/detail/lead-poisoning-and-health>. Accessed November 2022.
- World Organisation for Animal Health. 2022. *Reference laboratories*. <https://www.woah.org/en/what-we-offer/expertise-network/reference-laboratories/#ui-id-3>. Accessed November 2022.
- Zook BC, Eisenberg JF, McLanahan E. 1973. Some factors affecting the occurrence of lead poisoning in captive primates. *J Med Primatol* 2:206–217.
- Zook BC, Sauer RM, Garner FM. 1970. Lead poisoning in Australian fruit bats (*Pteropus poliocephalus*). *J Am Vet Med Assoc* 157:691–694.
- Zook BC, Sauer RM, Garner FM. 1972. Lead poisoning in captive wild animals. *J Wildl Dis* 8:264–272.

Submitted for publication 22 March 2023.

Accepted 25 October 2023.