

# Association Between Ocular Biomechanics Measured With Corvis ST and Glaucoma Severity in Patients With Untreated Primary Open Angle Glaucoma

Na Wu<sup>1,2</sup>, Yuhong Chen<sup>1,2</sup>, and Xinghuai Sun<sup>1-3</sup>

<sup>1</sup> Department of Ophthalmology and Visual Science, Eye and ENT Hospital, Shanghai Medical College, Fudan University, Shanghai, China

<sup>2</sup> NHC Key Laboratory of Myopia (Fudan University), Key Laboratory of Myopia, Chinese Academy of Medical Sciences, and Shanghai Key Laboratory of Visual Impairment and Restoration (Fudan University), Shanghai, China

<sup>3</sup> State Key Laboratory of Medical Neurobiology, Institutes of Brain Science and Collaborative Innovation Center for Brain Science, Fudan University, Shanghai, China

**Correspondence:** Yuhong Chen, Department of Ophthalmology, Eye and ENT Hospital of Fudan University, 83 Fenyang Road, Shanghai 200031, China. e-mail: [yuhongchen@fudan.edu.cn](mailto:yuhongchen@fudan.edu.cn)

**Received:** November 3, 2021

**Accepted:** May 20, 2022

**Published:** June 9, 2022

**Keywords:** ocular biomechanics; glaucoma severity; Corvis ST

**Citation:** Wu N, Chen Y, Sun X. Association between ocular biomechanics measured with corvis ST and glaucoma severity in patients with untreated primary open angle glaucoma. *Transl Vis Sci Technol.* 2022;11(6):10, <https://doi.org/10.1167/tvst.11.6.10>

**Purpose:** To compare the ocular biomechanical differences between normal controls and patients with untreated primary open angle glaucoma, including normal-tension glaucoma (NTG) and high-tension glaucoma (HTG), and to investigate the association between ocular biomechanics and glaucoma severity in each group.

**Methods:** One hundred fifty-three eyes of 153 subjects, including 51 controls, 47 NTG, and 55 HTG cases, were enrolled in this cross-sectional study. Each participant underwent biomechanical measurements by using the Corneal Visualization Scheimpflug Technology. Glaucoma severity was evaluated by mean deviation (MD), pattern standard deviation (PSD), ganglion cell complex (GCC), and retinal nerve fiber layer (RNFL) thickness.

**Results:** Deformation amplitude ( $P = 0.001$ ) significantly increased, whereas first applanation time ( $P < 0.0001$ ), highest concavity time ( $P = 0.001$ ), stiffness parameter at first applanation ( $P = 0.009$ ), and time of whole eye movement (WEM,  $P = 0.008$ ) decreased significantly in NTG eyes compared with controls. Besides, NTG had the highest first applanation velocity than controls ( $P < 0.0001$ ) and HTG ( $P = 0.044$ ). Shorter time of WEM was independently correlated with worse MD ( $P = 0.02$ ) and higher values of PSD ( $P = 0.03$ ) in NTG. Axial length was positively related to PSD ( $P = 0.02$ ) and negatively related to GCC ( $P < 0.0001$ ) and RNFL ( $P < 0.0001$ ) thickness in HTG.

**Conclusions:** NTG corneas are more deformable than healthy ones and HTG. Time of WEM, which relates to orbital compliance, is significantly associated with glaucomatous visual field defect in NTG, whereas axial length is correlated with glaucoma severity in HTG.

**Translational Relevance:** Ocular biomechanics may partly account for the differences of pathogenic mechanisms between NTG and HTG.

## Introduction

Glaucoma is a leading cause of irreversible visual impairment and blindness worldwide.<sup>1</sup> It is accompanied by retinal nerve fibers loss and optic disc excavation, eventually leading to visual field (VF) defects. Intraocular pressure (IOP) is the most significant risk factor for glaucoma development and progression.<sup>2,3</sup>

Corneal biomechanical properties are of increasing interest in glaucoma because of their influences on corneal resistance to applanation and therefore IOP measurement obtained by Goldmann Applanation Tonometry.<sup>4,5</sup> In addition, it has been reported that corneal biomechanics is associated with optic nerve surface compliance during transient IOP elevation.<sup>6</sup> Corneal hysteresis (CH), measured with the ocular response analyzer, is a corneal biomechanical

parameter related to viscoelastic dampening.<sup>7</sup> CH is found to be lower in glaucomatous compared to healthy eyes<sup>8</sup> and is a risk factor for VF progression.<sup>9,10</sup>

Most of the available studies about the relationship between corneal biomechanics and glaucoma severity were conducted using the ocular response analyzer. CH is reported to be positively correlated with both of visual field index and mean deviation (MD) in primary open angle glaucoma (POAG) patients.<sup>11,12</sup> In addition, lower CH is associated with worse VF eyes in asymmetric POAG patients, which is independent of its effect on IOP measurements.<sup>7</sup> However, Hirneiss et al.<sup>13</sup> reported there was no difference of corneal biomechanics in both eyes of patients with unilateral POAG when CH was corrected for IOP. These results indicate that the relationship between corneal biomechanics and glaucoma severity is controversial and needs to be studied further.

The Corneal Visualization Scheimpflug Technology (Corvis ST) is another non-contact device that has the advantage of dynamic cross-sectional imaging during the corneal deformation, which give additional information about corneal biomechanics.<sup>14</sup> Only a few studies reported the relationship between corneal biomechanics measured by Corvis ST and glaucoma severity.<sup>15,16</sup> However, these studies did not exclude the influence of antiglaucomatous medications on corneal biomechanics, which has been demonstrated by several studies.<sup>17–19</sup>

Normal-tension glaucoma (NTG) and high-tension glaucoma (HTG) are two subtypes of POAG. IOP always remains within normal range in NTG, whereas elevated IOP is presented in HTG. Different pathogeneses may exist between NTG and HTG.<sup>20</sup> Previous literature had reported there were differences in ocular biomechanics between NTG and HTG. The corneas were significantly softer and more deformable in medically controlled NTG patients than those in HTG.<sup>16,21</sup> However, whether ocular biomechanics contribute differently to glaucomatous damage between NTG and HTG are still unknown. Therefore the aim of the current study was to investigate the correlation between Corvis ST measured ocular biomechanics and glaucoma severity in untreated NTG and HTG patients and in normal controls. The differences of biomechanical factors were compared among the groups as well.

## Materials and Methods

### Subjects


Newly diagnosed NTG and HTG patients were consecutively recruited in Eye & ENT Hospital, Fudan

University. Normal controls were enrolled from those who visited our hospital for regular eye checkups. Written informed consent was obtained from all participants before the enrollment. This study was approved by the hospital's Ethics Committee and was conducted in accordance with the tenets of the Declaration of Helsinki.

Healthy controls were defined as those with no history of eye diseases, IOP  $\leq 21$  mm Hg, open anterior chamber angles, normal appearances of the optic nerve head, and normal VF. A normal VF test result was defined as one with MD and pattern standard deviation (PSD) within 95% confidence limits of normal reference and glaucoma hemifield test within normal limits.

POAG was diagnosed as typical glaucomatous disc cupping and compatible VF defects in at least one eye and with the presence of open angles with Shaffer grading  $>2$  on gonioscopy. In addition, patients were diagnosed with HTG when they had at least one measurement of IOP  $> 21$  mm Hg with a Goldmann applanation tonometer taken at three different time points before treatment. On the contrary, patients were diagnosed as NTG when they had IOP  $\leq 21$  mm Hg at all time points in a 24-hour IOP test. Exclusion criteria included histories of intraocular surgery, laser, trauma, secondary glaucoma and corneal abnormalities. Patients with any VF loss caused by nonglaucomatous diseases were also excluded from this study. Subjects with concurrent or prior use of antiglaucoma medications were excluded as well.

All subjects underwent a comprehensive ophthalmologic examination, including visual acuity, best corrected visual acuity, slit-lamp biomicroscopy, fundus evaluation of the optic disc with a 90-diopter lens, IOP measurement using the Goldmann applanation tonometer and gonioscopy examination by an experienced glaucoma specialist. The VF tests were performed by the 30-2 SITA Standard program of the Humphrey Field Analyzer (Carl Zeiss Meditec, Dublin, CA, USA) to obtain MD and PSD. Only reliable VF measurements, which was defined as fixation loss rate less than 20%, false-positive and -negative rates less than 15%, were used for analysis. The retinal nerve fiber layer (RNFL) and ganglion cell complex (GCC) thickness were measured by spectral domain optical coherence tomography (SD-OCT; RTVue OCT; Optovue Inc., Fremont, CA, USA). VF and SD-OCT tests were conducted within one month before the Corvis ST measurements. Glaucoma severity was evaluated by the VF indexes (MD and PSD) and the structural parameters of optic disc and macular features (RNFL and GCC thickness). Axial length was measured by IOL Master (Zeiss, Oberkochen, Germany).



**Figure 1.** Schematic diagrams of the ocular biomechanical parameters provided by Corvis ST. (A) Cornea deformation during the Corvis ST measurement. From left to right: resting state before the measurement; first appplanation; highest concavity; second appplanation. (B) Graphs illustrating SP-A1, ARTh, DA ratio 2 mm, inverse radius and integrated radius. Lower values of SP-A1 and ARTh indicate a more deformable cornea, whereas higher values of DA ratio 2 mm, inverse radius and integrated radius indicate a more deformable cornea. (C) Correlation between deformation amplitude and whole eye movement. Deformation amplitude is a sum of whole eye movement and pure corneal deformation named deflection amplitude. *Yellow arrow:* whole eye movement; *Red arrow:* deflection amplitude; *Blue arrow:* deformation amplitude.

### Corvis ST Measurements

The Corvis ST (Oculus, GmbH, Wetzlar, Germany) was used to measure IOP, biomechanically corrected IOP, central corneal thickness (CCT), as well as ocular biomechanical parameters. Corvis ST monitors cornea’s dynamic reaction to an air impulse with a high-speed Scheimpflug camera, which captures 4330 images per second.<sup>22</sup> Under the pressure, the cornea bends inward from the resting state to the first appplanation point and continues to move until it reaches the

maximum deformation state, namely highest concavity (HC). When the pressure decreases, the cornea moves outward and passes the second appplanation point before reaching to its resting state.<sup>14,23</sup> After each measurement, corneal biomechanical parameters are produced as follows (Fig. 1A): time from start to the first and second applations (AT1 and AT2, respectively), velocity during the first and second applations (AV1 and AV2, respectively), time and maximum deformation amplitude (DA) from resting state to HC, peak distance (PD) between corneal

bending points and radius of HC. In addition, several new biomechanical parameters measured by the latest Corvis ST software (v. 1.3r1538) were included in our study (Figs. 1B, 1C):

Stiffness parameter at first applanation (SP-A1): the difference between the strength of the air puff at the corneal surface and biomechanically corrected IOP divided by deflection amplitude at the first applanation.<sup>24,25</sup>

Ambrosio relational thickness to the horizontal profile (ARTh): the quotient of corneal thickness at the thinnest point of the horizontal meridian and the thickness increase toward the periphery.<sup>25</sup>

DA ratio 2 mm: the ratio of deformation amplitude at corneal apex to that at 2 mm.<sup>25,26</sup>

Integrated radius: the integrated area under the radius of the inverted curvature (Fig. 1B) during the concave phase.<sup>26</sup>

Whole eye movement length (WEM length): the length of the linear anterior-posterior movement of the whole eye after maximum displacement of the cornea.<sup>25,27,28</sup>

WEM time: the time taken for the linear anterior-posterior movement of the whole eye after maximum displacement of the cornea.<sup>25,27,28</sup>

## Statistical Analysis

Only data of one eye for each subject was included for statistical analysis. The left eye was selected if both eyes had eligible Corvis ST measurements. Data were presented as mean  $\pm$  standard deviation (SD) or as median interquartile range. Analysis of variance or Kruskal Wallis rank sum tests were used to assess the differences of demographic and ocular parameters among the normal control, NTG and HTG groups. Biomechanical differences among the three groups were analyzed by general linear model. Furthermore,

the Bonferroni post hoc tests were used for pairwise comparisons. Spearman rank correlation tests were employed to identify the possible correlated factors of glaucoma severity indices since at least one variable in each group did not obey the normal distribution. Multivariate linear regression tests were performed to evaluate the association between glaucoma severity parameters with the demographic and ocular variables in each group. In this study, significant *P* value was set at  $<0.05$  (two tailed), and SPSS version 23.0 program was used for the whole analysis.

## Results

A total of 51 normal controls and 102 untreated POAG patients, including 47 NTG and 55 HTG patients were enrolled in the study. The demographic and ocular characteristics of the participants were shown in Table 1. Gender, age, IOP, and all glaucoma severity parameters (MD, PSD, GCC and RNFL) showed statistically significant differences among the groups of control, NTG and HTG (all  $P < 0.01$ ), whereas axial length and CCT revealed no significant differences ( $P = 0.143$  and  $0.155$ , respectively).

The differences of biomechanical parameters among the three groups were compared by the general linear model with adjustment for gender, age, axial length, IOP and CCT. There were statistically significant differences in several parameters between the controls and untreated glaucoma groups (Table 2). DA was significantly higher whereas AT1 and HC time were significantly lower in NTG (DA =  $1.01 \pm 0.01$  mm,  $P = 0.001$ ; AT1 =  $7.62 \pm 0.02$  ms,  $P < 0.0001$ ; HC time =  $16.91 \pm 0.10$  ms,  $P = 0.001$ ) and HTG eyes (DA =  $1.02 \pm 0.01$  mm,  $P = 0.002$ ; AT1 =  $7.64 \pm 0.03$  ms,  $P < 0.0001$ ; HC time =  $16.77 \pm 0.10$  ms,  $P = 0.0002$ ) than in controls (DA =  $0.95 \pm 0.01$  mm; AT1 =  $7.82 \pm 0.02$  ms; HC time =  $17.41 \pm 0.01$

**Table 1.** Demographic and Ocular Characteristics of Normal Control and Glaucoma Participants

Variables	Mean $\pm$ SD/Median (P25, P75)			<i>P</i>
	Normal Control (n = 51)	NTG (n = 47)	HTG (n = 55)	
Gender (Male/Female)	22/29	19/28	42/13	$<0.0001$
Age, years	37.00 (32.00, 40.00)	44.00 (36.00, 51.50)	37.50 (32.50, 46.00)	0.0012
Axial length, mm	24.87 (24.07, 26.77)	25.78 (23.77, 27.01)	26.08 (24.75, 27.13)	0.143
IOP, mmHg	16.50 (15.25, 18.75)	17.00 (15.75, 18.25)	20.75 (18.19, 23.31)	$<0.0001$
CCT, $\mu$ m	$546.74 \pm 31.72$	$547.10 \pm 33.73$	$557.45 \pm 31.27$	0.155
MD, dB	$-2.01 (-2.81, -0.77)$	$-5.33 (-8.98, -2.43)$	$-4.17 (-9.73, -2.92)$	$<0.0001$
PSD, dB	1.69 (1.52, 2.05)	5.33 (2.82, 10.72)	4.90 (2.21, 11.12)	$<0.0001$
GCC, $\mu$ m	$96.90 \pm 5.62$	$77.77 \pm 9.42$	$76.57 \pm 10.99$	$<0.0001$
RNFL, $\mu$ m	$100.43 \pm 8.32$	$82.23 \pm 12.96$	$78.38 \pm 13.41$	$<0.0001$

**Table 2.** Adjusted Ocular Biomechanical Parameters and Comparative Analysis Among Groups of Normal Control, NTG and HTG

Variables	Normal Control	NTG	HTG	<i>P</i>	P1	P2	P3
DA, mm	0.95 ± 0.01	1.01 ± 0.01	1.02 ± 0.01	<b>&lt;0.0001</b>	<b>0.001</b>	<b>0.002</b>	1.000
AT1, ms	7.82 ± 0.02	7.62 ± 0.02	7.64 ± 0.03	<b>&lt;0.0001</b>	<b>&lt;0.0001</b>	<b>&lt;0.0001</b>	1.000
AV1, m/s	0.12 ± 0.01	0.14 ± 0.01	0.13 ± 0.01	<b>&lt;0.0001</b>	<b>&lt;0.0001</b>	0.096	<b>0.044</b>
AT2, ms	21.83 ± 0.10	21.62 ± 0.10	21.49 ± 0.11	0.057	0.356	0.062	1.000
AV2, m/s	−0.24 ± 0.01	−0.25 ± 0.01	−0.24 ± 0.01	0.497	0.719	1.000	1.000
HC time, ms	17.41 ± 0.01	16.91 ± 0.10	16.77 ± 0.10	<b>&lt;0.0001</b>	<b>0.001</b>	<b>0.0002</b>	1.000
HC PD, mm	4.74 ± 0.04	4.80 ± 0.04	4.80 ± 0.04	0.539	0.925	1.000	1.000
HC radius, mm	7.32 ± 0.50	6.97 ± 0.53	7.97 ± 0.57	0.472	1.000	1.000	0.672
SP-A1, mm Hg/mm	132.85 ± 3.02	119.55 ± 3.19	126.09 ± 3.41	<b>0.012</b>	<b>0.009</b>	0.477	0.554
Integrated radius, mm <sup>−1</sup>	7.98 ± 0.14	8.42 ± 0.14	8.17 ± 0.15	0.089	0.084	1.000	0.881
ARTh, μm	418.52 ± 10.51	449.16 ± 11.16	451.72 ± 11.88	0.065	0.144	0.142	1.000
DA ratio 2 mm	4.44 ± 0.10	4.32 ± 0.12	4.19 ± 0.12	0.349	1.000	0.444	1.000
WEM length, mm	0.29 ± 0.01	0.28 ± 0.01	0.32 ± 0.01	0.130	1.000	0.357	0.152
WEM time, ms	22.86 ± 0.17	22.12 ± 0.18	22.34 ± 0.19	<b>0.008</b>	<b>0.008</b>	0.147	1.000

*P*, *P* values of the general linear model with adjustment for gender, age, axial length, IOP and CCT;

P1: *P* values by Bonferroni post hoc tests between control and NTG;

P2: *P* values by Bonferroni post hoc tests between control and HTG;

P3: *P* values by Bonferroni post hoc tests between NTG and HTG.

Bold *P* values are <0.05 with statistical significances.

ms). In addition, NTG showed the highest AV1 (0.14 ± 0.01 m/s) compared with controls (0.12 ± 0.01 m/s, *P* < 0.0001) and HTG (0.13 ± 0.01 m/s, *P* = 0.044). SP-A1 and time of WEM decreased statistically significantly in NTG eyes (SP-A1 = 119.55 ± 3.19 mm Hg/mm; WEM = 22.12 ± 0.18 ms) than in controls (SP-A1 = 132.85 ± 3.02 mm Hg/mm, *P* = 0.009; WEM = 22.86 ± 0.17 ms, *P* = 0.008). The ocular biomechanical differences between the POAG patients (composed of NTG and HTG) and healthy controls were also analyzed by the general linear model and the results were shown in Supplementary Table S1. DA (1.02 ± 0.01 vs. 0.95 ± 0.01 mm, *P* < 0.0001), AV1 (0.13 ± 0.01 vs. 0.12 ± 0.01 m/s, *P* < 0.0001), integrated radius (8.35 ± 0.09 vs. 7.97 ± 0.14 mm<sup>−1</sup>, *P* = 0.028) and ARTh (449.65 ± 7.11 vs. 420.84 ± 10.44 μm, *P* = 0.031) were significantly higher, whereas AT1 (7.62 ± 0.02 vs. 7.82 ± 0.02 ms, *P* < 0.0001), AT2 (21.57 ± 0.06 vs. 21.82 ± 0.09 ms, *P* = 0.035), HC time (16.87 ± 0.06 vs. 17.40 ± 0.09 ms, *P* < 0.0001), SP-A1 (121.97 ± 2.05 vs. 132.82 ± 3.01 mm Hg/mm, *P* = 0.005) and time of WEM (22.20 ± 0.11 vs. 22.85 ± 0.17 ms, *P* = 0.002) were significantly lower in POAG eyes compared to those in normal controls.

In further analyses on the associated factors of glaucoma severity, we found a number of demographic and ocular variables were significantly associated with both of VF and SD-OCT indices by Spearman rank

correlation (Supplementary Tables S2 and S3). The biomechanical parameters which showed statistically significant differences in the univariate analysis in each group as well as gender, age, axial length, CCT and IOP, were entered into the multivariate linear regression models. In normal controls (Supplementary Table S4), there was a significant negative correlation between DA ratio 2mm and MD ( $\beta = -0.43$ , 95% confidence interval [CI], −0.75 to −0.10, *P* = 0.01). CCT was significantly associated with PSD ( $\beta = -0.01$ , 95% CI, −0.01 to 0.01, *P* = 0.03). Longer axial length was associated with decreased thickness of GCC ( $\beta = -1.28$ , 95% CI, −2.38 to −0.19, *P* = 0.02) and RNFL ( $\beta = -2.65$ , 95% CI, −4.22 to −1.08, *P* = 0.0007). In addition, healthy females showed reduced GCC thickness compared with males ( $\beta = -3.72$ , 95% CI, −7.04 to −0.40, *P* = 0.03). No statistically significant variables were detected in POAG subjects except axial length showed a positive relationship with PSD ( $\beta = 0.64$ , 95% CI, 0.08 to 1.20, *P* = 0.03, Supplementary Table S4).

Interestingly, when NTG and HTG were analyzed separately, shorter time of WEM was significantly associated with worse MD ( $\beta = 1.57$ , 95% CI, 0.21 to 2.94, *P* = 0.02) and higher values of PSD ( $\beta = -1.55$ , 95% CI, −2.98 to −0.12, *P* = 0.03) in NTG (Table 3). However, only axial length revealed a positive correlation with PSD ( $\beta = 0.95$ , 95% CI, 0.15 to 1.75, *P* =

**Table 3.** Association Between Demographic and Ocular Variables With Glaucoma Severity Parameters by Multivariate Linear Regression Analysis in NTG and HTG Groups

	MD		PSD		GCC		RNFL	
	$\beta$ (95% CI)	P	$\beta$ (95% CI)	P	$\beta$ (95% CI)	P	$\beta$ (95% CI)	P
<b>NTG</b>								
Gender	0.55 (-2.28 to 3.39)	0.70	0.57 (-2.17 to 3.32)	0.67	6.62 (-0.08 to 13.32)	0.05	3.63 (-5.03 to 12.29)	0.40
Age	-0.09 (-0.19 to 0.01)	0.08	0.05 (-0.05 to 0.15)	0.28	0.01 (-0.24 to 0.25)	0.95	-0.25 (-0.58 to 0.08)	0.13
Axial length	0.12 (-0.63 to 0.88)	0.74	-0.04 (-0.82 to 0.74)	0.92	0.59 (-1.09 to 2.27)	0.48	-0.47 (-2.72 to 1.79)	0.68
IOP	-0.08 (-0.62 to 0.47)	0.78	0.30 (-0.29 to 0.89)	0.31	-0.34 (-1.83 to 1.15)	0.65	0.21 (-1.62 to 2.05)	0.81
CCT	-0.01 (-0.05 to 0.03)	0.49	0.00 (-0.04 to 0.04)	0.93	-0.02 (-0.11 to 0.08)	0.73	-0.09 (-0.21 to 0.03)	0.15
WEM time	1.57 (0.21-2.94)	<b>0.02</b>	-1.55 (-2.98 to -0.12)	<b>0.03</b>	—	—	—	—
WEM length	17.88 (-1.15 to 36.91)	0.06	-19.61 (-39.24 to 0.02)	0.05	—	—	—	—
ARTh	-0.01 (-0.02 to 0.01)	0.21	—	—	—	—	—	—
AT2	—	—	-0.17 (-2.07 to 1.72)	0.85	—	—	—	—
<b>HTG</b>								
Gender	0.75 (-2.54 to 4.03)	0.65	-1.69 (-4.57 to 1.19)	0.24	-1.15 (-7.90 to 5.60)	0.73	-0.18 (-6.85 to 6.49)	0.96
Age	-0.02 (-0.16 to 0.11)	0.73	0.01 (-0.10 to 0.13)	0.82	-0.02 (-0.32 to 0.27)	0.88	-0.04 (-0.34 to 0.25)	0.77
Axial length	-0.72 (-1.73 to 0.28)	0.16	0.95 (0.15-1.75)	<b>0.02</b>	-3.21 (-5.02 to -1.40)	<b>&lt;0.0001</b>	-3.12 (-4.95 to -1.29)	<b>&lt;0.0001</b>
IOP	-0.31 (-0.65 to 0.03)	0.08	0.13 (-0.17 to 0.44)	0.38	-0.49 (-1.22 to 0.24)	0.19	-0.59 (-1.32 to 0.13)	0.11
CCT	0.05 (0.00-0.10)	0.07	-0.02 (-0.06 to 0.02)	0.38	0.09 (-0.01 to 0.20)	0.09	0.09 (-0.02 to 0.20)	0.10
DA ratio 2mm	0.72 (-0.54 to 1.98)	0.26	—	—	—	—	—	—
WEM time	—	—	—	—	1.83 (-0.80 to 4.47)	0.17	—	—

Bold P values are <0.05 with statistical significances.




Figure 2. Scatterplots of the correlation between time of WEM and glaucoma severity parameters in NTG patients.

0.02) and negative correlation with GCC ( $\beta = -3.21$ , 95% CI,  $-5.02$  to  $-1.40$ ,  $P < 0.0001$ ) and RNFL ( $\beta = -3.12$ , 95% CI,  $-4.95$  to  $-1.29$ ,  $P < 0.0001$ ) thickness

in HTG (Table 3). Scatterplots showing the relationship between these ocular parameters and glaucoma severity indexes were presented in Figures 2 and 3.




Figure 3. Scatterplots of the correlation between axial length and glaucoma severity parameters in HTG patients.

## Discussion

Corneal biomechanical properties are gaining increasingly research attention and have been demonstrated differences across various ocular situations. Our results found that the ocular biomechanics were significantly different between untreated POAG (including NTG and HTG separately) patients and normal controls. More interestingly, our findings revealed shorter time of WEM was independently associated with more severe visual field defect in NTG, whereas longer axial length was related to glaucoma damage in HTG, suggesting different involvement of biomechanics in glaucoma severity between NTG and HTG.

Comparative analysis by the general linear model suggested that the corneal deformability is higher in untreated POAG patients than normal controls as indicated by increased values of DA, AV1, integrated radius, and decreased values of AT1, HC time and SP-A1 (Supplementary Table S1). Different results were obtained about the corneal biomechanical differences between POAG and normal controls in previous studies. Some reported corneas of POAG were less deformable compared to controls,<sup>17,29</sup> whereas others reported that no biomechanical differences were found between the two groups.<sup>30,31</sup> The possible reasons for this disagreement could be due to the influences of antiglaucoma medications on corneal biomechanics, differences in sample size and variables used for adjustment. When NTG and HTG were analyzed separately and compared with controls (Table 2), DA, AT1 and HC time showed similar changes in the two glaucoma groups. Additionally, SP-A1 decreased and AV1 increased significantly in NTG than in controls. This finding revealed that NTG corneas were more deformable than healthy ones and HTG, which were in agreement with previous studies.<sup>16,21,32,33</sup> More deformable corneas were more likely to show glaucomatous visual field defect.<sup>9,16</sup> It has been reported that in untreated NTG patients with asymmetric visual field damage, the worse eyes were the ones with a larger degree of corneal deformability compared to the better eyes.<sup>34</sup> More compliant corneas may reflect higher deformability in sclera and lamina cribrosa to some extent since they are in structural continuum with the cornea, thus making the optic nerve head more vulnerable to glaucomatous damage.<sup>9,35,36</sup>

After the air impulse applying on the cornea, the kinetic energy of the puff is absorbed by the cornea as well as extraocular tissues such as fat and orbital muscles.<sup>37</sup> When the cornea reaches its maximum displacement, the whole eye displays a slow linear motion in the anterior-posterior direction,<sup>25</sup> namely

WEM, which reflects orbital compliance,<sup>38</sup> and the pure corneal deformation is called deflection amplitude. As shown in Figure 1C, WEM is a part of DA (deformation amplitude). Our study showed that the time of WEM was significantly decreased in NTG patients compared with controls (Table 2), which was in line with a previous study.<sup>32</sup> Decreased WEM indicated decreased orbital compliance<sup>28,39</sup> in these patients. Unlike other corneal biomechanical parameters, only a few literatures studied WEM, and its clinical relevance need to be fully established in future.

Furthermore, we evaluated the potential contribution of ocular biomechanics to glaucoma severity in each group. In total POAG patients, only axial length showed a significantly positive correlation with PSD (Supplementary Table S4). To our knowledge, studies about the relationship between ocular biomechanics measured by Corvis ST and glaucoma severity were very few. Hirasawa et al reported that eyes with fast AV1 and shorter AT2 were related to more severe VF damage in POAG patients.<sup>15</sup> Vinciguerra et al.<sup>16</sup> found there was a significant negative correlation between MD and DA ratio in addition to its positive correlation with SP-A1 in POAG patients. The most possible reason for this inconsistency could be that these two studies did not exclude patients under usage of hypotensive eye drops. IOP lowering medications, particularly the prostaglandin analogues, have been shown to influence corneal biomechanics significantly.<sup>17-19</sup> Our results were in agreement with the study from Bolivar et al.,<sup>40</sup> which found that the initial Corvis ST biomechanical parameters (DA, AT1, AV1, AT2, AV2, HC time, HC PD, HC radius) had no significant correlation with MD in treatment naïve POAG eyes.

Interestingly, when analyzed separately, shorter time of WEM was demonstrated to be associated with worse glaucoma damage in NTG, but not in HTG (Table 3). Decreased WEM in NTG patients indicated decreased orbital compliance, perhaps suggesting the buffering ability of the orbit tissues to the whole eyeball was reduced. That made the optic nerve head more vulnerable to forces, finally leading to increased susceptibility to glaucomatous damage. More studies should be conducted on the significant relevance of WEM in clinical setting. Regarding HTG, only axial length was associated with glaucoma severity in our study. The possible reason for the association between longer axial length and more severe glaucoma damage may be partly due to increased deformability of the eyeball with axial elongation. Eyeball elongation is often accompanied by a reduction of sclera collagen fiber bundles, and thus a thinner sclera<sup>41</sup> and lamina cribrosa,<sup>42</sup> leading to scleral rigidity decreased



and greater compliance of the eyeball.<sup>43</sup> Different involvement of biomechanics in glaucoma severity between NTG and HTG further suggested different pathogenic mechanisms existing between these two glaucoma subtypes. Our results added more biomechanical evidence for the underlying mechanism differences between NTG and HTG.

IOP and age have been demonstrated as important risk factors for visual field progression in glaucoma.<sup>44</sup> However, neither showed statistically significant association with glaucoma severity in our study. Several previous studies reported the lack of association between IOP and MD in untreated POAG patients as well.<sup>40,45</sup> Also, there was a study that observed that age was not correlated with glaucoma severity in POAG patients.<sup>15</sup> The lack of association between IOP and age with glaucoma severity in our study may be explained by the cross-sectional design and small sample size. Another possible reason for the nonsignificant association between age and glaucoma damage could be due to relatively young age of glaucoma patients in our study, in which the median age of NTG and HTG was 44 and 37.5 years, respectively.

Our study has several limitations. First, the sample size was small, and the participants were recruited from a single tertiary hospital, which may not necessarily represent for a more general normal and POAG population. In addition, the cross-sectional nature of the study made it only assess association but not causality. Longitudinal studies are necessary to further explore the relationship between baseline ocular biomechanics and glaucoma progression.

In conclusion, our study demonstrated NTG corneas were more deformable than those of normal controls and HTG. More importantly, our results showed glaucoma severity was correlated with different factors between NTG and HTG. Shorter time of WEM was independently associated with more severe visual field defect in NTG, whereas longer axial length was associated with glaucoma damage in HTG. Our findings may have implications for understanding the etiological differences between NTG and HTG. In future, more studies should be conducted on the biomechanical relevance of WEM in glaucoma.

## Acknowledgments

The authors are grateful to the normal and glaucoma subjects who participated in the study.

Supported by funds from the National Science Foundation of China (no. 81870692 and no. 81600730); the Shanghai Committee of Science and Technology

(no. 17410712500 and no. 20s31905800); the Clinical Research Plan of Shanghai Hospital Development Center (SHDC2020CR6029) and the top priority of clinical medicine center of Shanghai (no. 2017ZZ01020). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Disclosure: **N. Wu**, None; **Y. Chen**, None; **X. Sun**, None

## References

1. Martucci A, Nucci C. Evidence on neuroprotective properties of coenzyme Q10 in the treatment of glaucoma. *Neural Regen Res.* 2019;14:197–200.
2. Kass MA, Heuer DK, Higginbotham EJ, et al. The Ocular Hypertension Treatment Study: a randomized trial determines that topical ocular hypotensive medication delays or prevents the onset of primary open-angle glaucoma. *Arch Ophthalmol.* 2002;120:701–713; discussion 829–730.
3. Leske MC, Heijl A, Hyman L, Bengtsson B, Dong L, Yang Z. Predictors of long-term progression in the early manifest glaucoma trial. *Ophthalmology.* 2007;114:1965–1972.
4. Medeiros FA, Weinreb RN. Evaluation of the influence of corneal biomechanical properties on intraocular pressure measurements using the ocular response analyzer. *J Glaucoma.* 2006;15:364–370.
5. Liu J, Roberts CJ. Influence of corneal biomechanical properties on intraocular pressure measurement: quantitative analysis. *J Cataract Refract Surg.* 2005;31:146–155.
6. Wells AP, Garway-Heath DF, Poostchi A, Wong T, Chan KC, Sachdev N. Corneal hysteresis but not corneal thickness correlates with optic nerve surface compliance in glaucoma patients. *Invest Ophthalmol Vis Sci.* 2008;49:3262–3268.
7. Anand A, De Moraes CG, Teng CC, Tello C, Liebmann JM, Ritch R. Corneal hysteresis and visual field asymmetry in open angle glaucoma. *Invest Ophthalmol Vis Sci.* 2010;51:6514–6518.
8. Kaushik S, Pandav SS, Banger A, Aggarwal K, Gupta A. Relationship between corneal biomechanical properties, central corneal thickness, and intraocular pressure across the spectrum of glaucoma. *Am J Ophthalmol.* 2012;153:840–849.e842.
9. Congdon NG, Broman AT, Bandeen-Roche K, Grover D, Quigley HA. Central corneal thickness

- and corneal hysteresis associated with glaucoma damage. *Am J Ophthalmol.* 2006;141(5):868–875.
10. Medeiros FA, Meira-Freitas D, Lisboa R, Kuang TM, Zangwill LM, Weinreb RN. Corneal hysteresis as a risk factor for glaucoma progression: a prospective longitudinal study. *Ophthalmology.* 2013;120:1533–1540.
  11. Dascalescu D, Corbu C, Constantin M, et al. Correlations between corneal biomechanics and glaucoma severity in patients with primary open angle glaucoma. *Maedica (Bucur).* 2015;10:331–335.
  12. Detry-Morel M, Jamart J, Hautenauven F, Pourjavan S. Comparison of the corneal biomechanical properties with the Ocular Response Analyzer (ORA) in African and Caucasian normal subjects and patients with glaucoma. *Acta Ophthalmol.* 2012;90(2):e118–124.
  13. Hirneiss C, Neubauer AS, Yu A, Kampik A, Kernt M. Corneal biomechanics measured with the ocular response analyser in patients with unilateral open-angle glaucoma. *Acta Ophthalmol.* 2011;89(2):e189–192.
  14. Chansangpetch S, Panpruk R, Manassakorn A, et al. Impact of myopia on corneal biomechanics in glaucoma and nonglaucoma patients. *Invest Ophthalmol Vis Sci.* 2017;58:4990–4996.
  15. Hirasawa K, Matsuura M, Murata H, et al. Association between corneal biomechanical properties with ocular response analyzer and also CorvisST tonometry, and glaucomatous visual field severity. *Transl Vis Sci Technol.* 2017;6(3):18.
  16. Vinciguerra R, Rehman S, Vallabh NA, et al. Corneal biomechanics and biomechanically corrected intraocular pressure in primary open-angle glaucoma, ocular hypertension and controls. *Br J Ophthalmol.* 2020;104:121–126.
  17. Wu N, Chen Y, Yu X, Li M, Wen W, Sun X. Changes in corneal biomechanical properties after long-term topical prostaglandin therapy. *PLoS One.* 2016;11(5):e0155527.
  18. Bolivar G, Sanchez-Barahona C, Teus M, et al. Effect of topical prostaglandin analogues on corneal hysteresis. *Acta Ophthalmol.* 2015;93(6):e495–498.
  19. Tsikripis P, Papaconstantinou D, Koutsandrea C, Apostolopoulos M, Georgalas I. The effect of prostaglandin analogs on the biomechanical properties and central thickness of the cornea of patients with open-angle glaucoma: a 3-year study on 108 eyes. *Drug Des Devel Ther.* 2013;7:1149–1156.
  20. Park IK, Kim KW, Moon NJ, Shin JH, Chun YS. Comparison of superior and inferior visual field asymmetry between normal-tension and high-tension glaucoma. *J Glaucoma.* 2021;30:648–655.
  21. Wu N, Chen Y, Yang Y, Sun X. The changes of corneal biomechanical properties with long-term treatment of prostaglandin analogue measured by Corvis ST. *BMC Ophthalmol.* 2020;20:422.
  22. Hong J, Xu J, Wei A, et al. A new tonometer—the Corvis ST tonometer: clinical comparison with noncontact and Goldmann applanation tonometers. *Invest Ophthalmol Vis Sci.* 2013;54:659–665.
  23. Salvetat ML, Zeppieri M, Tosoni C, Felletti M, Grasso L, Brusini P. Corneal deformation parameters provided by the Corvis-ST pachy-tonometer in healthy subjects and glaucoma patients. *J Glaucoma.* 2015;24:568–574.
  24. Roberts CJ, Mahmoud AM, Bons JP, et al. Introduction of two novel stiffness parameters and interpretation of air puff-induced biomechanical deformation parameters with a dynamic scheimpflug analyzer. *J Refract Surg.* 2017;33:266–273.
  25. Vinciguerra R, Ambrosio R, Jr., Elsheikh A, et al. Detection of keratoconus with a new biomechanical index. *J Refract Surg.* 2016;32:803–810.
  26. Hirasawa K, Nakakura S, Nakao Y, et al. Changes in corneal biomechanics and intraocular pressure following cataract surgery. *Am J Ophthalmol.* 2018;195:26–35.
  27. Ziaei M, Gokul A, Vellara H, Lu LM, Patel DV, McGhee CNJ. Measurement of in vivo biomechanical changes attributable to epithelial removal in keratoconus using a noncontact tonometer. *Cornea.* 2020;39:946–951.
  28. Leszczynska A, Moehler K, Spoerl E, et al. Measurement of orbital biomechanical properties in patients with thyroid orbitopathy using the dynamic scheimpflug analyzer (Corvis ST). *Curr Eye Res.* 2018;43:289–292.
  29. Wang W, Du S, Zhang X. Corneal deformation response in patients with primary open-angle glaucoma and in healthy subjects analyzed by Corvis ST. *Invest Ophthalmol Vis Sci.* 2015;56:5557–5565.
  30. Pradhan ZS, Deshmukh S, Dixit S, et al. A comparison of the corneal biomechanics in pseudoexfoliation glaucoma, primary open-angle glaucoma and healthy controls using Corvis ST. *PLoS One.* 2020;15(10):e0241296.
  31. Silva N, Ferreira A, Baptista PM, et al. Corneal biomechanics for ocular hypertension, primary open-angle glaucoma, and amyloidotic glaucoma: a comparative study by Corvis ST. *Clin Ophthalmol.* 2022;16:71–83.
  32. Miki A, Yasukura Y, Weinreb RN, et al. Dynamic scheimpflug ocular biomechanical parameters in

- untreated primary open angle glaucoma eyes. *Invest Ophthalmol Vis Sci.* 2020;61(4):19.
33. Chen YY, Wang TH, Huang JY, Su CC. Relationship of axial length and corneal biomechanical properties with susceptibility to unilateral normal-tension glaucoma. *Graefes Arch Clin Exp Ophthalmol.* 2022;260:255–264.
  34. Li BB, Cai Y, Pan YZ, et al. Corneal biomechanical parameters and asymmetric visual field damage in patients with untreated normal tension glaucoma. *Chin Med J (Engl).* 2017;130:334–339.
  35. Bochmann F, Ang GS, Azuara-Blanco A. Lower corneal hysteresis in glaucoma patients with acquired pit of the optic nerve (APON). *Graefes Arch Clin Exp Ophthalmol.* 2008;246:735–738.
  36. Qassim A, Mullany S, Abedi F, et al. Corneal stiffness parameters are predictive of structural and functional progression in glaucoma suspect eyes. *Ophthalmology.* 2021;128:993–1004.
  37. Francis M, Pahuja N, Shroff R, et al. Waveform analysis of deformation amplitude and deflection amplitude in normal, suspect, and keratoconic eyes. *J Cataract Refract Surg.* 2017;43:1271–1280.
  38. Jung Y, Park HL, Yang HJ, Park CK. Characteristics of corneal biomechanical responses detected by a non-contact scheimpflug-based tonometer in eyes with glaucoma. *Acta Ophthalmol.* 2017;95(7):e556–e563.
  39. Hwang HS, Kim EC, Kim MS, Yang SW. A novel method for quantifying the biomechanical parameters of orbital soft tissue using a corneal dynamic Scheimpflug analyser: a retrospective study. *BMC Ophthalmol.* 2019;19:53.
  40. Bolivar G, Sanchez-Barahona C, Ketabi S, Kozobolis V, Teus MA. Corneal factors associated with the amount of visual field damage in eyes with newly diagnosed, untreated, open-angle glaucoma. *Ophthalmol Ther.* 2021;10:669–676.
  41. Funata M, Tokoro T. Scleral change in experimentally myopic monkeys. *Graefes Arch Clin Exp Ophthalmol.* 1990;228:174–179.
  42. Jonas JB, Berenshtein E, Holbach L. Lamina cribrosa thickness and spatial relationships between intraocular space and cerebrospinal fluid space in highly myopic eyes. *Invest Ophthalmol Vis Sci.* 2004;45:2660–2665.
  43. Yang Y, Ng TK, Wang L, et al. Association of 24-hour intraocular pressure fluctuation with corneal hysteresis and axial length in untreated Chinese primary open-angle glaucoma patients. *Transl Vis Sci Technol.* 2020;9(12):25.
  44. Kim JH, Rabiolo A, Morales E, et al. Risk factors for fast visual field progression in glaucoma. *Am J Ophthalmol.* 2019;207:268–278.
  45. Teus MA, Castejon MA, Calvo MA, Perez-Salaices P, Marcos A. Intraocular pressure as a risk factor for visual field loss in pseudoexfoliative and in primary open-angle glaucoma. *Ophthalmology.* 1998;105:2225–2229; discussion 2229–2230.