

Controlling invasive species with biologically-inspired robots

Giovanni Polverino¹ and Maurizio Porfiri^{2,3,4}

¹Centre for Evolutionary Biology, School of Biological Sciences, University of Western Australia, Perth, Australia

²Department of Mechanical and Aerospace Engineering, Tandon School of Engineering, New York University, Brooklyn, USA

³Department of Biomedical Engineering, Tandon School of Engineering, New York University, Brooklyn, USA

⁴Center for Urban Science and Progress, Tandon School of Engineering, New York University, Brooklyn, USA

gio.polverino@gmail.com, giovanni.polverino@uwa.edu.au

Abstract

Robotics is emerging as a promising approach to study animal behavior. Biologically-inspired robots can manipulate the behavior of live animals and are increasingly employed to uncover the underpinnings of sociality and mate choice in the animal realm. But behavioral variation between animals plays a critical role for their ecology and evolution, and ultimately it determines variation in the survival, growth, and reproduction of individuals. While the study of behavioral responses of animals toward their robotic counterparts dominates the literature, it remains largely untested whether the life-history strategies of live animals can be artificially manipulated with biologically-inspired robots. Recently, predator-mimicking robots allowed to successfully study antipredator responses of highly invasive fish in detail, revealing that costs of behavioral alternations induced by robotic predators can impact the health and survival of invaders. The evidence that biologically-inspired robots can undermine the ecological success of invasive animals opens the door to novel experimental analyses at the interface between robotics, ecology, and invasion biology.

Invasive species: a serious environmental problem

Human activities have offered unprecedented opportunities to some animals for crossing geographical barriers and invading new habitats (Elton, 2000; Richardson, 2011). As a result, nowadays invasive species are spread across the planet, driving native species to extinction (Clavero and García-Berthou, 2005), compromising ecosystems (Lockwood, 2013), and costing our economy billions of dollars every year (Pimentel et al., 2005). Freshwater ecosystems host more than 25% of the world's vertebrates and are particularly vulnerable to competitive and aggressive invaders (Leprieur et al., 2008; Su et al., 2021). Despite the urgent environmental challenge that invasive species represent for freshwater ecosystems, methods to eradicate invasive species are largely unsuccessful (Byers et al., 2002).

The invasive mosquitofish (*Gambusia holbrooki*, Figure 1A) is a major threat to freshwater fish and amphibians (Pyke, 2008). Its impact on biodiversity has been recognized by the *International Union for Conservation of Nature*, which lists mosquitofish as amongst the world's top hundred worst

invasive species (Lowe et al., 2000). For example, the presence of mosquitofish is causing the extinction of the red-finned blue eye (*Scaturiginichthys vermeilipinnis*) and the Edgbaston goby (*Chlamydogobius squamigenus*)—two of the most critically endangered fish species in Australia (Faulks et al., 2017; Kerezy and Fensham, 2013; Nicol et al., 2015). To date, no successful procedure exists for eradicating mosquitofish and mitigating their impact on native fauna (Brookhouse and Coughran, 2010; Westhoff et al., 2013; Willis and Ling, 2000).




Figure 1 (A) Mosquitofish and (B) the body morphology and (C) swimming features that we measured in the native mosquitofish's primary predator (*Micropterus salmoides*) and that we integrated into the design of (D) the biologically-inspired robotic predator presented in Polverino et al. 2019a. A is adapted from Polverino and Porfiri 2013a, B is from Adicaza 3D models, and C and D are prepared by Polverino and Porfiri for this article.

Harnessing cutting-edge technology to understand animal invasions

New technologies have offered invaluable tools for studying the ecology and evolution of living creatures (Aguilar et al., 2014) and addressing the seemingly intractable problem of invasive species (Kerr et al., 2017; McCann et al., 2019). In particular, robotics is emerging as a promising approach to studying animal behavior (Krause et al., 2011; Romano et al., 2018; Webb, 2000) and animal invasions (Dufour et al., 2020; Polverino et al., 2019). State-of-the-art robots can move autonomously (Butail et al., 2013; Butail et al., 2014; Gribovskiy et al., 2018; Katzschmann et al., 2018), simulate

specific characteristics of live fish (Landgraf et al., 2016; Phamduy et al., 2014; Polverino et al., 2012; Romano et al., 2017a), infiltrate social groups (Bierbach et al., 2018; Butail et al., 2013; Marras and Porfiri, 2012), and interact with live animals in real time (Bonnet et al., 2019; Kopman et al., 2013; Polverino et al., 2019a).

Recent studies underscore the extraordinary opportunity of using robots inspired by live predators to control the behavior of pests such as locusts (Romano et al., 2017b, 2019) that pose a threat to human agricultural economies. Similarly, falcon-like robots are a powerful tool to deter other birds from flying in the vicinity of aircrafts (Folkertsma et al., 2017). Robots are inherently safer to use than live agents in the wild, since robots can be finely controlled and cannot breed.

Efforts from our group have shown that biologically-inspired robotic predators can repel invasive mosquitofish (Polverino and Porfiri, 2013a), undermine their health (Polverino et al., 2019a), and simultaneously attract non-invasive fish species (Polverino and Porfiri, 2013b).

Advancing the field

The unparalleled control offered by biologically-inspired robotics to successfully manipulate animal behavior leads to new exciting questions. We know that small behavioral adjustments have relatively large effects on the survival and reproduction of animals (Réale et al., 2007) and can determine the ecological success of invaders, including mosquitofish (Brodin et al., 2019; Rehage and Sih, 2004). What we do not know is whether and how the behavior of invasive animals can be artificially manipulated to undermine their ecological success.

Our previous analyses have demonstrated that variation in behavioral, physiological, and life-history traits in fish is associated at the individual level—individuals that are more active grow faster and have higher metabolic rates than less active ones (Polverino et al., 2018). We have uncovered the association between individual variation in risk taking and fertility, where males that take more risks have high number of sperm over risk-averse males (Gasparini et al., 2019). We also found empirical evidence that manipulating social experience during development alters the mating acquisition of fish in adulthood and, hence, their reproductive success (Polverino et al., 2019b).

The knowledge that altering the behavior of fish has repercussions on their physiology, growth, fertility, and reproduction is central for studying the developmental and evolutionary consequences of behavioral manipulation. Yet whether exposure to biologically-inspired robots has effects that extend beyond the behavior of animals remains unexplored (but see Polverino et al., 2019a).

We believe that state-of-the-art robots inspired by live predators offer an extraordinary opportunity to alter the behavior of invasive species and study the detrimental consequences of fear and anxiety on their survival and reproduction. World-leading experts in ecology and evolution indeed advocate for expanding the conceptual underpinning and practice of biological control (Culshaw-Maurer et al., 2020). In this vein, cutting-edge technologies can play a key role in revealing the evolutionary vulnerabilities of invasive

and pest species, and transfer the fundamental knowledge from predator-prey ecology to inform a new generation of biocontrol practices.

Acknowledgments

This work was supported by the Forrest Research Foundation (postdoctoral fellowship to G.P.), the University of Western Australia (Faculty of Science ‘Rising Stars Award’ 68000003 to G.P.), and the National Science Foundation (CMMI-1505832 and CMMI-1901697 to M.P.).

References

- Aguilar, W., Santamaría-Bonfil, G., Froese, T. and Gershenson, C. (2014). The past, present, and future of artificial life. *Frontiers in Robotics and AI*, 1:8.
- Bierbach, D., Lukas, J., Bergmann, A., Elsner, K., Höhne, L., Weber, C., . . . and Krause, J. (2018). Insights into the social behavior of surface and cave-dwelling fish (*Poecilia mexicana*) in light and darkness through the use of a biomimetic robot. *Frontiers in Robotics and AI*, 5:1–9.
- Bonnet, F., Mills, R., Szopek, M., Schönwetter-Fuchs, S., Halloy, J., Bogdan, S., . . . and Schmickl, T. (2019). Robots mediating interactions between animals for interspecies collective behaviors. *Science Robotics*, 4:eaa07897.
- Brodin, T., Fogarty, S., Sih, A. and Cote, J. (2019). Personality-dependent survival of the invasive mosquitofish: being social can be deadly. *Aquatic Invasions*, 14:1798–6540.
- Brookhouse, N. and Coughran, J. (2010). Exploring the potential for an ecology-specific, physical control method of the exotic pest mosquitofish, *Gambusia holbrooki*. *Ecological Management & Restoration*, 11:226–228.
- Butail, S., Bartolini, T. and Porfiri, M. (2013). Collective response of zebrafish shoals to a free-swimming robotic fish. *PLoS One*, 8:e76123.
- Butail, S., Polverino, G., Phamduy, P., Del Sette, F. and Porfiri, M. (2014). Influence of robotic shoal size, configuration, and activity on zebrafish behavior in a free-swimming environment. *Behavioural Brain Research*, 275:269–280.
- Byers, J. E., Reichard, S., Randall, J. M., Parker, I. M., Smith, C. S., Lonsdale, W. M., . . . and Hayes, D. (2002). Directing research to reduce the impacts of nonindigenous species. *Conservation Biology*, 16:630–640.
- Clavero, M. and García-Berthou, E. (2005). Invasive species are a leading cause of animal extinctions. *Trends in Ecology & Evolution*, 20:110.
- Culshaw-Maurer, M., Sih, A. and Rosenheim, J. A. (2020). Bugs scaring bugs: enemy-risk effects in biological control systems. *Ecology Letters*, 23:1693–1714.
- Dufour, C. M. S., Clark, D. L., Herrel, A. and Losos, J. B. (2020). Recent biological invasion shapes species recognition and

- aggressive behaviour in a native species: A behavioural experiment using robots in the field. *Journal of Animal Ecology*, 89:1604–1614.
- Elton, C. S. (2000). *The ecology of invasions by animals and plants*: University of Chicago Press.
- Faulks, L. K., Kerezszy, A., Unmack, P. J., Johnson, J. B. and Hughes, J. M. (2017). Going, going, gone? Loss of genetic diversity in two critically endangered Australian freshwater fishes, *Scaturiginichthys vermeilipinnis* and *Chlamydogobius squamigenus*, from Great Artesian Basin springs at Edgbaston, Queensland, Australia. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 27:39–50.
- Folkertsma, G. A., Straatman, W., Nijenhuis, N., Venner, C. H. and Stramigioli, S. (2017). Robird: a robotic bird of prey. *IEEE Robotics & Automation Magazine*, 24:22–29.
- Gasparini, C., Speechley, E. M. and Polverino, G. (2019). The bold and the sperm: positive association between boldness and sperm number in the guppy. *Royal Society Open Science*, 6:190474.
- Gribovskiy, A., Halloy, J., Deneubourg, J. L. and Mondada, F. (2018). Designing a socially integrated mobile robot for ethological research. *Robotics and Autonomous Systems*, 103:42–55.
- Katzschmann, R. K., DelPreto, J., MacCurdy, R. and Rus, D. (2018). Exploration of underwater life with an acoustically controlled soft robotic fish. *Science Robotics*, 3:eaar3449.
- Kerezszy, A. and Fensham, R. (2013). Conservation of the endangered red-finned blue-eye, *Scaturiginichthys vermeilipinnis*, and control of alien eastern gambusia, *Gambusia holbrooki*, in a spring wetland complex. *Marine and Freshwater Research*, 64:851–863.
- Kerr, P. J., Cattadori, I. M., Liu, J., Sim, D. G., Dodds, J. W., Brooks, J. W., . . . and Read, A. F. (2017). Next step in the ongoing arms race between myxoma virus and wild rabbits in Australia is a novel disease phenotype. *Proceedings of the National Academy of Sciences*, 114:9397–9402.
- Kopman, V., Laut, J. W., Polverino, G. and Porfiri, M. (2013). Closed-loop control of zebrafish response using a bioinspired robotic-fish in a preference test. *Journal of the Royal Society Interface*, 10:20120540.
- Krause, J., Winfield, A. F. T. and Deneubourg, J. L. (2011). Interactive robots in experimental biology. *Trends in Ecology & Evolution*, 26:369–375.
- Landgraf, T., Bierbach, D., Nguyen, H., Muggelberg, N., Romanczuk, P. and Krause, J. (2016). RoboFish: increased acceptance of interactive robotic fish with realistic eyes and natural motion patterns by live Trinidadian guppies. *Bioinspiration & Biomimetics*, 11:015001.
- Leprieur, F., Beauchard, O., Blanchet, S., Oberdorff, T. and Brosse, S. (2008). Fish invasions in the world's river systems: when natural processes are blurred by human activities. *PLoS Biology*, 6:0404–0410.
- Lockwood, J. L. (2013). *Invasion Ecology, 2nd Edition*: Wiley-Blackwell.
- Lowe, S., Browne, M., Boudjelas, S. and De Poorter, M. (2000). 100 of the world's worst invasive alien species a selection from the global invasive species database. *Aliens*, 12:1–12.
- Marras, S. and Porfiri, M. (2012). Fish and robots swimming together: attraction towards the robot demands biomimetic locomotion. *Journal of the Royal Society Interface*, 9:1856–1868.
- McCann, S., Crossland, M., Greenlees, M. and Shine, R. (2019). Invader control: factors influencing the attraction of cane toad (*Rhinella marina*) larvae to adult parotoid exudate. *Biological Invasions*, 21:1895–1904.
- Nicol, S., Haynes, T. B., Fensham, R. and Kerezszy, A. (2015). Quantifying the impact of *Gambusia holbrooki* on the extinction risk of the critically endangered red-finned blue-eye. *Ecosphere*, 6:1–18.
- Phamduy, P., Polverino, G., Fuller, R. C. and Porfiri, M. (2014). Fish and robot dancing together: bluefin killifish females respond differently to the courtship of a robot with varying color morphs. *Bioinspiration & Biomimetics*, 9:036021.
- Pimentel, D., Zuniga, R. and Morrison, D. (2005). Update on the environmental and economic costs associated with alien-invasive species in the United States. *Ecological Economics*, 52:273–288.
- Polverino, G., Abaid, N., Kopman, V., Macri, S. and Porfiri, M. (2012). Zebrafish response to robotic fish: preference experiments on isolated individuals and small shoals. *Bioinspiration & Biomimetics*, 7:036019.
- Polverino, G., Karakaya, M., Spinello, C., Soman, V. R. and Porfiri, M. (2019a). Behavioural and life-history responses of mosquitofish to biologically inspired and interactive robotic predators. *Journal of the Royal Society Interface*, 16:20190359.
- Polverino, G., Palmas, B. M., Evans, J. P. and Gasparini, C. (2019b). Individual plasticity in alternative reproductive tactics declines with social experience in male guppies. *Animal Behaviour*, 148:113–121.
- Polverino, G. and Porfiri, M. (2013a). Mosquitofish (*Gambusia affinis*) responds differentially to a robotic fish of varying swimming depth and aspect ratio. *Behavioural Brain Research*, 250:133–138.
- Polverino, G. and Porfiri, M. (2013b). Zebrafish (*Danio rerio*) behavioural response to bioinspired robotic fish and mosquitofish (*Gambusia affinis*). *Bioinspiration & Biomimetics*, 8:044001.
- Polverino, G., Santostefano, F., Díaz-Gil, C. and Mehner, T. (2018). Ecological conditions drive pace-of-life syndromes by shaping relationships between life history, physiology and behaviour in two populations of Eastern mosquitofish. *Scientific Reports*, 8:14673.
- Pyke, G. (2008). Plague minnow or mosquito fish? A review of the biology and impacts of introduced *Gambusia* species.

Annual Review of Ecology, Evolution and Systematics, 39:171–191.

- Réale, D., Reader, S. M., Sol, D., McDougall, P. T. and Dingemanse, N. J. (2007). Integrating animal temperament within ecology and evolution. *Biological Reviews*, 82:291–318.
- Rehage, J. S. and Sih, A. (2004). Dispersal behavior, boldness, and the link to invasiveness: a comparison of four *Gambusia* species. *Biological Invasions*, 6:379–391.
- Richardson, D. M. (2011). *Fifty years of invasion ecology: the legacy of Charles Elton*: Wiley-Blackwell.
- Romano, D., Benelli, G., Donati, E., Remorini, D., Canale, A. and Stefanini, C. (2017a). Multiple cues produced by a robotic fish modulate aggressive behaviour in Siamese fighting fishes. *Scientific Reports*, 7:4667.
- Romano, D., Benelli, G. and Stefanini, C. (2017b). Escape and surveillance asymmetries in locusts exposed to a Guinea fowl-mimicking robot predator. *Scientific Reports*, 7:12825.
- Romano, D., Benelli, G. and Stefanini, C. (2019). Encoding lateralization of jump kinematics and eye use in a locust via bio-robotic artifacts. *Journal of Experimental Biology*, 222:jeb187427.
- Romano, D., Donati, E., Benelli, G. and Stefanini, C. (2018). A review on animal–robot interaction: from bio-hybrid organisms to mixed societies. *Biological Cybernetics*, 1–25.
- Su, G., Logez, M., Xu, J., Tao, S., Villéger, S. and Brosse, S. (2021). Human impacts on global freshwater fish biodiversity. *Science*, 371:835–838.
- Webb, B. (2000). What does robotics offer animal behaviour? *Animal Behaviour*, 60:545–558.
- Westhoff, J. T., Watts, A. V. and Mattingly, H. T. (2013). Efficacy of artificial refuge to enhance survival of young Barrens topminnows exposed to western mosquitofish. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 23:65–76.
- Willis, K. and Ling, N. (2000). Sensitivities of mosquitofish and black mudfish to a piscicide: could rotenone be used to control mosquitofish in New Zealand wetlands? *New Zealand Journal of Zoology*, 27:85–91.