

Serum-Containing Medium Effect on Isolation Rate of Dental Pulp Cells from Cryopreserved Intact Deciduous Teeth

Bressan Werle S*/ Lindemann D**/ Machado J***/ Pranke P****/ Casagrande L*****

Objectives: To isolate cells from pulp of intact cryopreserved deciduous teeth. The null hypothesis raised here is to find no difference in the establishment of cell culture after cryopreservation (1) using culture medium supplemented with different concentrations of fetal bovine serum (FBS); and (2) between teeth with different stages of physiological root resorption. **Study design:** Intact deciduous teeth with different root resorption stages were cryopreserved using FBS and Dimethyl Sulfoxide (DMSO) medium (9:1) in a progressive freezing process, by placing the samples in the refrigerator (4°C/60 min) and subsequently transferring them to a -80°C freezer (controlled device -1°C/min/24 hours), and finally into liquid nitrogen (-196°C/30 days). After the thawing process, the cell isolation was performed by enzymatic digestion (type I collagenase). The cells were re-suspended into the culture medium with 10% (G1) or 20% (G2) of FBS. Microscopic analysis was performed after 30 days to visualize the cell attachment. **Results:** The culture establishment rate was higher in G2 (75%) than G1 (12.5%) ($p=0.041$). There was no difference between the different stages of root resorption. **Conclusions:** It was possible to establish cell cultures from the pulp of intact cryopreserved deciduous teeth. The use of 20% FBS after thawing improved the culture rate.

Key words: Deciduous teeth; Cryopreservation; Fetal Bovine Serum.

INTRODUCTION

Stem cells from deciduous human teeth exhibit a high proliferation rate and differentiation potential in multiple lineage, which promotes their potential use in research and pre-clinical trials^{1,2}. The option for these cells is also attractive because their obtainment causes minimal discomfort to the donor once deciduous teeth are physiologically exfoliated and pulp tissue, a stem cell niche, is usually discarded³. The exfoliation process occurs usually between 5 and 13 years of age, thus, after this period it is no longer possible to obtain these cells.

The culture and expansion of stem cells after their isolation would be one method to maintain them for a long-term period.

However, there are some complications, such as the decrease in the ability of differentiation, senescence, induced by consecutive passages⁴, the possibility of genetic alterations⁵ and the economic cost involved⁶. Thus, cryopreservation is presented as an alternative to maintain the viability of these cells until their use is required, allowing for long-term storage and safety for clinical application⁵.

Recent studies suggest that the cryopreservation of hDPSC (human dental pulp stem cells) is viable⁷. These hDPSC maintain the surface markers unaltered and they also have proliferation and differentiation potential^{4,8,9}, even when cryopreserved for long periods^{5,10}. Moreover, when the initial isolation processing is not possible, some studies have demonstrated the freezing capacity of the whole permanent tooth for future isolation of hDPSCs¹¹⁻¹⁴, showing the culture rates in a range of 20-70%.

Most of these studies utilized a culture medium supplemented with 15-20% of fetal bovine serum (FBS)^{4,5,9,13,14} or fetal calf serum (FCS)^{8,10} for the culture and expansion of the cells subjected to cryopreservation. Animal serum is an important additive for cell growth and essential for the early stages of isolation¹⁵. However, the use of lower concentrations or the absence of animal serum in the culture medium would be ideal because it can transmit zoonosis and it contains unspecific growth factors, reducing the possibilities for clinical application¹⁶.

The hDPSCs from deciduous teeth have the advantage of presenting higher proliferation rates and higher differentiation potential when compared to hDPSCs from permanent teeth¹⁷. However, few studies have reported the cryopreservation viability of these cells^{4,18} and there is still an absence in the literature regarding the cryopreservation of intact deciduous teeth without initial isolation processing.

The purpose of this study was to isolate cells from cryopreserved

From the Federal University of Rio Grande do Sul, Porto Alegre, Brazil

*Stefanie Bressan Werle, DDS, MS, PhD Student, Department of Pediatric Dentistry, School of Dentistry.

**Daniele Lindemann, DDS, MS Student, Department of Pediatric Dentistry, School of Dentistry.

***Janine Machado, Undergraduate Student, School of Pharmacy.

****Patricia Pranke, MS, PhD, Associate Professor, Head of Hematology and Stem Cell Laboratory and Stem Cell Research Institute, School of Pharmacy.

*****Luciano Casagrande, DDS, MS, PhD, Adjunct Professor, Department of Pediatric Dentistry, School of Dentistry.

Send all correspondence to Prof. Luciano Casagrande School of Dentistry, Pediatric Dentistry, Federal University of Rio Grande do Sul Ramiro Barcelos 2492, Porto Alegre, RS; ZIP: 90035-003, Brazil Phone: +(51)33085493

Fax: +(51)33085010

E-mail: luciano.casagrande@ufrgs.br


deciduous teeth and test the null hypothesis that there is no difference in the establishment of cell culture after cryopreservation of intact deciduous teeth using: (1) culture medium supplemented with different concentrations of FBS and (2) teeth at different root resorption stages.

MATERIALS AND METHOD

Sound deciduous teeth (n=16) with absence of trauma and no clinical or radiographic signs of pulpal disease, presenting different root resorption stages (Figure 1), were collected from patients aged from 7 to 13. Before tooth extraction, the parents were fully informed about the procedure and destination of the tooth, and signed a consent form approved by the Ethics Committee of Federal University of Rio Grande do Sul (n. 20865). The tooth extraction decision was part of the orthodontic treatment plan design for the patients by their dentists who were not related to this research.

Immediately after extraction, the teeth were placed in a flask with transport medium containing DMEM (Dulbecco’s modified Eagle medium – Sigma- Aldrich, St. Louis, MO) / HEPES, 10% fetal bovine serum (Laborclin, Pinhais, Brazil), 100U/mL penicillin, 100mg/mL streptomycin and 0.45mg/mL gentamicin (Gibco, Grand Island, NY) and transported to the laboratory within 6 hours. Under sterile conditions, the teeth were removed from the flask and cleaned with gauze and 0.12% chlorhexidine solution. The teeth were randomized into two groups: 10% FBS (supplemented medium containing 10% FBS for culture after cryopreservation) or 20% FBS (supplemented medium containing 20% FBS for culture after cryopreservation).

Figure 1. Root resorption stages according to groups: a) total root resorption; b) 1/3 root remaining and c) 2/3 root remaining.



Freezing

The deciduous teeth were displaced in a 2mL vial containing 1.5 mL cryoprotectant solution of 10% dimethylsulfoxide and 10% FBS (1:9), and kept in a 4°C environment for 60 minutes. Following, the vials were subjected to a dump-freeze method consisting of a vial suspension in an isopropanol bath (Mr. Frosty – NALGENE TM Cryo 10°C freezer Container, Cat. No 5100-0001) within a -80°C mechanical freezer for a -1C/min cooling rate for 24 hours. The following day, the vials were placed into -196°C liquid nitrogen.

Thawing

After 30 days of cryopreservation, the vials were recovered and partially thawed for less than 1 minute in a 37°C water bath. Once out of the vial, the teeth were washed with cold culture medium supplemented with 10 or 20% FBS (DMEM/HEPES, 10 or 20% FBS, 1 or 2% antibiotics) according to the groups.

Isolation and Cell Culture

For both groups, all the pulp tissue was removed from the dentin walls with a dentin excavator, placed in buffer containing 0.2% type I collagenase solution (Gibco, Grand Island, NY) for 60 minutes in a 37°C bath and centrifuged (10 min, 800 G). The remaining pellet was re-suspended and seeded onto a 12 well plate. The first medium change occurred after 72 hours and successively every 3 to 4 days. The identification of an agglomerate of cells within a period of 30 days was considered criteria for the success of the culture establishment. When the cells were not present after 30 days, the culture was inactivated and registered as a failure.

Statistical Analysis

The chi-square test was used to verify statistical differences in the establishment of the culture between serum groups and between resorption root stages. The statistical analysis was performed using the Statistical Package for Social Sciences version 20.0 (SPSS Inc., Chicago, IL, USA, 2011) at 5% significance level.

RESULTS

The success rate in establishing cell cultures was statistically significantly higher (p= 0.041) in 20% FBS group (Table 1). Figure 2 shows a microscopic view (40x) of the cells growing in culture after cryopreservation of the deciduous teeth.

Table 2 shows no differences in the culture rate, according to the different resorption stages of the deciduous teeth.

Table 1. Establishment of cell culture according to concentration of Fetal Bovine Serum in the culture medium.

	Culture success			%	p
	no	yes	total		
10% FBS	7	1	8	12.5	0.041
20% FBS	2	6	8	75	
Total	9	7	16		


Fisher’s exact test (p<0.05)

Table 2. Cell culture establishment regarding the root resorption stage of deciduous teeth.

		Culture success		p
		yes	total	
Root resorption stage	2/3 root remaining	3	5	0.66
	1/3 root remaining	2	5	
	Total root resorption	2	6	
	Total	7	16	

Pearson’s Chi-square (p>0.05)

Figure 2. A) Microscopic view (40x) of cell attachment (arrow) from intact cryopreserved deciduous tooth. B) Microscopic view (40x) of a confluence culture.



DISCUSSION

The dental pulp of deciduous tooth is a niche of stem cells and therefore, a potential source for tissue engineering². Usually, children lose their primary teeth when the permanent successor tooth starts the eruption process and this cell niche is then discarded³.

The results of this study have demonstrated the capacity of the establishment of cell culture from human intact deciduous teeth subjected to cryopreservation for 30 days. The success rate for the establishment of the culture varied according to the concentration of FBS used in the culture medium, rejecting the first null hypothesis. The different stages of root resorption of the deciduous teeth demonstrated no difference in cell culture establishment, so the second null hypothesis was confirmed.

Although the isolation of stem cells from pulp of deciduous teeth is related to the resorption process¹⁹, it seems that the different stages of resorption considered in this study did not affect the culture rate after cryopreservation. However, alterations of cell activities, such as proliferation and differentiation potential, may occur, and were not evaluated in this study.

When the Stem Cells from Human Exfoliated Deciduous teeth (SHED) were evaluated, Zhu and colleagues (2013) observed higher proliferation and mineralization potential in the final stage of root resorption compared to the initial stage. This data could be justified by the changes in the microenvironment, due to the resorption process as well as the activation of the stem cells during the repair process, which occurs in the final stage.

The 20% FBS group showed a 75% cell culture rate, corroborating with other studies in which permanent teeth were

cryopreserved intact¹¹⁻¹³, whereas 10% FBS group only showed a 12.5% cell culture rate, six times less. Chen and colleagues (2010) also reported only 20% of cell culture from cryopreserved permanent teeth; however the sample was comprised of teeth with pathological root resorption, inflammation, and periodontitis.

During the cryopreservation process, the formation of ice crystals in the cell cytoplasm leads to osmotic imbalance and possible dehydration²¹. In order to prevent this phenomenon, cryoprotectants are added to the freezing media⁸. When the pulp is frozen inside the hard tissue of the tooth, insulation occurs, which is caused by the dentin and enamel layers⁶ and the cryoprotectant only penetrates through the apical foramen or the area of resorption. In this case, dehydration can occur even with the use of cryoprotectants and a medium with more supplements (20% FBS) can support the cell growth.

Recent study demonstrated cell encapsulation and thawing techniques as an alternative to optimize the viability of cells from the pulp of deciduous teeth after cryopreservation¹⁸. However, when there is no time for initial processing, which is important when considering the natural exfoliation of the deciduous teeth, freezing the whole tooth may be an alternative for storage of the cells but it has not yet been studied.

Another technique demonstrated by Giuventù and colleagues (2011), utilized the laser to create micro-channels on non-exfoliated deciduous teeth roots. These teeth were subjected to freezing for 10 days at -80°C and showed a higher isolation success rate compared to the control (no laser). However, it is important to search for a method that does not require any prior processing for cryopreservation in liquid nitrogen (-196°C). Moreover, the use of laser to improve the establishment of cell culture involves time and high cost.

Thus, the use of the supplementation like recommended by the initial studies of characterization of dental pulp stem cell, with animal serum in higher concentrations (20% FBS)^{1,22}, is a way for optimizing cell isolation from intact cryopreserved deciduous teeth and can be used as an improvement in the process. The idea of banking stem cells for research from cryopreserved deciduous teeth may be considered because of the success rate of 75% in isolation and the recoverability of cells in all root resorption stages.

However, a significant number of stem cells are necessary for therapeutic purposes and in this study the proliferative capacity and differentiation potential of these cultures were not evaluated. Moreover, although stem cells from dental pulp cultured with animal serum have been used successfully in humans²³, there is a risk of immune response to animal products¹⁵.

According to the Mesenchymal and Tissue Stem Cell Committee of the International Society for Cellular Therapy, for any cells to be characterized as mesenchymal stem cells, the following criteria are necessary: (1) adherence to plastic, (2) presence of surface markers for mesenchymal cells (CD73, CD90 and CD105), and (3) differentiate into osteoblasts, chondroblasts and adipocytes²⁴. The present study is limited to the isolation of cells with plastic adherence similar to the stem cells from deciduous teeth isolated and characterized by a previous study from this laboratory¹⁹. Thus, future studies are required for the immunophenotypic characterization of cells from pulp of intact cryopreserved deciduous teeth, in relation to viability, differentiation potential and proliferation rate.

CONCLUSION

- It was possible to establish cell culture from intact cryopreserved teeth when using culture medium supplemented with 20% FBS.
- The establishment of pulp cell culture was not influenced by different stages of root resorption in deciduous teeth. Additional experiments will be required to establish the identity and usefulness of the cells obtained by this method.

REFERENCES

- Miura M, Gronthos S, Zhao M, Lu B, Fisher LW, Robey PG, Shi S. SHED: stem cells from human exfoliated deciduous teeth. *Proc Natl Acad Sci USA* 100: 5807-12, 2003.
- Kawashima N. Characterisation of dental pulp stem cells: a new horizon for tissue regeneration? *Arch Oral Biol* 11: 1439-58, 2012.
- Arora V, Arora P, Munshi AK. Banking stem cells from human exfoliated deciduous teeth (SHED): saving for the future. *J Clin Pediatr Dent Summer* 33: 289-94, 2009.
- Lizier NF, Kerkis A, Gomes CM, Hebling J, Oliveira CF, Caplan AI, Kerkis I. Scaling-up of dental pulp stem cells isolated from multiple niches. *PLoS One* 7: e39885, 2012.
- Ma L, Makino Y, Yamaza H, Akiyama K, Hoshino Y, Song G, Kukita T, Nonaka K, Shi S, Yamaza T. Cryopreserved dental pulp tissues of exfoliated deciduous teeth is a feasible stem cell resource for regenerative medicine. *PLoS One* 7: e51777, 2012.
- Gioventù S, Andriolo G, Bonino F, Frasca S, Lazzari L, Montelatici E, Santoro F, Rebullà P. A novel method for banking dental pulp stem cells. *Transfus Apher Sci* 47: 199-206, 2012.
- Collart Dutilleul PY, Thonat C, Jacquemart P, Cuisinier F, Levallois B, Chaubron F. Dental pulp stem cells: characteristics, cryopreservation and therapeutic potentialities. *Orthod Fr* 83: 209-16, 2012.
- Zhang W, Walboomers XF, Shi S, Fan M, Jansen JA. Multilineage differentiation potential of stem cells derived from human dental pulp after cryopreservation. *Tissue Eng* 12: 2813-23, 2006.
- Yalvaç ME, Ramazanoglu M, Tekguc M, et al. Human Tooth Germ Stem Cells Preserve Neuro-Protective Effects after Long-Term Cryo-Preservation. *Curr Neurovasc Res* 7: 49-58, 2010.
- Papaccio G, Graziano A, d'Aquino R, Graziano MF, Pirozzi G, Menditti D, De Rosa A, Carinci F, Laino G. Long-term cryopreservation of dental pulp stem cells (SBP-DPSCs) and their differentiated osteoblasts: a cell source for tissue repair. *J Cell Physiol* 208: 319-25, 2006.
- Perry BC, Zhou D, Wu X, Yang FC, Byers MA, Chu TM, Hockema JJ, Woods EJ, Goebel WS. Collection, cryopreservation, and characterization of human dental pulp-derived mesenchymal stem cells for banking and clinical use. *Tissue Eng Part C Methods* 14: 149-56, 2008.
- Woods EJ, Perry BC, Hockema JJ, Larson L, Zhou D, Goebel WS. Optimized cryopreservation method for human dental pulp-derived stem cells and their tissues of origin for banking and clinical use. *Cryobiology* 59: 150-7, 2009.
- Lee SY, Chiang PC, Tsai YH, Tsai SY, Jeng JH, Kawata T, Huang HM. Effects of cryopreservation of intact teeth on the isolated dental pulp stem cells. *J Endod* 36: 1336-40, 2010.
- Chen YK, Huang AH, Chan AW, Shieh TY, Lin LM. Human dental pulp stem cells derived from different cryopreservation methods of human dental pulp tissues of diseased teeth. *J Oral Pathol Med* 40: 793-800, 2011.
- Hirata TM, Ishkitiev N, Yaegaki K, Calenic B, Ishikawa H, Nakahara T, Mitev V, Tanaka T, Haapasalo M. Expression of multiple stem cell markers in dental pulp cells cultured in serum-free media. *J Endod* 36: 1139-44, 2010.
- Ishkitiev N, Calenic B, Aoyama I, Ii H, Yaegaki K, Imai T. Hydrogen sulfide increases hepatic differentiation in tooth-pulp stem cells. *J Breath Res* 6: 017103, 2012.
- Wang X, Sha XJ, Li GH, et al. Comparative characterization of stem cells from human exfoliated deciduous teeth and dental pulp stem cells. *Arch Oral Biol* 57: 1231-40, 2012.
- Umemura E, Yamada Y, Nakamura S, Ito K, Hara K, Ueda M. Viable cryopreserving tissue-engineered cell-biomaterial for cell banking therapy in an effective cryoprotectant. *Tissue Eng Part C Methods* 17: 799-807, 2011.
- Bernardi L, Luisi SB, Fernandes R, et al. The isolation of stem cells from human deciduous teeth pulp is related to the physiological process of resorption. *J Endod* 37: 973-9, 2011.
- Zhu Y, Shang L, Chen X, et al. Deciduous dental pulp stem cells are involved in osteoclastogenesis during physiologic root resorption. *J Cell Physiol* 228: 207-15, 2013.
- Janz Fde L, Debes Ade A, Cavaglieri Rde C, Duarte SA, Romão CM, Morón AF, et al. Evaluation of distinct freezing methods and cryoprotectants for human amniotic fluid stem cells cryopreservation. *J Biomed Biotechnol* 2: 649353, 2012.
- Gronthos S, Mankani M, Brahimi J, Robey PG, Shi S. Postnatal human dental pulp stem cells (DPSCs) in vitro and in vivo. *Proc Natl Acad Sci USA* 97: 13625-30, 2000.
- d'Aquino R, De Rosa A, Lanza V, et al. Human mandible bone defect repair by the grafting of dental pulp stem/progenitor cells and collagen sponge biocomplexes. *Eur Cell Mater* 12: 75-83, 2009.
- Dominici M, Le Blanc K, Mueller I, et al. Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. *Cytotherapy* 8: 315-7, 2006.