

Mitogenome of the Monotypic Genus *Rhaetus* (Coleoptera: Scarabaeidae: Lucanidae)¹

Liu Jing, Shi-Ju Zhou, Yong-Jing Chen, and Xia Wan²

Department of Ecology, School of Resources and Engineering, Anhui University, Anhui, Hefei, 23000, China

J. Entomol. Sci. 53(4): 503–513 (October 2018)

Abstract *Rhaetus westwoodi* Parry (Coleoptera: Lucanidae) is the type species of monotypic genus *Rhaetus* Parry, which is distributed only in the region of the Eastern Himalayas. Little is known about the biology, ecology, genetics, and conservation status of this rare beetle. In this study, the mitochondrial genome of *R. westwoodi* was sequenced for the first time to obtain useful genetic data for this monotypic genus. Like most of known mitogenome in beetles, the mitochondrial genome of *R. westwoodi* is a closed circular molecule 18,134 bp in length, comprising 13 protein-coding genes, 22 transfer RNA (tRNAs) genes, 2 ribosomal RNAs (rRNAs), and 1 control region. The nucleotide composition is 36.30% adenine (A), 31.20% thymine (T), 21.49% cytosine (C), and 11.01% glycine (G), with a highly biased AT content of 67.50%. The phylogenetic analysis showed that *R. westwoodi* and *Hexarthrius vitalis* Tsukamotoi were recovered as one clade with strong nodal support (1.00 Bayesian Phylogenetics and Phylogeography [BPP], 100% maximum likelihood bootstrap [MLB]). The topology also strongly supported that *R. westwoodi* appeared to be a sister species to *Pseudorhaetus sinicus* Boileau (1.00 BPP, 100% MLB).

Key Words monotypic genus, *Rhaetus westwoodi*, complete mitochondrial genome, phylogenomic analysis

Rhaetus Parry is a monotypic genus with the type species of *R. westwoodi* Parry from Darjeeling, N. India (Parry 1862). As the monotype of *Rhaetus*, *R. westwoodi* is an amazing beetle due to its large size, shiny black body, and striking mandibles in males.

Nagai (2000) published the subspecies *R. westwoodi kazumiae* from North Myanmar, with the description of a large male. This male had a longer medial tooth, possibly distinguishing it from the typical *R. westwoodi* male. However, some large specimens of *R. westwoodi* from N. India also had the same mandibular characteristics of this reported subspecies and are depicted in the illustration of Didier and Séguy (1952). The mandibular “difference” of the subspecies, in our opinion, should be regarded as intraspecific variation among males of *R. westwoodi*. Therefore, in this study, we treated *R. westwoodi* as the only valid taxon in *Rhaetus*.

¹Received 11 November 2017; accepted for publication 6 December 2017.

²Corresponding author (email: wanxia@ahu.edu.cn).

Table 1. PCR primers.

Gene	Primer Name	Sequence (5'–3')	Length (bp)	Reference
<i>COI</i>	<i>COI-F1</i>	CAACATTTATTTTGATTTTTTGG	23	Simon et al., 1994
	<i>COI-R1</i>	TCCAATGCACTAATCTGCCATATTA	25	Simon et al., 1994
<i>CYTB</i>	<i>CYTB-F2</i>	GAGGAGCAACTGTAATTACTAA	22	Balke et al., 2004
	<i>CYTB-R2</i>	AAAAGAAARTATCATTGAGGTTGAAT	26	Balke et al., 2004
<i>rRNAI</i>	<i>16SF1</i>	CCGGTTTGAAGCTCAGATCATG	21	Hosoya et al., 2001
	<i>16SR1</i>	TAATTTATTGTACCTTGTGTATCAG	25	Hosoya et al., 2001

Little is known about the biology, ecology, and conservation status of *R. westwoodi*. The species is limited in its distribution to the high mountain forests (elevations averaging over 1,500 m above sea level) of the Eastern Himalayas (N. India, N. Myanmar, Tibet, N. W. Yunnan) (Arrow 1943, 1950; Benesh, 1960; Didier and Séguéy 1952; Fujita 2010; Huang and Chen 2013; 2017; Krajcik 2001; Maes 1992; Parry 1862; Wan et al. 2010). Unfortunately, due to their fascinating appearance, specimens of this species have been much in demand as valuable collections, with an asking price of US\$450 to 1,999 per large male (>80 mm in length) and US\$90 to 350 per small male (approximately 60 mm in length). With this market, individuals are collecting the beetles for financial returns and, thus, may overhunt *R. westwoodi* in its restricted habitats to the point that the species may become increasingly endangered. In this study, we sequenced its complete mitogenome for the first time to provide genetic data for future conservation work and for establishing the phylogenetic relationships of *R. westwoodi* with other beetles.

Materials and Methods

Sample collection and DNA extraction. The voucher specimen of *R. westwoodi* was collected from Motuo, Tibet, in August 2006 by Ming Bai and deposited in the Museum of Anhui University (Hefei, China). Total genomic DNA was extracted from the muscle of *R. westwoodi* by using the Qiagen DNeasy Kit (Qiagen, Shanghai, China).

Polymerase chain reaction (PCR) amplification and sequencing. Complete mitogenomes were assembled from amplified fragments, with all primers used for amplification listed in Table 1. PCR amplification reactions were conducted in 25- μ l volumes containing 10 μ M of each primer (forward and reverse), 2 μ l template DNA,

12.5 μL $2\times$ EasyTaq SuperMix (+dye), and 8.5 μL sterile double-distilled water to a final volume of 25 μL . PCR amplifications were performed under the following conditions: an initial denaturation at 94°C for 2 min, followed by 35–37 cycles of denaturation at 94°C for 40 s, annealing at 52–58°C for 50 s, elongation at 70°C for 1 min, and then a final extension step at 72°C for 7 min. The temperature of annealing was determined by the length of the fragments. Sequencing was conducted with the Illumina HiSeq 2000 platform. Cluster strands created by bridge amplification were primed, and all four fluorescently labeled and 3-OH blocked nucleotides were added to the flow cell with DNA polymerase. The cluster strands were extended in single nucleotides. Following the incorporation step, the unused nucleotides and DNA polymerase molecules were rinsed away, a scan buffer was added to the flow cell, and then the optics system scanned each lane of the flow cell in imaging units (tiles). Once imaging was completed, chemicals that affect cleavage of the fluorescent labels and the 3-OH blocking groups were added to the flow cell, which prepares the cluster strands for another round of fluorescent nucleotide incorporation. The sequence was submitted to GenBank and assigned accession number MG159815.

Mitogenome assembly, annotation, and analysis. The mitogenomes were assembled using SOAPdenovo (BGI Company, Shenzhen, China), and preliminary annotations were made with the MITOS WebServer (<http://mitos.bioinf.uni-leipzig.de/index.py>). tRNA genes and their secondary structures were inferred using tRNAscan-SE version 1.21 (<http://lowelab.ucsc.edu/tRNAscan-SE/>). Those not identified by tRNAscan-SE, in addition to 16S rRNA (*rrnL*, *lrRNA*), and 12S rRNA (*rrnS*, *srRNA*), were determined according to sequence similarity with related species. The protein-coding genes (PCGs) were determined by ORF Finder (<http://www.ncbi.nlm.nih.gov/gorf/gorf.html>) under the invertebrate mitochondrial genetic code. Nucleotide compositions, codon usage, and relative synonymous codon usage (RSCU) values of PCGs were calculated with MEGA version 6.05 (Tamura et al. 2013). PCGs were translated with DNAMAN version 7.0.2.176 (Lynnsoft Biosoft, Vaudreuil-Dorion, Canada). Mitogenomes were mapped with CGView (Grant and Stothard 2008). Composition skew analysis was conducted according to the formulas $\text{AT skew} = \frac{[A - T]}{[A + T]}$ and $\text{GC skew} = \frac{[G - C]}{[G + C]}$ (Perna and Kocher 1995).

Phylogenetic analyses. In this study, we retrieved 14 mitogenome sequences from GenBank (Table 2) and, to this data set, we added a newly sequenced *R. westwoodi*, generating a dataset of 15 taxa. Each of the PCGs was aligned individually, based on codon-based multiple alignments by using the MEGA version 7.0.26. Models of nucleotide substitution were selected according to the Akaike Information Criterion with jModelTest version 2.1.4 (Posada 2008). Phylogenetic trees were generated from maximum likelihood (ML) analysis using RAxML (Gillett et al. 2014) and Bayesian inference (BI) with MrBayes version 3.2.5 (Huelsenbeck and Ronquist 2013), both under the GTR + I + G model. Node support in the ML tree was estimated through bootstrap analysis with 1,000 replicates. The BI was conducted with two simultaneous Markov chain Monte Carlo runs of 2 million generations, sampled every 1,000 steps, with the first 25% discarded as burn-in. Phylogenetic trees were viewed and edited in Figtree version 1.4.3. (Rambaut 2016).

Table 2. Coleopteran mitogenomes.

Species	GenBank Accession Number
<i>Rhaetus westwoodi</i>	MG159815
<i>Ageus angustus</i>	JX313668
<i>Lucanus mazama</i>	NC013578
<i>Dorcus parallelipedus</i>	JX412841
<i>Hemisodorcus rubrofemoratus</i>	JX313682
<i>Prosopocoilus gracilis</i>	KP735805
<i>Prosopocoilus blanchardi</i>	KF364622
<i>Prosopocoilus confucius</i>	KU552119
<i>Hexarthrius vitalis</i>	JX313676
<i>Pseudoechthistatus sinicus</i>	KP987575
<i>Nigidious parryi</i>	KP987576
<i>Sinodendron yunnanense</i>	KP735804
<i>Cheirotonus jansonii</i>	NC023246
<i>Protaetia brevitarsis</i>	NC023453
<i>Rhopaea magnicornis</i>	NC013252

Results and Discussion

Mitogenome. The complete mitogenome of *R. westwoodi* (GenBank accession number MG159815) measured 18,134 bp in length as a closed circular molecule (Fig. 1), consisting of 22 tRNA genes, two rRNA genes (*rrn-L* and *rrn-S*), 13 PCGs, and 1 control region. Four PCGs, 2 rRNAs, and 8 tRNAs were located on the N-strand, whereas the other 9 PCGs and 14 tRNAs were located on the J-strand (Table 3). Among the 13 PCGs, only 4 genes (*nad4*, *nad4l*, *nad5*, *nad1*) were located on the N-strand, whereas the other 9 (*cox1*, *cox2*, *cox3*, *atp8*, *atp6*, *nad2*, *nad3*, *nad6*, *cytb*) were located on the J-strand. These 13 PCGs comprised 3,705 codons, with the largest being the *nad5* gene (1,806 bp) and the smallest the *atp8* gene (155 bp). The average AT content of the PCGs was 65.65%, with the highest in *nad4l* (70.92%) and the lowest in *cox1* (61.33%) (Table 4). All PCG codons started with ATN except *cox1*, which started with AAT (Table 3). Nine PCGs ended with TAA and TAG codons, with the incomplete stop codons TA and T in the remaining genes. In general, incomplete codon structures signal a halt of protein translation in insects and other invertebrates (Cheng et al. 2016, Masta and Boore 2004, Wu et al. 2014).

The RSCU analysis showed a biased usage of A and T nucleotides in the genome (Fig. 2). The most frequently encountered codons in other insects have been UUA (Leu), UUU (Phe), and AUU (Ile) (Wei et al. 2016). The PCGs coded on

Table 3. Annotation for *Rhaetus westwoodi*.

Gene	Strand	Region	Length (bp)	Start Codon	Stop Codon	Anticodon	Intergenic Nucleotides (bp)
<i>trnI</i>	J	1–63	63	-	-	GAT	-1
<i>trnQ</i>	N	61–129	68	-	-	TTG	0
<i>trnM</i>	J	129–197	68	-	-	CAT	0
<i>nad2</i>	J	198–1211	1013	ATA	TAG	-	2
<i>trnW</i>	J	1214–1278	64	-	-	TCA	-8
<i>trnC</i>	J	1271–1331	60	-	-	GCA	0
<i>trnY</i>	N	1332–1394	62	-	-	GTA	1
<i>cox1</i>	J	1396–2931	1535	AAT	TAA	-	-5
<i>trnL(CUN)</i>	J	2927–2989	62	-	-	TAA	1
<i>cox2</i>	J	2991–3678	687	ATT	T	-	0
<i>trnK</i>	J	3679–3749	70	-	-	CTT	1
<i>trnD</i>	J	3749–3811	62	-	-	GTC	0
<i>atp8</i>	J	3812–3967	155	ATC	TAA	-	-7
<i>atp6</i>	J	3961–4632	671	ATG	TAA	-	-7
<i>cox3</i>	J	4632–5415	783	ATG	T	-	0
<i>trnG</i>	J	5416–5479	63	-	-	TCC	-12
<i>nad3</i>	J	5468–5833	365	ATA	ATG	-	-2
<i>trnA</i>	J	5832–5896	64	-	-	TGC	-1
<i>trnR</i>	J	5896–5959	63	-	-	TCG	-1
<i>trnN</i>	J	5959–6022	63	-	-	GTT	0
<i>trnS(AGN)</i>	J	6023–6089	67	-	-	TCT	0
<i>trnE</i>	J	6090–6153	63	-	-	TTC	-2
<i>trnF</i>	N	6152–6215	63	-	-	GAA	0
<i>nad5</i>	N	6216–7932	1806	ATT	T	-	0
<i>trnH</i>	N	7933–7996	63	-	-	GTG	0
<i>nad4</i>	N	8001–9332	1331	ATG	T	-	-7
<i>nad4l</i>	N	9325–9607	282	ATA	TAA	-	8
<i>trnT</i>	J	9619–9682	63	-	-	TGT	0
<i>trnP</i>	N	9683–9747	64	-	-	TGG	5
<i>nad6</i>	J	9750–10247	497	ATG	TAA	-	-1

Table 3. Continued.

Gene	Strand	Region	Length (bp)	Start Codon	Stop Codon	Anticodon	Intergenic Nucleotides (bp)
<i>cob</i>	J	10247–11389	1142	ATG	TAG	-	-2
<i>trnS(UCN)</i>	J	11391–11455	64	-	-	TGA	19
<i>nad1</i>	N	11472–12425	953	ATA	TAG	-	-3
<i>trnL(CUN)</i>	N	12426–12488	62	-	-	TAG	-3
<i>rrnL</i>	N	12486–13753	1267	-	-	-	1
<i>trnV</i>	N	13755–13823	68	-	-	TAC	-1
<i>rrnS</i>	N	13823–14585	762	-	-	-	0
Control region	-	14586–18131	3545	-	-	-	0

L2 and *tRNA-V*, whereas the *rRNA-S* gene (AT content, 68.94%) was between *tRNA-V* and the AT-rich region.

The mitogenome was found to have a 7-bp conserved sequence (TTGTTCA) in the upstream of *ND1*, which has been observed in other beetles and insects (Cameron and Whiting 2008, Cameron 2014b, Kim et al. 2015, Lin et al. 2017, Sheffield et al. 2009, Yang et al. 2016). Intergenic spacers appeared frequently in the mitogenome, were interspersed throughout the PCGs and RNA genes, and ranged from 1 to 97 bp in length (Table 3).

The AT-rich region was 3,545 bp in length (70.87% AT) and was locked between *rRNA-S* and *tRNA-M*. A 16-bp poly-TA was locked in 17,079 bp downstream of the AT-rich region, whereas a 12-bp poly-T, 10-bp poly-A, and 21-bp poly-A were locked in 437 bp, 750 bp, and 696 bp upstream, respectively, of the *tRNA-M*.

Phylogenetic analysis. The phylogenetic relationships of Scarabaeoidea were reconstructed based on the concatenated nucleotide sequences of 11 PCGs from 12 beetles as in-groups and other three scarabs as out-groups (14 of them downloaded from GenBank) (Table 2). BPP values ≥ 0.95 were considered as




Fig. 2. The RSCU of *R. westwoodi*.




Fig. 3. The ML and BI phylogenetic trees of *R. westwoodi* and 14 other species based on 11 PCGs. GTR + G and GTR + I + G were selected as the best models, and with three scarab species as outgroups.

strong support, whereas BPP values ≤ 0.75 as weak support, and the ML bootstrap (MLB) values $\geq 75\%$ as strong support, MLB values between 50% and 75% as moderate support, and $\leq 50\%$ as weak support (Kim et al. 2015).

The topologies of the BI and ML analyses were highly consistent (Fig. 3). *Rhaetus westwoodi* and *Hexarthrus vitalis* Tsukamotoi were recovered as one clade with strong nodal value (1.00 BPP, 100% MLB). The two species appeared to be a sister group to *Pseudorhaetus sinicus* Boileau, with robust support (1.00 BPP, 100% MLB). The results corresponded to the morphological work of Huang and Chen (2013) and the latest molecular study inferred from the fragments (Wu and Wan 2016). Furthermore, these three species were in the tribe Dorcini first described by Parry (1864) with the same relationships we described herein. In contrast, the resolution did not agree with their relationships that were shown by Bartolozzi and Sprecher-Uebersax (2006), Benesh (1960), and Krajcik (2001), in which *R. westwoodi* and *P. sinicus* were in the tribe Rhaetulini and *H. vitalis* was in the tribe Lucaini.

Table 4. The skew of *Rhaetus westwoodi*.

Genes	T(U)	C	A	G	A+T	G+C	AT-Skew	GC-Skew
PCGs	37.90	18.35	27.46	16.29	65.36	34.64	-0.16	-0.06
tRNA	34.51	12.82	36.78	15.90	71.28	28.72	0.03	0.11
rRNA	39.78	8.22	31.12	20.88	70.90	29.10	-0.12	0.43
A+T rich region	34.04	19.67	36.83	9.47	70.87	29.13	0.04	-0.35
Mitogenome	31.20	21.49	36.30	11.01	67.50	32.50	0.08	-0.32

Acknowledgments

We are grateful to Ming Bai (Institute of Zoology, Chinese Academy of Sciences) for donation specimens, Hu Li and Fan Song (Department of Entomology, China Agricultural University) for statistical advice, and laboratory colleagues for their kind help. This study was supported by the National Natural Science Foundation of China (no. 31201745, no. 31071954, and no. 31572311). Thanks for the funding provided by the National Natural Science Foundation.

References Cited

- Arrow, G.J. 1943.** On the genera and nomenclature of the lucanid Coleoptera, and descriptions of a few new species. *Proc. Royal Entomol. Soc. London (B)*. 12: 133–143.
- Arrow, G.J. 1950.** The Fauna of India Including Pakistan, Ceylon, Burma and Malaya. Coleoptera, Lamellicornia, Lucanidae and Passalidae. Volume IV. Taylor & Francis Ltd., London.
- Balke, M., I. Ribera and A.P. Vogler. 2004.** MtDNA phylogeny and biogeography of Copelatinae, a highly diverse group of tropical diving beetles (Dytiscidae). *Mol. Phy. Evol.* 32: 866–880.
- Bartolozzi, L. and E. Sprecher-Uebersax. 2006.** Lucanidae, Pp. 63–76. *In* Lobl I. and A. Smetana (eds.), *Catalogue of Palaearctic Coleoptera*, Vol. 3: Scarabaeoidea – Scirtoidea – Dascilloidea – Buprestoidea – Byrrhoidea. Apollo Books, Brooklyn, NY.
- Benesh, B. 1960.** Lucanidae. *Coleopterorum Catalogus Suppl.* 8: 1–178.
- Cameron, S.L. and M.F. Whiting. 2008.** The complete mitochondrial genome of the tobacco hornworm, *Manduca sexta*, (Insecta: Lepidoptera: Sphingidae), and an examination of mitochondrial gene variability within butterflies and moths. *Gene*. 408: 112–123.
- Cameron, S.L. 2014a.** Insect mitochondrial genomics: Implication for evolution and phylogeny. *Annu. Rev. Entomol.* 59: 95–117.
- Cameron, S.L. 2014b.** How to sequence and annotate insect mitochondrial genomes for systematic and comparative genomics research. *Syst. Entomol.* 39: 400–411.
- Cao, S.S. and Y.Z. Du. 2014.** Characterization of the complete mitochondrial genome of *Chilo auricilius* and comparison with three other rice stem borers. *Gene*. 548: 270–276.
- Cheng, X.F., L.P. Zhang, D.N. Yu, K.B. Storey and J.Y. Zhang. 2016.** The complete mitochondrial genomes of four cockroaches (Insecta: Blattodea) and phylogenetic analyses within cockroaches. *Gene*. 586: 115–122.
- Didier, R. and E. Séguéy. 1952.** Notes sur quelques espèces de Lucanides et descriptions de formes nouvelles. *Rev. Française d'Entomol.* 19: 220–233.
- Fujita, H. 2010.** The Lucanid Beetles of the World. Mushi-Sha, Japan, 6: 488 pp.
- Hosoya, T., M. Honda and K. Araya. 2001.** Genetic variations of 16S rRNA gene observed in *Ceruchus lignarius* and *Dorcus rectus rectus* (Coleoptera: Lucanidae). *Entomol. Sci.* 4: 335–344.
- Huang, H. and C.C. Chen. 2013.** Stag Beetles of China II. Formosa Ecological Company, Taiwan. 1–716 pp.
- Huang, H. and C.C. Chen. 2017.** Stag beetles of China III. Formosa Ecological Company, Taiwan. 1–524 pp.
- Huelsenbeck, J.P. and F.R. Ronquist. 2013.** MrBayes: A program for the Bayesian inference of phylogeny. *Biometrics*. 17: 754–755.
- Gillett, C.P., A. Crampton-Platt, M.J. Timmermans, B.H. Jordal, B.C. Emerson and A.P. Vogler. 2014.** Bulk de novo mitogenome assembly from pooled total DNA elucidates the phylogeny of weevils (Coleoptera: Curculionoidea). *Mol. Biol. Evol.* 31: 2223–2237.
- Grant, J.R. and P. Stothard. 2008.** The CGView Server: A comparative genomics tool for circular genomes. *Nucl. Acids Res.* 36: 181–184.

- Kim, M.J., K.G. Kim, S.R. Kim and I. Kim. 2013.** Complete mitochondrial genome of the two-spotted stag beetle, *Metopodontus blanchardi* (Coleoptera: Lucanidae). Mitochondr. DNA. 26: 307–309.
- Kim, S.I. and B.D. Farrell. 2015.** Phylogeny of world stag beetles (Coleoptera: Lucanidae) reveals a Gondwanan origin of Darwin's stag beetle. Mol. Phy. Evol. 86: 35–48.
- Krajcik, M. 2001.** Lucanidae of the World. Catalogue—Part I. Checklist of the stag beetles of the World (Coleoptera: Lucanidae). Published by the author, Plzen, Czech Republic. 108 pp.
- Lin, Z.Q., F. Song, T. Li, Y.Y. Wu and X. Wan. 2017.** New mitogenomes of two Chinese stag beetles (Coleoptera, Lucanidae) and their implications for systematics. J. Insect. Sci. 17: 63.
- Liu, J., Y.J. Chen, S.J. Zhou and X. Wan. 2017.** The first complete mitogenome of *Cyclommatus* stag beetles (Coleoptera: Lucanidae) with the phylogenetic implications. Entomol. 39: 294–299.
- Maes, J.M. 1992.** Lista de los Lucanidae del Mundo. R Nicaraguense Entomol. 22, A: 1–60; B: 61–121.
- Masta, S.E. and J.L. Boore. 2004.** The complete mitochondrial genome sequence of the spider *Habronattus oregonensis* reveals rearranged and extremely truncated tRNAs. Mol. Biol. Evol. 21: 893–902.
- Nagai, S. 2000.** Twelve new species, three new subspecies, two new status[es] and with the chick [check] list of the family Lucanidae of northern Myanmar. (Coleoptera: Lamellicornia). Not. Eur. Insect. 3: 73–108
- Parry, F.J.S. 1862.** Further descriptions and characters of undescribed lucanoid *Coleoptera*. Proc. Entomol. Soc. London. 3:107–113.
- Parry, F.J.S. 1864.** A catalogue of lucanoid Coleoptera; with illustrations and descriptions of various new and interesting species. Trans. Entomol. Soc. London. 2(3): 1–113.
- Perna, N.T. and T.D. Kocher. 1995.** Patterns of nucleotide composition at fourfold degenerate sites of animal mitochondrial genomes. J. Mol. Evol. 41: 353–358.
- Posada, D. 2008.** jModelTest: Phylogenetic model averaging. Mol. Biol. Evol. 25: 1253–1256.
- Rambaut, A. 2016.** FigTree: Tree figure drawing tool, Version 1.4.3. Inst. Evol. Biol. University of Edinburgh. <http://tree.bio.ed.ac.uk/>; accessed 16 June 2017.
- Sheffield, N.C., H.J. Song, S.L. Cameron and M.F. Whiting. 2009.** No stationary evolution and compositional heterogeneity in beetle mitochondrial phylogenomics. Syst. Biol. 58: 381–394.
- Simon, C., F. Frati, A. Beckenbach, B. Crespi, H. Liu and P. Flook. 1994.** Evolution, weighting, and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction primers. Ann. Entomol. Soc. Am. 87: 651–701.
- Tamura, K., G. Stecher, D. Peterson, A. Filipksi and S. Kumar. 2013.** MEGA6: Molecular evolutionary genetics analysis version 6.0. Mol. Biol. Evol. 30: 2725–2729.
- Wan, X., M. Bai, J.Z. Cui and X.K. Yang. 2010.** Six new species record of stag beetles in China. Zool. Syst. 35: 247–250.
- Wang, P., H. Li, Y. Wang, J.H. Zhang, X. Dai, J. Chang, B.W. Hu and W.Z. Cai. 2014.** The mitochondrial genome of the plant bug *Apolygus lucorum* (Hemiptera: Miridae): Presently known as the smallest in Heteroptera. Ins. Sci. 21:159–173.
- Wei, S.J., M. Shi, X.X. Chen, M.J. Sharkey, C. Achterberg, G.Y. Ye and J.H. He. 2010.** New views on strand asymmetry in insect mitochondrial genomes. PLoS One, 5: e12708.
- Wei, S.J., L.J. Cao, Q.L. Wu and C.D. Zhu. 2016.** Mitochondrial gene rearrangement within genus *Gasteruption* (Hymenoptera: Evaniioidea: Gasteruptionidae). Zool. Syst. 20: 840–853
- Wu, Q.L., Q. Li, Y. Gu, B.C Shi, C. Achterberg, S.J. Wei and X.X. Chen. 2014.** The complete mitochondrial genome of *Taeniogonalos taihorina* (Bischoff) (Hymenoptera: Trigonalidae) reveals a novel gene rearrangement pattern in the Hymenoptera. Gene 543: 76–84.

- Wu, Y.Y., Y.Y. Cao, J. Fang and X. Wan. 2016.** The first complete mitochondrial genome of stag beetle from China, *Prosopocoilus gracilis* (Coleoptera, Lucanidae). *Mitochondr. DNA.* 27(4): 2633.
- Wu, Y.Y. and X. Wan. 2016.** Molecular phylogeny of Lucanidae from China (Coleoptera: Scarabaeoidea). M.S. Thesis, Anhui University, Anhui, Hefei, China.
- Yang, J., F. Ye and Y. Huang. 2016.** Mitochondrial genomes of four katydids (Orthoptera: Phaneropteridae): New gene rearrangements and their phylogenetic implications. *Gene* 575: 702–711.
- Zhang, K.J., W.C. Zhu, X. Rong, J. Liu, X.L. Ding and X.Y. Hong. 2014.** The complete mitochondrial genome sequence of *Sogatella furcifera* (Horvath) and a comparative mitogenomic analysis of three predominant rice planthoppers. *Gene.* 533: 100–109.