

The Effect of Restricting Residents' Duty Hours on Patient Safety, Resident Well-Being, and Resident Education: An Updated Systematic Review

Lauren Bolster, MD
Liam Rourke, PhD

ABSTRACT

Background Despite 25 years of implementation and a sizable amount of research, the impact of resident duty hour restrictions on patients and residents still is unclear. Advocates interpret the research as necessitating immediate change; opponents draw competing conclusions.

Objective This study updates a systematic review of the literature on duty hour restrictions conducted 1 year prior to the implementation of the Accreditation Council for Graduate Medical Education's 2011 regulations.

Methods The review draws on reports catalogued in MEDLINE and PreMEDLINE from 2010 to 2013. Interventions that dealt with the duty hour restrictions included night float, shortened shifts, and protected time for sleep. Outcomes were patient care, resident well-being, and resident education. Studies were excluded if they were not conducted in patient care settings.

Results Twenty-seven studies met the inclusion criteria. Most frequently, the studies concluded that the restrictions had no impact on patient care (50%) or resident wellness (47%), and had a negative impact on resident education (64%). Night float was the most frequent means of implementing duty hour restrictions, yet it yielded the highest proportion of unfavorable findings.

Conclusions This updated review, including 27 recent applicable studies, demonstrates that focusing on duty hours alone has not resulted in improvements in patient care or resident well-being. The added duty hour restrictions implemented in 2011 appear to have had an unintended negative impact on resident education. New approaches to the issue of physician fatigue and its relationship to patient care and resident education are needed.

Introduction

Despite a large number of primary studies and numerous systematic reviews, the impact of resident duty hour restrictions on patients and residents remains unclear. In turn, reviewers find the impact to be favorable, unfavorable, neither, or inconclusive.¹⁻⁶ For programs that have implemented these changes, and for those about to, this literature is vexing.

From 2004 to 2013, 9 peer-reviewed systematic reviews of the duty hour literature were conducted (TABLE 1).¹⁻⁹ Eight of the reviews examined the impact of duty hour restrictions on patient safety. A total of 2 reported a positive impact,^{1,5} 2 reported no impact,^{3,6} and 4 found that the effect differed across studies.^{2,4,7,9} Five reviews examined the impact on resident well-being, with 4 finding positive changes.^{1,4,8,9} Resident education is examined in 5 reviews, with 1 concluding the impact is unfavorable,⁵ 2 concluding there is no impact,^{4,6} and 2 finding that the impact is inconclusive.^{8,9} Despite these equivocal findings, organizations that represent both patients and resi-

dents cite empirical support for action. The Bell Commission, which instituted New York State's limit on resident work hours¹⁰; the Institute of Medicine's Committee on Optimizing Graduate Medical Trainee (Resident) Hours and Work Schedule to Improve Patient Safety¹; the Accreditation Council for Graduate Medical Education (ACGME)¹¹; the Fédération des médecins résidents du Québec¹²; and Canada's National Steering Committee on Resident Duty Hours¹³ all point to the literature in their campaigns to limit duty hours. Others, notably residents,^{14,15} program directors,¹⁶ and other experts,¹⁷ read the same body of literature and concluded that restrictions may be detrimental.

Although much evidence has been produced for this debate, it has been inconclusive so far. The evidence includes systematic reviews; however, the most recent review⁵ assessed that the effect of night float, shortened shifts, or protected time for sleep is out of date. This review, by Reed and colleagues,⁵ was completed 1 year prior to when the ACGME's 2011 duty hour regulations took effect, and before the publication of hundreds of additional potentially pertinent studies. The purpose of this study is to update this systematic review of the duty hour literature.

DOI: <http://dx.doi.org/10.4300/JGME-D-14-00612.1>

TABLE 1
Results of 9 Systematic Reviews of the Impact of Resident Duty Hour Restrictions on 3 Categories of Outcomes

Source, y	Patient Care	Resident Wellness	Resident Education
Fletcher et al, ⁷ 2004	Inconclusive		
Fletcher et al, ⁸ 2005		Favorable	Inconclusive
Ulmer et al, ¹ 2009	Favorable	Favorable	
Fletcher et al, ⁹ 2011	Inconclusive	Favorable	Inconclusive
Moonesinghe et al, ⁶ 2011	No impact		No impact
Baldwin et al, ² 2011	Inconclusive		
Philibert et al, ⁴ 2013	Inconclusive	Favorable	No impact
Jamal et al, ³ 2012	No impact		
Reed et al, ⁵ 2010	Favorable	Inconclusive	Unfavorable

Methods

As an update, this review modeled the methodology of the original study, including the search query, data sources, eligibility criteria, study selection and data extraction, assessment of study quality, and data synthesis. Here we present a concise description of the methodology: the full description is available in the original study.⁵ Our review followed the PRISMA guidelines for systematic reviews.¹⁸

Eligibility Criteria, Study Selection, and Data Extraction

We included English-language original research studies published in MEDLINE and PreMEDLINE. The literature search for the original review included articles published between January 1, 1989, and May 21, 2010. The search for our review included all relevant literature between May 22, 2010, and February 4, 2014. We included studies that imple-

mented duty hour restrictions by reducing shift length, providing protected time for sleep, and implementing night float. As in the original review, *shift length* is defined as the number of consecutive hours worked by residents without protected sleep. *Protected time for sleep* is defined as a period during a work shift in which residents transfer all of their responsibilities (such as admitting and cross-covering patients, and performing procedures) to another individual, so that they

can obtain uninterrupted sleep. *Night float* is defined as a staffing system in which dedicated residents work during the night and not during the day.

As in the original study, we included shift length and night float studies only if the studies occurred in actual practice settings, and described at least 1 outcome related to patient care, residents' health, or residents' education. Studies with any outcome that examined protected time for sleep for residents were included.

The search strategy yielded a total of 1362 unique citations. Uncertainties regarding their fitness for inclusion were resolved through discussion among the investigators. The interrater reliability for study inclusion was excellent ($\kappa = 0.81$) as $\kappa > 0.80$ indicates near perfect agreement.¹⁹ A total of 203 abstracts were selected for full text review, and 27 articles were included in the study (FIGURE). Both authors independently reviewed the articles and performed data extraction. Disagreements were resolved by discussion and consensus, and the interrater agreement was calculated for study inclusion and for the qualitative characterization of each study's overall effects.




FIGURE
Summary of Evidence Search and Selection

TABLE 2
Summary of Studies

Intervention Type	Patient Care	Resident Wellness	Resident Education
Shift length			
Auger et al, ³⁴ 2012		No change	Unfavorable
Bismilla et al, ³³ 2011		Favorable	Unfavorable
Choma et al, ²⁸ 2013	No change		
DeLaroche et al, ³⁰ 2014			Unfavorable
Emler et al, ³¹ 2012	Favorable		No change
Rosenbluth et al, ²⁵ 2013	Favorable		
Schwartz et al, ²⁹ 2013			Unfavorable
Sen et al, ²⁷ 2013	Unfavorable	No change	
Stroud et al, ³⁵ 2012	Favorable	No change	Unfavorable
Theobald et al, ²⁶ 2013			Favorable
Wysocki and McGowan, ³² 2010		Favorable	No change
Yaghoubian et al, ²³ 2010	No change		
Yaghoubian et al, ²⁴ 2010	No change		
Night float			
Naughton et al, ³⁷ 2011		Unfavorable	
Chua et al, ³⁸ 2011		Unfavorable	
McCormick et al, ³⁹ 2012		Unfavorable	
Davenport et al, ⁴⁰ 2010	Favorable		
McCoy et al, ⁴¹ 2011	No change	No change	Unfavorable
Zahrai et al, ⁴² 2011		Unfavorable	No change
Luks et al, ⁴³ 2010		Unfavorable	Unfavorable
Borman et al, ⁴⁴ 2011		No change	
Kee et al, ⁴⁵ 2011		Favorable	Favorable
Schuh et al, ⁴⁶ 2011		No change	Unfavorable
Brandenberger et al, ⁴⁷ 2010		Unfavorable	
Desai et al, ³⁶ 2013			Unfavorable
Protected sleep time			
Volpp et al, ⁴⁸ 2012	No change	Favorable	
Amin et al, ⁴⁹ 2012		No change	

Data were entered into a structured form to facilitate the abstraction of information about study design, sample characteristics, type of intervention, outcomes, and study quality.

Assessment of Study Quality

The Medical Education Research Quality Study Instrument (MERSQI) was used to evaluate the studies' quality.²⁰ The MERSQI highlights 6 aspects of a study, including the type of outcomes it evaluates, the research design, the sampling strategy, and the

data collection and analysis procedures. Since its introduction in 2007, its psychometric properties have been investigated in multiple studies, and evidence is accruing for its validity and reliability.²⁰⁻²² These data were extracted in duplicate, and intraclass correlations were calculated to measure interrater reliability.

Data Synthesis

Consistent with the original review, the data were synthesized qualitatively. We grouped the studies by the intervention type (shift length, night float, protected sleep time), and recorded their results categorically as having a favorable impact, unfavorable impact, or no impact on the following 3 classes of outcomes: patient care, resident education, and resident well-being. A study was determined to be favorable if its intervention had achieved a positive outcome, unfavorable if it had a negative outcome, and no impact if the intervention had a neutral outcome. Both authors coded each study's outcome, and any disagreements were resolved by discussion. The interrater reliability for the qualitative categorization of study effects overall was substantial

($\kappa = 0.76$). When a study examined the impact on more than 1 outcome it was included in each group. We tallied the frequency of results for each category of outcome and each type of intervention. Due to the heterogeneity of the outcome measures, we did not synthesize the results quantitatively, and calculation of effect size was not performed.

Results

There were 27 studies included in the final review: 13 (48%) reviewed shift length,²³⁻³⁵ 12 (44%) examined

TABLE 3A
Outcomes Associated With Duty Hour Reduction Interventions—Reduce Shift Length

Source, y	Study Design	Study Size	Specialty	Shift Length, h	MERSQI Score
Yaghoubian et al, ²³ 2010	Nonrandomized 2-group	4634 surgical cases	Surgery	>16 h	10.36
Yaghoubian et al, ²⁴ 2010	Nonrandomized 2-group	1432 surgical cases	Surgery, trauma	>16 h	10.36
Rosenbluth et al, ²⁵ 2013	Single group pre and post	664 patients	Pediatrics	30 versus 13 h	11.25
Theobald et al, ²⁶ 2013	Single group pre and post	97 residents	IM	30 versus 13 h	9
Sen et al, ²⁷ 2013	Single group pre and post	2323 residents	IM, surgery, pediatrics, emergency	>16 h vs <16 h	12.5
Choma et al, ²⁸ 2013	Nonrandomized 2-group	3991 patients	IM	30 versus 16 h	12.5
Schwartz et al, ²⁹ 2013	Nonrandomized 2-group	249 residents	Surgery	30 versus 16 h	10.36
DeLaroche et al, ³⁰ 2014	Nonrandomized 2-group	825 residents	Pediatrics	>16 versus <16 h	8.5
Emler et al, ³¹ 2012	Nonrandomized 2-group	19 residents	ICU	30 versus 13 h	Education, 12.66 Patient care, 10.36

Downloaded from http://meridian.allenpress.com/jgme/article-pdf/7/3/349/2262417/jgme-d-14-00612_1.pdf by guest on 20 January 2021

TABLE 3A
Extended

Patient Care	Resident Wellness	Resident Education
No significant difference in complications for daytime procedures (shift length <16 h) compared to nighttime procedures (shift length >16 h)		
Nighttime surgeries (shift >16 h) had a lower complication rate (15.6%) compared to daytime surgeries (shift <16 h; 20%; $P = .040$) No significant difference in mortality between daytime and nighttime procedures		
Postintervention significant reduction (14%) in length of stay (RR = 0.86, 95% CI 0.77–0.96)		
	Average number of hours worked per week did not change (69.0 h/wk pre versus 67.8 h/wk post, $P = .24$)	No difference in morning report attendance (79% versus 78%, $P = .49$). Increase in weekly noon teaching conference (60% of expected attendees pre versus 68% post, $P = .001$) No change in procedural experience ($P = .94$) Increase in patient encounters (118 patients per 24 wk pre versus 140 patients per 24 wk post, $P = .005$)
Increase in residents reporting concern about making serious medical error (19% pre to 23.3% post, $P = .007$)	No significant change in hours slept per day (6.8 h/d pre versus 7.0 h/d post, $P = .17$) No significant change in depressive symptoms or well-being scores ($P = .55$, $P = .86$)	
No significant change in length of stay, mortality, 30-day readmission rates, complication rates, or ICU transfers		
		Significant decrease (25%) in operative cases per year per PGY-1 pre and post (88.8/y per PGY-1 versus 65.9/y per PGY-1, $P = .005$)
		No significant difference on written examination knowledge ($P = .34$) Significant decrease in attendance at didactic teaching sessions: grand rounds (50.9% versus 37.1%, $P < .001$), lectures (83.9% versus 70.5%, $P < .001$), and mock resuscitation (65.2% versus 55%, $P = .005$)
Continuity: no significant difference in perceived continuity of care Clinical outcomes: length of stay in ICU was significantly shorter for the intervention schedule, 5.65 versus 8.43 d for the control schedule ($P = .040$) There was no significant difference in mortality or ICU readmission rates between schedules		Education: no significant difference in lecture attendance between the intervention or control schedule

TABLE 3A
Continued

Source, y	Study Design	Study Size	Specialty	Shift Length, h	MERSQI Score
Wysocki and McGowan, ³² 2010	Nonrandomized 2-group	11 residents	Surgery	36 versus 19 h	12.66
Bismilla et al, ³³ 2011	Nonrandomized 2 group	51 residents	Pediatrics	28 h versus 24 + 2 h	Resident wellness, 13.03 Education, 14
Auger et al, ³⁴ 2012	Randomized control trial	15 residents	Pediatrics	30 versus 12 h	Resident wellness, 11 Education, 10
Stroud et al, ³⁵ 2012	Single group posttest	28 residents	IM	28 h versus 24 + 2 h	Patient care, 8 Education, 8

Abbreviations: MERSQI, Medical Education Research Quality Study Instrument; RR, relative risk; CI, confidence interval; IM, internal medicine; ICU, intensive care unit; PGY, postgraduate year.

night float,³⁶⁻⁴⁷ and 2 (7%) examined protected time for sleep.^{48,49} Some studies measured more than 1 outcome, with 10 studies (37%) assessing patient care,^{23-28,31,40,41,48} 17 studies (63%) reporting on resident well-being,^{32-39,41-49} and 14 studies (52%) assessing educational impact.^{26,29-36,41-43,45,46} Across all of the studies and interventions, the impact of duty hour restrictions most frequently had no impact on patient care (5 of 10, 50%),^{23,24,28,41,48} no impact on resident well-being (7 of 17,

41%),^{27,34,35,41,44,46,49} and an unfavorable impact on resident education (9 of 14, 64%).^{29,30,33-36,41,43,46} A minority of studies found a favorable impact on resident well-being (4 of 17, 24%; TABLE 2).^{32,33,45,48} When analyzed by intervention type, the most frequent result of shift length changes was no impact (8 of 20, 40%),^{23,24,27,28,31,32,34,35} that of night float was an unfavorable impact (10 of 18, 56%),^{35-39,41-43,46,47} and for protected time for sleep, 1 study reported a favorable outcome for resident wellness (1 of 2, 50%)⁴⁸

TABLE 3A
Continued, Extended

Patient Care	Resident Wellness	Resident Education
	Traditional 74 h/wk versus shift intervention 60 h/wk Fatigued for 1/3 rostered hours for traditional, versus fatigued less than 1/10 rostered hours for intervention	Caseload: 90 operations for trainees in traditional schedule versus 83 operations for intervention schedule
	Workload: no difference in daytime patient load, fewer nighttime patients with intervention, fewer overnight pages after duty hour regulation (27.5 versus 15.5, $P = .001$), fewer overnight admissions for resident after duty hour regulations (4 versus 2, $P = .001$) Distance walked on call after duty hour regulation: 7.3 versus 5.24 km ($P < .001$) No significant difference in the rate of urinary ketones Each patient admitted was associated with 38 fewer minutes of sleep ($P < .001$)	No significant difference in educational attendance Significantly decreased staff supervision after duty hour regulations: 87.5 min to 30 min ($P < .001$)
	Fewer hours per week worked in intervention (66.6 versus 75.9 h) No significant difference in total sleep time (7.5 versus 7.3 h) Proportion of sleepless work hours (no sleep in prior 24 h) was 1% for intervention and 15% for control ($P < .001$) No significant differences in fatigue outcomes No significant differences in overall well-being	Educational survey items ranked significantly lower for intervention compared to control, including attending assessment of amount of education, quality of education, amount of didactic teaching, and resident's ability to reflect on clinical details ($P < .05$)
Perceived improved patient care and patient safety Perceived fewer gaps in transition in care and less cross-coverage, perceived improved continuity of care Increased accountability in the new system	No difference perceived in workload	Perceived negative effects on education

but no impact on patient care, while the other study was unfavorable for resident well-being (1 of 2, 50%).⁴⁹

Study Quality

The mean (SD) MERSQI score for the 27 studies was 10.55 (2.38, maximum 18 points). The interclass correlation for the 2 authors' MERSQI scoring was excellent (intraclass correlation coefficient = 0.90). Thus, the quality of these studies is consistent with the quality of studies in the original review, which

reported a mean MERSQI score of 11.90.⁵ As a set, the studies had consistent shortcomings in 4 of the 6 dimensions of MERSQI quality measures. Most frequently, analyses were based on a single institution setting (19 of 27, 70%),^{23–25,28,31–43,45,49} outcomes were assessed through self-reporting (15 of 27, 56%),^{27,30,31,34–39,41–43,45,46,49} no information was provided on the psychometric properties of the measurement tools (19 of 27, 70%),^{23–26,29–36,41–43,45,46,48,49} and measurement focused on lower-level outcomes, such as user reaction or knowledge gains (10 of 27,

TABLE 3B
Outcomes Associated With Duty Hour Reduction Interventions—Night Float (NF)

Source, y	Study Design	Study Size	Specialty	NF (No. of Consecutive Nights)	MERSQI Score
Desai et al, ³⁶ 2013	Randomized control trial	43 residents	IM	6 nights	8.73
Naughton et al, ³⁷ 2011	Nonrandomized 2 group	20 residents	Surgery	7 nights	10.36
Chua et al, ³⁸ 2011	Single group pre-post	31 residents	Pediatrics	5 nights	9.93
McCormick et al, ³⁹ 2012	Nonrandomized 2-group	27 residents	Surgery, orthopedics	Not described	Resident wellness, 11.45
Davenport et al, ⁴⁰ 2010	Single group cross-sectional	20 residents	Radiology	9 h, 7 nights	13.03
McCoy et al, ⁴¹ 2011	Single group pre-post	6 residents	IM, GI	11 h (EOD × 12, total 6 shifts)	12
Zahrai et al, ⁴² 2011	Nonrandomized 2-group	16 residents	Surgery, orthopedics	14 h, 5 nights	Resident wellness, 7.09 Education, 8.18
Luks et al, ⁴³ 2010	Single group posttest	116 residents	IM	12 h, 14 days	7.5
Borman et al, ⁴⁴ 2011	Single group posttest	6161 residents	Surgery	Not described	9
Kee et al, ⁴⁵ 2011	Single group posttest	8 residents	IM	11.5 h, 5d	5

Downloaded from http://meridian.allenpress.com/jgme/article-pdf/7/3/349/2262417/jgme-d-14-00612_1.pdf by guest on 20 January 2021

TABLE 3B
Extended

Patient Care	Resident Wellness	Resident Education
Experimental schedule increased handoffs Quality of patient care perceived to be reduced such that NF schedule terminated early	NF residents slept more on call versus control schedule residents (mean, 5.1 versus 8.3 h, $P = .003$)	Decreased availability for teaching conferences, and reduced PGY-1 presence during daytime hours
	Less sleep prior to shift: NF 6 h versus day shift 7 h ($P < .05$) Significantly higher sleepiness on NF than day, ESS 13 for NF versus 6 for day shift ($P < .001$), Significantly decreased sleep quality on NF ($P = .001$)	
	NF associated with decreased shift length, decreased work hours, but also significantly decreased total sleep: 5.47 versus 7.5 h ($P < .001$)	
	Subgroup analysis: mean NF sleep 5.1 h, mean day shift 5.7 h ($P = .08$) Significantly higher levels of fatigue on NF	
Between traditional call and NF, diagnostic discrepancies between residents and staff physicians were lower on NF, overnight versus night float OR 1.5		
No significant difference in length of stay, 30-day readmission rates, 30-day mortality, codes, rapid response team calls, ICU transfers between control and intervention months	No significant reduction in work hours per week (64.3 versus 68.9, $P = .40$) Overall experience rated lower on intervention month ($P = .040$)	Residents felt less prepared to manage cross-covered patients during intervention month ($P = .006$) Residents less likely to attend educational conference on intervention months ($P = .02$) Residents reported higher hours per week for scholarly research on the intervention month (9.5 versus 2 h, $P < .001$)
	Lower scores on SF36 Stressor Inventory for role of physical, bodily pain, and social functioning for NF residents No difference in stress levels between NF and standard call, less time for physical activity in NF group	No significant difference in attendance of education sessions, hours reading, or case volumes
	Fewer hours of sleep per day at home on NF (6.3 versus 7.1 h, $P < .001$)	Significantly less time at education sessions on NF (0.1 versus 2.7 h, $P < .001$) Significantly fewer hours per week with attending on NF (0.57 versus 2.97 h, $P < .001$) Significantly less time reading on NF (2.63 versus 3.33 h, $P = .004$)
	No difference in sleep duration between extended call and NF systems	
Perceived improved continuity of care with NF intervention	Perceived better alertness, more energy for residents	Perceived more time for reading

TABLE 3B
Continued

Source, y	Study Design	Study Size	Specialty	NF (No. of Consecutive Nights)	MERSQI Score
Schuh et al, ⁴⁶ 2011	Nonrandomized 2-group	34 residents	Neurology	11.5 h, 5 d versus 11.5 h, 4 d; 14 h, 5 d versus 15 h, 1 in 4 (not NF); 12 h, 6 d versus 12 h, 4 d	Resident wellness, 10.36 Education, 7.63
Brandenberger et al, ⁴⁷ 2010	Nonrandomized 2-group	14 residents	Surgery	12 nights	7.63

Abbreviations: MERSQI, Medical Education Research Quality Study Instrument; IM, internal medicine; NF, night float; ESS, Epworth Sleepiness Scale; OR, operating room; GI, gastroenterology; EOD, every other day.

37%),^{26,27,29,30,34–36,42,45,46} instead of the higher-level outcomes, such as behavior or patient outcomes.

Shift Length

TABLE 3A outlines the 13 studies examining shift length and shows the study design, quality score, and outcomes.^{23–35} The length of shifts that were studied ranged from 13 to 36 hours. The populations of interest, and subsequently the study outcomes, were not consistent across all of the studies. There were 2 studies reviewing surgical cases, using complication rates as outcomes (total n = 6066)^{23,24}; 9 studies involving residents with a variety of patient care, resident wellness, and resident education outcomes (total n = 3618)^{26,27,29–35}; and 2 studies involving patients with length of stay, mortality, and readmission rates used as outcomes (total n = 4655).^{25,28} Of the 2 studies of surgical cases, 1 found no difference in mortality for surgeries performed on shifts shorter than 16 hours compared with shifts longer than 16 hours, but this study found a slightly higher complication rate for procedures performed during shifts longer than 16 hours.²⁴ The other study found no difference in complication rates for surgeries performed on shifts longer than 16 hours.²³ Two studies found shorter length of stay for patients with the shorter shift length,^{25,31} although another study found no difference in length of stay,²⁸ and 3 found no difference in mortality.^{24,28,31}

Regarding resident wellness outcomes, the interventions most frequently had no impact (3 of 5, 60%).^{27,34,35} Notably, the only randomized controlled trial in the shift length intervention group found no impact on resident well-being with shifts of 12 versus 30 hours.³⁴ Resident education outcomes were most frequently unfavorable (5 of 8, 63%),^{29,30,33–35} with decreases in staff supervision,

opportunities for assessment, attendance at educational activities, and decreased operative cases reported as negative outcomes.

Night Float

TABLE 3B shows the study design, quality score, and outcomes of the 12 studies examining night float.^{36–47} Studies predominantly examined resident well-being (10 of 12, 83%),^{37–39,41–47} and the majority reported unfavorable outcomes (6 of 10, 60%), such as decreased sleep,^{36,38,43} higher fatigue,^{39,47} and higher stress,⁴³ compared to traditional call. The majority of studies that examined educational outcomes demonstrated worse outcomes with night float (4 of 6, 67%), with decreased attendance at teaching conferences,^{36,41–43} less time per week spent with attending physician,⁴³ and less time for independent reading.⁴³ Only 2 studies examined patient care outcomes, and their results are contradictory. One demonstrated decreased diagnostic errors,⁴⁰ and the other reported no differences in length of stay, 30-day readmission rates, 30-day mortality, codes, rapid response team calls, or intensive care unit transfers.⁴¹

Protected Time for Sleep

Two studies with high-quality ratings examined the effect of protecting time for sleep.^{48,49} A randomized controlled trial of 103 residents examined the impact on patient care and reported no significant impact on mortality, length of stay, or readmission rates for the 2657 patients cared for by the residents.⁴⁸ Both studies examined the impact on sleep as a measure of resident well-being, and results are contradictory. One found that protected time improved the quality and quantity of residents' sleep,⁴⁸ while the other found no differences⁴⁹ (TABLE 3C).

TABLE 3B
Continued, Extended

Patient Care	Resident Wellness	Resident Education
	Higher score for depersonalization and emotional exhaustion for residents on restricted hours Fatigue was better on restricted hours No significant difference in sleepiness or sleep quantity	Faculty favored longer shifts
	Fatigue was higher in NF group	

Discussion

We reviewed the literature that was published during this period (2010–2014) to clarify the impact of duty hour restrictions on patient safety, resident education, and resident wellness. Across 27 studies with 41 separate analyses, the largest proportion of results pointed to an unfavorable impact (16 of 41, 39%) and the second largest to no impact (15 of 41, 37%). Of the 3 types of outcomes, resident education garnered the highest proportion of unfavorable outcomes, followed by resident well-being. Of the 3 approaches for adapting resident schedules to duty hour restrictions, night float was associated with the highest proportion of unfavorable results.

The majority of previous reviews have examined retrospective outcomes before and after duty hour implementation, based on when the ACGME regulations took effect, and the studies included in these reviews did not state how duty hour regulations were implemented.^{3–5,7} The review by Reed et al⁵ and our review highlight that not all strategies to comply with duty hour regulations are equally effective. Reduced shift length is the strategy with the best evidence for improved patient care, although this is still only modest evidence, as there are numerous studies that demonstrate no impact. For the modest improvements, at best, in patient care, the reduced shift length did improve resident wellness and had a negative impact on education in most studies (TABLE 3A).

Night float is an increasingly common strategy, yet only 1 study in our review demonstrated a positive impact on patient care⁴⁰ and 1 demonstrated a favorable impact on residents.⁴⁵ The remainder of the studies demonstrated either an unfavorable impact or no change to resident well-being. Further, 4 of 6 studies (67%) found a negative impact on resident education. This is in contrast to the review by Fletcher et al⁹ that concluded resident quality of life

may be improved by duty hour restrictions, and that the impact on education was unclear.

As more prospective studies with defined interventions become available, there is increasing evidence that some strategies (night float) for duty hour regulations may have a negative impact on resident well-being. The vast majority of interventions did not use the suggested 4-night maximum recommended by the Institute of Medicine; therefore, these findings may be attributable to the acute sleep deprivation that has been reported with increased numbers of consecutive nights on call.⁵⁰ Previous reviews, including the 1 we updated, have been supportive of duty hour restrictions, although many have determined that the evidence is unclear.^{1–3,6} The original intended purpose of duty hour changes was to enhance patient safety through enhanced supervision and sleep.^{1,9,11} Many of the reviews have found no conclusive evidence that duty hour restrictions have improved patient care; therefore, one might conclude that duty hour restrictions have not had their intended impact. There is also a growing body of evidence to suggest that there is no definitive improvement in resident well-being, and that there is a potential negative impact on resident education. It is worth noting that while the issue of supervision after the institution of the duty hour limits is relatively underresearched, 2 of the studies in our review found decreased time with attending physicians and decreased supervision as a result of duty hour regulations. Enhanced supervision was a key feature in the Institute of Medicine report and the ACGME standards; however, it appears to not have been addressed in the majority of interventions, and may actually be decreased by implementing duty hour restrictions.

Limitations of this review are similar to those of the original study. Few of the studies were randomized controlled trials.^{34,36,48} However, many used a nonrandomized 2-group design, which is a reasonable design for an educational intervention. Unfortunately,

TABLE 3c
Outcomes Associated With Duty Hour Reduction Interventions—Protected Sleep Time (Nap)

Source, y	Study Design	Study Size	Specialty	Nap (h)/ Shift (h) Duration	MERSQI Score	Patient Care	Resident Wellness	Resident Education
Volpp et al, ⁴⁸ 2012	RCT	103 residents	IM	5/30	16.14	No significant difference between control and intervention for any of the clinical outcomes, including mortality, length of stay, or readmission	Significantly more hours slept with intervention 1.98 versus 2.86 h (VA Medical; $P < .001$), 2.04 versus 3.04 (HUP; $P < .001$) Significantly fewer residents with 0 h of sleep on call ($P < .001$) Significantly lower KSS score in intervention groups	
Amin et al, ⁴⁹ 2012	Nonrandomized 2-group	29 residents	IM	0.33/9	14.28		No difference in PSQI scores Decreased attentional failures post nap for intervention group Intervention residents showed improved CPT II test results	

Abbreviations: MERSQI, Medical Education Research Quality Study Instrument; RCT, randomized controlled trial; IM, internal medicine; VA, Veterans Affairs; HUP, Hospital of the University of Pennsylvania; KSS, Karolinska sleepiness scale; PSQI, Pittsburgh Sleep Quality Index; CPT, Continuous Performance Test.

the studies in this review used heterogeneous outcome measures, precluding quantitative synthesis and comparisons. Further, while the most important patient care outcomes are arguably mortality and morbidity, only 60% (6 of 10) of studies that reviewed patient care looked at mortality and morbidity outcomes.^{23–25,28,41,48} Regarding education, the outcome measures were largely related to attendance at teaching sessions and time with faculty. While important, these are rough surrogates for more meaningful educational outcomes.

As an update to a previous literature review with strict inclusion criteria, the studies in our review do not represent all of the literature on this topic. As in the original review, we excluded studies that did not occur in actual practice settings. The purpose of this criterion is to accentuate results that are ecologically valid over results derived from laboratory studies on the effects of sleep deprivation. Further, by limiting our review to studies with defined interventions, the results of each method of limiting duty hours are more apparent. Without a clearly defined intervention to implement duty hours, it is difficult to determine cause and effect, and even associations are less clear due to confounders. Given these limitations, this review does not definitively answer the questions regarding the impact of duty hour restrictions. However, the studies included are the most applicable, current evidence on this topic, and the number of studies in our review is consistent with other reviews of issues in medical education.^{51–54}

Future directions for this field should include attempts to standardize outcome measures across studies so that there will be more generalizability and studies will be able to be compared more accurately. Additionally, duty hours are only 1 factor in the safety of patients cared for by residents; handover-related communication errors and supervision of residents have also been identified as contributing factors. Future studies should seek to assess the impact of a more comprehensive approach to improving patient care. Recently, a prospective handoff “bundle” intervention study demonstrated a significant reduction in medical errors, and demonstrated proof of the concept of studying the impact on patient care from combining simultaneous interventions.⁵⁵

Conclusion

An updated review of 27 recent and highly applicable studies demonstrates that focusing on interventions to limit resident duty hours alone has not had the expected, consistent improvements to patient care, resident education, or resident well-being. Further

duty hour restrictions appear to have had an unintended negative impact on resident education. It is time to reevaluate the profession's approach to the issue of resident hours and its impact on patient care and residency education.

References

1. Ulmer C, Wolman DM, Johns MM, eds. *Resident Duty Hours: Enhancing Sleep, Supervision, and Safety*. Washington, DC: National Academies Press; 2009.
2. Baldwin K, Namdari S, Donegan D, Kamath A, Mehta S. Early effects of resident work-hour restrictions on patient safety: a systematic review and plea for improved studies. *J Bone Joint Surg Am*. 2011;93(2):e5.
3. Jamal M, Doi SA, Rousseau M, Edwards M, Rao C, Barendregt J, et al. Systematic review and meta-analysis of the effect of North American working hours restrictions on mortality and morbidity in surgical patients. *Br J Surg*. 2012;99(3):336–344.
4. Philibert I, Nasca T, Brigham T, Shapiro J. Duty-hour limits and patient care and resident outcomes: can high-quality studies offer insight into complex relationships? *Ann Rev Med*. 2013;64:467–483.
5. Reed D, Fletcher K, Arora VM. Systematic review: association of shift length, protected sleep time, and night float with patient care, residents' health, and education. *Ann Intern Med*. 2010;153(12):829–842.
6. Moonesinghe SR, Lowery J, Shahi N, Millen A, Beard JD. Impact of reduction in working hours for doctors in training on postgraduate medical education and patients' outcomes: systematic review. *BMJ*. 2011;342:d1580.
7. Fletcher KE, Davis SQ, Underwood W, Mangrulkar RS, McMahon LF, Saint S. Systematic review: effects of resident work hours on patient safety. *Ann Intern Med*. 2004;141(11):851–857.
8. Fletcher KE, Underwood W 3rd, Davis SQ, Mangrulkar RS, McMahon LF, Saint S. Effects of work hour reduction on residents' lives: a systematic review. *JAMA*. 2005;294(9):1088–1100.
9. Fletcher KE, Reed DA, Arora VM. Patient safety, resident education and resident well-being following implementation of the 2003 ACGME duty hour rules. *J Gen Intern Med*. 2011;26(8):907–919.
10. New York State Department of Health. *Final Report of the New York State Ad Hoc Advisory Committee on Emergency Services*. Albany, NY: New York State Department of Health; 1987.
11. Philibert I, Amis S. The ACGME 2011 duty hour standards: enhancing quality of care, supervision, and resident professional development. <http://www.acgme.org/acgmeweb/Portals/0/PDFs/jgme-monograph%5B1%5D.pdf>. Accessed April 20, 2015.
12. Dussault C, Saad N, Carrier J. 16-hour call duty schedules: the Quebec experience. *BMC Med Educ*. 2014;14(suppl 1):10. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4304263>. Accessed May 27, 2015.
13. National Steering Committee on Resident Duty Hours. Fatigue, risk and excellence: towards a pan-Canadian consensus on resident duty hours. http://www.residentdutyhours.ca/documents/fatigue_risk_and_excellence.pdf. Accessed May 15, 2015.
14. Fargen KM, Chakraborty A, Friedman W. Results of a national neurosurgery resident survey on duty hour regulations. *Neurosurgery*. 2011;69(6):1162–1170.
15. Drolet B, Christopher D, Fischer S. Residents' response to duty-hour regulations—a follow-up national survey. *N Engl J Med*. 2012;366(24):e35.
16. Antiel RM, Thompson SM, Reed DA, James KM, Tilburt JC, Bannon MP, et al. ACGME duty-hour recommendations—a national survey of residency program directors. *N Engl J Med*. 2010;363(8):e12.
17. McNoble DJ. Expanded liability of hospitals for the negligence of fatigued residents. *J Leg Med*. 1990;11:427–449.
18. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(7):e1000097.
19. Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*. 1977;33(1):159–174.
20. Reed DA, Cook DA, Beckman TJ, Levine RB, Kern DE, Wright SM. Association between funding and quality of published medical education research. *JAMA*. 2007;298(9):1002–1009.
21. Yucha CB, Schneider BS, Smyer T, Kowalski S, Stowers E. Methodological quality and scientific impact of quantitative nursing education research over 18 months. *Nurs Educ Perspect*. 2011;32(6):362–368.
22. Reed DA, Beckman TJ, Wright SM. An assessment of the methodologic quality of medical education research studies published in *The American Journal of Surgery*. *Am J Surg*. 2009;198(3):442–444.
23. Yaghoubian A, Kaji AH, Ishaque B, Park J, Rosing DK, Lee S, et al. Acute care surgery performed by sleep deprived residents: are outcomes affected? *J Surg Res*. 2010;163(2):192–196.
24. Yaghoubian A, Kaji AH, Putnam B, deVirgilio C. Trauma surgery performed by “sleep deprived” residents: are outcomes affected? *J Surg Educ*. 2010;67(6):449–451.
25. Rosenbluth G, Fiore DM, Maselli JH, Vittinghoff E, Wilson SD, Auerbach AD. Association between adaptations to ACGME duty hour requirements, length of stay, and costs. *Sleep*. 2013;36(2):245–248.
26. Theobald CN, Stover DG, Choma NN, Hathaway J, Green JK, Peterson NB, et al. The effect of reducing

- maximum shift lengths to 16 hours on internal medicine interns' educational opportunities. *Acad Med.* 2013;88(4):512–518.
27. Sen S, Kranzler HR, Didwania AK, Schwartz AC, Amarnath S, Kolars JC, et al. Effects of the 2011 duty hour reforms on interns and their patients: a prospective longitudinal cohort study. *JAMA Intern Med.* 2013;173(8):657–662.
 28. Choma NN, Vasilevskis EE, Sponsler KC, Hathaway J, Kripalani S. Effect of the ACGME 16-hour rule on efficiency and quality of care: duty hours 2.0. *AMA Intern Med.* 2013;173(9):819–821.
 29. Schwartz SI, Galante J, Kaji A, Dolich M, Easter D, Melcher ML, et al. Effect of the 16-hour work limit on general surgery intern operative case volume a multi-institutional study. *JAMA Surg.* 2013;148(9):829–833.
 30. DeLaroche A, Riggs T, Maisels MJ. Impact of the new 16-hour duty period on pediatric interns' neonatal education. *Clin Pediatr (Phila).* 2014;53(1):51–59.
 31. Emler LL, Al-Khafaji A, Kim YH, Venkataraman R, Rogers PL, Angus DC. Trial of shift scheduling with standardized sign-out to improve continuity of care in intensive care units. *Crit Care Med.* 2012;40(12):3129–3134.
 32. Wysocki AP, McGowan B. Revising the surgical registrar on-call roster. *ANZ J Surg.* 2010;80(7–8):490–494.
 33. Bismilla Z, Breakey VR, Swales J, Kulik DM, Pai N, Singh N, et al. Prospective evaluation of residents on call: before and after duty hour reduction. *Pediatrics.* 2011;127(6):1080–1087.
 34. Auger KA, Landrigan CP, Gonzales del Rey JA, Sieplinga KR, Sucharew HJ, Simmons JM. Better rested, but more stressed? Evidence of the effects of resident work hour restrictions. *Acad Pediatr.* 2012;12(4):335–343.
 35. Stroud L, Oulanova O, Szecket N, Ginsburg S. The benefits make up for whatever is lost: altruism and accountability in a new call system. *Acad Med.* 2012;87(10):1421–1427.
 36. Desai S, Feldman L, Brown L, Dezube R, Yeh HC, Punjabi N, et al. Regulation-compliant models on sleep duration, trainee education, and continuity of patient care among internal medicine house staff: a randomized trial. *AMA Intern Med.* 2013;173(8):649–655.
 37. Naughton PA, Aggarwal R, Wang TT, Van Herzele I, Keeling AN, Darzi AW, et al. Skills training after night shift work enables acquisition of endovascular technical skills on a virtual reality simulator. *J Vasc Surg.* 2011;53(3):858–866.
 38. Chua KP, Gordon MB, Sectish T, Landrigan CP. Effects of a night-team system on resident sleep and work hours. *Pediatrics.* 2011;128(6):1142–1147.
 39. McCormick F, Kadzielski J, Landrigan CP, Evans B, Herndon J, Rubash HE. Surgeon fatigue: a prospective analysis of the incidence, risk and intervals of predicted fatigue-related impairment in residents. *Arch Surg.* 2012;147(5):430–435.
 40. Davenport MS, Ellis JH, Khalatbari SH, Myles JD, Klein KA. Effect of work hours, caseload, shift type and experience on resident call performance. *Acad Radiol.* 2010;17(7):921–927.
 41. McCoy CP, Halvorsen AJ, Connor CG, McDonald FS, Oxentenko AS. Effect of 16-hour duty periods on patient care and resident education. *Mayo Clin Proc.* 2011;86(3):192–196.
 42. Zahrai A, Chahal J, Stojimirovic D, Schemitsch EH, Yee A, Kraemer W. Quality of life and educational benefit among orthopedic surgery residents: a prospective, multicentre comparison of the night float and the standard call systems. *Can J Surg.* 2011;54(1):25–32.
 43. Luks AM, Smith CS, Robins L, Wipf JE. Resident perceptions of the educational value of night float rotations. *Teach Learn Med.* 2010;22(3):196–201.
 44. Borman KR, Biester TW, Jones AT, Shea JA. Sleep, supervision, education, and service: views of junior and senior residents. *J Surg Educ.* 2011;68(6):495–501.
 45. Kee CL, Goh WP, Yap ES, Chan YC. Impact of a newly introduced medical officer night-float on-call system in a medical department in Singapore. *Singapore Med J.* 2011;52(1):60–62.
 46. Schuh LA, Khan MA, Harle H, Southerland AM, Hicks WJ, Falchook A, et al. Pilot trial of IOM duty hour recommendations in neurology programs: unintended consequences. *Neurology.* 2011;77(9):883–887.
 47. Brandenberger J, Kahol K, Feinstein AJ, Ashby A, Smith M, Ferrara JJ. Effects of duty hours and time of day on surgery resident proficiency. *Am J Surg.* 2010;200(6):814–819.
 48. Volpp KG, Shea JA, Small DS, Basner M, Zhu J, Norton L, et al. Effect of a protected sleep period on hours slept during extended overnight in-hospital duty hours among medical interns: a randomized trial. *JAMA.* 2012;308(21):2208–2217.
 49. Amin M, Graber M, Ahmad K, Manta D, Hossain S, Belisova Z, et al. The effects of a mid-day nap on the neurocognitive performance of first-year medical residents: a controlled interventional pilot study. *Acad Med.* 2012;87(10):1428–1433.
 50. Landrigan, CP, Rosthschild JM, Cronin JW, Kaushal R, Burdick E, Katz JT, et al. Effect of reducing interns' work hours on serious medical errors in intensive care units. *N Engl J Med.* 2004;351(18):1838–1848.
 51. De Jong J, Visser M, Van Dijk N, van der Vleuten C, Wieringa-de Waard M. A systematic review of the relationship between patient mix and learning in work-based clinical settings. A BEME systematic review: BEME Guide No. 24. *Med Teach.* 2013;35(6):e1181–e1196.

52. O'Dunn-Orto A, Hartling L, Campbell S, Oswald AE. Teaching musculoskeletal clinical skills to medical trainees and physicians: a best evidence in medical education systematic review of strategies and their effectiveness: BEME Guide No. 18. *Med Teach*. 2012;34(2):93–102.
53. Nelson C, Hartling L, Campbell S, Oswald AE. The effects of audience response systems on learning outcomes in health professions education. A BEME systematic review: BEME Guide No. 21. *Med Teach*. 2012;34(6):e386–e3405.
54. Thistlewaite JE, Davies D, Ekeocha S, Kidd JM, Macdougall C, Matthews P, et al. The effectiveness of case-based learning in health professional education. A BEME systematic review: BEME Guide No. 23. *Med Teach*. 2012;34(6):e421–e444.
55. Starmer AJ, Sectish TC, Simmon DW, Keohane C, McSweeney ME, Chung EY, et al. Rates of medical errors and preventable adverse events among

hospitalized children following implementation of a resident handoff bundle. *JAMA*. 2013;310(21):2262–2270.



Both authors are in the Department of Medicine, University of Alberta, Edmonton, Alberta, Canada. **Lauren Bolster, MD**, is Assistant Clinical Professor, Division of Hematology; and **Liam Rourke, PhD**, is Associate Professor.

Funding: The authors report no external funding source for this study.

Conflict of interest: The authors declare they have no competing interests.

Corresponding author: Lauren Bolster, MD, University of Alberta, Department of Medicine, 13-103 Clinical Sciences Building, 11350-83 Avenue, Edmonton, AB T6G 2P4 Canada, 780.407.1584, lauren.bolster@ualberta.ca

Received November 2, 2014; revisions received February 6, 2015, and March 22, 2015; accepted April 6, 2015.