

Erratum for ‘Retro-wedge foreland basin evolution along the ECORS line, eastern Pyrenees, France,’ *Journal of the Geological Society*, 173, 419–437

Mary Ford^{1,2*}, Louis Hemmer^{1,2*}, Arnaud Vacherat³, Kerry Gallagher⁴ & Frédéric Christophoul⁵

¹ CRPG, UMR 7358, 15 Rue Notre Dame des Pauvres, 54501 Vandoeuvre-lès-Nancy, France

² Université de Lorraine, ENSG, INP, Rue Doyen Marcel Roubault, 54501 Vandoeuvre-lès-Nancy, France

³ ISTEP, Université Pierre et Marie Curie, Paris 6, France

⁴ Géosciences Rennes, Université de Rennes 1, Campus de Beaulieu, 35042 Rennes, France

⁵ GET, UMR 5563, Université de Toulouse–CNRS–IRD–OMP, 14 Avenue Edouard Belin, 31400 Toulouse, France

* Correspondence: mary.ford@univ-lorraine.fr

The authors would like to apologise for the following error on page 422 in the paragraph beginning ‘Lithologically, the Grey Flysch Group...’:

In borehole MZ1 the lower levels of the Grey Flysch Group contain a fossil assemblage typical of inner platform (*Nummulites* sp., *Fallotia* sp., *Vidalina* sp., *Cribrstromoïdes* sp., *Rotalia* sp. and miliolids), whereas the upper levels contain outer platform assemblages including planktonic foraminifera (*Globotruncana* sp., keeled band morphotype and Heterohelicidae, biserial morphotype).

The sentence should instead read:

In borehole MZ1 the lower levels of the Grey Flysch group contain a fossil assemblage typical of inner platform (*Nummofallotia* sp., *Vidalina* sp., *Cribrstromoïdes* sp., *Rotalia* sp. and miliolids), while the upper levels contain outer platform assemblages including planktic foraminifera (*Globotruncana* sp., keeled band morphotype and Heterohelicidae, biserial morphotype).