

CO₂ emissions and their changes with H₂O emissions, soil moisture, and temperature during the wetting–drying process of the soil mixed with different biochar materials

Caner Yerli ^{a,*}, Talip Cakmakci ^a and Ustun Sahin ^b

^a Department of Biosystem Engineering, Faculty of Agriculture, Yuzuncu Yil University, Van, Turkey

^b Department of Agricultural Structures and Irrigation, Faculty of Agriculture, Ataturk University, Erzurum, Turkey

*Corresponding author. E-mail: caneryerli@yyu.edu.tr

 CY, 0000-0002-8601-8791; TC, 0000-0001-5815-1256; US, 0000-0002-1924-1715

ABSTRACT

Biochar is an organic regulator that improves crop yield by regulating soil properties. In addition, this organic regulator is also effective in reducing CO₂ emissions from soil. However, considering the management of CO₂ emissions together with many factors and the different properties of soil depending on the biochar content, CO₂ emissions can vary. Thus, the study investigated the soil moisture and temperature and H₂O emissions, which affect the emission, and CO₂ emission of biochars with different raw materials applied to the soil in the wetting–drying cycle of the soil. It was determined that biochar applications decreased CO₂ emissions, but the share of each biochar material in reduction differed, and CO₂ emissions were 82, 51, 20, and 13% lower in straw, hazelnut, apple, and sawdust biochar applications than in soil without biochar, respectively, and significant positive linear relationships of CO₂ emissions with soil moisture–temperature and H₂O emissions were determined. In addition, in biochar applications, H₂O and soil temperature decreased depending on the moisture retention in the soil. In the findings, it can be suggested that straw biochar application to soil is more effective in reducing the severity of increasing global warming, and that soil moisture and temperature should be managed to reduce CO₂ emissions.

Key words: biochar, CO₂ emission, soil moisture, soil temperature, wetting–drying

HIGHLIGHTS

- Biochar treatment decreased the CO₂ emissions from the soil.
- Straw biochar caused lower CO₂ emissions compared to hazelnut, apple, and sawdust biochar.
- Biochar treatment decreased H₂O emissions and soil temperature, while increased soil moisture.
- The relationships of CO₂ emissions with soil moisture, H₂O emissions, and soil temperature were positive linear.

INTRODUCTION

Agriculture is responsible for a quarter of global warming (Bennetzen *et al.* 2016). Soil can act as a reserve for organic carbon as well as a source of CO₂ emissions from the soil. Under faulty soil management conditions, the soil loses organic carbon and CO₂ is thus emitted into the atmosphere (Yerli *et al.* 2019). In this case, the increased CO₂ in the atmosphere absorbs the heat and causes global warming. Climate change, which is the effect of global warming, is regarded as the most worrying issue today (WMO 2019). In this context, studies investigating the effects of CO₂ emissions and thus global warming on agricultural activities are noteworthy.

Biochar can be defined as a charred organic improver as a result of the pyrolysis of biomass under oxygen-limited conditions at elevated temperatures. With its low bulk density, high porosity and surface area properties, biochar increases water retention in the soil and enables more profitable management of irrigation in today's conditions, in which effective management of water resources is very important (Tufenkci *et al.* 2022). Biochar both facilitates the crop's access to water and contributes to irrigation water savings by keeping soil moisture easier and preserving it for a long time, not only inside the pores but also between the particles due to its micropores (Cakmakci & Sahin 2022).

Biochar increases soil and crop productivity due to its low decomposition rate and long-term persistence in the soil (Tufenkci *et al.* 2022). The bulk density of biochar applied soil may decrease, and the soil porosity and soil hydraulic

This is an Open Access article distributed under the terms of the Creative Commons Attribution Licence (CC BY 4.0), which permits copying, adaptation and redistribution, provided the original work is properly cited (<http://creativecommons.org/licenses/by/4.0/>).

properties may improve (Ahmad Bhat *et al.* 2022). Thus, more effective irrigation management can be achieved. The biochar applied to the soil is effective in developing and improving the physical (Mukherjee & Lal 2013; Blanco-Canqui 2017; Zhang *et al.* 2020), chemical (Chintala *et al.* 2014; Rechberger *et al.* 2017; Jing *et al.* 2020), and biological (Luo *et al.* 2013; Dong *et al.* 2015; Somerville *et al.* 2020) properties of soil. In this context, it also provides positive reflections on the increase in the yield and quality of the crop (Zoghi *et al.* 2019). These contributions of biochar allow more economical and sustainable agricultural management by decreasing the need for synthetic fertilizers. Laghari *et al.* (2015) found that biochar increased phosphorus up to 70% and potassium up to 11% in the soil. Similarly, Apori *et al.* (2021) stated that nitrogen increased by 36% and phosphorus increased by 17% with biochar application. In addition, biochar plays an active role in biotic and abiotic stress conditions, increasing the resistance of the crop against stress conditions (Kul *et al.* 2021).

In addition to these positive reflections, considering biochar can increase the carbon storage capacity of the soil and stabilize the soil's organic carbon pools (Wang *et al.* 2014). As the effect of global warming is increasing day by day, biochar can also decrease the CO₂ emissions from agricultural soils (Deng *et al.* 2017). Thus, it can contribute to the increase in soil and crop yield by providing organic carbon increases and sequestration in the soil, and agricultural and environmental sustainability can be achieved by decreasing the emission values. However, considering that CO₂ emissions from soils can vary with soil moisture and soil temperature (Mancinelli *et al.* 2015; Yerli & Sahin 2021) and that the physical, chemical, and biological properties of soils can directly affect CO₂ emissions (Haddaway *et al.* 2017), it is possible to manage CO₂ emissions from soils effectively only with a good understanding of the CO₂ emissions relationships (Yerli *et al.* 2022a). Increasing moisture and temperature in the soil can trigger microorganism activities and increase CO₂ emissions by oxidizing organic carbon. After organic carbon encounters O₂, it turns into CO₂ and spreads from the soil to the atmosphere (Yerli *et al.* 2019). Zhao *et al.* (2020) reported a positive correlation between CO₂ emissions and soil temperature, and Buragiene *et al.* (2019) reported a positive correlation between CO₂ emissions and soil moisture. However, this is not always the case. Because situations such as water ingress into the soil increase the soil moisture and decrease the soil temperature by creating a cooling effect on the soil (Mancinelli *et al.* 2015), microorganism activities and thus oxidation can be effective between certain soil moisture and temperature values, and the increase in soil temperature due to a chemical reaction that occurs as a result of the oxidation of organic carbon additives mixed into the soil (Yerli & Sahin 2021) makes it difficult to understand and manage the CO₂ emissions from the soil. Yerli *et al.* (2022b) determined that CO₂ emissions from the soil showed positive correlations with H₂O emissions and soil moisture at 5-, 10-, and 20-cm soil depths, but the correlation with soil temperature was negative at the same soil depths.

Many studies have shown that biochar decreases CO₂ emissions from the soil (Liu *et al.* 2011; Wang *et al.* 2014; Zhang *et al.* 2017; Wu *et al.* 2018; Azeem *et al.* 2019; Akinyemi & Adesina 2020; Lehmann *et al.* 2021). Lehmann (2007) reported that biochar absorbs organic carbon and stabilizes soil carbon stocks due to its high surface area and porosity, thus decreasing CO₂ emissions. However, CO₂ emissions may also increase depending on the raw material content of biochar and preparation processes, such as stability, pyrolysis, and temperature, as well as the characteristics of the soil in which the biochar is mixed and its interaction with the soil (Zimmerman *et al.* 2011; Singh *et al.* 2012). In a study examining the effects of pine wood and grass-derived biochar applications in loamy soil on CO₂ emissions, it was determined that pine wood biochar did not affect CO₂ emissions, while grass-derived biochar increased CO₂ emissions (Hilscher *et al.* 2009). Karhu *et al.* (2011) reported that birch biochar mixed with soil has a negative effect on decreasing CO₂ emissions. Mohamed *et al.* (2015) evaluated the effects of woody waste mixed with sandy soil and its biochar on CO₂ emissions, and they found that both applications increased CO₂ emissions, but this increase was 3 times higher than biochar in the woody waste application and 6 times more than the control.

The insight shows that the mechanism of CO₂ emissions from the soil has a complex structure and that the effect of biochar on decreasing emissions varies. The studies have gained momentum to evaluate the effects of biochar applications on decreasing CO₂ emissions. However, there are no studies in the literature that compare the effects of different biochar sources on CO₂ emissions during the wetting–drying process of the soil and evaluate the soil moisture, H₂O emissions, and soil temperature that affect CO₂ emissions. Therefore, the objectives of this study were (i) to determine whether biochar applications would decrease the CO₂ emissions from the soil in the wetting–drying process, (ii) to evaluate the changes in CO₂ emissions depending on the raw material contents of the biochar, and (iii) to discuss the relationships of CO₂ emissions with the soil moisture, H₂O emissions, and soil temperature measured during the wetting–drying process. To achieve these objectives, this study investigated CO₂ emissions during the wetting–drying process of soil including biochars with different raw materials, and evaluated the relationships of CO₂ emissions with soil moisture and temperature and H₂O emissions.

MATERIALS AND METHODS

Study area and experimental design

The experiment was carried out in the laboratory of Van Yuzuncu Yil University, Faculty of Agriculture, Department of Bio-system Engineering. During the experiment, the mean air temperature was 24 ± 2 °C and the air humidity was $38 \pm 5\%$. The experiment was conducted with three replications using a completely randomized factorial design with hazelnut, straw, apple, and sawdust biochar materials and as a control in the soil in which biochar was not applied. Each wetting–drying cycle was completed in 1 week and repeated three times in total for 3 weeks.

Properties of soil and biochar materials

Prior to the experiment, the particle size distribution (texture), pH, EC, total N, organic matter and organic carbon contents of the experimental soil were analyzed (Table 1). The particle size distribution was determined by the Bouyoucos hydrometer method (Gee & Bauder 1986), and the soil texture was sandy loam (sand: 66.3%, silt: 15.6%, clay: 18.1%) according to the USDA classification. According to the pH and EC determined by pH-meter and EC-meter in 1:2.5 saturation extract, the experimental soil was very slightly alkaline and nonsaline. The total N determined by the Kjeldahl method (Bremner & Mulvaney 1982) was insufficient, and the organic matter content determined by the Walkley-Black method (Nelson & Sommers 1982) and the organic carbon calculated from the organic matter content (Avramidis *et al.* 2015) were low.

All biochar materials were prepared using the same procedures. First, the materials were dried, sieved, and homogenized, and finally, subjected to pyrolysis at 400 °C. Prior to the experiment, analyses and calculations were made to determine the pH, EC, total N, organic matter, and organic carbon contents of the biochar materials in the same way as the analysis and calculations of the soil (Table 1).

Applications and irrigation practices

Air-dried soil, sieved through a 2-mm sieve, was mixed with 1% biochar on a weight basis and tapped into pots with volumes of 1.5 l (diameter: 13 cm, height: 11 cm) by preserving the soil bulk density (1.31 Mg m^{-3}). All pots were incubated for two weeks at moisture level at field capacity. In the planning of irrigation events, the dry weights of control application pots with the soil were determined. To determine the water retained at field capacity (pot capacity), the pots without biochar (control) were first saturated with water and then surface covered to prevent evaporation. When the drainage completely stopped, the pots were weighed, and this moisture after conversion by bulk density, wet and dry weights was expressed as field capacity as the volume ($0.320 \text{ m}^3 \text{ m}^{-3}$). The irrigations were applied with the same water amounts in all applications to complete missing moisture to the field capacity considering the weights in control pots when each wetting–drying process was completed.

CO₂ emission measurement process

The CO₂ emission from the soil was measured daily during three wetting–drying periods with the EGM-5 infrared gas analyzer device (CFX-2, PPSystems, Stotfold, UK) by taking three readings from each pot. Soil moisture, H₂O emissions, and soil temperature were also measured simultaneously with CO₂ emissions. Soil moisture was determined by weighing the pots, while a CO₂ measuring device was used for H₂O emission and soil temperature measurements. The H₂O emission was automatically recorded by a CO₂ measuring device during CO₂ measurement. The STP-1 temperature probe connected to the CO₂ emission device was used to measure soil temperature at a 5-cm soil depth (Yerli & Sahin 2021; Yerli *et al.* 2022a, 2022b).

Table 1 | The properties of soil and biochar materials used in the experiment

Properties	Soil	Hazelnut biochar	Straw biochar	Apple biochar	Sawdust biochar
pH	7.17	8.11	8.56	7.88	8.88
EC (dS m ⁻¹)	0.62	1.16	1.85	0.82	0.80
Total N (%)	0.04	0.19	0.22	0.17	0.20
Organic matter (%)	0.81	56.1	58.2	42.3	45.2
Organic carbon (%)	0.47	32.5	33.8	24.5	26.2

Statistical analysis

Data analyses were performed using the General Linear Model in the SPSS program (Ver. 23). Duncan multiple comparison test at the 5% level of probability was used to compare significant means. Pearson correlation analysis was used to determine the relationships of CO₂ emissions with soil moisture, H₂O emissions, and soil temperature.

RESULTS AND DISCUSSION

Soil moisture increased in biochar applications in the experimental period compared to the control, and the highest soil moisture was determined in the straw biochar application (Figure 1). Mean soil moisture was found to be 7.6, 9.8, 2.2, and 4.0% higher in hazelnut, straw, apple, and sawdust biochar applications than in the control, respectively. The lowest H₂O emission was determined in the straw biochar application, and the highest soil moisture was obtained (Figure 2). Mean H₂O emissions were 6.1, 8.0, 2.2, and 3.4% lower in hazelnut, straw, apple, and sawdust biochar applications than in the control, respectively. These results showed that increased soil moisture decreased H₂O emissions considering a significant ($p < 0.01$) negative linear correlation ($r = -0.654$) between soil moisture and H₂O emissions. Similarly, Yerli & Sahin (2021) indicated that H₂O emissions decrease with increased moisture retention in the soil.

The decrease in H₂O emissions in parallel with the increase in moisture retention in biochar applications can be explained by the porous structure of biochar. Biochar retains moisture not only inside the pores but also between the particles and micropores, so it takes a longer time for the biochar applied soil to lose water and dry out (Cakmakci & Sahin 2022). Ahmad Bhat *et al.* (2022) stated that the improved hydraulic properties of the soil with biochar increased moisture retention in the soil. Ahmed *et al.* (2019) stated that biochar application improved the soil water holding capacity, thus preserving soil moisture and decreasing H₂O emissions, and reported that this contribution was due to the spongy structure of the biochar. Chen *et al.* (2010) also indicated that biochar application increases water storage in pores by improving soil aggregation and soil pore size distribution. The high soil moisture of the straw biochar among biochar applications can be related to its higher organic matter content than other biochar materials (Table 1). Lal (2020) reported that soil organic matter is very important in soil water-holding ability and H₂O emissions from soil. Charles Gould (2015) stated that by increasing the soil organic matter from 1 to 2%, the soil water storage can be increased by approximately 3 l for each 0.0283 m³ of soil. Organic



Figure 1 | Soil moisture contents in (I) biochar treatments daily, (II) daily in the mean of all weeks, and (III) mean of all measurements.




Figure 2 | H₂O emission from soil in (I) biochar treatments daily, (II) daily in the mean of all weeks, and (III) mean of all measurements.

matter additives decrease H₂O emissions from the soil by increasing the pore number and size and distribution of the soil and also the specific surface area of the soil and provide moisture to the soil (Devi *et al.* 2019; Yerli & Sahin 2021).

Soil temperature decreased in biochar applications in the experimental period compared to the control, and the lowest soil temperature was determined in straw and hazelnut biochar applications (Figure 3). The mean soil temperature was found to be 3.9, 4.4, 1.4, and 1.6% lower in hazelnut, straw, apple, and sawdust biochar applications than in the control, respectively. This can be explained by increasing soil moisture, which reduced soil temperature in biochar applications based on a significant ($p < 0.01$) negative linear correlation ($r = -0.625$) between soil moisture and temperature.

Longer preservation of soil moisture can significantly affect the heat capacity and thermal conductivity of the soil, resulting in a decrease in soil temperature. Licht & Kaisi (2005) reported that increased soil moisture causes lower soil temperatures. Soil temperature tends to decrease depending on the cooling of the surface soil and the increase in moisture values in the soil by supplying water to the soil, thus changing its thermal conductivity (Yerli *et al.* 2022a). Similarly, Mancinelli *et al.* (2015) also stated that water ingress into the soil can create a cooling effect on the soil by increasing soil moisture.

CO₂ emissions decreased in biochar applications in the experimental period compared to the control, and the lowest CO₂ emission was determined in straw biochar application (Figure 4). Mean CO₂ emissions were 51.0, 82.2, 20.3, and 13.2% lower in hazelnut, straw, apple, and sawdust biochar applications than in the control, respectively.

Biochar is a very effective material to decrease the emission of greenhouse gases from the soil, especially CO₂ (He *et al.* 2017). Organic carbons in the soil applied biochar become labile, thereby stabilizing the organic carbon in the soil (Wang *et al.* 2014), and the emission from soils decreases (Deng *et al.* 2017). Biochar can stabilize the carbon stocks of the soil for hundreds of years with its recalcitrant organic carbon (Schmidt *et al.* 2002). Due to the porosity of the biochar integrated with the soil, it absorbs organic carbon in the environment and decreases CO₂ emissions from the soil (Lehmann 2007). Case *et al.* (2014) stated that biochar traps the CO₂ emissions from the soil in its structure. Ge *et al.* (2020) indicated that biochar applied to soil limits CO₂ emissions from the soil by being effective on the labile carbon fraction, soil aggregates and soil respiration components. In addition, as a result of many studies, it was found that different biochar materials applied to the soil have a positive effect on decreasing CO₂ emissions from the soil (Liu *et al.* 2011; Wang *et al.* 2014; Zhang *et al.* 2017; Wu *et al.* 2018; Azeem *et al.* 2019; Akinyemi & Adesina 2020; Lehmann *et al.* 2021).




Figure 3 | Soil temperature contents in (I) biochar treatments daily, (II) daily in the mean of all weeks, and (III) mean of all measurements.




Figure 4 | CO₂ emission from soil in (I) biochar treatments daily, (II) daily in the mean of all weeks, and (III) mean of all measurements.




Figure 5 | The relationships of CO₂ emissions with soil moisture, H₂O emissions, and soil temperature.

Mukherjee & Lal (2013) reported that the CO₂ emissions from the soil decreased with the application of different biochar materials to the soil, but the emission rates can differ because biochars change the soil properties in different ways depending on their different characteristics. He *et al.* (2017) indicated that the effect of biochar on decreasing CO₂ emissions from the soil may vary depending on the properties of the soil to which the biochar is mixed and its interaction with the soil. Therefore, the lower CO₂ emissions in straw biochar among biochar applications can be explained by its higher EC value (Table 1). Yerli & Sahin (2021) reported that CO₂ emissions vary depending on the EC value of organic residues mixed into the soil. Increasing EC negatively affects soil microorganism activity and weakens oxidation processes (Zhai *et al.* 2011). Sakin & Yanardag (2019) stated that CO₂ emissions decreased with increasing EC values in the soil. In addition, due to the higher organic matter content of the straw biochar (Table 1), changing soil environmental conditions based on changed soil properties with a decrease in mineralization may have decreased CO₂ emissions. Chaudhari *et al.* (2013) reported that bacteria, fungi, and actinomycetes in the soil decreased by 40% with the addition of organic matter to the soil, decreasing the bulk density of the soil and thus, oxidation processes in the soil were adversely affected.

The strong ($p < 0.01$) positive linear relationships of CO₂ emissions with soil moisture, H₂O emissions, and soil temperature (Figure 5) show that emissions can be managed with soil moisture, H₂O emissions, and soil temperature. The relationship of CO₂ emissions with soil moisture and temperature can be explained by the increase in organic carbon mineralization due to increased soil moisture and temperature, while the relationship of CO₂ emissions with H₂O emissions can be evaluated indirectly by inducing H₂O emissions from increased soil moisture. The moisture balance of the soil greatly affects the oxidation of organic carbon in the soil (Shi & Marschner 2014). It is known that re-wetting dried soil triggers microbial activities and increases the emissions (Lamparter *et al.* 2009). Li *et al.* (2010) reported that the amount of irrigation changed the organic matter dynamics in the soil. Entry *et al.* (2008) stated that the continuous supply of water to the soil increases emissions by increasing organic carbon decomposition. Yerli *et al.* (2022a) stated in their study that examined the CO₂ emissions at different irrigation water levels that the soil microorganism activity slowed down with the decreasing amount of irrigation water, and thus the CO₂ emissions also decreased. In addition, as a result of many studies, positive linear relationships were determined between CO₂ emissions and soil temperature, similar to this study (Chen *et al.* 2018; Du *et al.* 2019; Zhao *et al.* 2020). Fan *et al.* (2021) reported that soil temperature has a direct effect on the population and number of microorganisms. Gonzalez-Mendez *et al.* (2015) determined that CO₂ emissions increased as a result of the acceleration of microorganism activities that carry out mineralization processes depending on the increasing soil temperature. Similarly, Jabro *et al.* (2008) stated that soil temperature increased CO₂ emissions by 59%.

CONCLUSION

As a result of this study, it was determined that (i) all biochar applications decreased CO₂ emissions, but the biochar material that decreased the emissions the most was straw biochar; (ii) with the biochar applications, the moisture retention in the soil

increased, and accordingly, the H₂O emissions and soil temperature decreased, and the straw biochar was more effective; and (iii) CO₂ emissions have strong positive linear relationships with soil moisture, H₂O emissions, and soil temperature. It has been concluded that the content and properties of the biochar applied to the soil are very important to decrease the CO₂ emissions from the soil and that the CO₂ emissions can be managed with soil moisture, H₂O emissions, and soil temperature. Therefore, it can be suggested to apply biochar to the soil by evaluating these strategies and to monitor the soil moisture and temperature to decrease emissions for more effective agricultural and environmental management, and it can be recommended to develop studies in this context.

DATA AVAILABILITY STATEMENT

All relevant data are included in the paper or its Supplementary Information.

CONFLICT OF INTEREST

The authors declare there is no conflict.

REFERENCES

- Ahmad Bhat, S., Kuriqi, A., Dar, M. U. D., Bhat, O., Sammen, S. S., Towfiqul Islam, A. R. M., Elbeltagi, A., Shah, O., Al-Ansari, N., Ali, R. & Heddami, S. 2022 Application of biochar for improving physical, chemical, and hydrological soil properties: a systematic review. *Sustainability* **14** (17), 11104.
- Ahmed, R., Li, Y., Mao, L., Xu, C., Lin, W., Ahmed, S. & Ahmed, W. 2019 Biochar effects on mineral nitrogen leaching, moisture content, and evapotranspiration after 15N urea fertilization for vegetable crop. *Agronomy* **9** (6), 331.
- Akinyemi, B. A. & Adesina, A. 2020 Recent advancements in the use of biochar for cementitious applications: a review. *Journal of Building Engineering* **32**, 101705.
- Apori, S. O., Byalebeka, J., Murongo, M., Ssekandi, J. & Noel, G. L. 2021 Effect of coapplied corncob biochar with farmyard manure and NPK fertilizer on tropical soil. *Resources, Environment and Sustainability* **5**, 100034.
- Avramidis, P., Nikolaou, K. & Bekiari, V. 2015 Total organic carbon and total nitrogen in sediments and soils: a comparison of the wet oxidation-titration method with the combustion-infrared method. *Agricultural Science Procedia* **4**, 425–430.
- Azeem, M., Hayat, R., Hussain, Q., Ahmed, M., Pan, G., Tahir, M. I., Imran, M., Irfan, M. & Hassan, M. 2019 Biochar improves soil quality and N₂-fixation and reduces net ecosystem CO₂ exchange in a dryland legume–cereal cropping system. *Soil and Tillage Research* **186**, 172–182.
- Bennetzen, E. H., Smith, P. & Porter, J. R. 2016 Agricultural production and greenhouse gas emissions from world regions-The major trends over 40 years. *Global Environmental Change* **37**, 43–55.
- Blanco-Canqui, H. 2017 Biochar and soil physical properties. *Soil Science Society of America Journal* **81** (4), 687–711.
- Bremner, J. M. & Mulvaney, C. S., 1982 Nitrogen-Total 1. In: *Methods of Soil Analysis, Part 2, Physical and Mineralogical Methods* (Klute, A., ed.). Agronomy Society of America and Soil Science Society America, Madison, WI.
- Buragiene, S., Sarauskis, E., Romaneckas, K., Adamaviciene, A., Kriauciuniene, Z., Avizienyte, D., Marozas, V. & Naujokiene, V. 2019 Relationship between CO₂ emissions and soil properties of differently tilled soils. *Science of the Total Environment* **662**, 786–795.
- Cakmakci, T. & Sahin, U. 2022 Yield, physiological responses and irrigation water productivity of capia pepper (*Capsicum annuum* L.) at deficit irrigation and different biochar levels. *Gesunde Pflanzen*, 1–11.
- Case, S. D., McNamara, N. P., Reay, D. S. & Whitaker, J. 2014 Can biochar reduce soil greenhouse gas emissions from a *M. iscanthus* bioenergy crop? *Gcb Bioenergy* **6** (1), 76–89.
- Charles Gould, M. 2015 *Compost Increases Water Holding Capacity of Droughty Soils*. Available from: <https://www.manuremanager.com/compost-increases-water-holding-capacity-of-droughty-soils-18241/> (accessed 20 July 2022).
- Chaudhari, P. R., Ahire, D. V., Ahire, V. D., Chkravarty, M. & Maity, S. 2013 Soil bulk density as related to soil texture, organic matter content and available total nutrients of Coimbatore soil. *International Journal of Scientific and Research Publications* **3** (2), 1–8.
- Chen, Y., Shinogi, Y. & Taira, M. 2010 Influence of biochar use on sugarcane growth, soil parameters, and groundwater quality. *Soil Research* **48** (7), 526–530.
- Chen, H., Hou, H., Wang, X., Zhu, Y., Saddique, Q., Wang, Y. & Cai, H. 2018 The effects of aeration and irrigation regimes on soil CO₂ and N₂O emissions in a greenhouse tomato production system. *Journal of Integrative Agriculture* **17** (2), 449–460.
- Chintala, R., Mollinedo, J., Schumacher, T. E., Malo, D. D. & Julson, J. L. 2014 Effect of biochar on chemical properties of acidic soil. *Archives of Agronomy and Soil Science* **60** (3), 393–404.
- Deng, W., Van Zwieten, L., Lin, Z., Liu, X., Sarmah, A. K. & Wang, H. 2017 Sugarcane bagasse biochars impact respiration and greenhouse gas emissions from a latosol. *Journal of Soils and Sediments* **17** (3), 632–640.
- Devi, N. S., Nongmeikakpam, G. & Devi, T. S., 2019 Organic manures for improving soil physical properties. Chapter 2. In: *Current Research in Soil Science* (Kumar, N., ed.). AkiNik Publications, Faizabad, India.

- Dong, D., Feng, Q., Mcgrouter, K., Yang, M., Wang, H. & Wu, W. 2015 Effects of biochar amendment on rice growth and nitrogen retention in a waterlogged paddy field. *Journal of Soils and Sediments* **15** (1), 153–162.
- Du, Y., Gu, X., Wang, J. & Niu, W. 2019 Yield and gas exchange of greenhouse tomato at different nitrogen levels under aerated irrigation. *Science of the Total Environment* **668**, 1156–1164.
- Entry, J. A., Mills, D., Mathee, K., Jayachandran, K., Sojka, R. E. & Narasimhan, G. 2008 Influence of irrigated agriculture on soil microbial diversity. *Applied Soil Ecology Journal* **40**, 146–154.
- Fan, L., Tarin, M. W. K., Zhang, Y., Han, Y., Rong, J., Cai, X., Chen, L., Shi, C. & Zheng, Y. 2021 Patterns of soil microorganisms and enzymatic activities of various forest types in coastal sandy land. *Global Ecology and Conservation* **28**, e01625.
- Ge, X., Cao, Y., Zhou, B., Xiao, W., Tian, X. & Li, M. H. 2020 Combined application of biochar and N increased temperature sensitivity of soil respiration but still decreased the soil CO₂ emissions in moso bamboo plantations. *Science of the Total Environment* **730**, 139003.
- Gee, G. W. & Bauder, J. W., 1986 Particle-size analysis. In: *Methods of Soil Analysis, Part 1, Physical and Mineralogical Methods* (Klute, A., ed.). Agronomy Society of America and Soil Science Society America, Madison, WI.
- Gonzalez-Mendez, B., Webster, R., Fiedler, S., Loza-Reyes, E., Hernandez, J. M., Ruiz-Suarez, L. G. & Siebe, C. 2015 Short-term emissions of CO₂ and N₂O in response to periodic flood irrigation with waste water in the Mezquital Valley of Mexico. *Atmospheric Environment* **101**, 116–124.
- Haddaway, N. R., Hedlund, K., Jackson, L. E., Katterer, T., Lugato, E., Thomsen, I. K., Jorgensen, H. B. & Isberg, P. E. 2017 How does tillage intensity affect soil organic carbon? A systematic review. *Environmental Evidence* **5** (1), 1–8.
- He, Y., Zhou, X., Jiang, L., Li, M., Du, Z., Zhou, G., Shao, J., Wang, X., Xu, Z., Bai, S. H., Wallance, H. & Xu, C. 2017 Effects of biochar application on soil greenhouse gas fluxes: a meta-analysis. *Gcb Bioenergy* **9** (4), 743–755.
- Hilscher, A., Heister, K., Siewert, C. & Knicker, H. 2009 Mineralisation and structural changes during the initial phase of microbial degradation of pyrogenic plant residues in soil. *Organic Geochemistry* **40** (3), 332–342.
- Jabro, J. D., Sainju, U., Stevens, W. B. & Evans, R. G. 2008 Carbon dioxide flux as affected by tillage and irrigation in soil converted from perennial forages to annual crops. *Journal of Environmental Management* **88** (4), 1478–1484.
- Jing, F., Chen, X., Wen, X., Liu, W., Hu, S., Yang, Z., Guo, B., Luo, Y., Yu, Q. & Xu, Y. 2020 Biochar effects on soil chemical properties and mobilization of cadmium (Cd) and lead (Pb) in paddy soil. *Soil Use and Management* **36** (2), 320–327.
- Karhu, K., Mattila, T., Bergström, I. & Regina, K. 2011 Biochar addition to agricultural soil increased CH₄ uptake and water holding capacity-Results from a short-term pilot field study. *Agriculture, Ecosystems & Environment* **140** (1–2), 309–313.
- Kul, R., Arjumend, T., Ekinci, M., Yildirim, E., Turan, M. & Argin, S. 2021 Biochar as an organic soil conditioner for mitigating salinity stress in tomato. *Soil Science and Plant Nutrition* **67** (6), 693–706.
- Laghari, M., Mirjat, M. S., Hu, Z., Fazal, S., Xiao, B., Hu, M., Chen, Z. & Guo, D. 2015 Effects of biochar application rate on sandy desert soil properties and sorghum growth. *Catena* **135**, 313–320.
- Lal, R. 2020 Soil organic matter and water retention. *Agronomy Journal* **112** (5), 3265–3277.
- Lamparter, A., Bachmann, J., Goebel, M. O. & Woche, S. K. 2009 Carbon mineralization soil: impact of wetting-drying, aggregation and water repellency. *Geoderma* **150**, 324–333.
- Lehmann, J. 2007 A handful of carbon. *Nature* **447**, 143–144.
- Lehmann, J., Cowie, A., Masiello, C. A., Kammann, C., Woolf, D., Amonette, J. E., Cayuela, L. M., Camps-Arbestain, M. & Whitman, T. 2021 Biochar in climate change mitigation. *Nature Geoscience* **14** (12), 883–892.
- Li, X., Liu, F., Li, G., Lin, Q. & Jensen, C. R. 2010 Soil microbial response, water and nitrogen use by tomato under different irrigation regimes. *Agriculture Water Management* **98**, 414–418.
- Licht, M. A. & Al-Kaisi, M. 2005 Strip-tillage effect on seedbed soil temperature and other soil physical properties. *Soil and Tillage Research* **80** (1–2), 233–249.
- Liu, Y., Yang, M., Wu, Y., Wang, H., Chen, Y. & Wu, W. 2011 Reducing CH₄ and CO₂ emissions from waterlogged paddy soil with biochar. *Journal of Soils and Sediments* **11** (6), 930–939.
- Luo, Y., Durenkamp, M., De Nobili, M., Lin, Q., Devonshire, B. J. & Brookes, P. C. 2013 Microbial biomass growth, following incorporation of biochars produced at 350 C or 700 C, in a silty-clay loam soil of high and low pH. *Soil Biology and Biochemistry* **57**, 513–523.
- Mancinelli, R., Marinari, S., Brunetti, P., Radicetti, E. & Campiglia, E. 2015 Organic mulching, irrigation and fertilization affect soil CO₂ emission and C storage in tomato crop in the Mediterranean environment. *Soil and Tillage Research* **152**, 39–51.
- Mohamed, E. M., El-Naggar, A. H., Usman, A. R. & Al-Wabel, M. 2015 Dynamics of CO₂ emission and biochemical properties of a sandy calcareous soil amended with *Conocarpus* waste and biochar. *Pedosphere* **25** (1), 46–56.
- Mukherjee, A. & Lal, R. 2013 Biochar impacts on soil physical properties and greenhouse gas emissions. *Agronomy* **3** (2), 313–339.
- Nelson, D. W. & Sommers, L. E., 1982 Total carbon, organic carbon, and organic matter. In: *Methods of Soil Analysis, Part 2, Physical and Mineralogical Methods* (Klute, A., ed.). Agronomy Society of America and Soil Sci. Society America, Madison, WI.
- Rechberger, M. V., Kloss, S., Rennhofer, H., Tintner, J., Watzinger, A., Soja, G., Lichtenegger, H. & Zehetner, F. 2017 Changes in biochar physical and chemical properties: accelerated biochar aging in an acidic soil. *Carbon* **115**, 209–219.
- Sakin, E. & Yanardag, I. H. 2019 Effect of application of sheep manure and its biochar on carbon emissions in salt affected calcareous soil in Sanliurfa Region SE Turkey. *Fresenius Environmental Bulletin* **28** (4), 2553–2560.
- Schmidt, M. W., Skjemstad, J. O. & Jäger, C. 2002 Carbon isotope geochemistry and nanomorphology of soil black carbon: Black chernozemic soils in central Europe originate from ancient biomass burning. *Global Biogeochemical Cycles* **16** (4), 1123.

- Shi, A. D. & Marschner, P. 2014 Drying and rewetting frequency influences cumulative respiration and its distribution over time in two soils with contrasting management. *Soil Biology and Biochemistry* **72**, 172–179.
- Singh, B. P., Cowie, A. L. & Smernik, R. J. 2012 Biochar carbon stability in a clayey soil as a function of feedstock and pyrolysis temperature. *Environmental Science & Technology* **46** (21), 11770–11778.
- Somerville, P. D., Farrell, C., May, P. B. & Livesley, S. J. 2020 Biochar and compost equally improve urban soil physical and biological properties and tree growth, with no added benefit in combination. *Science of the Total Environment* **706**, 135736.
- Tufenkci, S., Sahin, U., Cakmakci, T., Yerli, C., 2022 Biochar and drought. In: *Modern Agricultural Practices* (Yilmaz, A. & Soysal, S., eds). Iksad Publications, Ankara, Turkey.
- Wang, Z., Li, Y., Chang, S. X., Zhang, J., Jiang, P., Zhou, G. & Shen, Z. 2014 Contrasting effects of bamboo leaf and its biochar on soil CO₂ efflux and labile organic carbon in an intensively managed Chinese chestnut plantation. *Biology and Fertility of Soils* **50** (7), 1109–1119.
- WMO 2019 *World Meteorological Organization, Greenhouse Gas Bulletin*. Available from: https://library.wmo.int/doc_num.php?explnum_id=5455 (accessed 28 September 2022).
- Wu, D., Senbayram, M., Zang, H., Ugurlar, F., Aydemir, S., Brüggemann, N., Kuzyakov, Y., Bol, R. & Blagodatskaya, E. 2018 Effect of biochar origin and soil pH on greenhouse gas emissions from sandy and clay soils. *Applied Soil Ecology* **129**, 121–127.
- Yerli, C. & Sahin, U. 2021 Effect of different manure applications and wetting-drying cycles on CO₂ emissions from soil. *Environmental Engineering and Management Journal* **20** (9), 317–324.
- Yerli, C., Sahin, U., Cakmakci, T. & Tufenkci, S. 2019 Effects of agricultural applications on CO₂ emission and ways to reduce. *Turkish Journal of Agriculture-Food Science and Technology* **7** (9), 1446–1456.
- Yerli, C., Sahin, U. & Oztas, T. 2022a CO₂ emission from soil in silage maize irrigated with wastewater under deficit irrigation in direct sowing practice. *Agricultural Water Management* **271**, 107791.
- Yerli, C., Sahin, U., Kiziloglu, F. M., Oztas, T. & Ors, S. 2022b Deficit irrigation with wastewater in direct sowed silage maize reduces CO₂ emissions from soil by providing carbon savings. *Journal of Water and Climate Change* **13** (7), 2837–2846.
- Zhai, L. M., Liu, H. B., Zhang, J. Z., Huang, J. & Wang, B. R. 2011 Long-term application of organic manure and mineral fertilizer on N₂O and CO₂ emissions in a red soil from cultivated maize-wheat rotation in China. *Agricultural Sciences in China* **10** (11), 1748–1757.
- Zhang, A., Cheng, G., Hussain, Q., Zhang, M., Feng, H., Dyck, M., Sun, B., Zhao, Y., Chen, H., Chen, J. & Wang, X. 2017 Contrasting effects of straw and straw-derived biochar application on net global warming potential in the Loess Plateau of China. *Field Crops Research* **205**, 45–54.
- Zhang, C., Li, X., Yan, H., Ullah, I., Zuo, Z., Li, L. & Yu, J. 2020 Effects of irrigation quantity and biochar on soil physical properties, growth characteristics, yield and quality of greenhouse tomato. *Agricultural Water Management* **241**, 106265.
- Zhao, P., Pumpanen, J. & Kang, S. 2020 Spatiotemporal variability and controls of soil respiration in a furrow-irrigated vineyard. *Soil and Tillage Research* **196**, 104424.
- Zimmerman, A. R., Gao, B. & Ahn, M. Y. 2011 Positive and negative carbon mineralization priming effects among a variety of biochar-amended soils. *Soil Biology and Biochemistry* **43** (6), 1169–1179.
- Zoghi, Z., Hosseini, S. M., Kouchaksaraei, M. T., Kooch, Y. & Guidi, L. 2019 The effect of biochar amendment on the growth, morphology and physiology of *Quercus castaneifolia* seedlings under water-deficit stress. *European Journal of Forest Research* **138** (6), 967–979.

First received 9 August 2022; accepted in revised form 10 November 2022. Available online 24 November 2022