

## Developmental effects and genotoxicity of 10 water disinfection by-products in zebrafish

Elisabet Teixidó, Esther Piqué, Javier Gonzalez-Linares, Joan M. Llobet and Jesús Gómez-Catalán

### ABSTRACT

Disinfection by-products are contaminants produced during drinking water disinfection. Several DBPs have been implicated in a variety of toxic effects, mainly carcinogenic and genotoxic effects. Moreover, DBPs exposure has also been associated with an increased risk of developmental effects. In this study, the developmental toxicity and genotoxicity of 10 DBPs (four trihalomethanes [THMs], five haloacetic acids [HAAs] and sodium bromate) in the zebrafish embryo model were evaluated. Embryos exposed for 72 hours were observed for different endpoints such as growth, hatching success, malformations and lethality. THMs exposure resulted in adverse developmental effects and a significant reduced tail length. Two HAAs, tribromoacetic acid and dichloroacetic acid, along with sodium bromate were found to cause a significant increase in malformation rate. Chloroform, chlorodibromomethane and sodium bromate produced a weak induction of DNA damage to whole embryos. However, developmental effects occurred at a range of concentrations (20–100 µg/mL) several orders of magnitude above the levels that can be attained in fetal blood in humans exposed to chlorinated water. In conclusion, the teratogenic and genotoxic activity observed by some DBPs in zebrafish reinforce the view that there is a weak capacity of disinfection products to cause developmental effects at environmentally relevant concentrations.

**Key words** | comet assay, developmental toxicity, haloacetic acids, trihalomethanes, water disinfection by-products, zebrafish

Elisabet Teixidó (corresponding author)  
Esther Piqué  
Joan M. Llobet  
Jesús Gómez-Catalán  
GRET-CERETOX and Toxicology Unit,  
Public Health Department,  
School of Pharmacy,  
University of Barcelona. Av. Joan XXIII s/n,  
E-08028,  
Barcelona,  
Spain  
E-mail: [elisabet.teixido@gmail.com](mailto:elisabet.teixido@gmail.com)

Javier Gonzalez-Linares  
GRET-CERETOX and Experimental Toxicology and  
Ecotoxicology Unit,  
Barcelona Science Park. Baldiri i Reixac 10-12,  
E-08028,  
Barcelona,  
Spain

### INTRODUCTION

Disinfection by-products (DBPs) are contaminants formed as a consequence of chemical disinfection of public drinking waters. The disinfectants, such as chlorine, can react with natural organic matter in surface waters leading to the formation of a complex mixture of DBPs. So far, more than 600 DBPs have been identified and reported in the literature; nevertheless, they represent less than half of all possible environmental DBPs. The most prevalent DBPs include the four trihalomethanes (THMs), chloroform, bromoform, chlorodibromomethane (CDBM) and bromodichloromethane (BDCM), the group of haloacetic acids (HAAs) and bromate anion. DBPs in drinking water are generally present at sub-µg/L or low- to mid-µg/L levels (Richardson *et al.* 2007) and some countries have regulated

the levels of some DBPs in drinking water (Table 1, US Environmental Protection Agency (EPA) 2006). Moreover, World Health Organization (WHO) guidelines exist as well as European Union (EU) DBP standards (Table 1, EU Directive 98/83/EC 1998; WHO 2004).

Several DBPs have been confirmed as mutagenic, genotoxic and/or carcinogenic in different test systems (Richardson *et al.* 2007). There is an increasing concern about the association of DBPs exposure with adverse developmental effects. Some individual DBPs have been found to cause developmental toxicity in mammalian assays at high doses (Ruddick *et al.* 1983; Narotsky *et al.* 1996; Christian *et al.* 2001a). Some DBPs induce specific congenital malformations of the cardiovascular and neurological system, but,

doi: 10.2166/wh.2014.006

**Table 1** | DBP regulations and guideline values (highest concentration allowed in drinking water)**US EPA, Safe Drinking Water Act (SDWA)<sup>a</sup>**

DBP	MCL (mg/L)
Total THMs	0.08
Haloacetic acids (HAA5)	0.06
Bromate	0.01

**World Health Organization (WHO) guidelines<sup>b</sup>**

DBP	Guideline value (mg/L)
Chloroform	0.2
Bromoform	0.1
CDBM	0.1
BDCM	0.06
DCA	0.05 <sup>c</sup>
TCA	0.2
Bromate	0.01 <sup>c</sup>

**European Union Standards**

DBP	Standard value (mg/L)
Total THMs	0.1
Bromate	0.01 <sup>d</sup>

<sup>a</sup>The total THMs represent the sum of the concentrations of four THMs: chloroform, bromoform, BDCM, and CDBM. The HAAs represent the sum of monochloro-, dichloro-, trichloro-, monobromo-, and DBA (US EPA 2006).

MCL: maximum contaminant levels.

<sup>b</sup>WHO guidelines on THMs state that for authorities wishing to establish a total THM standard to account for additive toxicity the sum of the ratio of the concentration of each THM to its respective guideline value should not exceed unity.

<sup>c</sup>Provisional guideline value.

<sup>d</sup>Where possible, without compromising disinfection, EU member states should strive for a lower value. This value must be met, at the latest, 10 calendar years after the issue of Directive (EU Directive 98/83/EC 1998); within 5 years of the Directive, a value of 0.025 mg/L must be met.

in general, fetal body weight reduction is often reported as the major effect (Epstein *et al.* 1992; Hunter III *et al.* 1996). Recent studies had evaluated the developmental effects of complex mixtures and these appeared to exert no adverse developmental effects (Narotsky *et al.* 2012). The epidemiologic studies found inconsistent results or very weak associations for congenital anomalies/birth defects, central nervous system anomalies, neural tube defects and spontaneous abortion. However, these studies suggested a positive association with some measure of growth retardation (reviewed in Graves *et al.* 2001; Tardiff *et al.* 2006; Colman *et al.* 2011).

Chemical treatment of public water supplies is designed to kill pathogens that may exist in the drinking water, so the risk-benefit balance of water disinfection is considered positive. However some factors, such as the huge magnitude of the population affected, the distorted perception by the population of chemical risks, the availability of several alternative water treatment methodologies, the uncertainties in the hazard characterization of DBPs and the intrinsic limitations of the epidemiological studies, warrant the requirement of a better knowledge about DBPs toxicity and mechanisms of action (Colman *et al.* 2011). In this study, we have explored the capabilities of the zebrafish embryo model in order to identify and characterize the potential embryotoxic and genotoxic effects of some DBPs.

Zebrafish (*Danio rerio*) is a prominent model vertebrate in developmental genetics, toxicology and ecotoxicology (Postlethwait *et al.* 2000; Hill *et al.* 2005; Scholz *et al.* 2008). Size, easy husbandry, high fecundity and fast development represent the main benefits of using zebrafish over other vertebrate species. In addition, zebrafish embryos are transparent and develop outside the mother. Hence, morphological structures and internal organs can be easily visualized (Zhang *et al.* 2003). Moreover, the mechanisms of embryogenesis are well conserved along the vertebrates and there are several studies that confirm the ability of the zebrafish model to predict the teratogenic potential of chemicals in mammals (Brannen *et al.* 2010; Padilla *et al.* 2012; Selderslaghs *et al.* 2012). Hence, the zebrafish is increasingly used for assessing developmental toxicity of chemicals.

Currently, 'whole-mixture' approaches are being used to address concerns related to the potential adverse health effects of DBPs exposure. Scientists from the US EPA office of Research and Development proposed an experimental design for a multigenerational reproductive/developmental bioassay to optimize the probability of detecting adverse effects (Simmons *et al.* 2008). In the present study, the capability of the zebrafish embryo, as an alternative model, was explored in order to investigate the developmental effects and genotoxicity of DBPs. The selected DBPs – four THMs (CDBM and BDCM), five HAAs (dichloroacetic acid [DCA], trichloroacetic acid [TCA], dibromoacetic acid [DBA], tribromoacetic acid [TBA] and bromochloroacetic acid [BCA]) and sodium bromate – represented the most prevalent compounds and

data were available for comparison with other test systems and models.

## MATERIALS AND METHODS

### Chemicals and test media

DBA, TCA, chloroform, bromoform and sodium bromate were purchased from Sigma-Aldrich (St Louis, MO). DCA and TBA were delivered from Tokyo Chemical Industry (Tokyo, Japan). BCA, BDCM and CDBM were purchased from Alfa Aesar (Karlsruhe, Germany). Buffered embryo medium (17.4 mM NaCl; 0.23 mM KCl; 0.12 mM MgSO<sub>4</sub> · 7 H<sub>2</sub>O; 0.18 mM Ca(NO<sub>3</sub>)<sub>2</sub>; 1.5 mM HEPES; pH 7.4) was used as the medium for all solutions during the experiments to keep the pH stable and constant between assays (Gustafson *et al.* 2012).

### Embryo exposure

All stock solutions were prepared with buffered embryo medium except for THMs that were initially prepared in 100% dimethylsulfoxide (DMSO) and subsequently diluted in buffered embryo medium with a final DMSO concentration of 0.1% (v/v).

For all substances, a concentration range-finding experiment was conducted with a constant spacing factor 2. The range finding test allow us to select the final tested concentrations based on the presence of a 0 and 100% effect level (for both malformation and mortality). Each substance was tested in 5–7 concentrations with a negative control, test medium only or solvent control with 0.1% of DMSO. Exposure concentrations are anticipated to be stable throughout the test duration.

Zebrafish embryos were collected by natural spawning and staged according to Kimmel *et al.* (1995). Fertilization success was checked and only batches of eggs with a fertilization rate of at least 80% were used. Exposures of embryos began at 4 hours post-fertilization (hpf) and were incubated at 27 ± 1 °C on a 14-h light and 10-h dark cycle for 72 hours. The exposure was semi-static and solutions were renewed every 24 hours.

Embryos were exposed to HAAs and sodium bromate in a 6-well culture plate (Greiner Bio-one, Germany). Ten

embryos were randomly distributed into wells and filled with 5 mL of each solution. Each 6-well plate held five different concentrations of the test compound and the negative or solvent control. In order to prevent losses by volatilization, THMs were tested in 20 mL hermetically sealed glass vials. Ten embryos per vial and treatment were exposed with 10 mL of each test solution. For each substance, three independent exposure experiments were conducted using eggs from independent spawning events ( $n = 3$ ).

### Evaluation of developmental effects

At 8, 28, 52 and 76 hpf, mortality of embryos was checked using a stereo microscope (SMZ-168, Motic). According to Nagel (2002), four endpoints were considered as indicators of mortality: coagulation of eggs, non-development of somites, non-detachment of the tail and no presence of heartbeat. Dead embryos were removed daily after assessment of mortality rate. The fraction of dead embryos at the end of the test was used to calculate median lethal concentration ( $LC_{50}$ ) values.

The teratogenic effects were evaluated and recorded as described in Teixidó *et al.* (2013) (Table 2). The fraction of abnormal embryos was determined for each concentration and control group in order to calculate median effective concentration ( $EC_{50}$ ) values. The frequency of teratogenic effects in all tested groups of a test substance was also analyzed. If the following criteria were fulfilled, (a) concentration-response relationship and (b) the endpoint is observed in ≥50% of all embryos showing malformations, the effect was considered to be a distinctive or identifying malformation (fingerprint endpoint) for this substance (Weigt *et al.* 2011). Embryos were observed using a stereo microscope, and images were obtained with a camera (MotiCam 2000, Motic). Image processing was performed in Image J 1.41 (available at <http://rsb.info.nih.gov/ij/>) and Adobe Photoshop CS3 (Adobe Systems Inc., USA). Test concentrations are expressed in nominal concentrations.

### Hatching success and tail length measurement

From 48 hpf the embryos are able to hatch. Hatching success was recorded at 76 hpf as the percentage of embryos that hatched with respect to surviving embryos. The

**Table 2** | Lethal and teratogenic effects evaluated in zebrafish embryos at 76 hpf

Type	Physiological/dysmorphogenic effect	Description
Lethal	Coagulated egg	Denaturated fish egg. No clear structures of the embryo are observable anymore.
	Non-detachment of the tail from the yolk	From 16 hpf the tail begins to detach from the yolk and to extend.
	Non-development of somites	The somites are structures of the early trunk or tail segments that will eventually form the skeletal muscle, skin and cartilage.
	Lack of heartbeat	
Teratogenic	Malformation of the chorda	No tail, malformation of chorda or spinal cord.
	Malformation of the eyes	Abnormal pigmentation, small eyes or asymmetric eyes.
	Malformation of the ear	Formation of no, one or more than two otoliths per sacculus. Absence or abnormally shaped vesicles.
	Malformation of the head	Brain necrosis, hemorrhage or small head.
	Malformation of the heart	Pericardial edema, big heart, hemorrhage or abnormal chambers.
	Malformation of the tail	Hemorrhage, tail necrosis, bent tail, bent or twisted tip tail.
	Abnormal pigmentation	

malformations observed in each embryo could affect its movement and consequently reduce or delay the hatching. Therefore, motility was also assessed by touch evoked response as a complementary endpoint.

Embryos that had not yet hatched were dechorionated. All embryos were anesthetized with buffered tricaine methanesulfonate (MMS) (0.5 mM, Sigma-Aldrich, St Louis, MO) and photographed (Moticam 2000, Motic) positioned on their lateral side in order to measure the distance between the anus and the posterior end of the notochord, defined as tail length (Bachmann 2002). The minimum concentration to inhibit growth (MCIG) is defined as the minimum concentration to significantly produce a decrease in tail length.

### Cell isolation and alkaline comet assay

After exposure for 72 h, seven surviving fish embryos for the treatments and control groups were processed for cell isolation and comet assay. Cell isolation was carried out mechanically according to the protocol by Kosmehl *et al.* (2006). The  $EC_{50}$  values from teratogenic effects were used as the test concentrations for genotoxicity testing.

The alkaline single cell gel electrophoresis (SCGE) or comet assay was performed as described by Singh *et al.* (1988) with some modifications. Fifty microliters of cell suspension were added to a tube containing 100  $\mu$ L of 0.9% low

melting point agarose (37 °C). The suspension was added to an agarose pre-coated slide and gently covered with a cover slide to make a micro gel. The gel was allowed to solidify for 10 min at room temperature and 6 min at –20 °C. Immersion in lysis buffer (2.5 M NaCl, 100 mM disodium EDTA and 10 mM Tris, pH 10) containing lauryl sarcosine 1% (v/v), Triton 1% (v/v) and DMSO 10% (v/v) in the dark was performed for 1.5 h at 4 °C. The slides were then placed in a horizontal gel electrophoresis unit, immersed in cold alkaline electrophoresis buffer (300 mM NaOH and 1 Mm Na<sub>2</sub>EDTA, pH >13.5), and left in solution for 20 min at 4 °C. After electrophoresis in the same buffer at 25 V and 300 mA for 20 min, slides were neutralized by washing three times with 0.4 M Tris buffer at pH 7.5.

DNA was stained with 20  $\mu$ L of DAPI solution (4',6-diamidino-2-phenylindole) and immediately analyzed using a Nikon E600 fluorescence microscope. DNA damage as a percentage of DNA in the tail was measured using the Comet Assay IV software (Perceptive Instruments, Suffolk, UK). Zebrafish *in vivo* exposure to methyl MMS during 72 h (from 4 hpf to 76 hpf) was used as a positive control for comet assay.

### Data evaluation

Concentration-response curves for mortality and teratogenicity were plotted for compounds that showed a clear

concentration-response relationship. It has been demonstrated that ignoring control mortality, even at lower than 10%, can lead to biased estimation of  $LC_{50}$  (lethal effects) and  $EC_{50}$  (teratogenic effects) (Hoekstra 1987). Therefore, data were corrected for control mortality with Abbott's formula:  $P_c = (P - P_i)/(100 - P_i) \times 100$  where  $P_c$  is the corrected percentage,  $P$  is the percentage mortality of the treated embryos and  $P_i$  is the percentage mortality of the control embryos (Abbott 1987).

Concentration-response curves were calculated using probit analysis (SPSS 15.0). Confidence intervals (CI) were set at 95%. Based on  $LC_{50}$  and  $EC_{50}$  values, a teratogenic index (TI) was calculated as the ratio  $LC_{50}/EC_{50}$ . In the case where no TI could be calculated, the compound was considered to be a non-teratogen although it could still be embryotoxic. Effective concentration 20 ( $EC_{20}$ ) values were calculated for comparison between zebrafish embryo and human exposure data.

Statistical analysis was performed with SPSS 15.0. One-way analysis of variance followed by post hoc multi-comparison with the Bonferroni's test was used to analyze homogeneous data of the continuous variables. The Kruskal-Wallis test was used to analyze non-homogeneous data followed by Dunnett's post hoc test. Significance was accepted when  $p < 0.05$ .

## RESULTS AND DISCUSSION

### Exposure of zebrafish embryos to THMs

Exposure to the THMs selected in this study resulted in adverse developmental effects in zebrafish. The cumulative concentration-response curves for lethality and teratogenesis are shown in Figure 1(a). The vehicle control (0.1% DMSO) was not toxic to embryos. The estimated  $LC_{50}$ ,  $EC_{50}$ ,  $EC_{20}$  and TI values at 76 hpf are represented in Table 3. Malformations that occurred with the highest frequency included cardiac edema and tail developmental abnormalities (Table 4 and Figure 1(b)) and are in accordance with those reported by Brennan *et al.* (2005) for tadpoles. In our study, the  $EC_{20}$  values range between 0.11 mM (23 mg/L) for CDBM and 0.7 mM (84.7 mg/L) for chloroform. Using the  $EC_{20}$  to compare the chemicals


a ranking of the compounds in order of decreasing potency can be established as: CDBM > Bromoform  $\approx$  BDCM > Chloroform. CDBM ( $LC_{50} = 0.48$  mM, 100 mg/L and  $EC_{50} = 0.16$  mM, 33.3 mg/L) resulted to be three times more potent than chloroform ( $LC_{50} = 2.4$  mM, 286.5 mg/L and  $EC_{50} = 0.84$  mM, 100.3 mg/L). This order of toxicity was also found by Mattice *et al.* (1981) after exposure of common carp embryos to THMs. All THMs inhibited the growth of the embryo (Table 3), an effect that is often observed when there is a cardiovascular impairment (Billiard *et al.* 1999). Delayed hatching was also observed after THMs exposure and could be due to an inhibition of enzymes involved in hatching or a decreased mobility in the embryo (Von Westernhagen 1988). The developmental abnormalities and the concentration dependent sedative effect observed after THMs exposure partially or completely inhibit movement which may be required for proper hatching (Figure 2).

The  $EC_{50}$  values reported in this study for chloroform ( $EC_{50} = 0.85$  mM) and BDCM ( $EC_{50} = 0.26$  mM) were similar to those reported for the Frog Embryo Teratogenesis Assay with *Xenopus laevis* in Brennan *et al.* (2005) (0.92 mM and 0.4 mM respectively). Based on the TI values (from 2.5 to 3.6), all THMs were found to produce teratogenic effects in zebrafish. Studies in mammals suggest evidence of a fetotoxic response for THMs but not a teratogenic effect (Thompson *et al.* 1974; Ruddick *et al.* 1983). Other mammalian studies have reported a decreased fetal growth, delayed ossification and craniofacial abnormalities in rats treated by chloroform inhalation (Schwetz *et al.* 1974; Murray *et al.* 1979). Rats exposed to chloroform via drinking water have only showed to have an impaired post-natal growth (Lim *et al.* 2004). Brown-Woodman *et al.* (1998) reported a no effect concentration for chloroform of about 1.05 mM in rat whole embryo culture (WEC), similar to the  $EC_{20}$  concentration reported in our study ( $EC_{20} = 0.7$  mM).

### Exposure of zebrafish embryos to five HAAs

The only HAAs that induced some morphological effects were DCA and TBA at high concentrations (Figure 3(b) and Table 3). Embryos exposed to DCA exhibited moderate pericardial edema whereas no lethality was observed up to





**Figure 1** | (a) Concentration-response curves for mortality (o, full line) and teratogenesis (x, dotted line) for CDBM and BDCM exposure from 4–76 hpf. % effect (mean  $\pm$  S.D.) is shown versus the logarithm of concentration tested. (b) Representative images of fish embryos exposed to CDBM and BDCM at 76 hpf.

46.5 mM (Figure 3(a) and (b), Table 4). The estimated  $LC_{50}$ ,  $EC_{50}$ ,  $EC_{20}$  and TI-values are represented in Table 3. DCA were previously found to be developmental toxic in zebrafish by Hassoun *et al.* (2005). Craniofacial abnormalities (reduced mouth and jaw formation), skeletal muscle

deformations and yolk sac and cardiac edema were observed at 144 hpf after an exposure of DCA in a concentration range between 8 and 32 mM. However, Weber *et al.* (2004) did not find any embryotoxic effect up to a concentration of DCA of about 124 mM in *Xenopus laevis*.

**Table 3** | The  $LC_{50}$  and  $EC_{50}$  values are represented with their 95% CI and TI of all water DBPs tested. MCIG and  $EC_{20}$  effect concentrations at 76 hpf are also represented for all substances

Substance	CAS-No.	$LC_{50}$ mM (95% CI)	$EC_{50}$ mM (95% CI)	TI	$EC_{20}$ mM (95% CI)	MCIG mM
Chloroform	67-66-3	2.1 (1.75–2.31)	0.85 (0.75–0.97)	2.5	0.7 (0.6–0.8)	1.26
Bromoform	75-25-2	0.52 (0.45–0.60)	0.20 (0.17–0.23)	2.6	0.15 (0.12–0.17)	0.1
BDCM	75-27-4	0.93 (0.80–1.06)	0.26 (0.20–0.32)	3.6	0.17 (0.10–0.22)	0.3
CDBM	124-48-1	0.45 (0.38–0.54)	0.16 (0.13–0.19)	2.8	0.11 (0.08–0.13)	0.06
DCA	79-43-6	>46.5	28.9 (24.5–33.5)	–	22.1 (16.1–25.8)	46.5
TCA	76-03-9	>42.8	>42.8	–	>42.8	> 42.8
DBA	631-64-1	>20	>20	–	>20	10
TBA	75-96-7	12.7 (11.6–13.9)	5.7 (4.8–6.5)	2.2	4.4 (3.2–5.2)	2.6
BCA	5589-96-8	>30	>30	–	>30	> 30
Sodium bromate	7789-38-0	65.4 (57.6–73.0)	49.2 (42.0–56.6)	1.3	40.7 (25.3–46.0)	68.3

–, indicates that could not be calculated. Abbreviations used: CDBM (chlorodibromomethane), BDCM (bromodichloromethane), DCA (dichloroacetic acid), TCA (trichloroacetic acid), DBA (dibromoacetic acid), TBA (tribromoacetic acid) and BCA (bromochloroacetic acid).

**Table 4** | Frequency of endpoints observed at 76 hpf in zebrafish embryos exposed to DBPs.  $\Sigma_t$  represents the sum of embryos in all test concentrations showing teratogenic effects for each specific endpoint

Malformation	Chloroform		Bromoform		CDBM		BDCM		TBA		DCA		NaBr	
	$\Sigma_t$	(%)	$\Sigma_t$	(%)	$\Sigma_t$	(%)	$\Sigma_t$	(%)	$\Sigma_t$	(%)	$\Sigma_t$	(%)	$\Sigma_t$	(%)
Chorda	9	24.3	0	0.0	16	28.1	25	36.3	18	27.7	0	0.0	3	12.0
Ear	2	5.4	9	23.7	0	0.0	9	13.0	25	38.5	0	0.0	1	4.0
Head	4	10.8	14	36.8	19	33.3	9	13.0	19	29.2	0	0.0	1	4.0
Eyes	29	<b>78.4</b>	25	<b>65.8</b>	30	<b>52.6</b>	23	33.3	44	<b>67.7</b>	8	22.9	20	<b>80.0</b>
Heart	28	<b>75.7</b>	23	<b>60.5</b>	30	<b>52.6</b>	39	<b>56.5</b>	40	<b>61.5</b>	<b>31</b>	<b>88.6</b>	24	<b>96.0</b>
Tail	29	<b>78.4</b>	27	<b>71.0</b>	31	<b>54.4</b>	37	<b>53.6</b>	48	<b>73.8</b>	12	34.3	10	40.0

Abbreviations used: CDBM (chlorodibromomethane), BDCM (bromodichloromethane), TBA (tribromoacetic acid), DCA (dichloroacetic acid).


The percentage of embryos showing each endpoint was calculated with respect to the total number of embryos with teratogenic effects found in all test concentrations. Bold values indicate that the endpoint followed a concentration-response relationship and was observed in  $\geq 50\%$  of all embryos showing malformations.

DCA have been shown to produce cardiovascular defects when administered to pregnant dams at high dose levels (Epstein *et al.* 1992; Johnson *et al.* 1998).

In our study, TBA exposure caused curvatures of the spinal column, swelling of the pericardia and small eyes (Figure 3(b), Table 4) with an  $EC_{20}$  of 4.4 mM (1,300 mg/L, Table 3). The current data are in agreement with results from Hunter III *et al.* (1996) who reported that brominated acetic acids are potentially more toxic than chlorinated species to the developing embryo. Developmental toxicity studies with frogs (Bantle *et al.* 1999) reported a TI of about 3.86 after TBA treatment, a higher value compared

to our results with zebrafish embryos (TI = 2.2). *In vivo*, TBA exposure was found to not produce any adverse developmental effect (NTP (National Toxicology Program) 1998a). HAAs have been shown to produce neural tube defects, prosencephalic and pharyngeal arch hypoplasia, and heart and eye defects in rats and mice WEC (Hunter III *et al.* 1996; Andrews *et al.* 2004), but at a lower range concentration (0.05–2 mM) compared to zebrafish (10–30 mM).

BCA exposure did not produce any adverse developmental effects at the highest concentration tested (30 mM, Table 3). In BCA-exposed tadpoles, some gut abnormalities and decreased growth were observed only at higher



**Figure 2** | Percentage of embryos (mean  $\pm$  S.E.M) displaying reduced motility after THMs exposure and hatching success of these embryos at 76 hpf. Motility of the embryos was checked after dechlorination if the embryo had not hatched at 76 hpf. \*Significantly different from solvent control group (0),  $p < 0.05$ .

concentrations (57.7 mM and 46 mM, respectively) (Brennan *et al.* 2005). BCA was found to not produce any adverse developmental effect *in vivo* in mammals (NTP 1998b). In our study, TCA exposure did not cause developmental effects (Table 3). *In vivo*, administered TCA to pregnant dams at high dose levels has been shown to produce cardiovascular defects and low weight (Smith *et al.* 1989; Johnson *et al.* 1998). Zebrafish embryo exposure to DBA only produced a decrease in tail length (Table 3). DBA was found to not produce any adverse developmental effect *in vivo* in mammals (Christian *et al.* 2001b).

### Exposure of zebrafish embryos to sodium bromate


Sodium bromate showed embryotoxic effects on zebrafish only at very high concentration ( $LC_{50} = 65.4$  mM, Table 3), probably by an irrelevant non-specific effect. Small eyes and pericardial edema were the predominant sodium bromate-induced malformations in zebrafish embryos (Figure 3(b) and Table 4). As shown in Table 3, the  $EC_{20}$  value after exposure to sodium bromate was 51.7 mM. The

96-h  $LC_{50}$  (8.32 mM) reported for frog embryos after sodium bromate treatment resulted to be about 8 times lower than the  $LC_{50}$  reported for zebrafish embryos in this study (65.4 mM). Studies in mammals have shown that sodium bromate did not cause any adverse developmental effect (Wolf & Kaiser 1996).

### Genotoxicity of water DBPs

DNA-damage caused by water DBPs was investigated using the comet assay. Genotoxicity could represent a potential mechanism leading to developmental disorders and embryo mortality, and also it has been associated with carcinogenic effects provoked by DBPs. The four THMs, the TBA and sodium bromate were selected for the comet assays as they exhibit embryo toxicity in zebrafish. The test concentrations used were the  $EC_{50}$  values from teratogenic effects. Chloroform (0.85 mM) and CDBM (0.16 mM) produced significant DNA damage compared to the control solvent group (Figure 4(b)). If compared to the DNA damage produced by the positive control (Figure 4(a)),






**Figure 3** | (a) Concentration-response curves for mortality (o, full line) and teratogenesis (x, dotted line) for TBA, DCA and sodium bromate exposure from 4–76 hpf. % effect (mean  $\pm$  S.D.) is shown versus the logarithm of concentration tested. (b) Representative images of fish embryos exposed to DCA, TBA and sodium bromate at 76 hpf.

DBPs displayed a weak positive response. Exposure to the positive control (MMS) produced a concentration-dependent increase in the mean percentage of DNA in the tail ( $r = 0.89$ ,  $p < 0.01$ ; Figure 4(a)) and ranged between 4 and 37%. The highest tested concentration of MMS (25 mg/L) produced less than 10% embryo mortality, however all embryos showed developmental effects (data not shown). Many *in vitro* techniques have been used to investigate the mutagenic and genotoxic properties of THMs and HAAs (Richardson *et al.* 2007). These studies have shown that THMs are weak inducers of DNA damage (Landi *et al.* 2003; Zhang *et al.* 2012) and that glutathione S-transferase-theta (GSTT1-1) activity mediated transformation of brominated THMs to mutagenic intermediates (Pegram *et al.*

1997). Although the homolog GSTT1-1 gene for zebrafish has not yet been found, embryos possess a lower GST activity that could play a role not only in the weak mutagenicity observed but also in the low teratogenicity observed. HAAs caused DNA breaks in Chinese hamster ovary cells (Plewa *et al.* 2010) and in the human derived hepatoma cell line (Zhang *et al.* 2012). In our study, the strongest effect was found in sodium bromate-treated embryos with a median damage of 8% of DNA in the tail (Figure 4(c)). It has been reported that bromate induced DNA damage (SCGE assay) in mammalian cells through oxidative damage (Priestley *et al.* 2010).

Exposure to DBPs in humans has been quantitatively assessed by measuring the concentration of DBPs and its



**Figure 4** | (a) Genotoxicity following exposure of whole embryos to increasing concentrations of methyl MMS (from 4 to 76 hpf). DNA damage expressed as percentage of DNA in tail in primary cells derived from 76-h-old zebrafish embryos. (b) Genotoxicity following exposure to the  $EC_{50}$  of chloroform (0.85 mM), bromoform (0.2 mM), BDCM (0.26 mM) and CDBM (0.16 mM). (c) Genotoxicity following exposure to the  $EC_{50}$  of TBA (5.7 mM) and sodium bromate (49.2 mM). For each treatment and control groups, 100 cells were investigated, while  $2 \times 50$  cells on two replicate slides were tested for treatments plots. \*Significantly different from control group (Dunnett's test with  $p < 0.05$ ).

metabolites in blood, urine and exhaled breath after the oral intake of chlorinated water and after the dermal or inhalatory exposure during shower and bath. The reported blood levels in exposed humans are in the pg/mL range, with the highest levels of about 300 pg/mL for chloroform after a shower with a tap water surpassing the current standards (Nieuwenhuijsen *et al.* 2000; Nuckols *et al.* 2005). Transplacental crossing of some DBPs in blood at concentrations equal to or greater than in maternal blood has been demonstrated in rodents and humans (Dowty *et al.* 1976; Danielsson *et al.* 1986; Christian *et al.* 2001a). In this study, effect concentrations of 20% were between 0.11–0.7 mM (20–100  $\mu\text{g}/\text{mL}$ ) after DBPs exposure in zebrafish, several orders of magnitude above the levels to which human embryos would be exposed through their mothers' blood.

It should be noted that it has yet not been demonstrated that plasma concentrations in mammals relate to toxicity effect concentrations similarly to the way exposure concentrations relate to toxicity in the fish embryo test.

One of the alleged weaknesses of the zebrafish embryo as a model for teratogenicity in mammals is the difference in metabolic activity towards exogenous substances. This is especially relevant in the case of xenobiotics whose toxicity is mediated by their metabolites. It is well known that some of the toxic effects of halogenated short-chain hydrocarbons are mediated by intermediate electrophilic metabolites. Zebrafish have a total of 94 CYP genes, distributed among 18 gene families, most of which are direct orthologs of human CYPs. Most of these CYPs are expressed in embryos during various time courses along the first 48 hours after fertilization. Indeed,

some maternally-derived CYPs RNA transcripts are present in the unfertilized egg (Goldstone *et al.* 2010). Jones *et al.* (2010) have further demonstrated the expression of several xenobiotic metabolizing genes similar to human genes (CYP1A, CYP2B6, CYP3A5, UGT1A1) and their functional capacity metabolizing some model compounds during the early development. Therefore, the zebrafish embryo is endowed with a wide spectrum metabolic capacity, but the capability of metabolic transformation in comparison to mammalian models is still not fully understood (Hill *et al.* 2012). To our knowledge, there are no data about the capacity of the zebrafish embryo to metabolize THMs or HAAs. However, there is evidence of the capacity to bioactivate THMs by other fish species (Räbergh & Lipsky 1997; Vega-López *et al.* 2012).

## CONCLUSIONS

Effect concentrations in zebrafish embryos support previous observation of a weak teratogenic and genotoxic potential of DBPs. The proximity of effect concentration of lethality and malformations suggest that probably the teratogenic effects are related to unspecific embryo toxicity. The effects are observed only at concentration levels well above those that can be attained in the fetal blood in humans exposed to chlorinated water, providing further evidence for only a weak teratogenic potential of DBP products. However, more studies are needed to explore the involvement of metabolism in the potential DBP toxicity and to extend our knowledge about exposure to mixtures and the possible developmental effects and genotoxicity. Finally, our study indicates that zebrafish embryos are as sensitive as other test systems and can be used as a potential screening and prioritization tool to assess a large number of DBPs.

## REFERENCES

- Abbott, W. S. 1987 A method of computing the effectiveness of an insecticide. 1925. *J. Am. Mosq. Control Assoc.* **3** (2), 302–303.
- Andrews, J. E., Nichols, H. P., Schmid, J. E., Mole, L. M., Hunter III, E. S. & Klinefelter, G. R. 2004 Developmental toxicity of mixtures: The water disinfection by-products dichloro-, dibromo- and bromochloro acetic acid in rat embryo culture. *Reprod. Toxicol.* **19** (1), 111–116.
- Bachmann, J. 2002 Entwicklung eines Teratogenitäts – Screening – Tests mit Embryonen des Zebraärlings *Danio rerio*. Ph.D. Thesis, Technical University Dresden, Germany.
- Bantle, J. A., Finch, R. A., Fort, D. J., Stover, E. L., Hull, M., Kumsher-King, M. & Gaudet-Hull, A. M. 1999 Phase III interlaboratory study of FETAX. Part 3. FETAX validation using 12 compounds with and without an exogenous metabolic activation system. *J. Appl. Toxicol.* **19** (6), 447–472.
- Billiard, S. M., Querbach, K. & Hodson, P. V. 1999 Toxicity of retene to early life stages of two freshwater fish species. *Environ. Toxicol. Chem.* **18** (9), 2070–2077.
- Brannen, K. C., Panzica-Kelly, J. M., Danberry, T. L. & Augustine-Rauch, K. A. 2010 Development of a zebrafish embryo teratogenicity assay and quantitative prediction model. *Birth Defects Res. B Dev. Reprod. Toxicol.* **89** (1), 66–77.
- Brennan, L. M., Toussaint, M. W., Kumsher, D. M., Dennis, W. E., Rosencrance, A. B., Brown, C., Van Der Schalie, W. H. & Gardner, H. S. 2005 Developmental toxicity of drinking water disinfection by-products to embryos of the African clawed frog (*Xenopus laevis*). *Bull. Environ. Contam. Toxicol.* **75** (2), 361–367.
- Brown-Woodman, P. D. C., Hayes, L. C., Huq, F., Herlihy, C., Picker, K. & Webster, W. S. 1998 In vitro assessment of the effect of halogenated hydrocarbons: Chloroform, dichloromethane, and dibromoethane on embryonic development of the rat. *Teratology* **57** (6), 321–333.
- Christian, M. S., York, R. G., Hoberman, A. M., Diener, R. M. & Fisher, L. C. 2001a Oral (drinking water) developmental toxicity studies of bromodichloromethane (BDCM) in rats and rabbits. *Int. J. Toxicol.* **20** (4), 225–237.
- Christian, M. S., York, R. G., Hoberman, A. M., Diener, R. M., Fisher, L. C. & Gates, G. A. 2001b Biodisposition of dibromoacetic acid (DBA) and bromodichloromethane (BDCM) administered to rats and rabbits in drinking water during range-finding reproduction and developmental toxicity studies. *Int. J. Toxicol.* **20** (4), 239–253.
- Colman, J., Rice, G. E., Wright, J. M., Hunter, E. S., Teuschler, L. K., Lipscomb, J. C., Hertzberg, R. C., Simmons, J. E., Fransen, M., Osier, M. & Narotsky, M. G. 2011 Identification of developmentally toxic drinking water disinfection byproducts and evaluation of data relevant to mode of action. *Toxicol. Appl. Pharmacol.* **254** (2), 100–126.
- Danielsson, B. R. G., Ghantous, H. & Dencker, L. 1986 Distribution of chloroform and methyl chloroform and their metabolites in pregnant mice. *Biol. Res. Pregnancy Perinatol.* **7** (2), 77–83.
- Dowty, B. J., Laseter, J. L. & Storer, J. 1976 The transplacental migration and accumulation in blood of volatile organic constituents. *Pediatr. Res.* **10** (7), 696–701.
- Epstein, D. L., Nolen, G. A., Randall, J. L., Christ, S. A., Read, E. J., Stober, J. A. & Smith, M. K. 1992 Cardiopathic effects of dichloroacetate in the fetal long-evans rat. *Teratology* **46** (3), 225–235.
- EU Directive 98/83/EC 1998 The Drinking Water Directive (DWD). <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:1998:330:0032:0054:EN:PDF>.

- Goldstone, J. V., McArthur, A. G., Kubota, A., Zanette, J., Parente, T., Jönsson, M. E., Nelson, D. R. & Stegeman, J. J. 2010 Identification and developmental expression of the full complement of Cytochrome P450 genes in Zebrafish. *BMC Genomics* **11** (1).
- Graves, C. G., Matanoski, G. M. & Tardiff, R. G. 2001 Weight of evidence for an association between adverse reproductive and developmental effects and exposure to disinfection by-products: A critical review. *Regul. Toxicol. Pharmacol.* **34** (2), 103–124.
- Gustafson, A. L., Stedman, D. B., Ball, J., Hillegass, J. M., Flood, A., Zhang, C. X., Panzica-Kelly, J., Cao, J., Coburn, A., Enright, B. P., Tornesi, M. B., Hetheridge, M. & Augustine-Rauch, K. A. 2012 Inter-laboratory assessment of a harmonized zebrafish developmental toxicology assay – Progress report on phase I. *Reprod. Toxicol.* **33** (2), 155–164.
- Hassoun, E., Kariya, C. & Williams, F. E. 2005 Dichloroacetate-induced developmental toxicity and production of reactive oxygen species in zebrafish embryos. *J. Biochem. Mol. Toxicol.* **19** (1), 52–58.
- Hill, A., Mesens, N., Steemans, M., Xu, J. J. & Aleo, M. D. 2012 Comparisons between in vitro whole cell imaging and in vivo zebrafish-based approaches for identifying potential human hepatotoxicants earlier in pharmaceutical development. *Drug Metab. Rev.* **44** (1), 127–140.
- Hill, A. J., Teraoka, H., Heideman, W. & Peterson, R. E. 2005 Zebrafish as a model vertebrate for investigating chemical toxicity. *Toxicol. Sci.* **86** (1), 6–19.
- Hoekstra, J. A. 1987 Acute bioassays with control mortality. *Water Air Soil Pollut.* **35** (3–4), 311–317.
- Hunter III, E. S., Rogers, E. H., Schmid, J. E. & Richard, A. 1996 Comparative effects of haloacetic acids in whole embryo culture. *Teratology* **54** (2), 57–64.
- Johnson, P. D., Dawson, B. V. & Goldberg, S. J. 1998 Cardiac teratogenicity of trichloroethylene metabolites. *J. Am. Coll. Cardiol.* **32** (2), 540–545.
- Jones, H. S., Panter, G. H., Hutchinson, T. H. & Chipman, J. K. 2010 Oxidative and conjugative xenobiotic metabolism in zebrafish larvae in vivo. *Zebrafish* **7** (1), 23–30.
- Kimmel, C. B., Ballard, W. W., Kimmel, S. R. & Schilling, T. F. 1995 Stages of embryonic development of the zebrafish. *Dev. Dyn.* **203** (3), 253–310.
- Kosmehl, T., Hallare, A. V., Reifferscheid, G., Manz, W., Braunbeck, T. & Hollert, H. 2006 A novel contact assay for testing genotoxicity of chemicals and whole sediments in zebrafish embryos. *Environ. Toxicol. Chem.* **25** (8), 2097–2106.
- Landi, S., Naccarati, A., Ross, M. K., Hanley, N. M., Dailey, L., Devlin, R. B., Vasquez, M., Pegram, R. A. & DeMarini, D. M. 2003 Induction of DNA strand breaks by trihalomethanes in primary human lung epithelial cells. *Mutat. Res.* **538**, 41–50.
- Lim, G. E., Stals, S. I., Petrik, J. J., Foster, W. G. & Holloway, A. C. 2004 The effects of in utero and lactational exposure to chloroform on postnatal growth and glucose tolerance in male Wistar rats. *Endocrine* **25** (3), 223–228.
- Mattice, J. S., Tsai, S. C., Burch, M. B. & Beauchamp, J. J. 1981 Toxicity of trihalomethanes to common carp embryos. *Trans. Am. Fish. Soc.* **110** (2), 261–269.
- Murray, F. J., Schwetz, B. A., McBride, J. G. & Staples, R. E. 1979 Toxicity of inhaled chloroform in pregnant mice and their offspring. *Toxicol. Appl. Pharmacol.* **50** (3), 515–522.
- Nagel, R. 2002 DarT: The embryo test with the Zebrafish *Danio rerio* – a general model in ecotoxicology and toxicology. *ALTEX* **19**, 38–48.
- Narotsky, M. B., Hamby, B. T., Best, D. S. & Hunter, E. S. 1996 In vivo developmental effects of dibromoacetic acid (DBA) and dichloroacetic acid (DCA) in mice. *Teratology* **53** (2), 96–97.
- Narotsky, M. G., Pressman, J. G., Miltner, R. J., Speth, T. F., Teuschler, L. K., Rice, G. E., Richardson, S. D., Best, D. S., McDonald, A., Hunter, E. S. & Simmons, J. E. 2012 Developmental toxicity evaluations of whole mixtures of disinfection by-products using concentrated drinking water in rats: Gestational and lactational effects of sulfate and sodium. *Birth Defects Res. B Dev. Reprod. Toxicol.* **95** (3), 202–212.
- Nieuwenhuijsen, M. J., Toledano, M. B. & Elliott, P. 2000 Uptake of chlorination disinfection by-products; a review and a discussion of its implications for exposure assessment in epidemiological studies. *J. Expo. Anal. Environ. Epidemiol.* **10** (6 Pt 1), 586–599.
- NTP (National Toxicology Program) 1998a Short Term Reproductive and developmental Toxicity Study of Bromochloroacetic Acid (CAS No. 5589–96–8) when Administered to Sprague–Dawley Rats in the Drinking Water. National Toxicology Program, National Institute of Environmental Health Sciences, Research Triangle Park, NC. NTPRDGT96001. PB98172414.
- NTP (National Toxicology Program) 1998b Final Report on the Short Term Reproductive and Developmental Toxicity of Tribromoacetic Acid (CAS No. 75–96–7) Administered in Drinking Water to Sprague–Dawley Rats. Research Triangle Park, NC: National Toxicology Program, National Institute of Environmental Health Sciences. NTPRDGT94009. PB98165111.
- Nuckols, J. R., Ashley, D. L., Lyu, C., Gordon, S. M., Hinckley, A. F. & Singer, P. 2005 Influence of tap water quality and household water use activities on indoor air and internal dose levels of trihalomethanes. *Environ. Health Perspect.* **113** (7), 863–870.
- Padilla, S., Corum, D., Padnos, B., Hunter, D. L., Beam, A., Houck, K. A., Sipes, N., Kleinstreuer, N., Knudsen, T., Dix, D. J. & Reif, D. M. 2012 Zebrafish developmental screening of the ToxCast™ Phase I chemical library. *Reprod. Toxicol.* **33** (2), 174–187.
- Pegram, R. A., Andersen, M. E., Warren, S. H., Ross, T. M. & Claxton, L. D. 1997 Glutathione S-Transferase-mediated mutagenicity of trihalomethanes in *Salmonella typhimurium*: Contrasting results with bromodichloromethane and chloroform. *Toxicol. Appl. Pharmacol.* **144** (1), 183–188.
- Plewa, M. J., Simmons, J. E., Richardson, S. D. & Wagner, E. D. 2010 Mammalian cell cytotoxicity and genotoxicity of the haloacetic acids, a major class of drinking water disinfection by-products. *Environ. Mol. Mutagen.* **51** (8–9), 871–878.



- Postlethwait, J. H., Woods, I. G., Ngo-Hazelett, P., Yan, Y. L., Kelly, P. D., Chu, F., Huang, H., Hill-Force, A. & Talbot, W. S. 2000 Zebrafish comparative genomics and the origins of vertebrate chromosomes. *Genome Res.* **10** (12), 1890–1902.
- Priestley, C. C., Green, R. M., Fellows, M. D., Doherty, A. T., Hodges, N. J. & O'Donovan, M. R. 2010 Anomalous genotoxic responses induced in mouse lymphoma L5178Y cells by potassium bromate. *Toxicology* **267** (1–3), 45–53.
- Räbergh, C. M. I. & Lipsky, M. M. 1997 Toxicity of chloroform and carbon tetrachloride in primary cultures of rainbow trout hepatocytes. *Aquat. Toxicol.* **37** (2–3), 169–182.
- Richardson, S. D., Plewa, M. J., Wagner, E. D., Schoeny, R. & DeMarini, D. M. 2007 Occurrence, genotoxicity, and carcinogenicity of regulated and emerging disinfection by-products in drinking water: A review and roadmap for research. *Mutat. Res.* **636** (1–3), 178–242.
- Ruddick, J. A., Villeneuve, D. C., Chu, I. & Valli, V. E. 1983 A teratological assessment of four trihalomethanes in the rat. *J. Environ. Sci. Health B Pesticides, Food Contaminants Agric. Wastes* **18** (3), 333–349.
- Scholz, S., Fischer, S., Gündel, U., Küster, E., Luckenbach, T. & Voelker, D. 2008 The zebrafish embryo model in environmental risk assessment – Applications beyond acute toxicity testing. *Environ. Sci. Pollut. Res. Int.* **15** (5), 394–404.
- Schwetz, B. A., Leong, B. K. J. & Gehring, P. J. 1974 Embryo- and fetotoxicity of inhaled chloroform in rats. *Toxicol. Appl. Pharmacol.* **28**, 442–451.
- Selderslaghs, I. W. T., Blust, R. & Witters, H. E. 2012 Feasibility study of the zebrafish assay as an alternative method to screen for developmental toxicity and embryotoxicity using a training set of 27 compounds. *Reprod. Toxicol.* **33** (2), 142–154.
- Simmons, J. E., Richardson, S. D., Teuschler, L. K., Miltner, R. J., Speth, T. F., Schenck, K. M., Hunter III, E. S. & Rice, G. 2008 Research issues underlying the four-lab study: Integrated disinfection by-products mixtures research. *J. Toxicol. Environ. Health A* **71**, 1125–1132.
- Singh, N. P., McCoy, M. T., Tice, R. R. & Schneider, E. L. 1988 A simple technique for quantitation of low levels of DNA damage in individual cells. *Exp. Cell Res.* **175** (1), 184–191.
- Smith, M. K., Randall, J. L., Read, E. J. & Stober, J. A. 1989 Teratogenic activity of trichloroacetic acid in the rat. *Teratology* **40**, 445–451.
- Tardiff, R. G., Carson, M. L. & Ginevan, M. E. 2006 Updated weight of evidence for an association between adverse reproductive and developmental effects and exposure to disinfection by-products. *Regul. Toxicol. Pharmacol.* **45** (2), 185–205.
- Teixidó, E., Piqué, E., Gómez-Catalán, J. & Llobet, J. M. 2013 Assessment of developmental delay in the zebrafish embryo teratogenicity assay. *Toxicol. in Vitro* **27** (1), 469–478.
- Thompson, D. J., Warner, S. D. & Robinson, V. B. 1974 Teratology studies on orally administered chloroform in the rat and rabbit. *Toxicol. Appl. Pharmacol.* **29** (3), 348–357.
- US EPA (Environmental Protection Agency) 2006 National primary drinking water regulations: stage 2 disinfectants and disinfection by products rule. *Fed. Reg.* **71**, 387–493.
- Vega-López, A., Carrillo-Morales, C. I., Olivares-Rubio, H. F., Lilia Domínguez-López, M. & García-Latorre, E. A. 2012 Evidence of bioactivation of halomethanes and its relation to oxidative stress response in *Chirostoma riojai*, an endangered fish from a polluted lake in Mexico. *Arch. Environ. Contam. Toxicol.* **62** (3), 479–493.
- Von Westernhagen, H. 1988 Sublethal effects of pollutants on fish eggs and larvae. In: *Fish Physiology* (W.S. Hoar & D.J. Randall, eds). Academic Press, London, 11, pp. 253–346.
- Weber, N. M., Higuchi, T. T., Tessari, J. D. & Veeramachaneni, D. N. R. 2004 Evaluation of the effects of water disinfection by-products, bromochloroacetic and dibromoacetic acids, on frog embryogenesis. *J. Toxicol. Environ. Health A* **67** (12), 929–939.
- Weigt, S., Huebler, N., Strecker, R., Braunbeck, T. & Broschard, T. H. 2011 Zebrafish (*Danio rerio*) embryos as a model for testing proteratogens. *Toxicology* **281** (1–3), 25–36.
- WHO (World Health Organization) 2004 *Guidelines for Drinking Water Quality* (3rd edn) [http://www.who.int/water\\_sanitation\\_health/dwq/gdwq3rev/en](http://www.who.int/water_sanitation_health/dwq/gdwq3rev/en).
- Wolf, G. W. & Kaiser, L. 1996 *Final Report Sodium Bromate: Short Term Reproductive and Development Toxicity Study when Administered to Sprague Dawley Rats in the Drinking Water*. Submitted to National Toxicology Program, National Institute of Environmental Health Science, National Institutes of Health, Research Triangle Park, NC (NTP/NIEHS No. NOI-ES-15323).
- Zhang, C., Willett, C. & Fremgen, T. 2003 Unit 1.7 Zebrafish: An animal model for toxicological studies. *Curr. Protoc. Toxicol.* **1.7.1–1.7.18**.
- Zhang, L., Xu, L., Zeng, Q., Zhang, S. H., Xie, H., Liu, A. L. & Lu, W. Q. 2012 Comparison of DNA damage in human-derived hepatoma line (HepG2) exposed to the fifteen drinking water disinfection byproducts using the single cell gel electrophoresis assay. *Mutat. Res.* **741**, 89–94.

First received 23 October 2013; accepted in revised form 4 May 2014. Available online 29 May 2014