

Quantification and molecular characterisation of human sapoviruses in water sources impacted by highly polluted discharged wastewater in South Africa

Tanya Y. Murray and Maureen B. Taylor

ABSTRACT

Sapoviruses (SaVs) were detected and quantified in 8/10 water samples collected from wastewater treatment works (WWTWs) and water sources impacted by these WWTWs in Limpopo Province, South Africa. The median SaV concentration was 2.45×10^6 copies/L and SaV genotypes I.2 and IV were characterised. This study provides new data on the high concentrations of clinically relevant SaVs in rivers and dams impacted by poor-performing WWTWs.

Key words | genotypes, quantification, sapovirus, South Africa, water

Tanya Y. Murray (corresponding author)
Maureen B. Taylor
Department of Medical Virology,
University of Pretoria,
Pretoria,
South Africa
E-mail: tanyaymurray@gmail.com

INTRODUCTION

Sapovirus (SaV), a genus in the Caliciviridae (CV) family, is increasingly being recognised as a cause of gastroenteritis outbreaks worldwide, infecting both children and adults (Miyoshi *et al.* 2010; Svraka *et al.* 2010; Lee *et al.* 2012). The genus is divided into at least five genogroups (GI–GV) based on the capsid gene sequence (Farkas *et al.* 2004). Sapoviruses that infect humans belong to GI, GII, GIV and GV (Farkas *et al.* 2004) and GI and GII are each further divided into at least seven genotypes (Oka *et al.* 2012). Sapovirus has been implicated in a waterborne outbreak of gastroenteritis in France, along with other human CVs (Bon *et al.* 2005) and has also been associated with foodborne outbreaks, where bivalve shellfish (Iizuka *et al.* 2010) and food handlers (Usuku *et al.* 2008; Kobayashi *et al.* 2012) were identified as suspected sources of infection. Few countries have reported the presence of SaVs in water sources. In Japan and Europe (France, Italy and Spain), SaVs have been documented in influent and effluent wastewater (Hansman *et al.* 2007; Sano *et al.* 2011; Sima *et al.* 2011; Di Bartolo *et al.* 2013). In South Africa (SA), SaVs have also been detected in discharged wastewater in several provinces of the country (Murray *et al.* 2013a).

Malfunctioning wastewater treatment works (WWTWs) contribute significantly to the deteriorating water quality in SA (Mitchell *et al.* 2014). Consequently, in 2008 the

Department of Water Affairs (DWA) introduced the Green Drop system which aims to improve WWTW performance by providing an incentive in the form of a scoring system. This system rates aspects of the WWTW's performance and those which perform properly are awarded Green Drop status (Mitchell *et al.* 2014). One of the poorest-performing provinces is Limpopo (LP), which had the second lowest rating of 24% in the last Green Drop report (Department of Water Affairs 2011). This score indicates that the province's wastewater treatment systems are in a critical state and need urgent attention. Green Drop certification criteria include a microbiological compliance score which is determined from faecal coliform counts of <1,000 colony forming units (cfu) per 100 mL and is reported as the percentage of samples that have <1,000 cfu/100 mL over the total number of samples tested. Over a 1-year period from January to December 2012, which includes the duration of the study period, LP's average microbiological compliance was 5.75% (Department of Water and Sanitation 2012). This minimal functionality of WWTWs in the province is of particular concern for fresh produce farmers who use water sources downstream from the WWTWs to irrigate their crops. In this study, concentrations and genotypes of SaVs from water sources in a selected region of LP, including rivers

and dams impacted by poor-performing WWTWs, are reported.

METHODS

In January and March 2012, 10 water samples (10 L) were collected from several locations in close proximity to two WWTWs in LP, SA. Samples were collected directly downstream from the two WWTWs (LP5 and LP23) as well as from two rivers (LP1 and LP7) into which the discharged wastewater flows. Samples were also collected from dams located nearby (LP9, LP16B and LP17B) and from the irrigation pivot (LP PVT) on a fresh produce farm in the close vicinity. Water samples were transported to the laboratory in cooler bags with cold packs and the temperature and pH were recorded upon arrival. Samples were stored at 4 °C for no more than 24 hours, until processing.

Prior to virus recovery, mengovirus (5×10^5 genome copies) was added to each water sample as a process control (Saïd *et al.* 2014). Viruses were recovered from the water samples in a volume of 100 mL, using a glass wool adsorption-elution method which was developed by Vilaginès *et al.* (1993) and subsequently modified (Mans *et al.* 2013). Viruses were further concentrated to a final volume of 10 mL in phosphate-buffered saline (pH 7.4) (Sigma-Aldrich Co., St Louis, MO, USA) by polyethylene glycol 8000 (Amresco, Solon, OH, USA) and sodium chloride (Merck Schuchardt OHC, Hohenbrunn, Germany) (PEG8000/NaCl) precipitation as described in the ISO/TS 15216-1:2013 (2013). Nucleic acid was extracted from 1 mL virus concentrate using the semi-automated Nucli-SENS® EasyMAG® instrument (BioMérieux, Marcy l'Etoile, France) as previously described (Saïd *et al.* 2014). Mengovirus (5 µL extracted nucleic acid) was detected qualitatively by real-time reverse transcription-polymerase chain reaction (RT-PCR) using a commercial kit (mengo@ceeramTools™ Kit, Ceeram s.a.s, La Chappelle-Sur-Erdre, France) (Saïd *et al.* 2014). Norovirus GI and GII (5 µL extracted nucleic acid) were also detected qualitatively using commercial kits with internal amplification controls (IACs) (norovirusGI@ceeramTools™ Kit and norovirusGII@ceeramTools™ Kit). Sapovirus (2.5 µL extracted nucleic acid) was quantified in each sample using a real-time RT-PCR which targets 104 bp

of the conserved RNA polymerase and capsid gene junction in ORF1. The assay included a SaV RNA standard, which contains only the 104 bp target region of the SaV genome, as the positive control in the assay and quantification standard. The assay also incorporates a competitive IAC (201 bp), constructed from non-target *Theileria parva* DNA flanked with the SaV-specific primers, in each sample to monitor inhibition (Murray *et al.* 2013a). All samples were tested once and if PCR inhibition was observed, the assay was repeated at a 10-fold dilution of the extracted nucleic acid. Detected SaVs were further characterised by genotyping a 5' region (approximately 300 bp) of the capsid gene using a nested PCR as previously described (Murray *et al.* 2013b). Sapovirus sequences were submitted to GenBank (accession numbers: KC904508–KC904513).

Standard quality control precautions were applied to reduce the possibility of cross-contamination and false-positive results. Separate laboratories were used for sample processing and preparation, nucleic acid extraction, reaction mix preparation and manipulation of amplicons. An extraction negative control and RT-PCR negative control were included in the real-time assay and PCR negative controls were included in both rounds of the genotyping nested PCR to monitor for false-positive results (Bosch *et al.* 2011).

RESULTS AND DISCUSSION

Sapoviruses were detected in 80% (8/10) of the water samples (Table 1). Mengovirus was also detected in each sample (10/10), including samples from Dam 1 and Dam 2 that were SaV-negative. None of the samples required retesting with diluted nucleic acid, indicating successful virus recovery and nucleic acid extraction with the adequate removal of inhibitors. Seven of the eight SaV-positive samples were also positive for hepatitis A virus (HAV) and norovirus (NoV) (Table 1; Saïd *et al.* 2014), indicating the presence of a variety of enteric viruses in the water sources. Sapovirus concentrations ranged from 1.11×10^5 copies/L to 1.62×10^7 copies/L (median 2.45×10^6 copies/L). The highest concentrations were detected directly below the WWTWs (1.59×10^7 copies/L and 1.62×10^7 copies/L) and the lowest concentrations from Dam 3 (1.11×10^5 copies/L) and the irrigation pivot (8.32×10^5 copies/L) on the farm. The SaV

Table 1 | Water samples analysed for sapoviruses

Sample ID	Collection date	Sample description	SaV concentration (copies/L)	SaV genotype	HAV ^a	NoV
LP1	31 January 2012	River A	5.88×10^6	GI.2	+	+
LP5	31 January 2012	WWTW (discharged wastewater)	1.59×10^7	GI.2	+	+
LP7	31 January 2012	River B	1.70×10^6	GI.2	+	+
LP9	31 January 2012	Dam 1 (borehole water)	–	–	–	–
LP16B	31 January 2012	Dam 2 (surface water)	–	–	–	–
LP17B	31 January 2012	Dam 3 (surface water)	1.11×10^5	ND	–	–
LP23	31 January 2012	WWTW (discharged wastewater)	1.62×10^7	GI.2	+	+
LP7	27 March 2012	River B	3.21×10^6	GI.2	+	+
LP17B	27 March 2012	Dam 3 (surface water)	1.42×10^6	ND	+	+
LP PVT	27 March 2012	Irrigation pivot	8.32×10^5	GIV	+	+

ND = genotype was not determined, WWTW = wastewater treatment works, LP5 and LP23 are two different WWTWs in the same municipality.

^aData taken from Said *et al.* (2014).

concentrations from the WWTWs correlate to quantities previously reported in discharged wastewater in other provinces of SA (Murray *et al.* 2013a). The high concentrations also correlate to those reported in raw wastewater in Spain (Sano *et al.* 2011), suggesting the WWTWs are not successfully removing the virus prior to discharge. This reconfirms the poor performance of selected WWTWs in this area of SA, which impact water sources downstream from the WWTWs, as indicated by the low Green Drop score for the province.

Sapovirus strains were successfully genotyped from six of eight SaV-positive samples. Five of these strains clustered with GI.2 and one strain grouped with GIV. Genotype I.2 and GIV are two of the most prevalent genotypes implicated in SaV-associated outbreaks of gastroenteritis (Svraka *et al.* 2010; Kobayashi *et al.* 2012; Lee *et al.* 2012). Genotype I.2 is also frequently reported in water sources including in SA (Kitajima *et al.* 2011; Sano *et al.* 2011; Di Bartolo *et al.* 2013; Murray *et al.* 2013b). The GI.2 strains shared high nucleotide identity ($\geq 99\%$) to strains previously identified in water sources from other provinces of SA. In addition, the GI.2 strains were closely related (98–99% nucleotide identity) to strains found in water sources from Japan (Kitajima *et al.* 2011), Spain (Sano *et al.* 2011) and Italy (Di Bartolo *et al.* 2013). This is the first report of SaV GIV in the environment in SA. It was identified in a sample taken directly from an irrigation pivot on the farm, indicating the potential for direct contamination of fresh produce during irrigation. Sapovirus GIV has been reported in untreated sewage in Japan (Iwai

et al. 2009) and Spain (Sano *et al.* 2011). The SA GIV strain shared 98% nucleotide identity over the typed region with strains identified from patients with gastroenteritis in Venezuela (HM214146) and the United States (HM800905). The strain also shared 97% nucleotide identity with a SaV genotyped from clams in Japan (Iizuka *et al.* 2013).

CONCLUSION

This study has provided new and informative data on the high concentration of SaVs present in water sources impacted by poor-performing WWTWs in a selected region of SA. In addition, it has shown that clinically relevant SaV GI.2 and GIV strains are circulating in these water sources. Enteric viruses have a low infectious dose (10–100 infectious virions; Carter 2005; Teunis *et al.* 2008) and consequently, high titres of pathogens such as SaV in poorly treated wastewater discharge may have a negative impact on the health of populations using the water sources for domestic, recreational or agricultural purposes.

ACKNOWLEDGEMENTS

The authors thank Dr Erika Du Plessis, from the Department of Microbiology and Plant Pathology, University of Pretoria, for the collection of the water samples. The authors also

thank Gabriël de Ridder and Rachida Saïd for assisting with norovirus and mengovirus analysis. This study was funded, in part, by the Poliomyelitis Research Foundation, and in part by an on-going solicited Water Research Commission research project co-funded by Department of Agriculture, Forestry and Fisheries, SA. 'An investigation into the link between water quality and microbiological safety of fruit and vegetables from the farming to the processing stages of production and marketing' (Project No K5/1875//4, Water Research Commission Abridged Knowledge review 2009/10, Pretoria).

REFERENCES

- Bon, F., Ambert-Balay, K., Giraudon, H., Kaplon, J., Le Guyader, S., Pommepuy, M., Gallay, A., Vaillant, V., de Valk, H., Chikhi-Brachet, R., Flahaut, A., Pothier, P. & Kohli, E. 2005 Molecular epidemiology of caliciviruses detected in sporadic and outbreak cases of gastroenteritis in France from December 1998 to February 2004. *J. Clin. Microbiol.* **43** (9), 4659–4664.
- Bosch, A., Sánchez, G., Abbaszadegan, M., Carducci, A., Guix, S., Le Guyader, F. S., Netshikweta, R., Pintó, R. M., van der Poel, W. H. M., Rutjes, S., Sano, D., Taylor, M. B., van Zyl, W. B., Rodríguez-Lázaro, D., Kovač, K. & Sellwood, J. 2011 Analytical methods for virus detection in water and food. *Food Anal. Methods* **4** (1), 4–12.
- Carter, M. J. 2005 Enterically infecting viruses: pathogenicity, transmission and significance for food and waterborne infection. *J. Appl. Microbiol.* **98** (6), 1354–1380.
- Department of Water Affairs 2011 Green Drop Report, DWA. http://www.dwaf.gov.za/dir_ws/GDS/Docs/DocsDefault.aspx (accessed 16 September 2014).
- Di Bartolo, I., Ponterio, E., Battistone, A., Bonomo, P., Cicala, A., Mercurio, P., Triassi, M., Pennino, F., Fiore, L. & Ruggeri, F. M. 2013 Identification and genotyping of human sapoviruses collected from sewage water in Naples and Palermo, Italy, in 2011. *Food Environ. Virol.* **5** (4), 236–240.
- Department of Water and Sanitation. 2012 Green Drop System version 2.1. http://www.dwaf.gov.za/dir_ws/GDS/ (accessed 26 August 2014).
- Farkas, T., Zhong, W. M., Jing, Y., Huang, P. W., Espinosa, S. M., Martinez, N., Morrow, A. L., Ruiz-Palacios, G. M., Pickering, L. K. & Jiang, X. 2004 Genetic diversity among sapoviruses. *Arch. Virol.* **149** (7), 1309–1323.
- Hansman, G. S., Sano, D., Ueki, Y., Imai, T., Oka, T., Katayama, K., Takeda, N. & Omura, T. 2007 Sapovirus in water, Japan. *Emerg. Infect. Dis.* **13** (1), 133–135.
- Iizuka, S., Oka, T., Tabara, K., Omura, T., Katayama, K., Takeda, N. & Noda, M. 2010 Detection of sapoviruses and noroviruses in an outbreak of gastroenteritis linked genetically to shellfish. *J. Med. Virol.* **82** (7), 1247–1254.
- Iizuka, S., Takai-Todaka, R., Ohshiro, H., Kitajima, M., Wang, Q., Saif, L. J., Wakita, T., Noda, M., Katayama, K. & Oka, T. 2013 Detection of multiple human sapoviruses from imported frozen individual clams. *Food Environ. Virol.* **5** (2), 119–125.
- ISO/TS 15216-1:2013 2013 Microbiology of food and animal feed – horizontal method for determination of hepatitis A virus and norovirus in food using real-time RT-PCR – part 1: method for quantification.
- Iwai, M., Hasegawa, S., Obara, M., Nakamura, K., Horimoto, E., Takizawa, T., Kurata, T., Sogen, S. & Shiraki, K. 2009 Continuous presence of noroviruses and sapoviruses in raw sewage reflects infections among inhabitants of Toyama, Japan (2006 to 2008). *Appl. Environ. Microbiol.* **75** (5), 1264–1270.
- Kitajima, M., Haramoto, E., Phanuwat, C. & Katayama, H. 2011 Genotype distribution of human sapoviruses in wastewater in Japan. *Appl. Environ. Microbiol.* **77** (12), 4226–4229.
- Kobayashi, S., Fujiwara, N., Yasui, Y., Yamashita, T., Hiramatsu, R. & Minagawa, H. 2012 A foodborne outbreak of sapovirus linked to catered box lunches in Japan. *Arch. Virol.* **157** (10), 1995–1997.
- Lee, L. E., Cebelin, E. A., Fuller, C., Keene, W. E., Smith, K., Vinje, J. & Besser, J. M. 2012 Sapovirus outbreaks in long-term care facilities, Oregon and Minnesota, USA, 2002–2009. *Emerg. Infect. Dis.* **18** (5), 873–876.
- Mans, J., Netshikweta, R., Magwalivha, M., van Zyl, W. B. & Taylor, M. B. 2013 Diverse norovirus genotypes identified in sewage-polluted river water in South Africa. *Epidemiol. Infect.* **141**, 303–313.
- Mitchell, S. A., de Wit, M. P., Blignaut, J. N. & Crookes, D. 2014 Wastewater treatment plants: The financing mechanisms associated with achieving Green Drop rating. WRC Report no. 2085/1/14., Water Research Commission, Pretoria.
- Miyoshi, M., Yoshizumi, S., Kanda, N., Karino, T., Nagano, H., Kudo, S., Okano, M. & Ishida, S. 2010 Different genotypic sapoviruses detected in two simultaneous outbreaks of gastroenteritis among schoolchildren in the same school district in Hokkaido, Japan. *Jpn J. Infect. Dis.* **63** (1), 75–78.
- Murray, T. Y., Mans, J., van Zyl, W. B. & Taylor, M. B. 2013a Application of a competitive internal amplification control for the detection of sapoviruses in wastewater. *Food Environ. Virol.* **5** (1), 61–68.
- Murray, T. Y., Mans, J. & Taylor, M. B. 2013b Human calicivirus diversity in wastewater in South Africa. *J. Appl. Microbiol.* **114** (6), 1843–1853.
- Oka, T., Mori, K., Iritani, N., Harada, S., Ueki, Y., Iizuka, S., Mise, K., Murakami, K., Wakita, T. & Katayama, K. 2012 Human sapovirus classification based on complete capsid nucleotide sequences. *Arch. Virol.* **157** (2), 349–352.
- Saïd, R., Wolfaardt, M. & Taylor, M. B. 2014 Molecular characterisation of hepatitis A virus strains from water sources in South Africa. *Water Sci. Technol.* **69** (5), 923–933.
- Sano, D., Pérez-Sautu, U., Guix, S., Pintó, R. M., Miura, T., Okabe, S. & Bosch, A. 2011 Quantification and genotyping of human

- sapoviruses in the Llobregat river catchment, Spain. *Appl. Environ. Microbiol.* **77** (5), 1111–1114.
- Sima, L. C., Schaeffer, J., Le Saux, J. C., Parnaudeau, S., Elimelech, M. & Le Guyader, F. S. 2011 [Calicivirus removal in a membrane bioreactor wastewater treatment plant](#). *Appl. Environ. Microbiol.* **77** (15), 5170–5177.
- Svraka, S., Vennema, H., van der Veer, B., Hedlund, K. O., Thorhagen, M., Siebenga, J., Duizer, E. & Koopmans, M. 2010 [Epidemiology and genotype analysis of emerging sapovirus-associated infections across Europe](#). *J. Clin. Microbiol.* **48** (6), 2191–2198.
- Teunis, P. F., Moe, C. L., Liu, P., Miller, S. E., Lindesmith, L., Baric, R. S., Le Pendu, J. & Calderon, R. L. 2008 [Norwalk virus: how infectious is it?](#) *J. Med. Virol.* **80** (8), 1468–1476.
- Usuku, S., Kumazaki, M., Kitamura, K., Tochikubo, O. & Noguchi, Y. 2008 An outbreak of food-borne gastroenteritis due to sapovirus among junior high school students. *Jpn J. Infect. Dis.* **61** (6), 438–441.
- Vilaginès, Ph., Sarrette, B., Husson, G. & Vilaginès, R. 1993 Glass wool for virus concentration from water at ambient pH level. *Water Sci. Technol.* **27** (3–4), 299–306.

First received 25 September 2014; accepted in revised form 15 March 2015. Available online 10 April 2015