

## ***Staphylococcus aureus* isolated from wastewater treatment plants in Tunisia: occurrence of human and animal associated lineages**

Meriam Ben Said, Mohamed Salah Abbassi, Paula Gómez, Laura Ruiz-Ripa, Senda Sghaier, Chourouk Ibrahim, Carmen Torres and Abdennaceur Hassen

### **ABSTRACT**

The objective was to characterize *Staphylococcus aureus* isolated from two wastewater treatment plants (WWTPs) located in Tunis City (Tunisia), during the period 2014–2015. Genetic lineages, antibiotic resistance mechanisms and virulence factors were determined for the recovered isolates. *S. aureus* isolates were recovered from 12 of the 62 wastewater samples tested (19.35%), and one isolate/sample was characterized, all of them being methicillin-susceptible (MSSA). Six *spa* types (t587, t674, t224, t127, t701 and t1534) were found among the 12 isolates, and the *spa*-t587, associated with the new sequence type ST3245, was the most predominant one (7 isolates). The remaining isolates were assigned to five clonal complexes (CC5, CC97, CC1, CC6 and CC522) according to the sequence-type determined and/or the *spa*-type detected. *S. aureus* isolates were ascribed to *agrI* ( $n = 3$ ), *agrII* ( $n = 7$ ) and *agrIII* ( $n = 1$ ); however, one isolate was non-typeable. *S. aureus* showed resistance to (number of isolates): penicillin (12), erythromycin (7), tetracycline (one) and clindamycin (one). Among the virulence factors investigated, only one isolate harboured the *tst* gene, encoding the TSST-1 (toxic shock syndrome toxin 1). Despite the low number of studied isolates, the present study reports the occurrence of both human- and animal-associated *S. aureus* clonal complexes in WWTPs in Tunisia.

**Key words** | animal and human lineages, *S. aureus*, wastewater

**Meriam Ben Said**  
**Mohamed Salah Abbassi**  
**Senda Sghaier**  
Université de Tunis El Manar, Institut de la  
Recherche Vétérinaire de Tunisie,  
20 rue Jebel Lakhdar, Bab Saadoun,  
Tunis 1006,  
Tunisia

**Mohamed Salah Abbassi**  
Université de Tunis El Manar,  
Faculté de Médecine de Tunis,  
Laboratoire de résistance aux antibiotiques  
LR99ES09,  
Tunisie

**Meriam Ben Said**  
**Senda Sghaier**  
**Chourouk Ibrahim**  
**Abdennaceur Hassen**  
Laboratoire de Traitement des Eaux Usées,  
Centre de Recherche et des Technologies des Eaux  
(CERTE),  
Technopoles Borj-Cédria, BP 273,  
8020 Soliman,  
Tunisia

**Paula Gómez**  
**Laura Ruiz-Ripa**  
**Carmen Torres** (corresponding author)  
Área de Bioquímica y Biología Molecular,  
Universidad de La Rioja,  
26006 Logroño,  
Spain  
E-mail: carmen.torres@unirioja.es

### **INTRODUCTION**

The frequent use of antibiotics in hospitals, veterinary and agricultural settings has certainly contributed to the dissemination of resistant bacteria or their resistance genes in the environment. Furthermore, a large part of the antibiotics consumed are not absorbed and can end up in wastewater. It has been suggested that wastewater treatment plants (WWTPs) may play a role in the spread and the development of antibiotic-resistant bacteria (Börjesson *et al.* 2010; Rizzo *et al.* 2013). In addition, treated sewage water is reused in various ways, such as for agriculture, aquaculture and industrial purposes.

During these activities, individuals may come into contact with reused wastewater and can be potentially exposed to resistant and virulent bacteria that may remain in the treated wastewater. *Staphylococcus aureus* is commonly found in the skin and the nails of humans and various animal species; however, it may also act as an opportunistic pathogen that causes minor to severe infections due to the production of different virulence factors. Nowadays, methicillin-resistant *S. aureus* (MRSA) is one of the most important threats to human and animal health worldwide (Cuny *et al.* 2015). The

occurrence of *S. aureus* and MRSA in non-clinical environments such as wastewater has rarely been studied since the majority of wastewater studies are focused on microbial indicators of faecal contamination (George *et al.* 2002; Tran *et al.* 2015). Nonetheless, some studies have reported a low prevalence of *S. aureus* in hospital and municipal wastewater (Schwartz *et al.* 2003; Shannon *et al.* 2007). MRSA has been previously detected from wastewater (Börjesson *et al.* 2010; Kumar *et al.* 2015; Gómez *et al.* 2016).

The main aim of this study was to assess the occurrence of *S. aureus* in two WWTPs located in the suburbs of Tunis City (Tunisia), and to characterize the recovered isolates by determining their antibiotic susceptibility, the genes encoding antibiotic resistance and some virulence factors, and finally their clonal lineages.

## MATERIALS AND METHODS

### Sampling and microbiological identification

Sixty-two water samples were collected during the period of December 2014 to February 2015 from two WWTPs located in the north-east suburbs of Tunis City (Tunisia) (La Charguia: 59 samples and El Menzah: 3 samples). The two WWTPs received urban municipal wastewater from distinct areas in Tunis City. In the case of La Charguia WWTP, the sample collection was done at different points during the wastewater treatment process (at the entrance as influent, after the primary treatment, after the secondary treatment achieved by activated sludge, after the returned sludge, and lastly at the exit of the plant as effluent). In the case of El Menzah WWTP, only influent water was collected. Wastewater samples were recovered in a 500 mL sterile bottle, and transported under refrigeration condition of +4 °C to the laboratory for analysis.

Petri dishes containing Mannitol Salt Agar (MSA, Becton-Dickinson) were seeded by 1 mL of different wastewater samples, and incubated for 18–24 hours. One colony from each positive wastewater sample that showed typical *S. aureus* morphology was picked up for further study. Further identification of *S. aureus* was based mainly on colony morphology, Gram staining, the ability to coagulate rabbit plasma (BioMerieux, France) and the DNase activity. Identification of *S. aureus* was confirmed by a duplex polymerase chain

reaction (PCR) that amplifies both the *nuc* gene and the methicillin-resistance genetic determinant *mecA* (CRL-AR 2009).

### Antimicrobial susceptibility testing

Susceptibility to eight antimicrobial agents was performed using the disc-diffusion method as recommended by the Clinical and Laboratory Standards Institute (CLSI 2015). Antimicrobial agents tested were (charge in µg/disc): penicillin (10 units), cefoxitin (30), tetracycline (30), gentamicin (10), tobramycin (10), ciprofloxacin (5), erythromycin (15) and clindamycin (2).

### Detection of antimicrobial resistance genes

The presence of genes that confer resistance to penicillin (*blaZ*), tetracycline [*tet*(K), *tet*(L) and *tet*(M)], clindamycin [*lnu*(A), *lnu*(B), *vga*(A) and *vga*(C)] and erythromycin [*msr*(A)] was analysed by PCR (Gómez-Sanz *et al.* 2010; Lozano *et al.* 2012). Positive and negative controls from the collection of the University of La Rioja (Logroño, Spain) were used in all PCRs.

### Molecular typing methods of *S. aureus* isolates

All *S. aureus* isolates obtained were characterized by *spa* typing (Harmsen *et al.* 2003), and *agr* typing as previously described (Shopsin *et al.* 2003). Multi-locus sequence typing was performed in selected *S. aureus* isolates (Enright *et al.* 2000). The clonal complex (CC) of the isolates was ascribed according to the sequence-type (ST) determined and/or *spa*-type detected.

### Virulence factors

All isolates were tested by PCR for the presence of *tst* genes encoding the TSST-1 (toxic shock syndrome toxin), *eta* and *etb* genes encoding exfoliative toxins ETA and ETB, respectively (Jarraud *et al.* 2002) and *lukF/lukS-PV* genes encoding Panton–Valentine leucocidin (Lina *et al.* 1999).

## RESULTS AND DISCUSSION

Very few data concerning the presence and the genetic lineages of *S. aureus* in wastewater are available in Tunisia.

Therefore, this study was undertaken to expand the knowledge about *S. aureus* that might circulate among WWTPs. In this study, *S. aureus* was detected in 12 of the 62 wastewater samples collected (19.35%), and one isolate/sample was characterized: 11 isolates from La Chargaia WWTP and one from El Menzah WWTP (Table 1). The percentage of samples carrying *S. aureus* detected in wastewater in this study is different from previous studies conducted in the United States and in Spain, where high rates (50–55%) of *S. aureus* occurrence in wastewater were observed (Rosenberg Goldstein et al. 2012; Gómez et al. 2016). MRSA was not detected in this study, in dissimilarity with some European studies where MRSA has often been reported in wastewater (Börjesson et al. 2010; Porrero et al. 2014). This finding might be related to the low rate or the absence of MRSA from healthy humans or animals in Tunisia (Ben Slama et al. 2011; Gharsa et al. 2015). Regarding antimicrobial resistance, all the isolates showed resistance to penicillin, which was encoded by the *blaZ* gene in seven isolates. Indeed, penicillin resistance is widely disseminated among *S. aureus* isolates worldwide (Lowy 2003). The *msr(A)* gene was detected in the seven erythromycin-resistant isolates (Table 1). The *msr(A)* gene codes for an ABC transporter protein that confers co-resistance to macrolides and streptogramin B antibiotics, and it has been reported

from animal and human *S. aureus* isolates (Schmitz et al. 2000; Wendlandt et al. 2013, 2015). The *lnu(A)*, *lnu(B)*, *vga(A)* and *vga(C)* genes were not detected in the clindamycin-resistant isolate. Only one isolate was resistant to tetracycline and harboured the *tet(K)* gene. In fact, *S. aureus* resistance to tetracycline was more common in MRSA than in the MSSA isolates (Sharma et al. 2013). In general, tetracycline resistance in *S. aureus* is often encoded by the *tet(M)*, *tet(L)* and *tet(K)* genes (Wendlandt et al. 2013, 2015). Only one isolate harboured the *tst* gene, and other virulence genes were not detected. Similarly, low rates of TSST-1 were earlier observed in *S. aureus* of wastewater origin (Gómez et al. 2016), in contrast to animal and human *S. aureus* isolates, where TSST-1 and other virulence factors are more frequent, especially in some specific genetic lineages (Monecke et al. 2009; Gharsa et al. 2012a).

Six *spa* types (t587, t674, t224, t127, t701 and t1534) were detected in this study (Table 1), the *spa*-type t587 was the most predominant one (7 isolates, 58.3%). This *spa*-type was associated with *agr* type II in all but one of the isolates, and with a new ST named ST3245 (singleton) which differs from ST9 in *glpF* and *tpi* alleles. The ST9-CC9 is linked worldwide to livestock (Hasman et al. 2010). As far as we know, the *spa*-type t587 has not been previously described in wastewater; nevertheless, the lineage t587-CC9

**Table 1** | Characteristics of the 12 *S. aureus* isolates recovered in this study

Strains	WWTP samples <sup>a</sup>	<i>spa</i> -type	ST/CC <sup>b</sup>	<i>agr</i> -type	Phenotype of resistance <sup>c</sup>	Resistance genes	Virulence genes
C8300	C-AS	t587	<b>ST3245</b>	NT	PEN, ERY	<i>msr(A)</i>	–
C8302	C-APT	t587	ST3245	<i>agr</i> II	PEN, ERY	<i>blaZ</i> , <i>msr(A)</i>	–
C8303	C-AS	t587	ST3245	<i>agr</i> II	PEN, ERY	<i>blaZ</i> , <i>msr(A)</i>	–
C8304	C-E	t587	ST3245	<i>agr</i> II	PEN, ERY	<i>blaZ</i> , <i>msr(A)</i>	–
C8305	C-I	t587	ST3245	<i>agr</i> II	PEN, ERY	<i>msr(A)</i>	–
C8306	C-APT	t587	ST3245	<i>agr</i> II	PEN, ERY	<i>blaZ</i> , <i>msr(A)</i>	–
C8307	C-RS	t587	ST3245	<i>agr</i> II	PEN, ERY	<i>blaZ</i> , <i>msr(A)</i>	–
C8308	C-I	t674	ST15/CC5	<i>agr</i> II	PEN, TET	<i>blaZ</i> , <i>tet(K)</i>	–
C8309	C-APT	t224	(CC97)	<i>agr</i> I	PEN	–	–
C8310	M-I	t127	(CC1)	<i>agr</i> III	PEN, CLIN	<i>blaZ</i>	–
C8311	C-E	t701	(CC6)	<i>agr</i> I	PEN	–	–
C8313	C-APT	t1534	(CC522)	<i>agr</i> I	PEN	–	<i>tst</i>

<sup>a</sup>WWTP samples: the location of the WWTP (C: La Chargaia; M: El Menzah) is indicated as well as the type of the samples (I, influent; APT, after primary treatment; AS, activated sludge; RS, returned sludge; E, effluent).

<sup>b</sup>New ST is shown in bold. In some cases, the CC was assumed according to the *spa*-type (in this case it is shown in parentheses).

<sup>c</sup>PEN, penicillin; CLI, clindamycin; ERY, erythromycin; TET, tetracycline; NT, non-typeable.

has been detected in bloodstream infections in French patients, and in intensive care unit patients from 14 hospitals in the Netherlands (Rijnders *et al.* 2009; Lamamy *et al.* 2013). Furthermore, two lineages, t224-CC97 and t1534-CC522, have been reported in bovine and small ruminant isolates, respectively (Hata *et al.* 2010; Porrero *et al.* 2012). The lineage CC1, associated with *spa*-type t127 and *agr*III, was detected in one isolate (Table 1); this CC is frequently detected in human isolates (Monecke *et al.* 2009; Lozano *et al.* 2011), although it has also been frequently reported in animals, including healthy donkeys in Tunisia (Gharsa *et al.* 2012b). The lineage t701-CC6, detected in one isolate, was previously reported in Tunisia among both human and animal isolates (Kechrid *et al.* 2010; Gharsa *et al.* 2012b). The lineage t674-ST15-CC5, found in one isolate, was previously detected in patients in Spain, but with different *spa* types (Lozano *et al.* 2015). It is worth noting that the CC5 is a human-associated lineage, frequently associated with MRSA strains (Nübel *et al.* 2008).

## CONCLUSIONS

*S. aureus* was detected in approximately 20% of wastewater samples of two WWTPs tested in Tunisia; we cannot discount that this relatively low frequency of detection could be influenced by the low volume of the sample (1 mL) used in this study. Different CCs associated with either animals or humans have been found. High rates of isolates showed resistance to penicillin and erythromycin, although resistance to tetracycline was also reported. The detection of one *S. aureus* isolate of lineage t1534 carrying the *tst* gene is of relevance and its impact on public health should be evaluated. Future studies, including a larger number of WWTPs of different geographic areas of the country, should be performed to expand the results obtained in this study.

## ACKNOWLEDGEMENTS

Ben Said Meriam has a fellowship from the Tunisian Ministry of Higher Education and Scientific Research. Paula Gómez has a pre-doctoral fellowship from the Universidad de La Rioja (Spain). This work was partially

supported by project SAF2016-76571-R of the Agencia Estatal de Investigación (AEI) of Spain and Fondo Europeo de Desarrollo Regional (FEDER).

## REFERENCES

- Ben Slama, K., Gharsa, H., Klibi, N., Jouini, A., Lozano, C., Gómez-Sanz, E., Zarazaga, M., Boudabous, A. & Torres, C. 2011 Nasal carriage of *Staphylococcus aureus* in healthy humans with different levels of contact with animal in Tunisia: genetic lineage, methicillin resistance, and virulence factors. *Eur. J. Clin. Microbiol. Infect. Dis.* **30**, 499–508.
- Börjesson, S., Matussek, A., Melin, S., Löfgren, S. & Lindgren, P. E. 2010 Methicillin resistant *Staphylococcus aureus* (MRSA) in municipal wastewater: an uncharted threat? *J. Appl. Microbiol.* **108**, 1244–1251.
- CLSI (Clinical Laboratory Standards Institute) 2015 *Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Fifth Information Supplement. M100-S25*. National Committee for Clinical Laboratory Standards, Wayne, PA, USA.
- CRL-AR (Community Reference Laboratory for Antimicrobial Resistance) 2009 *Multiplex PCR for the Detection of the *MecA* Gene and the Identification of *Staphylococcus Aureus**. National Food Institute, Technical University of Denmark, Copenhagen, Denmark.
- Cuny, C., Wieler, L. H. & Witte, W. 2015 Livestock-associated MRSA: the impact on humans. *Antibiotics* **4**, 521–543.
- Enright, M. C., Day, N. P., Davies, C. E., Peacock, S. J. & Spratt, B. G. 2000 Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. *J. Clin. Microbiol.* **38**, 1008–1015.
- George, I., Crop, P. & Servais, P. 2002 Fecal coliform removal in wastewater treatment plants studied by plate counts and enzymatic methods. *Water Res.* **3**, 2607–2617.
- Gharsa, H., Ben Slama, K., Lozano, C., Gómez-Sanz, E., Klibi, N., Ben Sallem, R., Gómez, P., Zarazaga, M., Boudabous, A. & Torres, C. 2012a Prevalence, antibiotic resistance, virulence traits and genetic lineages of *Staphylococcus aureus* in healthy sheep in Tunisia. *Vet. Microbiol.* **156**, 367–373.
- Gharsa, H., Ben Sallem, R., Ben Slama, K., Gómez-Sanz, E., Lozano, C., Jouini, A., Klibi, N., Zarazaga, M., Boudabous, A. & Torres, C. 2012b High diversity of genetic lineages and virulence genes in nasal *Staphylococcus aureus* isolates from donkeys destined to food consumption in Tunisia with predominance of the ruminant associated CC133 lineage. *BMC Vet. Res.* **8**, 203.
- Gharsa, H., Ben Slama, K., Gómez-Sanz, E., Lozano, C., Zarazaga, M., Messadi, L., Boudabous, A. & Torres, C. 2015 Molecular characterization of *Staphylococcus aureus* from nasal samples of healthy farm animals and pets in Tunisia. *Vector Borne Zoonotic Dis.* **15**, 109–115.

- Gómez, P., Lozano, C., Benito, D., Estepa, V., Tenorio, C., Zarazaga, M. & Torres, C. 2016 Characterization of *Staphylococci* in urban wastewater treatment plants in Spain, with detection of methicillin-resistant *Staphylococcus aureus* ST398. *Environ. Pollut.* **212**, 71–76.
- Gómez-Sanz, E., Torres, C., Lozano, C., Fernandez-Perez, R., Aspiroz, C., Ruiz-Larrea, F. & Zarazaga, M. 2010 Detection, molecular characterization, and clonal diversity of methicillin-resistant *Staphylococcus aureus* CC398 and CC97 in Spanish slaughter pigs of different age groups. *Foodborne Pathog. Dis.* **7**, 1269–1277.
- Harmsen, D., Claus, H., Witte, W., Rothganger, J., Claus, H., Turnwald, D. & Vogel, U. 2003 Typing of methicillin-resistant *Staphylococcus aureus* in a university hospital setting by using novel software for *spa* repeat determination and database management. *J. Clin. Microbiol.* **41**, 5442–5448.
- Hasman, H., Moodley, A., Guardabassi, L., Stegger, M., Skov, R. L. & Aarestrup, F. M. 2010 *Spa* type distribution in *Staphylococcus aureus* originating from pigs, cattle and poultry. *Vet. Microbiol.* **141**, 326–331.
- Hata, E., Katsuda, K., Kobayashi, H., Uchida, I., Tanaka, K. & Eguchi, M. 2010 Genetic variation among *Staphylococcus aureus* strains from bovine milk and their relevance to methicillin-resistant isolates from humans. *J. Clin. Microbiol.* **48**, 2130–2139.
- Jarraud, S., Mougel, C., Thioulouse, J., Lina, G., Meugnier, H., Forey, F., Nesme, X., Etienne, J. & Vandenesch, F. 2002 Relationships between *Staphylococcus aureus* genetic background, virulence factors, *agr* groups (alleles), and human disease. *Infect. Immun.* **70**, 631–641.
- Kechrid, A., Pérez-Vázquez, M., Smaoui, H., Hariga, D., Rodríguez-Baños, M., Vindel, A., Baquero, F., Cantón, R. & Del Campo, R. 2010 Molecular analysis of community-acquired methicillin-susceptible and resistant *Staphylococcus aureus* isolates recovered from bacteraemia and osteomyelitis infections in children from Tunisia. *Clin. Microbiol. Infect.* **17**, 1020–1026.
- Kumar, H., Palaha, R., Kaur, N., Ratnakar, W. S., Sodi, A., Kaur, M., Katiyar, R., Sharma, M., Kaur, C. & Kumar, V. 2015 *Staphylococcus aureus* in nasal carriage, food, wastewater and paper currency in Jalandhar city (north-western), an Indian state of Punjab. *Environ. Assess.* **187**, 4134.
- Lamamy, C., Berthelot, A., Bertrand, X., Valentin, A. S., Dos Santos, S., Thiais, S., Morange, V., Girard, N., Donnio, P. Y., Quentin, R., Schrenzel, J., François, P. & van der Mee-Marquet, N. 2013 CC9 livestock-associated *Staphylococcus aureus* emerges in bloodstream infection in French patients unconnected with animal farming. *Clin. Infect. Dis.* **56**, 83–86.
- Lina, G., Piémont, Y., Godail-Gamot, F., Bes, M., Peter, M. O., Gauduchon, V., Vandenesch, F. & Etienne, J. 1999 Involvement of Panton-Valentine leukocidin-producing *Staphylococcus aureus* in primary skin infections and pneumonia. *Clin. Infect. Dis.* **29**, 1128–1132.
- Lowy, F. D. 2003 Antimicrobial resistance: the example of *Staphylococcus aureus*. *J. Clin. Invest.* **111**, 1265–1273.
- Lozano, C., Aspiroz, C., Charlez, L., Gomez-Sanz, E., Toledo, M., Zarazaga, M. & Torres, C. 2011 Skin lesion by methicillin-resistant *Staphylococcus aureus* ST398-t1451 in a Spanish pig farmer: possible transmission from animals to humans. *Vector Borne Zoonotic Dis.* **11**, 605–607.
- Lozano, C., Aspiroz, C., Rezusta, A., Gómez-Sanz, E., Simon, C., Gómez, P., Ortega, C., Revillo, M. J., Zarazaga, M. & Torres, C. 2012 Identification of novel *vga(A)*-carrying plasmids and a *tn5406*-like transposon in methicillin-resistant *Staphylococcus aureus* and *Staphylococcus epidermidis* of human and animal origin. *Int. J. Antimicrob. Agents* **40**, 306–312.
- Lozano, C., Mari, A., Aspiroz, C., Gómez-Sanz, E., Ceballos, S., Fortuno, B., Barcenilla, F., Jover-Sáenz, A. & Torres, C. 2015 Nasal carriage of coagulase positive staphylococci in patients of a Primary-Healthcare-Centre: genetic lineages and resistance and virulence genes. *Enferm. Infecc. Microbiol. Clin.* **33**, 391–396.
- Monecke, S., Luedicke, C., Slickers, P. & Ehrlich, R. 2009 Molecular epidemiology of *Staphylococcus aureus* in asymptomatic carriers. *Eur. J. Clin. Microbiol. Dis.* **28**, 1159–1165.
- Nübel, U., Roumagnac, P., Feldkamp, M., Song, J. H., Ko, K. S., Huang, Y. C., Coombs, G., Ip, M., Westh, H., Skov, R., Struelens, M. J., Goering, R. V., Strommenger, B., Weller, A., Witte, W. & Achtman, M. 2008 Frequent emergence and limited geographic dispersal of methicillin-resistant *Staphylococcus aureus*. *Proc. Natl Acad. Sci. USA* **105**, 14130–14135.
- Porrero, M. C., Hasman, H., Vela, A. I., Fernandez-Garayzabal, J. F., Dominguez, L. & Aarestrup, F. M. 2012 Clonal diversity of *Staphylococcus aureus* originating from the small ruminants goats and sheep. *Vet. Microbiol.* **156**, 157–161.
- Porrero, M. C., Valverde, A., Fernandez-Llario, P., Díez-Guerrier, A., Mateos, A., Lavin, S., Canton, R., Fernandez-Garayzabal, J. F. & Dominguez, L. 2014 *Staphylococcus aureus* carrying *mecC* gene in animals and urban wastewater, Spain. *Infect. Dis.* **20**, 899–901.
- Rijnders, M. I. A., Deurenberg, R. H., Boumans, M. L. L., Hoogkamp-Korstanje, J. A. A., Beisser, P. S. & the Antibiotic Resistance Surveillance Group, Stobberingh, E. E. 2009 Population structure of *Staphylococcus aureus* strains isolated from Intensive Care Unit patients in the Netherlands over an 11-year period (1996 to 2006). *J. Clin. Microbiol.* **47**, 4090–4095.
- Rizzo, L., Manaia, C., Merlin, C., Schwartz, T., Dagot, C., Ploy, M. C., Michael, I. & Fatta-Kassions, D. 2013 Urban wastewater treatment plants as hotspots for antibiotic resistant bacteria and genes spread into the environment: a review. *Sci. Total Environ.* **447**, 345–360.
- Rosenberg Goldstein, R. E., Micallef, S. A., Gibbs, S. G., Davis, J. A., He, X., George, A., Kleinfelter, L. M., Schreiber, N. A., Mukherjee, S., Sapkota, A., Joseph, S. W. & Sapkota, A. R. 2012 Methicillin-resistant *Staphylococcus aureus* (MRSA) detected at four U.S. wastewater treatment plants. *Environ. Health Perspect.* **120**, 1551–1558.
- Schmitz, F. J., Sadurski, R., Kary, A., Boos, M., Geisel, R., Khörer, K., Verhoef, J. & Fluit, A. C. 2000 Prevalence of

- macrolide-resistance genes in *Staphylococcus aureus* and *Enterococcus faecium* isolates from 24 European university hospitals. *J. Antimicrob. Chemother.* **45**, 891–894.
- Schwartz, T., Kohnen, W., Jansen, B. & Obst, U. 2003 Detection of antibiotic resistant bacteria and their resistance genes in wastewater, surface water, and drinking water biofilms. *FEMS Microbiol. Ecol.* **43**, 325–335.
- Shannon, K. E., Lee, D. Y., Trevors, J. T. & Beaudette, L. A. 2007 Application of real-time quantitative PCR for the detection of selected bacterial pathogens during municipal wastewater treatment. *Sci. Total Environ.* **382**, 121–129.
- Sharma, N. K., Garg, R., Baliga, S. & Bhat, K. G. 2013 Nosocomial infections and drug susceptibility patterns in methicillin sensitive and methicillin-resistant *Staphylococcus aureus*. *J. Clin. Diagn. Res.* **7**, 2178–2180.
- Shopsin, B., Mathema, B., Alcabes, P., Said-Salim, B., Lina, G., Matsuka, A., Martinez, J. & Kreiswirth, B. N. 2003 Prevalence of *agr* specificity groups among *Staphylococcus aureus* strains colonizing children and their guardians. *J. Clin. Microbiol.* **41**, 456–459.
- Tran, N. H., Yew-Hoong Gin, K. & Ngo, H. H. 2015 Faecal pollution source tracking toolbox for identification, evaluation and characterization of faecal contamination in receiving urban surface waters and groundwater. *Sci. Total Environ.* **538**, 38–57.
- Wendlandt, S., Feßler, A. T., Monecke, S., Ehricht, R., Schwarz, S. & Kadlec, K. 2015 The diversity of antimicrobial resistance genes among staphylococci of animal origin. *Int. J. Med. Microbiol.* **303**, 338–349.
- Wendlandt, S., Shen, J., Kadlec, K., Wang, Y., Li, B., Zhang, W. J., Feßler, A. T., Wu, C. & Schwarz, S. 2015 Multidrug resistance genes in staphylococci from animals that confer resistance to critically and highly important antimicrobial agents in human medicine. *Trends Microbiol.* **23**, 44–54.

First received 16 September 2016; accepted in revised form 26 February 2017. Available online 5 April 2017