

## Molecular analysis of *Blastocystis* sp. and its subtypes from treated wastewater routinely used for irrigation of vegetable farmlands in Iran

Ehsan Javanmard, Hanieh Mohammad Rahimi, Maryam Niyiyati, Hamid Asadzadeh Aghdaei, Meysam Sharifdini, Hamed Mirjalali, Mohammad Reza Zali and Panagiotis Karanis

### ABSTRACT

Treated wastewater samples were collected, filtered using sterile 47-mm cellulose nitrate membrane and DNA extracted from the filtered materials. The presence of *Blastocystis* sp. was confirmed via polymerase chain reaction (PCR) targeting the SSU rRNA gene of *Blastocystis* sp. in 5/12 of samples. Based on the subtype analysis after sequencing, 2, 2 and 1 of ST2, ST6 and ST8 were detected among the isolates, respectively. Furthermore, both ST6s were allele 139, alleles 11 and 138 were identified in ST2 and the only ST8 was allele 95. The phylogenetic tree showed that one of ST2 was clustered together with those ST2 that were already reported from humans and animals. The presence of *Blastocystis* sp. in treated wastewater can indicate the potential role of this type of water for irrigation in the transmission of pathogenic microorganisms to downstream farmlands.

**Key words** | allele discrimination, *Blastocystis* sp., phylogenetic analysis, subtypes, treated wastewater

**Ehsan Javanmard**  
**Hanieh Mohammad Rahimi**  
**Hamed Mirjalali** (corresponding author)  
Foodborne and Waterborne Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran  
E-mail: [hamedmirjalali@sbm.ac.ir](mailto:hamedmirjalali@sbm.ac.ir)

**Maryam Niyiyati**  
Department of Medical Parasitology and Mycology, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

**Hamid Asadzadeh Aghdaei**  
Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

**Meysam Sharifdini**  
Department of Medical Parasitology and Mycology, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran

**Mohammad Reza Zali**  
Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

**Panagiotis Karanis**  
University of Cologne, Faculty of Medicine and University Hospital Cologne, Cologne, Germany

### INTRODUCTION

Waterborne diseases are considered as one of the most important public health problems, all over the world. Together with bacteria and viruses, protozoan parasites are frequently reported from waterborne outbreaks in both developing and developed countries (Karanis *et al.* 2007;

Baldursson & Karanis 2011; Moss 2016; Efstratiou *et al.* 2017). Importantly, polluted water resources are a public health concern, which threaten millions of people, particularly those living in developing countries without access to safe water resources for drinking, bathing, and irrigation

proposes (WHO/UNICEF 2014; Aldeyarbi *et al.* 2016; Plutzer & Karanis 2016; Mahmoudi *et al.* 2017; Rosado-Garcia *et al.* 2017; Ahmed *et al.* 2018). Therefore, people inhabiting areas without easy access to clean water are at risk for intestinal infections caused by waterborne parasites (Andrade *et al.* 2018). Although *Cryptosporidium* species and *Giardia duodenalis* (*G. intestinalis*, *G. lamblia*) are the most prevalent reported protozoan parasites, other protozoan parasites such as *Entamoeba histolytica*, *Cyclospora cayotensis*, *Toxoplasma gondii*, *Isospora belli*, *Blastocystis* sp., *Acanthamoeba* spp., *Balantidium coli*, and *Naegleria fowleri* are also involved in waterborne outbreaks (Karanis *et al.* 2007; Baldursson & Karanis 2011; Plutzer & Karanis 2016; Efstratiou *et al.* 2017).

*Blastocystis* sp. has been found to be a highly prevalent intestinal parasite of humans that is mainly transmitted via contaminated food and water as well as close contact with animals (Parija & Jeremiah 2013; Wawrzyniak *et al.* 2013; Stensvold & Clark 2016). *Blastocystis* sp. is a non-invasive, unicellular, large intestinal protozoan parasite with multiple stages (Vassalos *et al.* 2010; Ahmed & Karanis 2018). According to results from clinical studies, *Blastocystis* sp. has been isolated from stool specimens of patients with diarrhea, constipation, vomiting, cutaneous rash, inflammatory bowel syndrome, and inflammatory bowel diseases (Carrascosa *et al.* 1996; Jimenez-Gonzalez *et al.* 2012; El Safadi *et al.* 2013; Casero *et al.* 2015; Mirjalali *et al.* 2017; Cifre *et al.* 2018; Rezaei Riabi *et al.* 2018). Wide distribution, high prevalence, and a broad spectrum of hosts have substantially increased the number of studies on this microorganism in recent years.

In recent decades, molecular tools have been applied to identify lineages of *Blastocystis* sp. Accordingly, based on the phylogenetic classification of discriminative fragment of the subunit ribosomal RNA (SSU rRNA) gene, 17 subtypes have so far been characterized (Stensvold *et al.* 2007; Alfellani *et al.* 2013a; Clark *et al.* 2013). Although correlation between clinical manifestations and certain subtypes is under question (Al-Fellani *et al.* 2007; Alinaghizade *et al.* 2017; Jalallou *et al.* 2017; Rezaei Riabi *et al.* 2017), subtype determination has mainly been applied for molecular epidemiology and source finding (Alfellani *et al.* 2013a; 2013b; Yoshikawa *et al.* 2016).

Although the interest in this parasite has rapidly increased in recent years, few studies have indicated the

importance of water transmission of *Blastocystis* sp. to human communities (Lee *et al.* 2012; Anuar *et al.* 2013; Angelici *et al.* 2018). Accordingly, *Blastocystis* sp. was detected in drinking water (Leelayoova *et al.* 2004, 2008), water catchment (Noradilah *et al.* 2016), recreational water (Ithoi *et al.* 2011) as well as river and sea waters (Koloren *et al.* 2018). More recently, a protocol was established and explained by Lee *et al.* (2019) for detection of *Blastocystis* sp. from water samples.

In the current study, we investigated the presence of *Blastocystis* sp. and its subtypes and alleles in treated wastewater (TW) that was routinely used for irrigation of vegetable farmlands.

## METHODS

### Sampling

In the current study, we performed investigations on 12 treated wastewater samples that were collected in our previous study by Javanmard *et al.* (2018a) from a one-year follow-up study on a wastewater treatment plant (WTP) located south of Tehran, Tehran province, Iran. This WTP is the biggest sewage treatment complex and collects a huge proportion of wastewaters in Tehran. Coagulation, flocculation, sedimentation, activated sludge, and disinfectant were used during the treatment process. The WTP collects approximately more than 70% of the sewage of Tehran. This study was a one-year follow-up of this WTP, thus, only 12 samples were investigated. All samples were collected from the WTP during monthly microbial quality control from a depth of 50 cm below the surface. Gathered samples were aseptically transported to the Parasitology and Water Research Laboratory of Foodborne and Waterborne Diseases Research Center in Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran for further processing.

### Processing samples

In the laboratory, samples were processed in the same way as our previous study by Javanmard *et al.* (2018a). Briefly,

samples were filtered using a six-branch filtration system (Sartorius, Gottingen, Germany) and sterile 47-mm cellulose nitrate membrane with pore size 0.4  $\mu\text{m}$  together with a sterile water as the negative control. Then, the filtered membranes were washed in 30 mL of sterile PBS and kept at 4–8 °C for 24 h. Finally, samples were centrifuged at 2,500  $\times g$  for 20 min at 4 °C, supernatant was discarded and the remaining pellet was stored at –20 °C until DNA extraction.

### DNA extraction and PCR amplification

DNA of samples were extracted using a YTA DNA extraction kit (Yekta Tajhiz, Tehran, Iran) according to the manufacturer's instructions with some modifications. Briefly, 250  $\mu\text{L}$  of pellet sample was washed three times with sterile PBS (pH = 7.5), and after final centrifugation, supernatant was discarded and the pellet was added to 2 mL sterile tubes containing glass beads. After 2 h incubation of samples at 60 °C together with 40  $\mu\text{g}/\text{mL}$  of proteinase K, DNA was extracted and purified. Isolated DNA was stored at –20 °C until use.

In order to amplify the discriminative fragment for subtyping, the primer set BhrDr (GAGCTTTTAACTGCAAC AACG) and the broad-specificity eukaryote-specific primer RD5 (ATCTGGTTGATCCTGCCAGT), described elsewhere (Scicluna *et al.* 2006), were employed to target ~620-bp fragment of the SSU rRNA gene of *Blastocystis* sp. according to the following conditions: an initial denaturing step of 95 °C for 5 min and 35 cycles consisting of 94 °C for 30 s, 60 °C for 30 s, and 30 s at 72 °C. A final extension at 72 °C was performed for 5 min. Subsequently, 10  $\mu\text{L}$  of each polymerase chain reaction (PCR) product was electrophoresed on a 1.5% agarose gel in TBE (0.09 M Tris, 0.09 M boric acid, 2 mM EDTA) stained with 0.5  $\mu\text{g}/\text{mL}$  ethidium bromide and visualized using a UV Transilluminator. Finally, an ABI (3130) sequencer (California, USA) sequenced 20  $\mu\text{L}$  of all PCR products. The generated sequences were edited and trimmed by Chromas software and the results were compared with those available sequences in GenBank database (BLAST; <http://blast.ncbi.nlm.nih.gov/>). Furthermore, in order to characterize alleles, edited sequences were compared (<http://pubmlst.org/blastocystis/>). All sequences were

submitted in the GenBank database with accession numbers MH915564–MH915567.

### Phylogenetic analysis

All *Blastocystis* sp. sequences obtained from the new isolates were multi-aligned, with ClustalW software, with those of humans and animals, which were previously deposited in GenBank. A phylogenetic tree was drawn using Molecular and Evolution Genetic Analysis Software (MEGA v.7) based on the maximum-likelihood model and Tamura-3-parameter option (Tamura *et al.* 2013). In order to calculate statistical significance, bootstrap 1,000 was employed.

## RESULTS

### PCR amplification, subtyping and allele discrimination

The targeted fragment of the SSU rRNA gene was successfully amplified in 5/12 of samples. All five *Blastocystis* sp.-positive samples were sequenced such that the result showed the presence of 2, 2, and 1 of ST2, ST6, and ST8 among treated water samples, respectively. The findings showed that *Blastocystis* sp. was present once in the investigated treated waters in all four seasons, while two *Blastocystis* sp.-positive isolates were identified in the summer period. Furthermore, allele discrimination showed that both ST6s were allele 139, while alleles 11 and 138 were identified in ST2 and the only ST8 was allele 95 (Table 1).

### Phylogenetic analysis

A phylogenetic tree was drawn that showed all obtained subtypes were clearly separated. Accordingly, ST2 obtained from the current study was placed together with other subtypes 2, which were isolated from humans and cattle, while it was molecularly closer to human isolates. Both subtypes 6 were grouped together and far from other ST6, which were isolated from chickens and humans even though molecular diversity was seen between them. Indeed, the only ST8 was separated from other subtypes and placed in a clade closer to ST6 than those ST8 that were characterized from non-human primates (Figure 1).



sequencing of the 'barcoding region' showed the majority of ST1–3 in both human and animals, some other less prevalent subtypes were recently reported from human cases. In a study conducted by Rezaei Riabi *et al.* (2018), ST6 and ST7 were isolated from human cases. However, they concluded that probably animal sources were involved in the transmission cycle of *Blastocystis* sp. Indeed, based on the non-host specificity feature of most human-infecting subtypes of *Blastocystis* sp. (Stensvold 2013; Stensvold & Clark 2016), it seems that not only close contact with animals, but also water resources that are polluted with feces of animals can play an important role in the transmission of *Blastocystis* sp. to human communities. Angelici *et al.* (2018) reported the case of a symptomatic patient who carried *Blastocystis* sp. subtype 1 and based on the history of the infected subject, they strongly suggested that drinking from an unpotable water source was the source of transmission.

The important role of water resources, particularly surface water in transmission of *Blastocystis* sp. to humans, was previously indicated (Table 2). Leelayoova *et al.* (2008) in a study on the prevalence of *Blastocystis* sp. in schoolchildren of a rural community in Thailand proposed that the water supply within a school was most

probably the source of infection. Researchers in Malaysia investigated the presence of *Blastocystis* sp. among primary schoolchildren and claimed that, based on the epidemiologic evidence, the source of drinking water most probably was infected with *Blastocystis* sp. (Abdulsalam *et al.* 2012). At the same time, another study in Nepal by Lee *et al.* (2012), which was performed on animals and river water, showed the majority of *Blastocystis* sp. ST4 in both animals and river waters and suggested high probability of zoonotic transmission of *Blastocystis* sp. via water resources.

The presence of *Blastocystis* in wastewater has also been reported. Suresh *et al.* (2005) investigated the presence of *Blastocystis* sp. in influent and effluent of a WTP and showed the presence of viable cysts of *Blastocystis* sp. in both types of sewage. In another study conducted by Banaticla & Rivera (2011), viable cysts of *Blastocystis* sp. were detected from both influent and effluent of a WTP. Although the number of positive samples in the discharge water was significantly lower than that observed in raw sewage, the authors concluded that the presence of viable cysts in effluent may indicate inefficiency of the usual wastewater treatment process. Therefore, it seems that *Blastocystis* sp. can not only be resistant to the water treatment process and be distributed in water environments,

**Table 2** | All reports of *Blastocystis* sp. and its subtypes isolated from different types of water regarding method of detection and characterization, worldwide

No.	Country	Sample	Subtype	Prevalence of <i>Blastocystis</i> sp.	Method of detection	References
1	Iran (Asia)	Treated wastewater	ST2, ST6, ST8	5/12	PCR sequencing	Current study
2	Malaysia (Asia)	River water	ST1, ST2, ST3, ST4, ST8, ST10	27.7%	PCR sequencing	Ithoi <i>et al.</i> (2011)
3	Malaysia (Asia)	Wastewater	–	Influent 50%, effluent 28%	Microscopy	Suresh <i>et al.</i> (2005)
3	Thailand (Asia)	Drinking water	ST1, ST2	–	PCR-RFLP	Leelayoova <i>et al.</i> (2008)
4	Philippines (Asia)	Wastewater treatment plants	ST1, ST2	Influent 23%, effluent 7%	PCR sequencing	Banaticla & Rivera (2011)
5	Turkey (Europe)	River water, drinking water	ST1, ST3	10%	PCR sequencing	Koloren <i>et al.</i> (2018)
6	Italy (Europe)	Non-potable drinking water	ST1	Positive	PCR-RFLP	Angelici <i>et al.</i> (2018)
7	Egypt (Africa)	Ponds and canal water	–	15.87%	Microscopy	Khalifa <i>et al.</i> (2014)
8	Egypt (Africa)	Potable water tanks, river	–	1%	Microscopy	Elshazly <i>et al.</i> (2007)
9	Venezuela (South America)	River water	–	–	Microscopy	Mora <i>et al.</i> (2010)

but also remain viable during the common processes that are employed for treatment of drinking water and sewage.

In the current study, three subtypes, ST2, ST6, and ST8, were characterized among the TW samples. Apart from ST2, which is usually reported from human subjects, ST6 and ST8 were mainly described from birds and non-human primates, respectively. Regarding the fact that Iran is not the habitat of non-human primates, it seems that together with ST2, ST8 came from a human source. In the case of ST6, it is necessary to mention that the WTP was not covered, and thus, most probably, the source of this subtype was free-range birds with access to the WTP.

Interestingly, the results of phylogenetic analysis confirmed the probable sources of each subtype. In the phylogenetic analysis, ST2 was placed together with other subtypes 2, which were previously obtained from human subjects, while ST8 was located in a separate branch from those ST8 that were reported from non-human primates. This observation indicated that, most probably, the presence of ST8 in TW was correlated with human sources. Inversely, both ST6 obtained from the current study were grouped together but separated from those subtypes 6 that were previously isolated from human subjects in Iran. The current finding strengthens the hypothesis that the source of ST6 in TW was probably birds.

Surprisingly, allele analysis of *Blastocystis* sp. showed that apart from one of the ST2 isolates (allele 11), all other subtypes revealed alleles that were not previously reported in humans and animals. Accordingly, ST2 allele 11 was previously characterized by Ramirez *et al.* (2016, 2017) in South America and Rezaei Riabi *et al.* (2008) in Iran. Presumably, the presence of this allele in TW indicated the insufficiency of the usual method for elimination of *Blastocystis* sp. cysts from raw wastewater. On the other hand, in the current study, we identified ST2 allele 138 together with ST6 allele 139 and ST8 allele 95 and there are no data about their distribution in humans. However, detection of these alleles may suggest secondary contamination of treated wastewater with non-human isolates.

## CONCLUSION

Epidemiological frequency of *Blastocystis* sp. subtypes in symptomatic and asymptomatic patients has been reported

in different countries and their prevalence is variable according to geographical distribution. Although knowledge on the prevalence and distribution of *Blastocystis* sp. and its subtypes has increased to a great extent, it seems that this parasite is largely underestimated and there are not enough data about the role of water resources in transmission of this protozoan parasite. Furthermore, there is no comprehensive study on the sufficiency of wastewater treatment processes in elimination of *Blastocystis* sp. from sewage. The current study showed that treated water can remain either contaminated with *Blastocystis* sp. after wastewater treatment or polluted with this protozoan via access of free-range animals. However, evaluation of small sample size was the limitation of the current research, therefore, more intensive studies on large water samples, axenization and/or genetic analyses of new subtypes will be beneficial to improve the knowledge about *Blastocystis* sp. species and its subtypes in water resources.

## ACKNOWLEDGEMENTS

The authors appreciate the valuable cooperation of all of the staff of Foodborne and Waterborne Diseases Research Center. The authors declare that they have no conflict of interest. The current study was financially supported by Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran (Grant no. 929).

## REFERENCES

- Abdulsalam, A. M., Ithoi, I., Al-Mekhlafi, H. M., Ahmed, A., Surin, J. & Mak, J. W. 2012 [Drinking water is a significant predictor of \*Blastocystis\* infection among rural Malaysian primary schoolchildren](#). *Parasitology* **139** (8), 1014–1020.
- Ahmed, S. A. & Karanis, P. 2018 *Blastocystis* spp., ubiquitous parasite of human, animals and environment. In: *Encyclopedia of Environmental Health*, 2nd edn (J. O. Nriagu, ed.). Elsevier Science Inc., Burlington, MA, USA, pp 1–6.
- Ahmed, S. A., Guerrero Florez, M. & Karanis, P. 2018 [The impact of water crises and climate changes on the transmission of protozoan parasites in Africa](#). *Pathogens Global Health* **112**, 281–293.

- Aldeyarbi, H. M., Abu El-Ezz, N. M. & Karanis, P. 2016 *Cryptosporidium* and cryptosporidiosis: the African perspective. *Environmental Science and Pollution Research International* **23** (14), 13811–13821.
- Al-Fellani, M. A., Khan, A. H., Al-Gazoui, R. M., Zaid, M. K. & Al-Ferjani, M. A. 2007 Prevalence and clinical features of *Blastocystis hominis* infection among patients in Sebha, Libya. *Sultan Qaboos University of Medical Journal* **7** (1), 35–40.
- Alfellani, M. A., Stensvold, C. R., Vidal-Lapiedra, A., Onuoha, E. S., Fagbenro-Beyioku, A. F. & Clark, C. G. 2013a Variable geographic distribution of *Blastocystis* subtypes and its potential implications. *Acta Tropica* **126** (1), 11–18.
- Alfellani, M. A., Taner-Mulla, D., Jacob, A. S., Imeede, C. A., Yoshikawa, H., Stensvold, C. R. & Clark, C. G. 2013b Genetic diversity of *Blastocystis* in livestock and zoo animals. *Protist* **164** (4), 497–509.
- Alinaghizade, A., Mirjalali, H., Mohebbali, M., Stensvold, C. R. & Rezaeian, M. 2017 Inter- and intra-subtype variation of *Blastocystis* subtypes isolated from diarrheic and non-diarrheic patients in Iran. *Infection, Genetics and Evolution* **50**, 77–82.
- Andrade, L., O'Dwyer, J., O'Neill, E. & Hynds, P. 2018 Surface water flooding, groundwater contamination, and enteric disease in developed countries: a scoping review of connections and consequences. *Environmental Pollution* **236**, 540–549.
- Angelici, M. C., Nardis, C., Scarpelli, R. & Ade, P. 2018 *Blastocystis hominis* transmission by non-potable water: a case report in Italy. *New Microbiology* **41** (2), 173–177.
- Anuar, T. S., Ghani, M. K., Azreen, S. N., Salleh, F. M. & Mokhtar, N. 2013 *Blastocystis* infection in Malaysia: evidence of waterborne and human-to-human transmissions among the Proto-Malay, Negrito and Senoi tribes of Orang Asli. *Parasites & Vectors* **6**, 40.
- Baldursson, S. & Karanis, P. 2011 Waterborne transmission of protozoan parasites: review of worldwide outbreaks – an update 2004–2010. *Water Research* **45**(20), 6603–6614.
- Banaticla, J. E. & Rivera, W. L. 2011 Detection and subtype identification of *Blastocystis* isolates from wastewater samples in the Philippines. *Journal of Water and Health* **9** (1), 128–137.
- Carrasco, M., Martinez, J. & Perez-Castrillon, J. L. 1996 Hemorrhagic proctosigmoiditis and *Blastocystis hominis* infection. *Annals Internal Medicine* **124** (2), 278–279.
- Casero, R. D., Mongi, F., Sanchez, A. & Ramirez, J. D. 2015 *Blastocystis* and urticaria: examination of subtypes and morphotypes in an unusual clinical manifestation. *Acta Tropica* **148**, 156–161.
- Cifre, S., Gozalbo, M., Ortiz, V., Soriano, J. M., Merino, J. F. & Trelis, M. 2018 *Blastocystis* subtypes and their association with irritable bowel syndrome. *Medical Hypotheses* **116**, 4–9.
- Clark, C. G., van der Giezen, M., Alfellani, M. A. & Stensvold, C. R. 2013 Recent developments in *Blastocystis* research. *Advance Parasitology* **82**, 1–32.
- Efstratiou, A., Ongerth, J. E. & Karanis, P. 2017 Waterborne transmission of protozoan parasites: review of worldwide outbreaks – an update 2011–2016. *Water Research* **114**, 14–22.
- El Safadi, D., Meloni, D., Poirier, P., Osman, M., Cian, A., Gaayeb, L., Wawrzyniak, I., Delbac, F., El Alaoui, H., Delhaes, L., Dei-Cas, E., Mallat, H., Dabboussi, F., Hamze, M. & Viscogliosi, E. 2013 Molecular epidemiology of *Blastocystis* in Lebanon and correlation between subtype 1 and gastrointestinal symptoms. *American Journal of Tropical Medicine and Hygiene* **88** (6), 1203–1206.
- Elsazly, A. M., Elsheikha, H. M., Soltan, D. M., Mohammad, K. A. & Morsy, T. A. 2007 Protozoal pollution of surface water sources in Dakahlia Governorate, Egypt. *Journal of Egypt Society of Parasitology* **37** (1), 51–64.
- Ithoi, I., Jali, A., Mak, J. W., Wan Sulaiman, W. Y. & Mahmud, R. 2011 Occurrence of *Blastocystis* in water of two rivers from recreational areas in Malaysia. *Journal of Parasitology Research* **2011**, 123916.
- Jalallou, N., Irvani, S., Rezaeian, M., Alinaghizade, A. & Mirjalali, H. 2017 Subtypes distribution and frequency of *Blastocystis* sp. isolated from diarrheic and non-diarrheic patients. *Iranian Journal of Parasitology* **12** (1), 63–68.
- Javanmard, E., Mirjalali, H., Niyayati, M., Jalilzadeh, E., Seyed Tabaei, S. J., Asadzadeh Aghdaei, H., Nazemalhosseini-Mojarad, E. & Zali, M. R. 2018a Molecular and phylogenetic evidences of dispersion of human-infecting microsporidia to vegetable farms via irrigation with treated wastewater: one-year follow up. *International Journal of Hygiene and Environmental Health* **221** (4), 642–651.
- Javanmard, E., Niyayati, M., Ghasemi, E., Mirjalali, H., Asadzadeh Aghdaei, H. & Zali, M. R. 2018b Impacts of human development index and climate conditions on prevalence of *Blastocystis*: a systematic review and meta-analysis. *Acta Tropica* **185**, 193–203.
- Jimenez-Gonzalez, D. E., Martinez-Flores, W. A., Reyes-Gordillo, J., Ramirez-Miranda, M. E., Arroyo-Escalante, S., Romero-Valdovinos, M., Stark, D., Souza-Saldivar, V., Martinez-Hernandez, F., Flisser, A., Olivo-Diaz, A. & Maravilla, P. 2012 *Blastocystis* infection is associated with irritable bowel syndrome in a Mexican patient population. *Parasitology Research* **110** (3), 1269–1275.
- Karanis, P., Kourenti, C. & Smith, H. 2007 Waterborne transmission of protozoan parasites: a worldwide review of outbreaks and lessons learnt. *Journal of Water and Health* **5** (1), 1–38.
- Khalifa, R. M., Ahmad, A. K., Abdel-Hafeez, E. H. & Mosllem, F. A. 2014 Present status of protozoan pathogens causing water-borne disease in northern part of El-Minia Governorate, Egypt. *Journal of Egypt Society of Parasitology* **44** (3), 559–566.
- Koloren, Z., Gulabi, B. B. & Karanis, P. 2018 Molecular identification of *Blastocystis* sp. subtypes in water samples collected from Black sea, Turkey. *Acta Tropica* **180**, 58–68.
- Lee, L. I., Chye, T. T., Karmacharya, B. M. & Govind, S. K. 2012 *Blastocystis* sp.: waterborne zoonotic organism, a possibility? *Parasites & Vectors* **5**, 130.

- Lee, I. L., Tan, T. C. & Govind, S. K. 2019 Establishing a protocol for water sample processing for the detection of *Blastocystis* sp. *Experimental Parasitology* **198**, 105–110.
- Leelayoova, S., Rangsin, R., Taamasri, P., Naaglor, T., Thathaisong, U. & Mungthin, M. 2004 Evidence of waterborne transmission of *Blastocystis hominis*. *American Journal of Tropical Medicine and Hygiene* **70** (6), 658–662.
- Leelayoova, S., Siripattanapipong, S., Thathaisong, U., Naaglor, T., Taamasri, P., Piyaraj, P. & Mungthin, M. 2008 Drinking water: a possible source of *Blastocystis* spp. subtype 1 infection in schoolchildren of a rural community in central Thailand. *American Journal of Tropical Medicine and Hygiene* **79** (3), 401–406.
- Mahmoudi, M. R., Ongerth, J. E. & Karanis, P. 2017 *Cryptosporidium* and cryptosporidiosis: the Asian perspective. *International Journal of Hygiene and Environmental Health* **220** (7), 1098–1109.
- Mirjalali, H., Abbasi, M. R., Naderi, N., Hasani, Z., Mirsamadi, E. S., Stensvold, C. R., Balaii, H., Asadzadeh Aghdaei, H. & Zali, M. R. 2017 Distribution and phylogenetic analysis of *Blastocystis* sp. subtypes isolated from IBD patients and healthy individuals in Iran. *European Journal of Clinical Microbiology and Infectious Diseases* **36** (12), 2335–2342.
- Mora, L., Martinez, I., Figuera, L., Segura, M. & Del Valle, G. 2010 [Protozoans in superficial waters and faecal samples of individuals of rural populations of the Montes municipality, Sucre state, Venezuela]. *Investigation Clinics* **51** (4), 457–466.
- Moss, J. A. 2016 Waterborne pathogens: the protozoans. *Radiologic Technology* **88** (1), 27–48.
- Noradilah, S. A., Lee, I. L., Anuar, T. S., Salleh, F. M., Abdul Manap, S. N., Mohd Mohtar, N. S., Azrul, S. M., Abdullah, W. O. & Mokhtar, N. 2016 Occurrence of *Blastocystis* sp. in water catchments at Malay villages and Aboriginal settlement during wet and dry seasons in Peninsular Malaysia. *PeerJ* **4**, e2541.
- Parija, S. C. & Jeremiah, S. 2013 *Blastocystis*: taxonomy, biology and virulence. *Tropical Parasitology* **3** (1), 17–25.
- Plutzer, J. & Karanis, P. 2016 Neglected waterborne parasitic protozoa and their detection in water. *Water Research* **101**, 318–332.
- Ramirez, J. D., Sanchez, A., Hernandez, C., Florez, C., Bernal, M. C., Giraldo, J. C., Reyes, P., Lopez, M. C., Garcia, L., Cooper, P. J., Vicuna, Y., Mongi, F. & Casero, R. D. 2016 Geographic distribution of human *Blastocystis* subtypes in South America. *Infections, Genetics and Evolution* **41**, 32–35.
- Ramirez, J. D., Florez, C., Olivera, M., Bernal, M. C. & Giraldo, J. C. 2017 *Blastocystis* subtyping and its association with intestinal parasites in children from different geographical regions of Colombia. *PLoS One* **12** (2), e0172586.
- Rezaei Riabi, T., Haghghi, A., Mirjalali, H., Mohammad Ali Gol, S., Karamati, S. A., Ghasemian, M., Bahadori Monfared, A., Aghamohammadi, E. & Zojaji, H. 2017 Study of prevalence, distribution and clinical significance of *Blastocystis* isolated from two medical centers in Iran. *Gastroenterology Hepatology From Bed to Bench* **10** (Suppl. 1), S102–S107.
- Rezaei Riabi, T., Mirjalali, H., Haghghi, A., Rostami Nejad, M., Pourhoseingholi, M. A., Poirier, P., Delbac, F., Wawrzyniak, I. & Zali, M. R. 2018 Genetic diversity analysis of *Blastocystis* subtypes from both symptomatic and asymptomatic subjects using a barcoding region from the 18S rRNA gene. *Infections, Genetics and Evolution* **61**, 119–126.
- Rosado-Garcia, F. M., Guerrero-Florez, M., Karanis, G., Hinojosa, M. D. C. & Karanis, P. 2017 Water-borne protozoa parasites: the Latin American perspective. *International Journal of Hygiene and Environmental Health* **220** (5), 783–798.
- Scicluna, S. M., Tawari, B. & Clark, C. G. 2006 DNA barcoding of *Blastocystis*. *Protist* **157** (1), 77–85.
- Stensvold, C. R. 2013 *Blastocystis*: genetic diversity and molecular methods for diagnosis and epidemiology. *Tropical Parasitology* **3** (1), 26–34.
- Stensvold, C. R. & Clark, C. G. 2016 Current status of *Blastocystis*: a personal view. *Parasitology International* **65** (6 Pt B), 763–771.
- Stensvold, C. R., Suresh, G. K., Tan, K. S., Thompson, R. C., Traub, R. J., Viscogliosi, E., Yoshikawa, H. & Clark, C. G. 2007 Terminology for *Blastocystis* subtypes—a consensus. *Trends in Parasitology* **23** (3), 93–96.
- Suresh, K., Smith, H. V. & Tan, T. C. 2005 Viable *Blastocystis* cysts in Scottish and Malaysian sewage samples. *Applied Environmental Microbiology* **71** (9), 5619–5620.
- Tamura, K., Stecher, G., Peterson, D., Filipiński, A. & Kumar, S. 2013 MEGA6: molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution* **30** (12), 2725–2729.
- Vassalos, C. M., Spanakos, G., Vassalou, E., Papadopoulou, C. & Vakalis, N. 2010 Differences in clinical significance and morphologic features of *Blastocystis* sp. subtype 3. *American Journal of Clinical Pathology* **133** (2), 251–258.
- Wawrzyniak, I., Poirier, P., Viscogliosi, E., Dionigia, M., Texier, C., Delbac, F. & Alaoui, H. E. 2013 *Blastocystis*, an unrecognized parasite: an overview of pathogenesis and diagnosis. *Therapeutic Advances in Infectious Diseases* **1** (5), 167–178.
- WHO 2017 *Guidelines for Drinking Water Quality*. World Health Organization, Geneva, Switzerland.
- WHO/UNICEF 2014 *Progress on Drinking Water and Sanitation: 2014 Update*. World Health Organization, Geneva, Switzerland.
- Yoshikawa, H., Tokoro, M., Nagamoto, T., Arayama, S., Asih, P. B., Rozi, I. E. & Syafruddin, D. 2016 Molecular survey of *Blastocystis* sp. from humans and associated animals in an Indonesian community with poor hygiene. *Parasitology International* **65** (6 Pt B), 780–784.

First received 2 March 2019; accepted in revised form 2 May 2019. Available online 19 July 2019