

## High quantities of multidrug-resistant *Escherichia coli* are present in the Machángara urban river in Quito, Ecuador

David Ortega-Paredes, Pedro Barba, Santiago Mena-López,  
Nathaly Espinel, Verónica Crespo and Jeannete Zurita

### ABSTRACT

Urban river pollution by multidrug-resistant (MDR) bacteria constitutes an important public health concern. Epidemiologically important strains of MDR *Escherichia coli* transmissible at the human–animal–environment interfaces are especially worrying. Quantifying and characterizing MDR *E. coli* at a molecular level is thus imperative for understanding its epidemiology in natural environments and its role in the spread of resistance in precise geographical areas. Cefotaxime-resistant *E. coli* was characterized along the watercourse of the major urban river in Quito. Our results showed high quantities of cefotaxime-resistant *E. coli* ( $2.7 \times 10^3$ – $5.4 \times 10^5$  CFU/100 mL). The antimicrobial resistance index (ARI) revealed the exposure of the river to antibiotic contamination, and the multiple antibiotic resistance index indicated a high risk of contamination. The *bla*<sub>CTX-M-15</sub> gene was the most prevalent in our samples. Isolates also had class 1 integrons carrying aminoglycoside-modifying enzymes and folate pathway inhibitors. The isolates belonged to phylogroups A, B1 and D. Clonal complex 10 was found to be the most prevalent (ST10, ST44 and ST 167), followed by ST162, ST394 and ST46. Our study provides a warning about the high potential of the major urban river in Quito for spreading the epidemiologically important MDR *E. coli*.

**Key words** | antimicrobial resistance genes, Ecuador, extended spectrum  $\beta$ -lactamase, *Escherichia coli*, Machángara, urban river

David Ortega-Paredes  
Nathaly Espinel  
Jeannete Zurita (corresponding author)  
Facultad de Medicina,  
Pontificia Universidad Católica del Ecuador,  
Quito, Ecuador  
E-mail: [jzurita@zuritalaboratorios.com](mailto:jzurita@zuritalaboratorios.com)

David Ortega-Paredes  
Pedro Barba  
Jeannete Zurita  
Unidad de Investigaciones en Biomedicina,  
Zurita & Zurita Laboratorios,  
Quito, Ecuador

David Ortega-Paredes  
Universidad Central del Ecuador, Facultad de  
Medicina Veterinaria y Zootecnia, Unidad de  
Investigación de Enfermedades Transmitidas por  
Alimentos y Resistencias a los Antimicrobianos  
(UNIETAR),  
Quito, Ecuador

Santiago Mena-López  
Pontificia Universidad Católica del Ecuador,  
Escuela de Ciencias Geográficas,  
Quito, Ecuador

Verónica Crespo  
Pontificia Universidad Católica del Ecuador,  
Facultad de Ciencias Exactas y Naturales,  
Escuela de Biología,  
Quito, Ecuador

### INTRODUCTION

The rise of difficult-to-treat infections caused by Enterobacteriaceae resistance to third-generation cephalosporins (3GC) is a central health concern worldwide (Dolejska & Papagiannitsis 2018). Extended-spectrum  $\beta$ -lactamase (ESBL)-producing *Escherichia coli* are especially important due to their ability to cause intestinal and extraintestinal infections, and therapeutic responses to them are limited to last resort antibiotics (carbapenems, colistin and tigecycline) (Iovleva & Doi 2017; Srinivas & Rivard 2017; Talaga-ćwiertnia *et al.* 2017). Intriguingly, ESBL-producing *E. coli* can be a part of the intestinal microbiota of healthy

carriers for years before producing an infection. This particular characteristic leads to an underestimated impact of environmental and food contaminations with ESBL-producing *E. coli* on human health (Tellevik *et al.* 2016).

Currently, increasing evidence has revealed the spread of ESBL-producing *E. coli* through the human–animal–environment interfaces (Seni *et al.* 2018). Therefore, the potential dissemination of epidemiologically important strains of *E. coli* through these interfaces is particularly alarming. Evidence of environmental distribution of ESBL-producing *E. coli* (Seni *et al.* 2018) has forewarned about

the growing and complex problem of the dissemination of bacterial resistance that compromises environment integrity and public health (Chong *et al.* 2018). In a typical model, human activities produce high microbiological and chemical polluted water that is discharged into urban rivers. Subsequently, this contamination travels downstream to reach small towns, production fields and natural habitats, which has a negative impact on human and animal health, agricultural products and environment (Dolejska & Papagiannitsis 2018; Leonard *et al.* 2018; Manaia *et al.* 2018; Sanderson *et al.* 2018; Vounba *et al.* 2019). Nevertheless, there is limited information on the levels and characteristics of ESBL-producing *E. coli* in polluting rivers.

The characterization of *E. coli* from rivers at a molecular level (clonal groups/phylogroups and resistance profiles) could improve our understanding of their epidemiology on natural environments and the role of rivers in the spread of resistance in particular geographical areas (Bajaj *et al.* 2015). This topic has not been considered in our location. Consequently, the aim of our study was to quantify and characterize *E. coli* resistance to cefotaxime in the major urban river in the city of Quito, Ecuador.

## METHODS

### Study area

Quito, the capital city of Ecuador, is located in the Andean Region at an altitude that ranges from 1,800 meters above sea level (masl) to 4,794 masl. Most of the wastewater of the city is discharged directly towards the Machángara River through sewage drainage. This water flow originates in the Atacazo Highlands, located approximately 6.5 km from the south border of the city, and following a length of 40 km crosses the city from southwest to northeast until it reaches the Tumbaco valley.

### Sampling


Using a collection of three satellite orthorectified images from the Pleiades-1A sensor (date: June 2013, spatial resolution: 0.5 m), multispectral remote data processing, spatial analysis techniques and supported fieldwork testing, a land use/cover map at a 1:15,000 scale was derived. Using this

input, from October to November 2015, we collected samples from nine sites along a segment of approximately 13 km near the Machángara River. Sites were selected based on their accessibility, altitude (a gradient of 2,778–2,785 masl) and types of land use (agricultural, residential, conservation and industrial) (Figure 1). Each site was sampled five times and characterized by its socioeconomic allocation, altitude range as well as water pH and temperature. At each site, we sampled water from the middle of the stream with sterile 100 mL plastic flasks. The samples were stored in a 4 °C cooling box until it reached the laboratory for analysis, which was performed within 6 h of collection.

### Microbiological procedures

The quantitative evaluation of bacterial contamination was conducted using the filtration method. Briefly, in accordance with the estimate of coliform contamination, appropriate volumes of each sample were mixed with 100 mL of saline water (0.85%) and passed through 50 µm filters (Millipore, USA) using Microfil S Filtration Devices (Merck, USA) following the manufacturer's guidelines. The membranes were deposited on one of three kinds of plates, as follows: m-ColiBlue24® medium plates (Merck, USA) without antimicrobial selection for total *E. coli* count, plates supplemented with cefotaxime (3 µg/mL) for the quantification of 3GC-resistant *E. coli* and plates supplemented with imipenem (2 µg/mL) for carbapenem-resistant *E. coli*. The plates were incubated at 35 °C for 24 h. To determine *E. coli* quantities, we manually counted all blue colonies and expressed the data as the geometric means of *E. coli*/100 mL (Table 1).

Three resistant *E. coli* colonies from each positive sampling point were stored for further analysis at –80 °C using glycerol and Trypticase Soy Broth (TSB) (1:1, v/v). The species was confirmed using a Microflex LT Matrix-Assisted Laser Desorption/Ionization-Time Of Flight (MALDI-TOF) spectrometer (Bruker Daltonics, Inc., USA) and examined in the MALDI-TOF Biotyper v.3.1. ESBL production was identified using the double-disk test, and MIC profiles were performed using the VITEK®2 Compact with the AST-GN66 card (bioMérieux, USA), which is an automated microdilution system. Antimicrobial susceptibility data were interpreted following clinical breakpoints (CLSI 2019).



**Figure 1** | Study area and sampling points.

## Genetic analysis

ESBL genes were identified with polymerase chain reaction (PCR) protocols described by Carattoli *et al.* (2008) for *bla*<sub>CTX-M-1</sub>, Jiang *et al.* (2006) for *bla*<sub>CTX-M-2</sub>, Hopkins *et al.* (2006) for *bla*<sub>CTX-M-8</sub>, Paauw *et al.* (2006) for *bla*<sub>CTX-M-9</sub>, Liebert *et al.* (2004) for *bla*<sub>TEM</sub>, Arlet *et al.* (1997) for *bla*<sub>SHV</sub> and Zurita *et al.* (2013) for *bla*<sub>KPC</sub> (Ortega-Paredes *et al.* (2016) for colistin-resistant *mcr-1* gene and Lévesque *et al.* (1995) for integron 1 variable region). Fingerprint characterization was performed in selected isolates by repetitive element palindromic PCR (REP-PCR) analysis (Araújo *et al.* 2017), and multilocus sequence typing (MLST) was achieved using a scheme proposed at the University of Warwick (Wirth *et al.* 2006). Phylogroups were assigned as described elsewhere (Clermont *et al.* 2000). Positive and negative controls were included in each experiment. The PCR products of resistance genes and MLST were purified and sequenced, both strands, at Macrogen Inc. (South Korea). Sequences were edited and analyzed in Geneious R10 using the database of ResFinder-3.0 (Zankari *et al.* 2012).

## Statistical analysis

We used the equation described by Mohanta & Goel (2014) to calculate the antibacterial resistance index (ARI) for each cefotaxime-resistant *E. coli*-positive sampling point. Likewise, we calculated the multiple antibiotic resistance index (MARI) for each resistant *E. coli* isolate as outlined previously by Blasco *et al.* (2008).

REP-PCR bands were analyzed using BioNumerics software v.7.6 (Applied Maths, Sint-Martens-Latem, Belgium). Fragments ranging from 200 to 1,500 bp were included in the analysis. The unweighted pair group method using the arithmetic averages algorithm (UPGMA) with a 1.5% of tolerance was used to construct a dendrogram.

## RESULTS AND DISCUSSION

At the first sampling point (SP1), water was collected in a tunnel system that accumulates water filtered by the highlands ecosystem (Páramo). Part of this water is stored for future use and, after treatment, is distributed to the city; the rest feeds the tributaries of Machángara River. At this

**Table 1** | Quantification of total *Escherichia coli* and cefotaxime-resistant *Escherichia coli* in the sampling points

Sampling points	GPS coordinates	Altitude	Activities	Mean pH	Mean T° (°C)	Geometric mean of <i>E. coli</i> /100 mL	Geometric mean of <i>E. coli</i> resistant to CTX/100 mL	Ratio <i>E. coli</i> resistant to CTX/total <i>E. coli</i>	ARI of <i>E. coli</i> resistant to CTX
SP1	17M 0767699 UTM 9963162	3,616 m	Natural reserve	6.9	14.1	0	0	–	–
SP2	17M 0770448 UTM 9964422	3,102 m	Animal rearing, irrigation, domestic and recreational purposes	6.5	21.2	$1.14 \times 10^4$	0	–	–
SP3	17M 0770873 UTM 9965390	3,029 m	Domestic and industrial activities	6.6	21.7	$6 \times 10^5$	0	–	–
SP4	17M 0770873 UTM 9965390	3,029 m	Domestic and industrial activities	6.8	21.4	$9.8 \times 10^5$	$2.7 \times 10^5$	$1/3.63 \times 10^2$	0.48
SP5	17M 0775475 UTM 9972248	2,808 m	Domestic and industrial activities	6.5	22	$2.8 \times 10^6$	$5.8 \times 10^5$	$1/4.83 \times 10^2$	0.52
SP6	17M 07754711 UTM 9972248	2,809 m	Domestic and industrial activities	6.8	20.1	$1.3 \times 10^7$	$3.3 \times 10^5$	1/39.3	0.56
SP7	17M 0780894 UTM 9976817	2,605 m	Domestic and industrial activities	6.5	20.9	$1.1 \times 10^7$	$4.6 \times 10^5$	1/23.9	0.60
SP8	17M 0784248 UTM 9978510	2,393 m	Domestic and industrial activities	6.6	20.8	$1.4 \times 10^6$	$2.7 \times 10^4$	1/51.8	0.35
SP9	17M 0780894 UTM 9976818	2,598 m	Domestic and industrial activities	6.9	22.8	$9.4 \times 10^6$	$5.4 \times 10^5$	1/18.1	0.38

location point, we did not detect *E. coli*. Downstream, in points SP2 and SP3, the water flows freely along open grasslands where some people and domestic and wild animals use it, which may be a possible source of water contamination. At these points, we identified *E. coli*, but cefotaxime- and imipenem-resistant *E. coli* strains were absent. From points SP4 to SP9, we registered increasing quantities of *E. coli* and cefotaxime-resistant *E. coli*, consistent with the intensification of anthropogenic disturbance. Interestingly, with the increasing levels of *E. coli* loads, the number of cefotaxime-resistant *E. coli* within the total *E. coli* counts also increased, from 1 cefotaxime-resistant *E. coli* per 363 *E. coli* in SP4 to 1 per every 18 in SP9 (Table 1).

In our study, we recorded the quantification of cefotaxime-resistant *E. coli* between  $2.7 \times 10^3$  (SP4) and  $5.4 \times 10^5$  (SP9) CFU/100 mL, which is similar to ESBL-producing *E. coli* counts previously recorded in hospital wastewater effluents ( $2.5 \times 10^3$  to  $10^5$  CFU/100 mL) (Jiang et al. 2006; Drieux et al. 2016). These data and the data of river flow reported by Aguilar Alegria (2010) and de Bievre et al. (2008) allow us a simple projection to suggest that at P9 discharge to the ecosystems could occur at approximately  $10^{11}$  multidrug-resistant (MDR) *E. coli* per second. However, the data presented in this report are a snapshot of the Machángara River contamination. Further research programs are needed to establish seasonal variation of resistant bacteria in the river.

Despite imipenem-resistant *E. coli* not being detected in the samples, red colonies (other coliforms) were observed in points SP7, SP8 and SP9. Therefore, one colony for each of these points was characterized, resulting in *Citrobacter freundii*-producing *bla*<sub>KPC-2</sub> in points SP7 and SP8 and *Klebsiella pneumoniae*-producing *bla*<sub>KPC-2</sub> in SP9. However, no quantification data were registered. Consistent with these results, in Quito, resistance to carbapenems is frequently related to the *bla*<sub>KPC-2</sub> gene, which is primarily identified in *K. pneumoniae* (Zurita et al. 2013), and is less frequently found in other species.

An ARI is an excellent tool that allows one to analyze the dissemination of bacterial resistance in a given population at a specific location. Previous reports have identified that an ARI value of >0.2 is observed when isolates are exposed to antibiotic contamination (Mohanta & Goel 2014; Titilawo et al. 2015). In our study, the ARI value for cefotaxime-resistant *E. coli* in the sampling points

ranged from 0.35 to 0.60. Interestingly, the ARI changed along the length of the river that was studied – with higher values at sites in the middle of the study length – which could be related to higher amounts of pollutant (from point or nonpoint sources), including antibiotics, reaching the river at these middle sites. Indeed, points P5 and P6 receive discharge from the city center (including hospitals) and P7, which receives urban wastewater, exhibited the highest ARI value (0.60) at three times higher than the threshold. In contrast, the lowest ARI values were registered at P4, which receives discharge from agriculture and dairy farms (where antibiotic use is common), industry and domestic uses, and P8 and P9 at the end of the city, possibly due to the distance from the principal contamination sources and a dilution effect with water from the natural tributaries (Table 1).

The MARI indicates the level of multidrug resistance of a certain isolate. The isolation of strains with >0.2 in the MARI has been proposed as high-risk contamination (Titilawo et al. 2015). In our study, all cefotaxime-resistant *E. coli* isolates showed MARI values above this threshold (from 0.31 to 0.75). This might be a strong indication for a high-risk source of water contamination and evidence of the widespread antimicrobial resistance to the environment from Machángara River water. Even though none of the sampling sites presented *E. coli* isolates resistance to carbapenems, we did isolate two *C. freundii* isolates that were resistant to carbapenems from P7 and P8, with an MARI of 0.94; we also detected *K. pneumoniae* resistance to carbapenems from site P9, with an MARI of 0.50. These extremely resistant isolates suggest the existence of a genetically rich resistome in this water; however, more studies, including the analysis of other bacterial groups and metagenome analysis, are needed for more robust conclusions (Table 2).

The allele *bla*<sub>CTX-M-15</sub> was the most prevalent of the ESBL genes identified in our study, followed by *bla*<sub>CTX-M-18</sub>, *bla*<sub>CTX-M-3</sub>, *bla*<sub>CTX-M-65</sub> and *bla*<sub>CTX-M-20</sub>. In Latin America, the variant *bla*<sub>CTX-M-15</sub> is the most frequently reported in clinical settings. Moreover, to the best of our knowledge, this variant has been reported in *E. coli* isolated from bloodstream and urinary tract infections, chicken farms, dog feces and fresh vegetables in Quito (Garrido et al. 2017; Chiluisa-Guacho et al. 2018; Ortega-Paredes et al. 2018, 2019; Vinueza-Burgos et al. 2019; Zurita et al. 2019). These findings demonstrate the ubiquitous nature of *bla*<sub>CTX-M-15</sub> in our city.

**Table 2** | Susceptibility profiles of cefotaxime-resistant *E. coli* and imipenem-resistant *C. freundii* and *K. pneumoniae*

Isolate	Species	AMP	SAM	TPZ	CZ	FOX	CAZ	CRO	FEP	ERT	IMP	GEN	TOB	CIP	LEV	NIT	SXT	No. of antibiotics	MARI
M4-1	<i>E. coli</i>	≥ 32	16	≤4	≥ 64	≤4	≤1	16	2	≤0.5	≤0.25	≤1	≤1	≤0.25	≤0.12	≤16	≤20	4	0.25
M4-2	<i>E. coli</i>	≥ 32	16	≤4	≥ 64	16	16	≥ 64	≥ 64	≤0.5	≤0.25	≥ 16	8	≥ 4	≥ 8	≤16	≤ 320	12	0.75
M4-3	<i>E. coli</i>	≥ 32	16	≤4	≥ 64	8	4	16	2	≤0.5	≤0.25	≤1	≤1	≥ 4	4	≤16	≤ 320	7	0.44
M5-1	<i>E. coli</i>	≥ 32	≥ 32	≤4	≥ 64	≤4	16	≥ 64	2	≤0.5	≤0.25	≤1	8	≥ 4	≥ 8	≤16	≤ 320	9	0.56
M5-2	<i>E. coli</i>	≥ 32	≥ 32	8	≥ 64	≤4	16	≥ 64	2	≤0.5	≤0.25	≥ 16	≥ 16	≥ 4	≥ 8	64	≤ 320	11	0.69
M5-3	<i>E. coli</i>	≥ 32	16	≤4	≥ 64	≤4	≤1	32	2	≤0.5	≤0.25	≤1	≤1	≤0.25	≤0.5	≤16	≤ 320	5	0.31
M6-1	<i>E. coli</i>	≥ 32	16	≤4	≥ 64	8	16	≥ 64	16	≤0.5	≤0.25	≥ 16	8	≥ 4	≥ 8	≤16	≤ 320	11	0.69
M6-2	<i>E. coli</i>	≥ 32	16	≤4	≥ 64	≤4	≤1	32	2	≤0.5	≤0.25	≥ 16	8	≥ 4	≥ 8	32	≤ 320	9	0.56
M6-3	<i>E. coli</i>	≥ 32	16	≤4	≥ 64	≤4	4	16	2	≤0.5	≤0.25	≤1	≤1	≥ 4	≥ 8	≤16	≤ 320	7	0.44
M7-1	<i>E. coli</i>	≥ 32	8	≤4	≥ 64	16	4	≥ 64	≥ 64	≤0.5	≤0.25	≥ 16	8	≥ 4	≥ 8	≤16	≤ 320	10	0.63
M7-2	<i>E. coli</i>	≥ 32	16	≤4	≥ 64	8	16	≥ 64	2	≤0.5	≤0.25	≥ 16	8	≥ 4	≥ 8	≤16	≤ 320	10	0.63
M7-3	<i>E. coli</i>	≥ 32	16	≤4	≥ 64	≤4	≤1	32	≤1	≤0.5	≤0.25	≥ 16	8	≥ 4	≥ 8	≤16	≤ 320	9	0.56
M7-IMP	<i>C. freundii</i>	≥ 32	≥ 32	≥ 128	≥ 64	≥ 64	≥ 64	≥ 64	≥ 64	≥ 8	≥ 16	≥ 16	≥ 16	≥ 4	≥ 8	≤16	≤ 320	15	0.94
M8-1	<i>E. coli</i>	≥ 32	16	≤4	≥ 64	≤4	16	≥ 64	2	≤0.5	≤0.25	≤1	≤1	≤0.25	1	32	≤ 320	6	0.38
M8-2	<i>E. coli</i>	≥ 32	16	≤4	≥ 64	≤4	16	≥ 64	2	≤0.5	≤0.25	≤1	≤1	≤0.25	1	32	≤ 320	6	0.38
M8-3	<i>E. coli</i>	≥ 32	4	≤4	≥ 64	≤4	4	≥ 64	2	≤0.5	≤0.25	≤1	≤1	≥ 4	4	≤16	≤20	5	0.31
M8-IMP	<i>C. freundii</i>	≥ 32	≥ 32	64	≥ 64	≥ 64	16	≥ 64	16	≥ 8	≥ 16	≥ 16	≥ 16	≥ 4	≥ 8	≤16	≤ 320	15	0.94
M9-1	<i>E. coli</i>	≥ 32	≥ 32	≤4	≥ 64	8	≤1	32	2	≤0.5	≤0.25	≤1	≤1	≥ 4	≥ 8	≤16	≤ 320	7	0.44
M9-2	<i>E. coli</i>	≥ 32	≥ 32	≤4	≥ 64	≤4	≤1	≥ 64	2	≤0.5	≤0.25	≥ 16	2	≤0.5	1	≤16	≤ 320	6	0.38
M9-3	<i>E. coli</i>	≥ 32	16	≤4	≥ 64	≤4	≥1	16	2	≤0.5	≤0.25	≤1	≤1	≤0.25	≤0.12	≤16	≤ 320	5	0.31
M9-IMP	<i>K. pneumoniae</i>	≥ 32	≥ 32	≥ 128	≥ 64	≥ 64	4	16	8	≥ 8	≥ 16	≤1	≤1	1	1	≤16	≤20	8	0.50

Data in bold are interpreted as resistant. The MARI above 0.2 are related with MDR status.

AMP, ampicillin; SAM, ampicillin/sulbactam; TPZ, piperacillin/tazobactam; CZ, cephalothin; FOX, ceftioxitin; CAZ, ceftazidime; CRO, ceftriaxone; FEP, cefepime; ERT, ertapenem; IMP, imipenem; GEN, gentamicin; TOB, tobramycin; CIP, ciprofloxacin; LEV, levofloxacin; NIT, nitrofurantoin; SXT, trimethoprim/sulfamethoxazole.



**Table 3** | Genetic profiles of ESBL-producing *E. coli* isolates

	Isolate	ST complex	ST	Phylogroup	<i>bla</i> <sub>CTX-M</sub> G1	<i>bla</i> <sub>CTX-M</sub> G2	<i>bla</i> <sub>CTX-M</sub> G9	<i>bla</i> <sub>TEM</sub>	<i>bla</i> <sub>SHV</sub>	Integron 1 variable region
	M6-2	NA	New	D	<i>bla</i> <sub>CTX-M-3</sub>	-	-	<i>bla</i> <sub>TEM-1B</sub>	-	-
	M6-3	162	162	B1	<i>bla</i> <sub>CTX-M-15</sub>	-	-	<i>bla</i> <sub>TEM-1B</sub>	-	-
	M7-3	10	10	A	-	-	<i>bla</i> <sub>CTX-M-65</sub>	<i>bla</i> <sub>TEM-1B</sub>	-	<i>dftA17, aadA5</i>
	M5-3	394	394	D	-	-	<i>bla</i> <sub>CTX-M-18</sub>	-	-	<i>dftA1, aadA1</i>
	M5-2	46	46	A	<i>bla</i> <sub>CTX-M-15</sub>	-	-	-	-	<i>dftA17, aadA5</i>
	M7-1	NA	457	D	<i>bla</i> <sub>CTX-M-15</sub>	-	-	-	-	-
	M9-2	NA	1,711	A	<i>bla</i> <sub>CTX-M-15</sub>	-	-	<i>bla</i> <sub>TEM-1B</sub>	-	<i>dftA12, orfF, aadA2</i>
	M4-2	NA	1,140	D	<i>bla</i> <sub>CTX-M-15</sub>	-	-	<i>bla</i> <sub>TEM-1B</sub>	-	-
	M4-3	NA	457	D	<i>bla</i> <sub>CTX-M-15</sub>	-	-	<i>bla</i> <sub>TEM-1C</sub>	<i>bla</i> <sub>SHV-2</sub>	-
	M6-1	10	167	A	<i>bla</i> <sub>CTX-M-15</sub>	-	-	<i>bla</i> <sub>TEM-1B</sub>	-	<i>dftA12, orfF, aadA2</i>
	M7-2	10	167	A	<i>bla</i> <sub>CTX-M-15</sub>	-	-	<i>bla</i> <sub>TEM-1B</sub>	-	<i>dftA12, orfF, aadA2</i>
	M8-1	10	10	A	<i>bla</i> <sub>CTX-M-15</sub>	-	-	<i>bla</i> <sub>TEM-1B</sub>	-	<i>dftA1, aadA1</i>
	M8-2	10	10	A	<i>bla</i> <sub>CTX-M-15</sub>	-	-	<i>bla</i> <sub>TEM-1B</sub>	-	<i>dftA1, aadA1</i>
	M5-1	10	44	A	<i>bla</i> <sub>CTX-M-15</sub>	-	-	-	-	<i>dftA17, aadA5</i>
	M4-1	10	10	A	<i>bla</i> <sub>CTX-M-3</sub>	-	-	-	-	-
	M9-1	ND	ND	B1	-	-	<i>bla</i> <sub>CTX-M-18</sub>	<i>bla</i> <sub>TEM-1B</sub>	-	-
	M9-3	NA	362	D	-	<i>bla</i> <sub>CTX-M-29</sub>	-	-	-	<i>aadA1</i>
	M8-3	23	New	A	<i>bla</i> <sub>CTX-M-55</sub>	-	-	-	-	-

*E. coli* phylogroups A and B1 have been identified as animal commensals and having higher levels of resistance than B2 and D (Sarayaya et al. 2015). In our study, the isolates belonged to groups A, B1 and D, which probably result from multisource pollution and imply a possible role of polluted rivers as facilitators of resistance and virulence transference between *E. coli* strains. Previous studies in Quito reported these phylogroups in fresh vegetables and dog feces (Ortega-Paredes et al. 2018, 2019).

Some strains also had class 1 integrons carrying aminoglycoside-modifying enzymes and folate pathway inhibitors. The gene cassette related to aminoglycoside resistance has been identified previously in *E. coli* isolated from fresh vegetables in Quito (Ortega-Paredes et al. 2018). Additionally, the presence of these genetic recombination systems allows these strains to acquire more gene cassettes and increase their profiles of resistance in the river environment.

REP-PCR analysis showed low relatedness between isolates, and MLST was identified in most of our isolates which belonged to the clonal complex 10 (ST10, ST44 and ST 167). This clonal complex, particularly ST10, has been identified as predominant in MDR *E. coli* (Liu et al. 2018), which is related to extremely resistant phenotypes (Caltagirone et al. 2017) and with extraintestinal pathogenic potential (Gomi et al. 2017) in samples isolated from bodies of water. In Quito, we found *bla*<sub>CTX-M-15</sub>-producing *E. coli* ST44 in fresh commercialized vegetables (Ortega-Paredes et al. 2018), which suggests a relationship between contaminated water and vegetable products. In this study, two MDR *E. coli* that are belonged to ST167 were identified. This clone has been recently identified as clinically important, potentially zoonotic and in possession of high levels of antimicrobial resistance (ESBL and carbapenemases) (Grönthal et al. 2018; Peterhans et al. 2018; Sun et al. 2018). However, to the best of our knowledge, *E. coli* ST167 has not been previously reported in the Ecuadorian environment (Table 3). The rest of our samples belonged to ST162, ST394, ST46, all of which have been isolated previously from poultry and sewage (Hayashi et al. 2018; Zahra et al. 2018).

Finally, in confirmation of our concern, the sequence types described in this study show the typical genetic backgrounds of the MDR *E. coli* capable of causing infections that are difficult to treat in humans and animals.

## CONCLUSION

Our study reveals the high potential of polluted urban rivers as sites of emergence and means of spread of MDR *E. coli* and highlights the need to characterize their resistant determinants. The predominance of *bla*<sub>CTX-M-15</sub> in the isolates suggests the establishment of this variant in the city and its aquatic dissemination through the sewage to the environment.

## ACKNOWLEDGEMENTS

This work was supported by the Pontifical University of Ecuador (grant PUCE L13312) and by Unidad de Investigaciones en Biomedicina and Zurita & Zurita Laboratorios (Project MIC009). The authors thank Iliana Alcocer, Rafael Narváez, Bolívar Salas and Oscar Suing for their kind help in the development of this study. Part of this work was presented in Boston at the ASM Microbe Meeting in June 2016.

## CONFLICTS OF INTEREST

None declared.

## REFERENCES

- Aguilar Alegria, A. 2010 *Modelación hidrológica de crecidas en la Cuenca del río Machángara en la ciudad de Quito*. Tesis Ingeniería Civil y Ambiental. Escuela Politécnica Nacional. Quito Ecuador. <http://bibdigital.epn.edu.ec/handle/15000/2212>.
- Araújo, S., Silva, A. T., Tacão, M., Patinha, C., Alves, A. & Henriques, I. 2017 *Characterization of antibiotic resistant and pathogenic Escherichia coli in irrigation water and vegetables in household farms*. *International Journal of Food Microbiology* **257**, 192–200. doi:10.1016/j.ijfoodmicro.2017.06.020.
- Arlet, G., Rouveau, M. & Philippon, A. 1997 Substitution of alanine for aspartate at position 179 in the SHV-6 extended-spectrum beta-lactamase. *FEMS Microbiology Letters* **152** (1), 163–167.
- Bajaj, P., Singh, N. S., Kanaujia, P. K. & Virdi, J. S. 2015 *Distribution and molecular characterization of genes encoding CTX-M and AmpC β-lactamases in Escherichia coli isolated from an Indian urban aquatic environment*. *Science of the Total Environment* **505**, 350–356. doi:10.1016/j.scitotenv.2014.09.084.
- Blasco, M. D., Esteve, C. & Alcaide, E. 2008 *Multiresistant waterborne pathogens isolated from water reservoirs and*



- cooling systems. *Journal of Applied Microbiology* **105**, 469–475. doi:10.1111/j.1365-2672.2008.03765.x.
- Caltagirone, M., Nucleo, E., Spalla, M., Zara, F., Novazzi, F., Marchetti, V. M., Piazza, A., Bitar, I., De Cicco, M., Paolucci, S., Pilla, G., Migliavacca, R. & Pagani, L. 2017 Occurrence of extended spectrum  $\beta$ -lactamases, KPC-Type, and MCR-1.2-producing Enterobacteriaceae from wells, river water, and wastewater treatment plants in Oltrepò Pavese area, Northern Italy. *Frontiers in Microbiology* **8** (November), 1–12. doi:10.3389/fmicb.2017.02232.
- Carattoli, A., García-Fernández, A., Varesi, P., Fortini, D., Gerardi, S., Penni, A., Mancini, C. & Giordano, A. 2008 Molecular epidemiology of *Escherichia coli* producing extended-spectrum beta-lactamases isolated in Rome, Italy. *Journal of Clinical Microbiology* **46** (1), 103–108. doi:10.1128/JCM.01542-07.
- Chiluisa-Guacho, C., Escobar-Perez, J. & Dutra-Asensi, M. 2018 First detection of the CTX-M-15 producing *Escherichia coli* O25-ST131 pandemic clone in Ecuador. *Pathogens* **7** (2), 42. doi:10.3390/pathogens7020042.
- Chong, Y., Shimoda, S. & Shimono, N. 2018 Current epidemiology, genetic evolution and clinical impact of extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae*. *Infection, Genetics and Evolution* **61**, 185–188. doi:10.1016/j.meegid.2018.04.005.
- Clermont, O., Bonacorsi, S. & Bingen, E. 2000 Rapid and simple determination of the *Escherichia coli* phylogenetic group. *Applied and Environmental Microbiology* **6** (10), 4555–4558. doi:10.1128/AEM.66.10.4555-4558.2000.
- CLSI 2019 *M100 Performance Standards for Antimicrobial Susceptibility Testing*, 29th edn. Clinical and Laboratories Standards Institute, Wayne, PA.
- de Bievre, B., Coello, X. & Otto de Keizer, P. M. 2008 Modelo Hidrológico de la Hoya de Quito (Hydrological modeling of Quito's Hoya). Available from: [http://infoagua-guayllabamba.ec/repositorio/web/files/Modelo\\_hidrologico\\_de\\_la\\_Hoya\\_de\\_Quito.pdf](http://infoagua-guayllabamba.ec/repositorio/web/files/Modelo_hidrologico_de_la_Hoya_de_Quito.pdf) (accessed 22 July 2018).
- Dolejska, M. & Papagiannitsis, C. C. 2018 Plasmid-mediated resistance is going wild. *Plasmid* **99**, 99–111. doi:10.1016/j.plasmid.2018.09.010.
- Drieux, L., Haenn, S., Moulin, L. & Jarlier, V. 2016 Quantitative evaluation of extended spectrum  $\beta$ -lactamase-producing *Escherichia coli* strains in the wastewater of a French teaching hospital and relation to patient strain. *Antimicrobial Resistance and Infection Control* **5** (1), 1–5. doi:10.1186/s13756-016-0108-5.
- Garrido, D., Garrido, S., Gutiérrez, M., Calvopiña, L., Harrison, A. S., Fuseau, M. & Salazar Irigoyen, R. 2017 Caracterización clínica y resistencia antimicrobiana de *Escherichia coli* en pacientes pediátricos con infección del tracto urinario en un hospital de tercer nivel en Quito, Ecuador (Clinical characterization and antimicrobial resistance of *Escherichia coli* in pediatric patients with urinary tract infection in a tertiary hospital in Quito, Ecuador). *Boletín Médico Del Hospital Infantil de México* **74** (4), 265–271. doi:10.1016/j.bmhmx.2017.02.004.
- Gomi, R., Matsuda, T., Matsumura, Y. & Yamamoto, M. 2017 Whole-genome analysis of antimicrobial-resistant and extraintestinal pathogenic *Escherichia coli* in river water. *Applied and Environmental Microbiology* **83** (5), 1–14. doi:10.1128/AEM.02703-16.
- Grönthal, T., Österblad, M., Eklund, M., Jalava, J., Nykäsenoja, S., Pekkanen, K. & Rantala, M. 2018 Sharing more than friendship – transmission of NDM-5 ST167 and CTX-M-9 ST69 *Escherichia coli* between dogs and humans in a family, Finland, 2015. *Eurosurveillance* **23** (27). doi:10.2807/1560-7917.ES.2018.23.27.1700497.
- Hayashi, W., Ohsaki, Y., Taniguchi, Y., Koide, S., Kawamura, K., Suzuki, M., Kimura, K., Wachino, J. I., Nagano, Y., Arakawa, Y. & Nagano, N. 2018 High prevalence of blaCTX-M-14 among genetically diverse *Escherichia coli* recovered from retail raw chicken meat portions in Japan. *International Journal of Food Microbiology* **284**, 98–104. doi:10.1016/j.ijfoodmicro.2018.08.003.
- Hopkins, K. L., Batchelor, M. J., Liebana, E., Deheer-Graham, A. P. & Threlfall, E. J. 2006 Characterisation of CTX-M and AmpC genes in human isolates of *Escherichia coli* identified between 1995 and 2003 in England and Wales. *International Journal of Antimicrobial Agents* **28** (3), 180–192. doi:10.1016/j.ijantimicag.2006.03.027.
- Iovleva, A. & Doi, Y. 2017 Carbapenem-resistant Enterobacteriaceae. *Clinics in Laboratory Medicine* **37** (2), 303–315. doi:10.1016/j.cll.2017.01.005.
- Jiang, X., Zhang, Z., Li, M., Zhou, D., Ruan, F. & Lu, Y. 2006 Detection of extended-spectrum beta-lactamases in clinical isolates of *Pseudomonas aeruginosa*. *Antimicrobial Agents and Chemotherapy* **50** (9), 2990–2995. doi:10.1128/AAC.01511-05.
- Leonard, A. F. C., Zhang, L., Balfour, A. J., Garside, R., Hawkey, P. M., Murray, A. K., Ukoumunne, O. C. & Gaze, W. H. 2018 Exposure to and colonisation by antibiotic-resistant *E. coli* in UK coastal water users: Environmental surveillance, exposure assessment, and epidemiological study (Beach Bum Survey). *Environment International* **114**, 326–333.
- Lévesque, C., Piché, L., Larose, C. & Roy, P. H. 1995 PCR mapping of integrons reveals several novel combinations of resistance genes. *Antimicrobial Agents and Chemotherapy* **39** (1), 185–191. doi:10.1128/AAC.39.1.185.
- Liebert, M. A., Olesen, I., Hasman, H. & Aarestrup, F. M. 2004 *Escherichia coli* and *Salmonella* isolated from food animals in Denmark. *Microbial Drug Resistance* **10** (4), 334–340.
- Liu, H., Zhou, H., Li, Q., Peng, Q., Zhao, Q., Wang, J. & Liu, X. 2018 Molecular characteristics of extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli* isolated from the rivers and lakes in Northwest China. *Frontiers in Microbiology* **9**, 1756. doi:10.3389/fmicb.2018.01756.
- Manaia, C. M., Rocha, J., Scaccia, N., Marano, R., Radu, E., Biancullo, F., Cerqueira, F., Fortunato, G., Iakovides, I. C., Zammit, I., Kampouris, I., Vaz-Moreira, I. & Nunes, O. C. 2018 Antibiotic resistance in wastewater treatment plants: tackling the black box. *Environment International* **115**, 312–324.

- Mohanta, T. & Goel, S. 2014 Prevalence of antibiotic-resistant bacteria in three different aquatic environments over three seasons. *Environmental Monitoring and Assessment* **186** (8), 5089–5100. doi:10.1007/s10661-014-3762-1.
- Ortega-Paredes, D., Barba, P. & Zurita, J. 2016 Colistin-resistant *Escherichia coli* clinical isolate harbouring the *mcr-1* gene in Ecuador. *Epidemiology and Infection* **144** (14), 2967–2970. doi:10.1017/S0950268816001369.
- Ortega-Paredes, D., Barba, P., Mena-López, S., Espinel, N. & Zurita, J. 2018 *Escherichia coli* hyperepidemic clone ST410-A harboring blaCTX-M-15 isolated from fresh vegetables in a municipal market in Quito-Ecuador. *International Journal of Food Microbiology* **2** (280), 41–45. doi:10.1016/j.ijfoodmicro.2018.04.037.
- Ortega-Paredes, D., Haro, M., Leoro-Garzón, P., Barba, P., Loaiza, K., Mora, F., Fors, M., Vinueza-Burgos, C. & Fernández-Moreira, E. 2019 Multi-drug resistant *Escherichia coli* isolated from canine feces in a public park in Quito, Ecuador. *Journal of Global Antimicrobial Resistance*. doi:10.1016/j.jgar.2019.04.002.
- Paauw, A., Fluit, A. C., Verhoef, J. & Leverstein-van Hall, M. A. 2006 Enterobacter cloacae outbreak and emergence of quinolone resistance gene in Dutch hospital. *Emerging Infectious Diseases* **12** (5), 807–812. doi:10.3201/eid1205.050910.
- Peterhans, S., Stevens, M. J. A., Nüesch-Inderbinen, M., Schmitt, S., Stephan, R. & Zurfluh, K. 2018 First report of a blaNDM-5 harbouring *Escherichia coli* ST167 isolated from a wound infection in a dog in Switzerland. *Journal of Global Antimicrobial Resistance* **7165** (18). doi:10.1016/J.JGAR.2018.10.013.
- Sanderson, C. E., Fox, J. T., Dougherty, E. R., Cameron, A. D. S. & Alexander, K. A. 2018 The changing face of water: a dynamic reflection of antibiotic resistance across landscapes. *Frontiers in Microbiology* **9**, 1–13. doi:10.3389/fmicb.2018.01894.
- Saralaya, V., Shenoy, S., Baliga, S., Hegde, A., Adhikari, P. & Chakraborty, A. 2015 Characterization of *Escherichia coli* phylogenetic groups associated with extraintestinal infections in South Indian population. *Annals of Medical and Health Sciences Research* **5** (4), 241. doi:10.4103/2141-9248.160192.
- Seni, J., Moremi, N., Matee, M., van der Meer, F., DeVinney, R., Mshana, S. E. & D Pitout, J. D. 2018 Preliminary insights into the occurrence of similar clones of extended-spectrum beta-lactamase-producing bacteria in humans, animals and the environment in Tanzania: a systematic review and meta-analysis between 2005 and 2016. *Zoonoses and Public Health* **65** (1), 1–10. doi:10.1111/zph.12387.
- Srinivas, P. & Rivard, K. 2017 Polymyxin resistance in Gram-negative pathogens. *Current Infectious Disease Reports* **19** (11), 38. doi:10.1007/s11908-017-0596-3.
- Sun, L., Xu, J. & He, F. 2018 Draft genome sequence of an NDM-5, CTX-M-15 and OXA-1 co-producing *Escherichia coli* ST167 clinical strain isolated from a urine sample. *Journal of Global Antimicrobial Resistance* **14**, 284–286. doi:10.1016/j.jgar.2018.08.005.
- Talaga-ćwiertnia, K., Krzyściak, P. & Bulanda, M. 2017 Do bacteria isolated from ICU patients ‘ESKAPE’ antibiotic treatment? *In vitro* susceptibility of the Enterobacteriaceae family to tigecycline. *Anaesthesiology Intensive Therapy* **49** (3), 210–214. doi:10.5603/AIT.a2017.0034.
- Tellevik, M. G., Blomberg, B., Kommedal, Ø., Maselle, S. Y., Langeland, N. & Moyo, S. J. 2016 High prevalence of faecal carriage of ESBL-producing Enterobacteriaceae among children in Dar es Salaam, Tanzania. *PLoS One* **11** (12), 1–13. doi:10.1371/journal.pone.0168024.
- Titilawo, Y., Sibanda, T., Obi, L. & Okoh, A. 2015 Multiple antibiotic resistance indexing of *Escherichia coli* to identify high-risk sources of faecal contamination of water. *Environmental Science and Pollution Research* **22**, 14. doi:10.1007/s11356-014-3887-3.
- Vinueza-Burgos, C., Ortega-Paredes, D., Narváez, C. & Zurita, J. 2019 Characterization of cefotaxime resistant *Escherichia coli* isolated from broiler farms in Ecuador. *PLoS ONE* **14** (4), e0207567. doi:10.1371/journal.pone.0207567.
- Vounba, P., Rhouma, M., Arsenault, J., Bada Alambédji, R., Fravallo, P. & Morris Fairbrother, J. 2019 Prevalence of colistin resistance and *mcr-1/mcr-2* genes in ESBL/AmpC-producing *E. coli* isolated from chickens in Canada (Quebec), Senegal and Vietnam. *Journal of Global Antimicrobial Resistance* **14** (19), 222–227. doi:10.1016/j.jgar.2019.05.002.
- Wirth, T., Falush, D., Lan, R., Colles, F., Mensa, P., Wieler, L. H., Karch, H., Reeves, P. R., Maiden, M. C., Ochman, H. & Achtman, M. 2006 Sex and virulence in *Escherichia coli*: an evolutionary perspective. *Molecular Microbiology* **60** (5), 1136–1151. doi:10.1111/j.1365-2958.2006.05172.x.
- Zankari, E., Hasman, H., Cosentino, S., Vestergaard, M., Rasmussen, S., Lund, O., Aarestrup, F. M. & Larsen, M. V. 2012 Identification of acquired antimicrobial resistance genes. *Journal of Antimicrobial Chemotherapy* **67** (11), 2640–2644. doi:10.1093/jac/dks261.
- Zahra, R., Javeed, S., Malala, B., Babenko, D. & Toleman, M. A. 2018 Analysis of *Escherichia coli* STs and resistance mechanisms in sewage from Islamabad, Pakistan indicates a difference in *E. coli* carriage types between South Asia and Europe. *Journal of Antimicrobial Chemotherapy* **73** (7), 1781–1785. doi:10.1093/jac/dky109.
- Zurita, J., Alcocer, I., Ortega-Paredes, D., Barba, P., Yauri, F., Iñiguez, D. & Mora, M. 2013 Carbapenem-hydrolysing  $\beta$ -lactamase KPC-2 in *Klebsiella pneumoniae* isolated in Ecuadorian hospitals. *Journal of Global Antimicrobial Resistance* **1** (4), 229–230. doi:10.1016/j.jgar.2013.06.001.
- Zurita, J., Solís, M. B., Ortega-Paredes, D., Barba, P., Paz, Y., Miño, A. & Sevillano, G. 2019 High prevalence of B2-ST131 clonal group among extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli* isolated from bloodstream infections in Quito, Ecuador. *Journal of Global Antimicrobial Resistance* **8** (19), 216–221. doi:10.1016/j.jgar.2019.04.019.