


Inside environmental *Clostridium perfringens* genomes: antibiotic resistance genes, virulence factors and genomic features

Johannes Cornelius Jacobus Fourie, Cornelius Carlos Bezuidenhout , Tomasz Janusz Sanko, Charlotte Mienie and Rasheed Adeleke

ABSTRACT

Until recently, research has focused on *Clostridium perfringens* in clinical settings without considering environmental isolates. In this study, environmental genomes were used to investigate possible antibiotic resistance and the presence of virulence traits in *C. perfringens* strains from raw surface water. *In silico* assembly of three *C. perfringens* strains, DNA generated almost complete genomes setting their length ranging from 3.4 to 3.6 Mbp with GC content of 28.18%. An average of 3,175 open reading frames was identified, with the majority associated with carbohydrate and protein metabolisms. The genomes harboured several antibiotic resistance genes for glycopeptides, macrolide–lincosamide–streptogramin B, β -lactam, trimethoprim, tetracycline and aminoglycosides and also the presence of several genes encoding for polypeptides and multidrug resistance efflux pumps and 35 virulence genes. Some of these encode for haemolysins, sialidase, hyaluronidase, collagenase, perfringolysin O and phospholipase C. All three genomes contained sequences indicating phage, antibiotic resistance and pathogenic islands integration sites. A genomic comparison of these three strains confirmed high similarity and shared core genes with clinical *C. perfringens* strains, highlighting their health security risks. This study provides a genomic insight into the potential pathogenicity of *C. perfringens* present in the environment and emphasises the importance of monitoring this niche in the future.


Key words | antibiotic resistance genes, *Clostridium perfringens*, genetic features, virulence genes, water environment, whole-genome sequencing

Johannes Cornelius Jacobus Fourie
Cornelius Carlos Bezuidenhout 
(corresponding author)
Tomasz Janusz Sanko
Charlotte Mienie
Rasheed Adeleke
Unit for Environmental Science and Management,
North-West University,
Potchefstroom,
South Africa
E-mail: carlos.bezuidenhout@nwu.ac.za

HIGHLIGHTS

- Genome sequences of three environmental *Clostridium perfringens* strains are reported.
- Several antibiotic resistance genes and virulence factors (VFs) were identified within all three genomes. Several VFs associated with other Gram-positive bacteria were also present.
- Genomes showed evidence of prophages and genomic islands (GIs). GIs in each strain contained genes contributing to virulence and drug resistance.
- Comparative genomics showed a close affinity between environmental strains and clinical strains.

GRAPHICAL ABSTRACT



INTRODUCTION

Clostridium perfringens (previously known as *Bacillus welchii*) is a strictly anaerobic, Gram-positive, rod-shaped bacterial pathogen (Hassan *et al.* 2015). This rapidly-growing bacterium has the ability to form endospores that are extremely resilient to toxic chemical and environmental stressors, such as heat or radiation (Paredes-Sabja *et al.* 2008). As a result, *C. perfringens* inhabits diverse environments including soil, water, sewage, food and even the gastrointestinal tract of mammals (Sathish & Swaminathan 2009). In the gastrointestinal tract, *Clostridium perfringens* does not infect healthy cells directly but rather acts by producing toxins and enzymes that cause a number of enteric and systemic diseases in the infected host (Gross *et al.* 1989; Kiu & Hall 2018).

Pathogenic *C. perfringens* strains have been found to produce and secrete more than 20 different extracellular toxins and/or hydrolytic enzymes (Revitt-Mills *et al.* 2015). They are classified into seven groups (toxintypes A–G) according to the combination toxin types they produce (α -toxin, β -toxin, ϵ -toxin and ι -toxin, enterotoxin (CPE) and NetB) (Kiu & Hall 2018). Clinically, the toxinotype A strain is the most common disease-causing agent responsible for gas gangrene (clostridial myonecrosis), necrotic enteritis, as well as mild diarrhoea (Brynstad & Granum 2002). Although these toxins and enzymes each play a specific role in the disease process, it is the synergistic actions of both on the host that could potentially be the key virulence factors involved in its pathophysiology (Rood 1998; Revitt-Mills *et al.* 2015).

Beside virulence factors in *Clostridium perfringens*, another trait such as antibiotic resistance poses a danger to clinically treated patients, mainly due to their ability to render the antibiotic treatment of severe and life-threatening infections ineffective (Kiu & Hall 2018). Therefore, extensive research has been conducted on antibiotic resistance profiles in *C. perfringens* using phenotypic methods (e.g. testing of minimal inhibitory concentration (MIC); Koch *et al.* 1998; Akhi *et al.* 2015). However, there are only a few studies that have used genomic methods (i.e. WGS, whole-genome sequencing) to investigate antibiotic resistance genes (ARGs; Kim *et al.* 2017; Li *et al.* 2017).

It is considering the clinical relevance of *Clostridium perfringens* holds and its wide distribution in nature; the current state of knowledge on the pathogenicity of *C. perfringens* or its non-clinical environment is still poor when compared with other more well-known pathogens (Shimizu *et al.* 2002). New technologies, such as next-generation sequencing (NGS), have become the new gold standard in *in silico* analysis. This is because NGS helps to understand bacterial pathogenesis, identify and characterise genes coding for virulence factors, toxins or antibiotic resistance in pathogens, especially *C. perfringens* strains which are hard to grow under laboratory conditions (Bakour *et al.* 2016). Thus, the aim of this study was to expand on the existing knowledge of *Clostridium perfringens* through the analysis of assembled genomes from surface water and identify the presence of virulence and ARGs.

MATERIAL AND METHODS

Bacterial isolation, antibiotic susceptibility testing and genomic DNA extraction

Clostridium perfringens was isolated from water of the Schoonspruit River (26°40'46.0''S, 26°34'58.7''E) in the North West Province (South Africa) in March 2016. A modified Fung's double tube method, along with tryptose-sulfite-cycloserine agar (Oxoid, UK), was used for the bacterial isolation from collected samples (Fourie *et al.* 2019). The isolates were purified through streaking and grown under anaerobic conditions at 44 °C and subjected to antibiotic susceptibility testing as recommended by the Clinical and Laboratory Standards Institute (CLSI 2016). The MICs of five clinically relevant antibiotics (ampicillin, tetracycline, clindamycin, chloramphenicol and metronidazole) were determined by the agar dilution method. Thereafter, multidrug-resistant (MDR) isolates were selected for genomic DNA extraction. gDNA extraction from overnight cultures was done using the NucleoSpin[®] Tissue kit (Macherey-Nagel, Germany) according to the instructions

provided by the manufacturer. Species confirmation was done by amplifying the 16S rRNA gene, followed by Sanger sequencing of the gene fragment. Subsequently, three MDR *C. perfringens* isolates with high DNA purity were selected for whole-genome sequencing.

Genome sequencing, assembly and annotation

DNA libraries were performed using the Nextera XT kit (Illumina Inc., San Diego, CA) according to the instructions provided by the manufacturer and sequenced using Illumina's MiSeq300 in paired-end reads. All short (<50 bp) or low quality value (QV <15) fragments were removed. The quality assessment and trimming were done in Trimmomatic (v.0.36; Bolger *et al.* 2014), followed by *de novo* assembly in SPAdes (v.3.9.0) to generate scaffolds and contigs (Bankevich *et al.* 2012). Gene prediction and annotation were conducted using the NCBI Prokaryotic Genome Annotation Pipeline (v.4.3) along with the Rapid Annotation using Subsystem Technology (RAST) server (Overbeek *et al.* 2014; Tatusova *et al.* 2016). Subsequently, the obtained draft genome sequences used in this study were deposited into the NCBI database (Fourie *et al.* 2019).

In silico analysis of antibiotic resistance, virulence factors, genomic islands and prophages

ARGs and virulence factors were identified within the genomes of all three *C. perfringens* strains by using BLASTx comparison to the virulence factors database (VFDB) and the ARGs database in deepARG (v.2.0; Chen *et al.* 2016; Arango-Argoty *et al.* 2018). The presence of genomic islands (GIs) within the assembled genomes was predicted using the VRprofile (v.2.0; Li *et al.* 2018). Prophage regions were identified and characterised and located using PHASTER (PHAge Search Tool – Enhanced Release; Arndt *et al.* 2016). Default parameters were used for all software unless otherwise specified.

Genomic comparison

Average nucleotide identity (ANI) was determined by using the OrthoANI algorithm (v.1.4) to analyse genomic relatedness between the three analysed and referenced *C. perfringens*

strains. OrthoANI percentages were calculated, and a UPGMA dendrogram was constructed (Lee *et al.* 2015). Annotated protein sequences were submitted to OrthoVenn2 to identify unique and/or shared orthologous clusters among three *C. perfringens* genomes, as well as three reference genomes (Xu *et al.* 2019). Reference strains used for comparison were *Clostridium perfringens*: ATCC 13124 (Genbank: NC_008261), Str 13 (Genbank: BA000016) and FORC_025 (Genbank: NZ_CP013101).

RESULTS AND DISCUSSION

Genomic annotations

The three assembled *Clostridium perfringens* strains (SC4-C13, SC4-C17 and SC4-C24) that were originally obtained from surface water produced genome sizes of 3.6, 3.5 and 3.4 Mbp, respectively (Table S1) and were organised in 205 (both SC4-C13 and SC4-C17) and 110 (SC4-C24) contigs (Fourie *et al.* 2019). The GC content between the three strains averaged at around 28.1%. Annotation of the genomes indicated the presence of open reading frames (ORFs) ranging between 3,079 and 3,245 for all three strains (Table S1), which is consistent with published data on this bacteria. The average genome size known for *C. perfringens* varies from 3.0 to 4.1 Mbp, with a GC content ranging between 27 and 28% (Kiu *et al.* 2017; Kiu & Hall 2018). The number of coding sequences present in their genome averaged between 2,500 and 3,600 (Kiu & Hall 2018).

RAST was used to functionally annotate the open reading frames of each genome into 25 subsystem categories (Figure 1). Based on the categories generated, all three strains showed similar amounts of assigned ORFs for each category. The majority was allocated to 'carbohydrates', thus showing the importance of carbohydrates for these pathogens, since they play a vital role in their growth and sporulation (Sacks 1983). This was then followed by 'protein metabolism', 'amino acids and derivatives' and 'cofactors, vitamins, prosthetic groups and pigments'. The less abundant ORFs belonged to the 'metabolism of aromatic compounds' category, along with 'motility and chemotaxis', 'transposable elements' and 'nitrogen metabolism'.




Figure 1 | ORFs from three analysed *C. perfringens* genomes (blue SC4-C13; orange SC4-C17 and grey SC4-C24) annotated and sorted into subsystems by the RAST together with counts per each category for each genome. Please refer to the online version of this paper to see this figure in colour: <http://dx.doi.org/10.2166/wh.2020.029>.

Antibiotic resistance

The deepARG tool applies a deep-learning approach to identifying and annotating ARGs from genomic data. It does this by creating a dissimilarity matrix based on ARG categories and comparing it against the three major datasets currently available, namely Comprehensive Antibiotic Resistance Database (CARD), Antibiotic Resistance Genes Database (ARDB) and Universal Protein Resource (UNIPROT; Arango-Argoty *et al.* 2018). This approach identified a total of 11 ARGs in the genomes of the three *Clostridium perfringens* strains (Table 1). Strains SC4-C13 and SC4-C17 both possess genes that encode for resistance against seven different classes of antibiotics (glycopeptide, β -lactamase, macrolide-lincosamide-streptogramin B (MLS_B), tetracycline, trimethoprim,

kasugamycin and bacitracin), whereas strain SC4-C24 only harboured resistance genes against six classes, with the absence of the β -lactam resistance gene (*bla*₂). Furthermore, all three strains hold two or more genes that encode for multi-drug resistance.

β -lactam antibiotics are the most commonly used class of antibiotics in human health medicine and agriculture (Price *et al.* 2019). The prolonged misuse of these antibiotics have resulted in the more frequent occurrence of β -lactam resistance in bacteria. This, in turn, has diminished its effectiveness in fighting against bacterial infections that were once easily treated. The bacterial mechanism of β -lactamase resistance is primarily due to the production of β -lactamase enzymes that hydrolyse the β -lactam ring, thereby inactivating the antibiotic (Bush &

Table 1 | ARGs predicted in three *C. perfringens* strains (SC4-C13; SC4-C17 and SC4-C23) using DeepARG. '+' marks if the gene was present in the genome, while '-' marks the absence of the gene in the genome

Antibiotic			SC4-C13	SC4-C17	SC4-C24
Class	Name	Gene			
Glycopeptide	Vancomycin	<i>vanHD</i>	-	-	+
		<i>vanTrL</i>	-	-	+
		<i>vanRB</i>	-	+	-
		<i>vanRG</i>	+	+	+
		<i>vanRI</i>	+	+	+
β-Lactam	Penicillin	<i>bla2</i>	+	+	-
MLSB ^a	Erythromycin	<i>ermQ</i>	+	+	+
Tetracyclines	Tetracycline	<i>tetB(P)</i>	+	+	+
Trimethoprim	Trimethoprim	<i>dfrK</i>	+	+	+
Aminoglycosides	Kasugamycin	<i>ksgA</i>	+	+	+
Polypeptides	Bacitracin	<i>bacA</i>	+	+	+
Multidrug	System regulators ATP-binding cassette transporters Efflux pump	<i>arIR</i>	+	+	+
		<i>vgaB</i>	+	+	-
		<i>mepA</i>	+	+	+

^aMacrolide-lincosamide-streptogramin B.

Bradford 2016). The presence of the *bla2* gene which encodes for β-lactamase in both SC4-C13 and SC4-C17 is worrisome, since most β-lactam resistance is not commonly associated with *Clostridium* species, with only a few reported incidences of resistance in clinical settings (Finegold et al. 2005; Mishra et al. 2016). Based on the UNIPROT database, the *bla2* gene was originally identified in *Firmicutes* bacterium CAG:212, which could suggest that the two *C. perfringens* strains acquired this gene from an external source and incorporated it into their genomes.

Glycopeptides encompass a group of antibiotics that are vital in combating infections caused by antibiotic-resistant bacteria, with vancomycin still the first choice of treatment against Gram-positive bacteria since its introduction 50 years ago (Blaskovich et al. 2018). Vancomycin works by binding to the D-Ala-D-Ala C-terminal of the growing pentapeptide chain during cell wall synthesis and inhibits further elongation and cross-linking of the peptidoglycan chain blocking through transglycosylation and transpeptidation (Sanakal & Kaliwal 2011). However, resistance to this antibiotic has become a common occurrence, especially in vancomycin-resistant enterococci and even in methicillin-

resistant *Staphylococcus aureus* (Willems et al. 2005). The resistance in these bacteria has been confirmed by the presence of several vancomycin resistance genes (*van*). Our strains possessed many of these genes (*vanHD*, *vanTrL*, *vanRB*, *vanRG* and/or *vanRI*), which encode for enzymes that produce synthetic precursors that replace the D-Ala-D-Ala C-terminal, altering the vancomycin-binding site (Sanakal & Kaliwal 2011). Quite a few studies have reported vancomycin resistance genes in *Clostridium* species (Peltier et al. 2013; Chia et al. 2017). However, *Clostridium perfringens* has not been one of the few to show such resistance since they are normally susceptible to vancomycin (Citron et al. 2005; Camacho et al. 2008; Chia et al. 2017). The presence of these genes could suggest a new resistance trait in these species if they are expressed.

A further study of the results presented in Table 1 shows that all three *C. perfringens* genomes harboured the same gene that encodes resistance for MLSB (*ermQ*), tetracycline (*tetB(P)*), trimethoprim (*dfrK*), kasugamycin (*ksgA*) and bacitracin (*bacA*). MLSB antibiotics work by inhibiting protein synthesis in both Gram-positive and Gram-negative bacteria. Resistance to this antibiotic is due to 23S rRNA methylation (encoded by *erm* genes) that prevents MLSB antibiotics from binding to the ribosome. Several *erm* genes have been detected in *Clostridium* species, such as *C. perfringens* and *C. difficile* (Spigaglia & Mastrantonio 2002). However, the *ermQ* gene was first identified in *C. perfringens* and found to be a distinct class of MLSB resistance determinant (Berryman et al. 1994; Shoemaker et al. 2001).

Tetracyclines are broad-spectrum antibiotics and the second most used antibiotics group after β-lactams. Bacterial resistance to tetracycline is achieved by harbouring one or more of the 36 known *tet* genes (Sheykhsaran et al. 2019). These *tet* genes follow one of three resistance mechanisms: energy-dependent efflux pumps, ribosomal protection proteins, or drug target alteration and enzymatic inactivation (van Hoek et al. 2011). The *tetB(P)* genes, found in our strains along with *tetA(P)* (not present), are originally two overlapping genes within the Tet P determinant of *C. perfringens* (Sloan et al. 1994). However, because the Tet P determinant has only been found in *Clostridium* species and is often associated with conjugative and non-conjugative plasmids, it has shown the capacity to spread rapidly across the whole genus (Roberts 2011; Vidor et al. 2019).

Clostridium species are known to show endogenous resistance to trimethoprim (Huovinen *et al.* 1995). However, the presence of the *drfK* gene in the three *C. perfringens* genomes suggests that it could have been acquired from an external source since it has mainly been associated with enterococci and staphylococci species (López *et al.* 2012). While the *drfK* gene may not be vital for trimethoprim resistance in *Clostridium*, its presence may have implications for further dissemination in the environment, or to other bacteria that are primarily susceptible to trimethoprim.

Antibiotics such as kasugamycin and bacitracin are widely used in agriculture and animal husbandry as growth promoters, with bacitracin also being used as a topical ointment for skin infections (Duffin & Seifert 2009; Charlebois *et al.* 2012). Resistance to the aminoglycoside antibiotic kasugamycin is due to the presence of the *KsgA* gene, which produces a predicted dimethyltransferase, whereas an ATP-dependent ABC-type efflux system consisting of three BcrA, -B and -C proteins is responsible for bacitracin resistance (El Ghachi *et al.* 2004; Duffin & Seifert 2009). Studies have shown that the use of antibiotics in agriculture and growth promoters in animal breeding have a direct correlation to the development and dissemination of antibiotic-resistant bacteria and genes into the surrounding water sources. This happens by means of surface runoff, where antibiotic resistance can be further intensified via horizontal gene transfer (HGT; Kümmerer 2004).

Genes for efflux pumps (*mepA*), ATP-binding cassette transporters (*vgaB*) and system regulators (*arlR*), which are associated with multidrug resistance, were found in our genomes as well. The *arlR* is part of a two-component regulatory system, a transmembrane sensor and its associated response regulator (*arlS-arlR*). This regulator was found to increase the expression of efflux pumps associated with *Staphylococcus aureus* (Fournier *et al.* 2000). The *vgaB* gene is one of the ATP transporter genes and can increase the level of resistance to pristinamycin, a mixture of streptogramin A and streptogramin B compounds via efflux pumps (Roberts 2002; Chesneau *et al.* 2005). The *mepA* encodes for a multidrug resistance efflux pump in *S. aureus*, which has reported low-level resistance to quaternary ammonium compounds, such as chlorhexidine, pentamidine and tigecycline (Costa *et al.* 2013).

When comparing the genotypic characteristics with the phenotypic profiles (Table S2 in Supplementary Materials)

of these three strains, some conclusions can be drawn. All the strains were shown to be susceptible to chloramphenicol, along with the absence of any associated resistance genes to support this. Two of the strains, SC4-C13 and SC4-C17, showed phenotypical resistance to ampicillin (β -lactam) and possessed the *bla2* gene, whereas SC4-C24 showed intermediate resistance with no genes to confirm resistance in its genome. Mobile genetic elements such as plasmids have been shown to carry β -lactam resistance genes (Partridge *et al.* 2018). If such plasmid-mediated resistance genes were to be present in *C. perfringens* and expressed, it could translate to phenotypical resistance to β -lactam antibiotics. Alternatively, this low level of resistance might also suggest an increase in the CLSI-defined susceptibility breakpoint for ampicillin in *C. perfringens*. Noticeably, all three strains also showed phenotypic resistance to tetracycline, clindamycin and metronidazole. The presence of genes such as *ermQ* and *tetB(P)* could explain the phenotypical resistance to clindamycin and tetracycline, respectively. However, no genes associated with metronidazole were identified in any of the assembled genomes. Previous studies have reported phenotypic resistance to metronidazole in *C. perfringens* (Tansuphasiri *et al.* 2005; Akhi *et al.* 2015). However, the understanding of its resistance mechanism in *Clostridium* species is still unclear (Lynch *et al.* 2013). It is also important to add that although the assembly delivered almost near genomes, the lack of genetic confirmation for phenotypic resistance may be due to incomplete gene reconstruction of the region where the gene occurs.

Virulence factors

The genes that are associated with virulence in *Clostridium perfringens* strains were analysed using the VFDB as a reference. A total of 35 different virulence genes were successfully annotated with this database, of which 79% are typically associated with *Clostridium perfringens* strains. These include genes that are responsible for adherence, such as fibronectin-binding protein, chaperonin GroEL and type IV pili, as well as the two members of the double-component VirR/VirS regulon (sensor histidine kinase and DNA-binding response regulator). There are also 19 genes present in all three strains that are capable of producing eight different toxins, namely α -clostripain,

phospholipase C, collagenase, hyaluronidase, perfringolysin O, sialidase and haemolysin (Table 2). Interestingly, the remaining 21% of virulence genes are usually associated with other Gram-positive bacteria, namely *Listeria ivanovii*, *Streptococcus pyogenes*, *Mycoplasma penetrans* and *Enterococcus faecium*. This might imply their potential influence or contribution to the expanding pathogenicity in these three strains as bile-resistance and adherence factors (Table 2). Among the three *C. perfringens* strains described in this study, SC4-C13 and SC4-C17 only possess 94.7% of the above-mentioned virulence factors, while for SC4-C24 it is 100%.

Clostridium perfringens can produce a variety of more than 20 different extracellular toxins and hydrolytic enzymes, giving it the ability to cause various histotoxic infections in humans and animals (Kiu & Hall 2018). However, not all strains are able to produce all these toxins. This is mainly due to some toxins being strain-specific (Kiu et al. 2017). The *C. perfringens* strains in this study possess genes able to code a vast array of toxins and enzymes. However, only a combination of several typing toxins is traditionally used to determine the toxinotype of a species. All three strains were identified as toxinotype A due to the presence of the *plc* gene, which encodes for α -toxin and

Table 2 | Virulence factors identified in *C. perfringens* strains based on the VFDB

Virulence factor

Class	Name	Gene	Reference species	Genbank	
Adherence	Fibronectin-binding protein	<i>fbp</i>	*	WP_011590967	
		<i>fbpA</i>	*	WP_011010006	
	Chaperonin GroEL	<i>groEL</i>	*	WP_003462314	
	Elongation Factor Tu	<i>tuf</i>	**	WP_011076858	
	Enterococcal surface protein Type IV pili	<i>esp</i>	***	WP_014387145	
		<i>pilA1</i>	*	WP_011010863	
		<i>pilB</i>	*	WP_011010636	
		<i>pilB2</i>	*	WP_011010862	
		<i>pilC</i>	*	WP_011010635	
		<i>pilC2</i>	*	WP_01101086	
		<i>pilD</i>	*	WP_003462279	
		<i>pilM</i>	*	WP_011010859	
	Regulation	Streptococcal plasmin receptor/GAPDH	<i>pilN</i>	*	WP_011010858
			<i>pilT</i>	*	WP_003451114
Sensor histidine kinase DNA-binding response regulator		<i>plr/gapA</i>	+	WP_002986042	
		<i>virS</i>	*	WP_011590863	
		<i>virR</i>	*	WP_003449818	
		Toxin	α -Clostripain	<i>cloSI</i>	*
Phospholipase C (α -toxin)	<i>plc</i>		*	WP_011590041	
κ -Toxin (collagenase)	<i>colA</i>		*	YP_697499	
μ -Toxin (hyaluronidase)	<i>nagH</i>		*	NP_561107	
	<i>nagJ</i>		*	NP_562150	
	<i>nagK</i>		*	NP_562195	
	<i>nagL</i>		*	NP562439	
	Perfringolysin O (θ -toxin/PFO)		<i>pfoA</i>	*	NP_561079
Sialidase	<i>nanH</i>		*	YP_695432	
	<i>nanI</i>		*	WP_011590331	
	<i>nanJ</i>		*	NP_561469	
	Haemolysin		<i>hlyB</i>	*	NP_561353
			<i>hlyC</i>	*	WP_003454634
<i>hlyD</i>		*	NP_562734		
<i>hlyE</i>		*	WP_011010677		
Bile resistance	Bile-salt hydrolase	<i>bsh</i>	++	YP_004855791	

* *Clostridium perfringens*; ** *Mycoplasma penetrans*; *** *Enterococcus faecium*; + *Streptococcus pyogenes*; ++ *Listeria ivanovii* subsp. *Ivanovii*.

the absence of the other typing toxin genes. Furthermore, *C. perfringens* toxinotype A strains are known to be human pathogens, causing diseases such as gas gangrene (clostridial myonecrosis), necrotic enteritis, as well as mild diarrhoea (Brynstad & Granum 2002). The presence of perfringolysin O (θ -toxin) has also shown to have a synergistic effect with α -toxin in the pathology of gas gangrene. Even on its own, this cholesterol-dependent cytolysin can form pores on cell membranes and lyse red blood cells, further highlighting the significant role perfringolysin O has in disease development (Awad *et al.* 2001; Kiu & Hall 2018). However, these two toxins, along with the κ -toxin gene (*colA*), are tightly regulated by specific regulatory systems in *C. perfringens* (Ohtani & Shimizu 2016). The VirS/VirR two-component system is one of the most important systems and consists of two genes: the response regulator (*virR*) and the sensor histidine kinase (*virS*; Ma *et al.* 2011). Still, this system assumes responsibility for coordinating the pathogenicity of *C. perfringens* type A strains.

The hydrolytic enzymes produced by *C. perfringens* also play an important role during *C. perfringens*-mediated tissue infection. Sialidase is encoded by several genes that contribute to internal virulence by improving the adherence of *C. perfringens* to host cells, enhancing the production, binding and activity of certain toxins that are responsible for intestinal infections, and contributes to intestinal colonisation by *C. perfringens* (Li *et al.* 2016). Mu-toxins (hyaluronidase), on the other hand, are enzymes that facilitate the breakdown of hyaluronate substrates, improving contact between *C. perfringens* and the specific cell wall receptors. It has also been found to promote the spread of α -toxin and, in doing so, potentiate its cytolytic activity (Hynes & Walton 2000).

As in most pathogens, adherence plays an essential role in the pathogenesis of *Clostridium perfringens*. Several adherence genes were identified in the genomes of *C. perfringens*, each of which encodes for a different approach of adherence and promotes colonisation. Fibronectin-binding protein (*fbpA*), for instance, makes it possible for *C. perfringens* to bind to fibronectin found in all human tissues and organs, including intestinal epithelial cells, and in doing so, invade the host cells (Katayama *et al.* 2009). Whereas type IV pili are elongated, flexible filaments extend from the bacterial cell surface and are implicated in attaching to and invading host cells, attachment to abiotic

surfaces, biofilm formation and bacteriophage susceptibility (Rodgers *et al.* 2011).

Genomic islands

Genomic islands (GIs) are regions of the bacterial genome that are acquired through HGT (Dobrindt *et al.* 2004; Klein *et al.* 2018). Antibiotic resistance islands (ARIs) and pathogenicity islands (PAIs) are two subgroups of GIs that aid and contribute to the pathogenesis of an organism (Yoon *et al.* 2015). Figure 2(a) shows the distribution of ARIs and PAIs present in the three *Clostridium perfringens* strains. All strains analysed possessed PAIs and ARIs homologous to those found in most enteropathogens. A total of 65 ARIs were identified in all three strains that were related to genes from *Staphylococcus aureus* (24/65), *Acinetobacter baumannii* (19/65), *Pseudomonas aeruginosa* (10/65), *Campylobacter coli* (3/65), *Shigella flexneri* (3/65), *Klebsiella pneumoniae* (2/65), *Proteus mirabilis* (2/65), *Escherichia coli* O157H7 (1/65) and *Staphylococcus hominis* (1/65) (Table S3). This included genes which encode for putative enzymes, such as acyl-CoA dehydrogenase, DNA topoisomerase III and enoyl-CoA hydratase/isomerase, as well as regulatory proteins such as KDP operon transcriptional regulatory protein (*KdpE*), propionate catabolism operon regulator (*prpR*), putative transcriptional and response (*ArmR*) regulators, to name a few. Genes encoding for proteins and enzymes contributing to antibiotic resistance (*TetA* class A, truncated tetracycline-resistant protein and streptothricin acetyltransferase) and tolerance to heavy metals (mercuric reductase and cadmium efflux system proteins B and C) were also identified. Heavy metals such as mercury and cadmium are commonly used for antimicrobial metal compounds in healthcare (Pal *et al.* 2017). However, the presence of genes encoding for resistance to antibiotics and heavy metal might suggest possible adverse implications for infectious cases of *C. perfringens*. Iron plays a key role in the growth and survival of many pathogens in any niche. It is utilised for important biological processes such as DNA synthesis, generating energy and protection against reactive oxygen species (Choo *et al.* 2016). All the *C. perfringens* strains possess ARIs related to *Shigella flexneri* that encodes for proteins responsible for the binding and transport of ferric




Figure 2 | Venn diagrams showing the ARIs (a) and PAIs (b) identified in the genomes of the three *C. perfringens* strains.

iron (III) (*FecC*, *FecD* and *FecE*; Table S3). Genes associated with iron acquisition have previously been linked to GIs in *Clostridium perfringens* (Myers et al. 2006).

The distribution of all PAIs identified among the *C. perfringens* strains are visualised in Figure 2(b). The various PAIs shared among the three strains ranged from hypothetical or unnamed proteins related to *Yersinia pestis*, *Escherichia coli*, *Edwardsiella tarda* and *Streptomyces lividans* (Table S4). Other examples include PAIs encoding for *Epd*, integrase, Orf17, Orf29, Orf34, Orf41, Orf59, *Pgk* and *TktA* from *Photothabdus luminescens*; dihydrofolate reductase type Ib (plasmid encoded), dihydropteroate synthase, *GlcD* protein, *GspE* type II secretion protein, haemolysin B, phosphoglycerate transport protein, putative ABC transporter ATP-binding and membrane proteins, putative lysyl-tRNA synthetase (*LysU*) and short-chain dehydrogenase from *Escherichia coli*; ATPases with chaperone activity ATP-binding subunit, serine/threonine protein phosphatase and transcriptional regulator from *Vibrio alginolyticus* and glutamate racemase from *Helicobacter pylori*. Additionally, sequences similar to ones encoding for *HrcN* and *HrpB6* in *Xanthomonas* species were also present. These two genes encode for the type III secretion system (TTSS) proteins, a requirement for the pathogenicity of several plant pathogens (Villela et al. 2019).

Among the three *Clostridium perfringens* strains, SC4-C13 and SC4-17 had the most unique PAIs and ARIs

(Tables S3 and S4). Noticeably, both these strains harboured genes encoding for putative transposase. Strain SC4-C13 possessed multiple transposases for IS285 (*Y. pseudotuberculosis*), ISSau4-like and IS256/Tn4001 (*Staphylococcus aureus*), whereas SC4-C17 harboured a transposase protein for IS30 (*Staphylococcus aureus*). Additionally, all strains possessed two or more GIs containing integrase genes from *Photothabdus luminescens*, *Acinetobacter baumannii* or *Staphylococcus aureus*. The presence of genes associated with transposases and integrases has shown to infer high potential for genetic gain or loss in *C. perfringens* genomes (Kiu et al. 2017).

A previous study identified over 300 GIs in the genomes of several clinical *C. perfringens* type A strains, of which almost all were chromosomally encoded (Myers et al. 2006). Less is known about the occurrence of GIs in strains isolated from environmental sources (Klein et al. 2018). Pathogens such as *C. perfringens* are found in various ecological niches, which also include diverse microbial communities that can contribute to HGT events (Kiu et al. 2017).

Prophages

The Phage Search Tool Enhanced Release (PHASTER) program identifies and characterises prophage regions into three categories based on the completeness scores, namely intact (>90), questionable (70–90) or incomplete (<70). All

three *Clostridium perfringens* strains used in this study contained two or more prophage regions in their genome (Table 3). Strain SC4-C24 were shown to harbour three prophage regions, of which one was confirmed to be intact, and the other two showed incomplete and questionable phage-related sequences. Another intact prophage was identified in strain SC4-C17, along with an additional incomplete region. Strain SC4-C13 showed putative fragments of two phages, one of which scored relatively high (90). Each prophage region varies in size, the number of CDS and GC content. The two intact prophages in SC4-C17 and SC4-C24 strains have sizes of 49.6 and 42.2 Kbp, respectively, with both containing the second-highest number of CDS (57). Functional annotation of the genes present in these prophage regions showed protein sequences encoding for phage-related elements such as integrase, terminase, portal, capsid, head, tail and/or transposase (Table S5). Phages play a very important role in the evolution of bacterial genomes (Ramisetty & Sudhakari 2019). Their interactions with each other have impacted both their survival and persistence. This is because prophages have several advantages to offer in the way of inversion, deletion and insertion via HGT of genetic material (Darmon & Leach 2014). This is because prophages are known to be ‘hotspots’ for carrying significant genes such as those involved in virulence, antibiotic resistance and metabolic pathways (Ramisetty & Sudhakari 2019). This led to further investigation into functional protein sequences in the identified prophage regions of the three *C. perfringens* genomes. However, only the presence of hypothetical proteins was discovered in these regions and could suggest that these genes encode for unknown functions from remote sources.

Genomic comparisons

The average nucleotide identity for all *Clostridium perfringens* genomes was successfully calculated using OrthoANI (Figure 3). ANI analyses of the three strains revealed them to be almost identical to each other as indicated by the ANI of >99% between the strains. Although the differences between the three strains appear negligible, based on the dendrogram, strains SC4-13 and SC4-C17 share relatively more similarities between them, as opposed to strain SC4-24 (Figure 3). When comparing the three *C. perfringens* genomes with that of the reference genomes (ATCC 13124, Str 13 and FORC_025), another high similarity was observed among the genomes, with our genomes sharing ANI values of 98.49–98.63% with the reference genomes. The dendrogram also showed that the three strains were clustered more closely to FORC_025, followed by the ATCC 13124 strain which was derived from clinical environments (Myers et al. 2006; Kiu et al. 2017), whereas only strain 13 was originally isolated from a natural environment (soil; Shimizu et al. 2002).

Orthologous genes originate from a single gene present in the last common ancestor through a series of speciation events and usually still have the same biological function in the present-day organism (Xu et al. 2019). Therefore, for further analysis of the three *C. perfringens* strains, a multi-species comparison of the shared core orthologous gene clusters was performed without (Figure 4(a)) and with reference genomes (Figure 4(b)). Results show a large number of (2,779) core orthologous genes shared between the three *C. perfringens* strains (Figure 4(a)). These clusters were all revealed to be associated with biological

Table 3 | Prophages regions identified in *C. perfringens* strains using PHASTER

Strain	Size (kb)	CDS	GC content (%)	Completeness	Score	Phage	
						Name	Genbank
SC4-C13	54.4	53	27.75	+ / –	90	Clostr_vB_CpeS_CP51	NC_021325
	6.1	14	33.54	–	30	Clostr_c_st	NC_007581
SC4-C17	43.6	40	28.10	–	50	Clostr_PhiS63	NC_017978
	49.6	57	28.10	+	100	Clostr_vB_CpeS_CP51	NC_021325
SC4-C24	40.6	59	28.41	+ / –	90	Clostr_vB_CpeS_CP51	NC_021325
	42.2	57	28.26	+	104	Clostr_phiSM101	NC_008265
	7.6	9	30.11	–	40	Sphing_PAU	NC_019521

+ Complete; – Incomplete; + / – Questionable.




Figure 3 | Dendrogram of *C. perfringens* strains relatedness calculated with the OrthoANI algorithm, clustered using UPGMA, and shown with the corresponding pairwise identity heat map.




Figure 4 | Venn diagram showing the core genome of (a) the three *C. perfringens* strains (SC4-C13, SC4-C17 and SC4-C24) and (b) the three *C. perfringens* strains and reference strains (ATCC 13124, FORC_025 and Str.3). The numbers represent the orthologous protein clusters shared by corresponding genomes.

processes, molecular functions and cellular components within their core genomes (Table S6). The orthologous analysis also supported the higher similarity between SC4-C13 and SC4-C17 based on them sharing over 130 orthologous clusters, while SC4-C24 has the highest amount of unique orthologous clusters of the three strains. When comparing the core orthologous clusters of the three strains with those of the reference strains, they were shown to share 2,300 clusters (Figure 4(b)). Altogether, only eight orthologous clusters were identified as unique among all the genomes. Interestingly, these clusters were found in strains SC4-C13 and SC4-C24. Strain SC4-C13 had two of the eight unique clusters, one of which was responsible for glycosyltransferase and the other for the biological response (movement, secretion, enzyme production or gene expression) to external cold stimuli of the cell. Strain SC4-C24 possessed the other six unique clusters, three clusters were responsible for the initiation of DNA-template transcription, hydrogen peroxide catabolic and antigen biosynthetic processes, while the other three clusters did not have assigned functions. Although only two of the three *C. perfringens* strains showed unique orthologous clusters when compared with the reference genomes, this may be attributed to the possibility of isolates niche adaptation (Braga *et al.* 2016).

CONCLUSION

The current study has shown that *Clostridium perfringens* strains from a water environment possess similar genomic features to published clinical *C. perfringens*. The *in silico* analysis of the three genomes revealed putative genes encoding for virulence factors normally responsible for pathogen adherence to host cells, production of toxins and extracellular enzymes and the presence of key system regulators managing the pathogenicity of *C. perfringens*. The strains were also classified as toxinotype A, which commonly consists of human pathogens only. Furthermore, the presence of ARGs encoding for several classes of antibiotics and multidrug resistance efflux pumps could contribute to the emergence of pathogenic multidrug-resistant *C. perfringens* strains in this environment. Interestingly, the presence of several genes normally associated with other pathogenic genera, as well as a few intact prophages and genomic

islands (ARIs and PAIs), might suggest the transfer of genetic material between bacteria through HGT within the water reservoir, thus indirectly implicating the presence of toxic substances in the water (cadmium, copper, mercury etc.), which are associated with HGT in bacteria. Moreover, expression analyses should be the next step of the investigation. Furthermore, comparative genomics showed great similarities between the environmental *C. perfringens* strains with those of clinical origin, therefore highlighting the importance of monitoring these bacteria for effective epidemiological surveillance purposes. Genomic studies on *Clostridium perfringens* in natural environments are currently very limited and should be explored further.

ACKNOWLEDGEMENTS

This work is based on research supported in part by the Water Research Commission (WRC) of South Africa (contract no. K5/2347//3), National Research Foundation (NRF) of South Africa (UID no. 109207), some NWU funding. We also thank the Centre for High Performance Computing (programme no. CBB10890) in Cape Town (South Africa) for access to computing resources. Views expressed are those of the authors and not of the funders.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this paper is available online at <https://dx.doi.org/10.2166/wh.2020.029>.

REFERENCES

- Akhi, M. T., Asl, S. B., Pirzadeh, T., Naghili, B., Yeganeh, F., Memar, Y. & Mohammadzadeh, Y. 2015 Antibiotic sensitivity of *Clostridium perfringens* isolated from faeces in Tabriz, Iran. *Jundishapur Journal of Microbiology* 8 (7), 6–9. <https://doi.org/10.5812/jjm.20863v2>.
- Arango-Argoty, G., Garner, E., Pruden, A., Heath, L. S., Vikesland, P. & Zhang, L. 2018 DeepARG: a deep learning approach for predicting antibiotic resistance genes from metagenomic data. *Microbiome* 6 (1), 1–15. <https://doi.org/10.1186/s40168-018-0401-z>.
- Arndt, D., Grant, J. R., Marcu, A., Sajed, T., Pon, A., Liang, Y. & Wishart, D. S. 2016 PHASTER: a better, faster version of the

- PHAST phage search tool. *Nucleic Acids Research* **44** (W1), W16–W21. <https://doi.org/10.1093/nar/gkw387>.
- Awad, M. M., Ellemor, D. M., Boyd, R. L., Emmins, J. J. & Rood, J. I. 2001 Synergistic effects of alpha-toxin and perfringolysin O in *Clostridium perfringens*-mediated gas gangrene. *Infection and Immunity* **69** (12), 7904–7910. <https://doi.org/10.1128/IAI.69.12.7904-7910.2001>.
- Bakour, S., Sankar, S. A., Rathored, J., Biagini, P., Raoult, D. & Fournier, P. E. 2016 Identification of virulence factors and antibiotic resistance markers using bacterial genomics. *Future Microbiology* **11** (3), 455–466. <https://doi.org/10.2217/fmb.15.149>.
- Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., Lesin, V. M., Nikolenko, S. I., Pham, S., Prjibelski, A. D., Pyshkin, A. V., Sirotkin, A. V., Vyahhi, N., Tesler, G., Alekseyev, M. A. & Pevzner, P. A. 2012 SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology* **19**, 455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Berryman, D. I., Lyristis, M. & Rood, J. I. 1994 Cloning and sequence analysis of ermQ, the predominant macrolide-lincosamide-streptogramin B resistance gene in *Clostridium perfringens*. *Antimicrobial Agents and Chemotherapy* **38** (5), 1041–1046. <https://doi.org/10.1128/AAC.38.5.1041>.
- Blaskovich, M. A. T., Hansford, K. A., Butler, M. S., Jia, Z., Mark, A. E. & Cooper, M. A. 2018 Developments in glycopeptide antibiotics. *ACS Infectious Diseases* **4** (5), 715–735. <https://doi.org/10.1021/acsinfecdis.7b00258>.
- Bolger, A. M., Lohse, M. & Usadel, B. 2014 Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Braga, R. M., Dourado, M. N. & Araújo, W. L. 2016 Microbial interactions: ecology in a molecular perspective. *Brazilian Journal of Microbiology* **47**, 86–98. <https://doi.org/10.1016/j.bjm.2016.10.005>.
- Brynstad, S. & Granum, P. E. 2002 *Clostridium perfringens* and foodborne infections. *International Journal of Food Microbiology* **74** (3), 195–202. [https://doi.org/10.1016/S0168-1605\(01\)00680-8](https://doi.org/10.1016/S0168-1605(01)00680-8).
- Bush, K. & Bradford, P. A. 2016 β -Lactams and β -lactamase inhibitors: an overview. *Cold Spring Harbor Perspectives in Medicine* **6**, a025247. <https://doi.org/10.1101/cshperspect.a025247>.
- Camacho, N., Espinoza, C., Rodríguez, C. & Rodríguez, E. 2008 Isolates of *Clostridium perfringens* recovered from Costa Rican patients with antibiotic-associated diarrhoea are mostly enterotoxin-negative and susceptible to first-choice antimicrobials. *Journal of Medical Microbiology* **57** (3), 343–347. <https://doi.org/10.1099/jmm.0.47505-0>.
- Charlebois, A., Jalbert, L. A., Harel, J., Masson, L. & Archambault, M. 2012 Characterization of genes encoding for acquired bacitracin resistance in *Clostridium perfringens*. *PLoS ONE* **7** (9). <https://doi.org/10.1371/journal.pone.0044449>
- Chen, L., Zheng, D., Liu, B., Yang, J. & Jin, Q. 2016 VFDB 2016: hierarchical and refined dataset for big data analysis—10 years on. *Nucleic Acids Research* **44**, 694–697. <https://doi.org/10.1093/nar/gkv1239>.
- Chesneau, O., Ligeret, H., Hosan-Aghaie, N., Morvan, A. & Dassa, E. 2005 Molecular analysis of resistance to streptogramin A compounds conferred by the Vga proteins of staphylococci. *Antimicrobial Agents and Chemotherapy* **49** (3), 973–980. <https://doi.org/10.1128/AAC.49.3.973-980.2005>.
- Chia, J. H., Su, L. H., Wu, T. L., Chia, J. H., Su, L. H., Wu, T. L. & Chiu, C. H. 2017 *Clostridium innocuum* is a significant vancomycin-resistant pathogen for extraintestinal clostridial infection. *Clinical Microbiology and Infection* **23** (8), 560–566. <https://doi.org/10.1016/j.cmi.2017.02.025>.
- Choo, J. M., Cheung, J. K., Wisniewski, J. A., Steer, D. L., Bulach, D. M., Hiscox, T. J., Chakravorty, A., Smith, A. I., Gell, D. A., Rood, J. I. & Awad, M. M. 2016 The NEAT domain-containing proteins of *Clostridium perfringens* bind heme. *PLoS ONE* **11** (9), e0162981. <https://doi.org/10.1371/journal.pone.0162981>.
- Citron, D. M., Kwok, Y. Y. & Appleman, M. D. 2005 In vitro activity of oritavancin (LY333328), vancomycin, clindamycin, and metronidazole against *Clostridium perfringens*, *Propionibacterium acnes*, and anaerobic Gram-positive cocci. *Anaerobe* **11** (1–2), 93–95. <https://doi.org/10.1016/j.anaerobe.2004.10.005>.
- Clinical and Laboratory Standards Institute (CLSI) 2016 *Performance Standards for Antimicrobial Susceptibility Testing*, 26th edn. CLSI supplement M100S. Clinical and Laboratory Standards Institute, Wayne, PA.
- Costa, S. S., Viveiros, M., Amaral, L. & Couto, I. 2013 Multidrug efflux pumps in *Staphylococcus aureus*: an update. *The Open Microbiology Journal* **7**, 59–71. <https://doi.org/10.2174/1874285801307010059>.
- Darmon, E. & Leach, D. R. F. 2014 Bacterial genome instability. *Microbiology and Molecular Biology Reviews* **78** (1), 1–39. <https://doi.org/10.1128/mmr.00035-13>.
- Dobrindt, U., Hochhut, B., Hentschel, U. & Hacker, J. 2004 Genomic islands in pathogenic and environmental microorganisms. *Nature Reviews Microbiology* **2** (5), 414–424. <https://doi.org/10.1038/nrmicro884>.
- Duffin, P. M. & Seifert, H. S. 2009 ksgA mutations confer resistance to kasugamycin in *Neisseria gonorrhoeae*. *International Journal of Antimicrobial Agents* **33** (4), 321–327. <https://doi.org/10.1016/j.ijantimicag.2008.08.030>.
- El Ghachi, M., Bouhss, A., Blanot, D. & Mengin-Lecreulx, D. 2004 The bacA gene of *Escherichia coli* encodes an undecaprenyl pyrophosphate phosphatase activity. *Journal of Biological Chemistry* **279** (29), 30106–30113. <https://doi.org/10.1074/jbc.M401701200>.
- Finegold, S. M., Song, Y., Liu, C., Hecht, D. W., Summanen, P., Könönen, E. & Allen, S. D. 2005 *Clostridium clostridioforme*: a mixture of three clinically important species. *European Journal of Clinical Microbiology and Infectious Diseases* **24** (5), 319–324. <https://doi.org/10.1007/s10096-005-1334-6>.

- Fourie, J. C. J., Sanko, T. J., Bezuidenhout, C. C., Mienie, C. & Adeleke, R. A. 2019 Draft genome sequences of potentially pathogenic *Clostridium perfringens* strains from environmental surface water in the North West Province of South Africa. *Microbiology Resource Announcements* **8** (32), 18–20. <https://doi.org/10.1128/mra.00407-19>.
- Fournier, B., Aras, R. & Hooper, D. C. 2000 Expression of the multidrug resistance transporter NorA from *Staphylococcus aureus* is modified by a two-component regulatory system. *Journal of Bacteriology* **182** (3), 664–671. <https://doi.org/10.1128/JB.182.3.664-671.2000>.
- Gross, T. P., Kamara, L. B., Hatheway, C. L., Powers, P., Libonati, J. P., Harmon, S. M. & Israel, E. 1989 *Clostridium perfringens* food poisoning: use of serotyping in an outbreak setting. *Journal of Clinical Microbiology* **27** (4), 660–663.
- Hassan, K. A., Elbourne, L. D. H., Tetu, S. G., Melville, S. B., Rood, J. I. & Paulsen, I. T. 2015 Genomic analyses of *Clostridium perfringens* isolates from five toxinotypes. *Research in Microbiology* **166** (4), 255–263. <https://doi.org/10.1016/j.resmic.2014.10.005>.
- Huovinen, P., Sundstrom, L., Swedberg, G. & Skold, O. 1995 Trimethoprim and sulfonamide resistance. *Antimicrobial Agents and Chemotherapy* **39** (2), 279–289. <https://doi.org/10.1128/aac.39.2.279>.
- Hynes, W. L. & Walton, S. L. 2000 Hyaluronidases of Gram-positive bacteria. *FEMS Microbiology Letters* **183** (2), 201–207. [https://doi.org/10.1016/S0378-1097\(99\)00669-2](https://doi.org/10.1016/S0378-1097(99)00669-2).
- Katayama, S., Nozu, N., Okuda, M., Hirota, S., Yamasaki, T. & Hitsumoto, Y. 2009 Characterization of two putative fibronectin-binding proteins of *Clostridium perfringens*. *Anaerobe* **15** (4), 155–159. <https://doi.org/10.1016/j.anaerobe.2009.03.001>.
- Kim, Y. B., Kim, J. Y., Song, H. S., Lee, C., Kwon, J., Kang, J. & Roh, S. W. 2017 Complete genome sequence of *Clostridium perfringens* CBA7123 isolated from a faecal sample from Korea. *Gut Pathogens* **9** (1), 1–6. <https://doi.org/10.1186/s13099-017-0181-1>.
- Kiu, R. & Hall, L. J. 2018 An update on the human and animal enteric pathogen *Clostridium perfringens*. *Emerging Microbes and Infections* **7** (1). <https://doi.org/10.1038/s41426-018-0144-8>.
- Kiu, R., Caim, S., Alexander, S., Pachori, P. & Hall, L. J. 2017 Probing genomic aspects of the multi-host pathogen *Clostridium perfringens* reveals significant pangenome diversity, and a diverse array of virulence factors. *Frontiers in Microbiology* **8** (Dec). <https://doi.org/10.3389/fmicb.2017.02485>.
- Klein, S., Pipes, S. & Lovell, C. R. 2018 Occurrence and significance of pathogenicity and fitness islands in environmental vibrios. *AMB Express* **8**, 177. <https://doi.org/10.1186/s13568-018-0704-2>.
- Koch, C. 1998 In-vitro antibiotic susceptibility and molecular analysis of anaerobic bacteria isolated in Cape Town, South Africa. *Journal of Antimicrobial Chemotherapy* **42** (2), 245–248. <https://doi.org/10.1093/jac/42.2.245>.
- Kümmerer, K. 2004 Resistance in the environment. *Journal of Antimicrobial Chemotherapy* **54** (2), 311–320. <https://doi.org/10.1093/jac/dkh325>.
- Lee, I., Kim, Y. O., Park, S. C. & Chun, J. 2015 OrthoANI: an improved algorithm and software for calculating average nucleotide identity. *International Journal of Systematic and Evolutionary Microbiology* **66**, 1100–1105. <https://doi.org/10.1099/ijsem.0.000760>.
- Li, J., Uzal, F. A. & McClane, B. A. 2016 *Clostridium perfringens* sialidases: potential contributors to intestinal pathogenesis and therapeutic targets. *Toxins* **8** (11). <https://doi.org/10.3390/toxins8110341>.
- Li, C., Yan, X. & Lillehoj, H. S. 2017 Complete genome sequences of *Clostridium perfringens* del1 strain isolated from chickens affected by necrotic enteritis. *Gut Pathogens* **9** (1), 1–7. <https://doi.org/10.1186/s13099-017-0217-6>.
- Li, J., Tai, C., Deng, Z., Zhong, W., He, Y. & Ou, H. Y. 2018 VRprofile: gene-cluster-detection-based profiling of virulence and antibiotic resistance traits encoded within genome sequences of pathogenic bacteria. *Briefings in Bioinformatics* **19** (4), 566–574. <https://doi.org/10.1093/bib/bbw141>.
- López, M., Kadlec, K., Schwarz, S. & Torres, C. 2012 First detection of the staphylococcal trimethoprim resistance gene *dfcK* and the *dfcK*-carrying transposon *tn559* in enterococci. *Microbial Drug Resistance* **18** (1), 13–18. <https://doi.org/10.1089/mdr.2011.0073>.
- Lynch, T., Chong, P., Zhang, J., Hizon, R., Du, T., Graham, M. R. & Mulvey, M. R. 2013 Characterization of a stable, metronidazole-resistant *Clostridium difficile* clinical isolate. *PLoS ONE* **8** (1). <https://doi.org/10.1371/journal.pone.0053757>.
- Ma, M., Vidal, J., Saputo, J., McClane, B. A. & Uzal, F. 2011 The VirS/VirR two-component system regulates the anaerobic cytotoxicity, intestinal pathogenicity, and enterotoxemic lethality of *Clostridium perfringens* type C isolate CN3685. *Health* **2** (1), 1–9. <https://doi.org/10.1128/mBio.00338-10>.
- Mishra, R., Sinha, N. & Duncalf, R. 2016 B lactamase producing *Clostridium perfringens* bacteremia in an elderly man with acute pancreatitis. *Case Reports in Critical Care* **2016**, 1–4. <https://doi.org/10.1155/2016/7078180>.
- Myers, G. S. A., Rasko, D. A., Cheung, J. K., Ravel, J., Seshadri, R., DeBoy, R. T. & Paulsen, I. T. 2006 Skewed genomic variability in strains of the toxigenic bacterial pathogen, *Clostridium perfringens*. *Genome Research* **16** (8), 1031–1040. <https://doi.org/10.1101/gr.5238106>.
- Ohtani, K. & Shimizu, T. 2016 Regulation of toxin production in *clostridium perfringens*. *Toxins* **8** (7), 1–14. <https://doi.org/10.3390/toxins8070207>.
- Overbeek, R., Olson, R., Pusch, G. D., Olsen, G. J., Davis, J. J., Disz, T. & Stevens, R. 2014 The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Research* **42**, 206–214. <https://doi:10.1093/nar/gkt1226>.
- Pal, C., Asiani, K., Arya, S., Rensing, C., Stekel, D. J., Larsson, D. G. J. & Hobman, J. L. 2017 Metal resistance and its association with antibiotic resistance. *Advances in Microbial*

- Physiology* **70** (April 2018), 261–313. <https://doi.org/10.1016/bs.ampbs.2017.02.001>.
- Paredes-Sabja, D., Torres, J. A., Setlow, P. & Sarker, M. R. 2008 *Clostridium perfringens* spore germination: characterization of germinants and their receptors. *Journal of Bacteriology* **190** (4), 1190–1201. <https://doi.org/10.1128/JB.01748-07>.
- Partridge, S. R., Kwong, S. M., Firth, N. & Jensen, S. O. 2018 Mobile genetic elements associated with antimicrobial resistance. *Clinical Microbiology Reviews* **31** (4), e00088-17. <https://doi.org/10.1128/CMR.00088-17>.
- Peltier, J., Courtin, P., El Meouche, I., Catel-Ferreira, M., Chapot-Chartier, M. P., Lemée, L. & Pons, J. L. 2013 Genomic and expression analysis of the vanG-like gene cluster of *Clostridium difficile*. *Microbiology* **159** (Pt 7), 1510–1520. <https://doi.org/10.1099/mic.0.065060-0>.
- Price, N. P. J., Jackson, M. A., Singh, V., Hartman, T. M., Blackburn, J. A. & Dowd, P. F. 2019 Synergistic enhancement of β -lactam antibiotics by modified tunicamycin analogs *tunr1* and *tunr2*. *Journal of Antibiotics* **72** (11), 807–815. <https://doi.org/10.1038/s41429-019-0220-x>.
- Ramisetty, B. C. M. & Sudhakari, P. A. 2019 Bacterial ‘Grounded’ prophages: hotspots for genetic renovation and innovation. *Frontiers in Genetics* **10** (65). <https://doi.org/10.3389/fgene.2019.00065>.
- Revitt-Mills, S. A., Rood, J. I. & Adams, V. 2015 *Clostridium perfringens* extracellular toxins and enzymes: 20 and counting. *Microbiology Australia* **9–12**. <https://doi.org/10.1071/ma15039>.
- Roberts, M. C. 2002 Resistance to tetracycline, macrolide-lincosamide-streptogramin, trimethoprim, and sulfonamide drug classes. *Applied Biochemistry and Biotechnology – Part B Molecular Biotechnology* **20** (3), 261–283. <https://doi.org/10.1385/MB:20:3:261>.
- Roberts, M. C. 2011 Mechanisms of bacterial antibiotic resistance and lessons learned from environmental tetracycline-resistant bacteria. In: *Antimicrobial Resistance in the Environment*. pp. 93–121. <https://doi.org/10.1002/9781118156247.ch7>
- Rodgers, K., Arvidson, C. G. & Melville, S. 2011 Expression of a *Clostridium perfringens* type IV pilin by *Neisseria gonorrhoeae* mediates adherence to muscle cells. *Infection and Immunity* **79** (8), 3096–3105. <https://doi.org/10.1128/IAI.00909-10>.
- Rood, J. I. 1998 Virulence genes of *Clostridium Myonecrosis* or gas Gangrene. *Annual Review of Microbiology* **52**, 333–360.
- Sacks, L. E. 1983 Influence of carbohydrates on growth and sporulation of *Clostridium perfringens* in a defined medium with or without guanosine. *Applied and Environmental Microbiology* **46** (5), 1169–1175.
- Sanakal, R. D. & Kaliwal, B. B. 2011 Vancomycin resistance genes in various organisms—an in silico study. *International Journal of Biometric and Bioinformatics* **5** (2), 111–129.
- Sathish, S. & Swaminathan, K. 2009 Genetic diversity among toxigenic clostridia isolated from soil, water, meat and associated polluted sites in South India. *Indian Journal of Medical Microbiology* **27** (4), 311–320. <https://doi.org/10.4103/0255-0857.55443>.
- Sheykhsaran, E., Baghi, H. B., Soroush, M. H. & Ghotaslou, R. 2019 An overview of tetracyclines and related resistance mechanisms. *Reviews in Medical Microbiology* **30** (1), 69–75. <https://doi.org/10.1097/mrm.000000000000154>.
- Shimizu, T., Ohtani, K., Hirakawa, H., Ohshima, K., Yamashita, A., Shiba, T. & Hayashi, H. 2002 Complete genome sequence of *Clostridium perfringens*, an anaerobic flesh-eater. *Proceedings of the National Academy of Sciences of the United States of America* **99** (2), 996–1001. <https://doi.org/10.1073/pnas.022493799>.
- Shoemaker, N. B., Vlamakis, H., Hayes, K., Salyers, A. A., Jensen, L. B., Hammerum, A. M. & Westh, H. 2001 Evidence for extensive resistance gene transfer among. *Applied and Environmental Microbiology* **67** (2), 561–568. <https://doi.org/10.1128/AEM.67.2.561>.
- Sloan, J., McMurry, L. M., Lyras, D., Levy, S. B. & Rood, J. I. 1994 The *Clostridium perfringens* Tet P determinant comprises two overlapping genes: *tetA(P)*, which mediates active tetracycline efflux, and *tetB(P)*, which is related to the ribosomal protection family of tetracycline-resistance determinants. *Molecular Microbiology* **11** (2), 403–415. <https://doi.org/10.1111/j.1365-2958.1994.tb00320.x>.
- Spigaglia, P. & Mastrantonio, P. 2002 Analysis of macrolide-lincosamide-streptogramin B (MLS_B) resistance determinant in strains of *Clostridium difficile*. *Microbial Drug Resistance* **8** (1), 45–53. <https://doi.org/10.1089/10766290252913755>.
- Tansuphasiri, U., Matra, W. & Sangsuk, L. 2005 Antimicrobial resistance among *Clostridium perfringens* isolated from various sources in Thailand. *Southeast Asian Journal of Tropical Medicine and Public Health* **36** (4), 954–961.
- Tatusova, T., Dicuccio, M., Badretdin, A., Chetvernin, V., Nawrocki, P., Zaslavsky, L., Lomsadze, A., Pruitt, K., D., Borodovsky, M. & Ostell, J. 2016 NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Research* **44**, 6614–6624. <https://doi:10.1093/nar/gkw569>.
- Van Hoek, A. H. A. M., Mevius, D., Guerra, B., Mullany, P., Roberts, A. P. & Aarts, H. J. M. 2011 Acquired antibiotic resistance genes: an overview. *Frontiers in Microbiology* **2** (Sep), 1–27. <https://doi.org/10.3389/fmicb.2011.00203>.
- Vidor, C. J., Bulach, D., Awad, M. & Lyras, D. 2019 *Paenoclostridium sordellii* and *Clostridioides difficile* encode similar and clinically relevant tetracycline resistance loci in diverse genomic locations. *BMC Microbiology* **19** (1), 1–12. <https://doi.org/10.1186/s12866-019-1427-5>.
- Villela, J. G. A., Ritschel, P., Barbosa, M. A. G., Baccin, K. M. S., Rossato, M., Maia, J. D. G. & Ferreira, M. A. S. V. 2019 Detection of *Xanthomonas citri* pv. *viticola* on grapevine by real-time PCR and BIO-PCR using primers designed from pathogenicity and xanthomonadin gene sequences. *European Journal of Plant Pathology* **155** (2), 445–459. <https://doi.org/10.1007/s10658-019-01779-y>.
- Willems, R. J. L., Top, J., Van Santen, M., Robinson, D. A., Coque, T. M., Baquero, F. & Bonten, M. J. M. 2005 Global spread of vancomycin-resistant *Enterococcus faecium* from

distinct nosocomial genetic complex. *Emerging Infectious Diseases* **11** (6), 821–828. <https://doi.org/10.3201/1106.041204>.

Xu, L., Dong, Z., Fang, L., Luo, Y., Wei, Z., Guo, H. & Wang, Y. 2019 *OrthoVenn2: a web server for whole-genome comparison and annotation of orthologous clusters across*

multiple species. Nucleic Acids Research **47** (W1), W52–W58. <https://doi.org/10.1093/nar/gkz333>.

Yoon, S. H., Park, Y. K. & Kim, J. F. 2015 *PAIDB v2.0: exploration and analysis of pathogenicity and resistance islands. Nucleic Acids Research* **43** (D1), D624–D630. <https://doi.org/10.1093/nar/gku985>.

First received 27 January 2020; accepted in revised form 4 May 2020. Available online 26 May 2020