

A systematic review of microorganisms as indicators of recreational water quality in natural and drinking water systems

José Luis Lugo, Elkyn Rafael Lugo and Mario de la Puente

ABSTRACT

The purpose of this review was to highlight the most frequent biological indicators used to estimate the microbiological quality of drinking and recreational water. It was observed that the incorporation of other microbiological indicators should be considered to strengthen the decision-making process on water quality management and guarantee its safe consumption in recreational activities.

Key words | drinking water, microbiological indicators, recreational water

José Luis Lugo

Udecino Group for Environmental Research,
Universidad de Cundinamarca,
19th Street N° 24-209,
Girardot-Cundinamarca,
Colombia

Elkyn Rafael Lugo

Corporación Universitaria Minuto de Dios,
Uniminuto, Cra. 53rd Street # 74-110,
Barranquilla,
Colombia

Mario de la Puente (corresponding author)

Universidad del Norte,
Barranquilla, Km 5,
Puerto Colombia,
Colombia
E-mail: mdelapuate@uninorte.edu.co

HIGHLIGHTS

- The incorporation of other microbiological indicators should be considered to strengthen the decision-making process on water quality management.
- The presence and concentration of bacterial, viral, and parasitic indicators are similar in several countries with diverse environmental conditions.

INTRODUCTION

Water is one of the most valuable natural resources, an essential element for life development and human activities (Romeu-Álvarez *et al.* 2012). It is intended for different uses: human consumption, industrial processes, recreational activities of primary and secondary contact, ecological conservation, among others (Baird & Cann 2014).

Due to alterations in natural and anthropogenic processes, it is necessary to guarantee its quality so that there is no risk for humans and the environment (Samboni-Ruiz *et al.* 2007; Lugo & Lugo 2018). Water is suitable for human consumption and domestic use, including personal hygiene and recreational activities, when it has low physical and chemical contaminant concentrations (WHO 2011; Lugo-Arias *et al.* 2020). Although the use of water in recreational activities can bring health benefits, it

can also generate adverse effects when it is contaminated (WHO 2000). To assess the microbiological water quality, organisms are used as single indicators since their presence reveals contamination (Silva-Iñiguez *et al.* 2007).

Microbiological indicators of fecal contamination expose pathogenic organisms. They must be in higher concentration than pathogens and have similar survival characteristics (Noble *et al.* 2003; Pulido *et al.* 2005). Fecal contamination indicators have proven to be an alternative to the difficulty of identifying and quantifying pathogens that cause water-origin diseases (Campos-Pinilla *et al.* 2008). Microbiological quality control of water for human consumption and recreational use (ACH) requires pathogenic microorganisms' analysis (Silva-Iñiguez *et al.* 2007).

These analyses are hard to execute due to the great variety of cultivable pathogenic bacteria, isolation tests, the low concentration of aggressive species, the need for specialized laboratories, high economic costs, and time-consuming. To identify the presence of pathogens in a reliable way, water

This is an Open Access article distributed under the terms of the Creative Commons Attribution Licence (CC BY 4.0), which permits copying, adaptation and redistribution, provided the original work is properly cited (<http://creativecommons.org/licenses/by/4.0/>).

doi: 10.2166/wh.2020.179

quality monitoring should be done by searching for fecal contamination indicators approved by international and national standards.

Microorganisms must meet the following requirements: (1) be a normal constituent of the intestinal microbiota of healthy individuals; (2) be present exclusively in the feces of homeothermic animals; (3) be present where pathogenic intestinal microorganisms are; (4) appear in high numbers; (5) facilitate isolation and identification; (6) be unable to reproduce outside the intestine of homeothermic animals. Their survival time must be equal to or greater than a pathogenic bacterium, and their resistance to environmental factors must be equal to or greater than fecal pathogens (Fernández *et al.* 2001). The objective of this review is to highlight the most common microbiological indicators that are used to evaluate the microbiological quality of drinking and recreational water to identify new methods that optimize microbiological monitoring of water quality.

METHOD

The research was conducted in the Science Direct, Redalyc, Scielo, and in the Google Scholar search engine selecting peer-reviewed documents and published research articles in the last ten years, based on the following keywords: microbiological indicators, drinking water, and recreational water. Articles' selection criteria were as follows: (1) biological indicators to assess the microbiological quality of drinking and recreational water quality; (2) water bodies that were affected by outflow wastewater; (3) the impact of microbiological water quality on individuals' health; and (4) recreational water affected by bathers.

Recreational waters in artificial systems such as swimming pools were not considered as they have different environmental conditions, compared to those of interest in this work. For example, swimming pool water quality criteria are like those for the quality of drinking water, including microbiological parameters that differ from the water quality characteristics of natural bodies (Carrasquero Ferrer *et al.* 2020). Also, swimming pool water is treated with different treatment methods, for example, disinfection with chlorine, UV-C radiation, ozone, membrane processes, among others (Dudziak *et al.* 2019; Skibinski *et al.* 2019).

The above is not feasible from a technical and economic point of view for natural water bodies such as seas and rivers.

Articles that did not meet the above criteria were excluded from the analysis. The guidelines were also based on microbiological quality standards for drinking and recreational water of the following organizations: the World Health Organization (WHO), the United States Environmental Protection Agency (EPA), and the standards of water quality of the European Union (EU). These served as the basis for defining the microbiological water quality criteria in Latin America and the world.

RESULTS AND DISCUSSION

The most damaging effect of contaminated water has been the transmission of diseases by microorganisms that can inhabit humans (see Table 1) (Romero *et al.* 2009).

Table 1 | Main microorganisms that transmit diseases in the water

Type of microorganism	Name of the microorganism	Disease
Bacterias	<i>Escherichia coli</i>	Diarrhea and stomach pain
	<i>Shigella</i> spp.	Shigellosis
	<i>Vibrio cholerae</i>	Cholera
	<i>Salmonella typhi</i>	Typhoid fever
	<i>Salmonella</i> spp.	Salmonellosis
	<i>Yersinia enterocolitica</i>	Yersiniosis
	<i>Campylobacter jejuni</i>	Enteritis
Viruses	Enterovirus	Various diseases: intestinal, respiratory, among others
	Rotavirus	Various diseases: intestinal, fever, among others
	Adenovirus	Various diseases: colds, conjunctivitis, among others
Protozoa	<i>Giardia lamblia</i>	Giardiasis
	<i>Cryptosporidium parvum</i>	Cryptosporidiosis
	<i>Entamoeba histolytica</i>	Dysentery
	<i>Ascaris lumbricoides</i>	Ascariasis

Source: Modified from WHO (2003).

Bioindicators serve as a complementary tool to evaluate water microbiological quality. Their application requires organism identification based on diversity indices, adjusting intervals that quantify water quality. For example, in Japan, the authorities in charge of water monitoring have illustrated guides of the organisms (bacteria, fungi, insects, amphibians, and fish) that can be found in water sources, including information on the tolerance of pollutants (heavy metals, dioxins, among others) to chemical compounds. This outlines information on the state of the water in real-time (Koschelov & Briedis 2013). An organism is considered a bioindicator when its degree of tolerance is known, since not all of them offer information due to their eating habits, life cycle, and frequency with which they are found (Miravet Sánchez et al. 2016).

It is found that the presence and concentration of bacterial, viral, and parasitic indicators are similar in several countries with diverse environmental conditions (Liberatore et al. 2015), proving their use in different types of water (Campos-Pinilla et al. 2008). Domestic contamination generates high health risks due to high concentrations of fecal microorganisms (Wade et al. 2015). Therefore, it is important to have laboratory tests, exposing bacteria, viruses, and parasite concentrations in a precise moment with low economic costs. The most common bacteria used for drinking water control, wastewater, and recreational uses are total coliforms, fecal coliforms, *Escherichia coli*, and *Enterococcus faecalis*.

Clostridium perfringens has also been proposed as an indicator of distant fecal contamination in groundwater and as an indicator of the presence of protozoan cysts (Campos-Pinilla et al. 2008). The presence of coliform bacteria is considered the best indicator of fecal contamination, and in the long term, it serves to monitor the effectiveness of control programs. Its use is advantageous because it gives a quick response to environmental changes such as pollution based on domestic discharges (Koschelov & Briedis 2013). Numerous epidemiological studies in different aquatic environments have shown a relationship between coliform counts and the appearance of infectious diseases in humans.

However, various studies reveal that there is no significant relationship between these indicators and diseases related to seawater bathing (Vergaray et al. 2007). Total coliforms and *Escherichia coli* (*E. coli*) have been suggested

as reliable indicators of wastewater contamination. Recreational waters such as seawater are susceptible to fecal contamination, which can increase the health risk associated with swimming in contaminated beach water (Praveena et al. 2013). In water analysis, the presence of *E. coli* indicates fecal contamination, with a positive correlation between the concentration of organisms and the amount of contamination (Romero et al. 2009).

Also, there is a trend towards linearity between the concentrations of thermotolerant coliforms and *E. coli*, which shows that such indicators could be used as a tool for river water microbiological quality (Romeu-Álvarez et al. 2012).

This includes fecal enterococci (FE) as a bacteriological indicator in marine waters for recreational use due to its resistance to seawater conditions, temperature, and relationship with gastrointestinal, respiratory, and dermatological diseases (Silva-Iñiguez et al. 2007). Although no indicator can accurately determine water quality, enterococci have a predictive bent in marine environments, while somatic bacteriophages are useful bioindicators in seawater ecosystems (Janelidze et al. 2011). In contrast, a study by Atoyan et al. (2011) suggests that FE and fecal coliforms (FC) are not reliable indicators of human fecal contamination in the Pettaquamscutt River due to inconsistencies found in high rates of false-positive and negative tests. Furthermore, there was a considerable amount of spatial and temporal variability in the reliability of the tests.

The inclusion of chemical nucleic acid-based methods for evaluating the microbiological quality of water can improve the identification of sources of human fecal contamination in water sources (Ahmed et al. 2015). The United States Environmental Protection Agency requires the use of FE as an indicator of the risk of contracting a gastrointestinal illness from swimming activities in waters contaminated by FC. However, none of these indicators are definite for human fecal contamination because there is a wide variety of warm-blooded animals (Atoyan et al. 2011). That is why several techniques have recently emerged for monitoring human fecal contamination sources based on indicators related to human beings.

Adolescentis bifidobacterium has been suggested as a specific bacterium of the human intestinal tract that can accurately identify human fecal contamination in

freshwater. It is an anaerobic Gram-negative bacillus that grows exclusively in human intestines and has been suggested as an alternative fecal contamination indicator, since it does not survive in seawater, so it cannot reproduce itself in the environment. Its evaluation is carried out using a nested polymerase chain reaction (PCR) method for the exclusive detection of *A. bifidobacterium* in human feces.

It has been used in water and sediment samples (Atoyan *et al.* 2011). On the other hand, the presence of pathogenic viruses has gained importance in recent decades due to epidemiological studies and improvements in environmental detection techniques.

However, these techniques require molecular procedures that are not within the reach of most water analysis laboratories (Moresco *et al.* 2012). As an alternative, bacteriophages such as the ones that infect *Bacteroides fragilis* have been proposed. It was observed that its behavior is like some viruses that cause diseases of hydric origin, and its detection can be done in a few hours. Depending on the type of water and environmental conditions, the use of phages is recommended (Campos-Pinilla *et al.* 2008). Monitoring coastal waters with total coliforms, FC, and *E. faecalis* phage indicators are recommended to prevent potential waterborne diseases (Janelidze *et al.* 2011).

Also, water sample testing is essential to verify the presence of indigenous microflora, such as pathogenic *Vibrio* species. It can be useful to predict and prevent waterborne diseases, such as cholera and *Vibrio*-related gastroenteritis (Janelidze *et al.* 2011). The risk of infection by enteric viruses and other pathogens in the use of recreational waters has been described in some studies. For example, the coastal water quality in Brazil was evaluated through base fecal indicators, such as coliforms or *Enterococcus* spp. (Prasad *et al.* 2015). However, bacterial contamination did not correlate with the presence of human enteric viruses (Wong *et al.* 2009).

Human viruses, such as adenoviruses (HAdV), hepatitis A (HAV), polyomaviruses (JCPyV), and human noroviruses (HuNoV), are associated with waterborne diseases. HAdV, HAV, and HuNoV can replicate inside the gastrointestinal tract and are excreted in high concentrations in feces of infected individuals (Moresco *et al.* 2012). JCPyV is found in the urine of approximately 80% of adults, and the infection is spread during childhood. These viruses are usually

spread into the water by industrial or agricultural activities (Lyons *et al.* 2015; Rusiñol *et al.* 2015). Once in the water, these viruses are very stable and resistant to chemical agents such as UV radiation and chlorine.

The presence of viruses in various water environments has been revealed, highlighting the need to include them in the analysis of water quality. In a study carried out by Moresco *et al.* (2012), no correlation was observed between virus detection and FC presence. It can be assumed that these common bioindicators (FC) are not accurate in associating the risk of disease contraction with water environments (Rusiñol *et al.* 2015). In the case of parasites, the presence of helminth eggs and protozoan cysts is the main risk to human health. The parasites most studied are giardia cysts and *Cryptosporidium* oocysts. The detection and counting of these parasites are an alternative health indicator since analysis is easy to carry out in laboratories with basic equipment (Campos-Pinilla *et al.* 2008).

Recreational water quality criteria

Water quality standards have been used through microbiological parameters that measure the degree of contamination (Bonamano *et al.* 2015). Table 2 shows the microbiological quality criteria for primary contact with recreational water. Table 3 shows studies of microbiological indicators for recreational waters.

Microbiological water quality analysis must consider the sources of microbial contamination of coastal areas since it has been found that fecal indicators, coliphages, and

Table 2 | Criteria for evaluating the microbiological quality of recreational water

Fecal indicator	Allowable limit	Normative
Total coliforms (TC)	<1,000 MPN/100 ml	Decree 1594 of 1984 (Colombia)
Fecal coliforms (FC)	<200 MPN/100 ml	Decree 1594 of 1984 (Colombia)
Enterococci	40 CFU/100 ml 200 CFU/100 ml 185 CFU/100 ml	NTS-TS 001-2 WHO (2003) Directive 2006/7/EC of the European Council
<i>Escherichia coli</i>	500 CFU/100 ml	Directive 2006/7/EC of the European Council

Source: Campos-Pinilla *et al.* (2008).

Table 3 | Studies carried out to determine the microbiological quality of recreational water

Water type	Indicator	Determination technique	Geographic location	References
Seawater	Total coliforms	Membrane filtration (CFU/100 ml)	(Black Sea, Georgia); (Teluk Kemang Beach, Malaysia); (Kuwait Bay)	Janelidze et al. (2011); Praveena et al. (2013); Lyons et al. (2015)
		Most probable number (MPN)	8 beaches of Peru	Vergaray et al. (2007)
	Fecal coliforms	Membrane filtration (CFU/100 ml)	(Black Sea, Georgia); (Kuwait Bay)	Janelidze et al. (2011); Lyons et al. (2015)
		Most probable number (MPN)	(8 beaches in Peru); (2 beaches in Costa Rica)	Vergaray et al. (2007); Badilla-Aguilar & Mora-Alvarado (2019)
	Fecal enterococci	Membrane filtration (CFU/100 ml)	(5 beaches in Brazil); (Kuwait Bay)	Lamparelli et al. (2015); Lyons et al. (2015)
		Most probable number (MPN)	(Playa la Boquita-México); (Black Sea, Georgia); (8 beaches in Peru); (Adriatic Sea, Italy); (2 beaches in Costa Rica)	Silva-Iñiguez et al. (2007); Janelidze et al. (2011); Vergaray et al. (2007); Liberatore et al. (2015); Badilla-Aguilar & Mora-Alvarado (2019)
	<i>Escherichia coli</i>	Most probable number (MPN)	(8 beaches in Peru); (2 beaches in Costa Rica)	Vergaray et al. (2007); Badilla-Aguilar & Mora-Alvarado (2019)
		Membrane filtration (CFU/100 ml)	(Santa Catarina Island-Brazil); (5 beaches in Brazil); (Teluk Kemang Beach, Malaysia)	Moresco et al. (2012); Lamparelli et al. (2015); Praveena et al. (2013)
	<i>Clostridium perfringens</i>	Membrane filtration (CFU/100 ml)	Atlantic Beaches (Colombia)	Moreno et al. (2019)
	Vibriopathogens	Membrane filtration	Black Sea, Georgia	Janelidze et al. (2011)
	Coliphages	Membrane filtration	Black Sea, Georgia	Janelidze et al. (2011)
	Adenovirus (HAdV)	Flocculation	Santa Catarina Island-Brazil	Moresco et al. (2012)
	Polyomavirus (JCPyV)	Flocculation	Santa Catarina Island-Brazil	Moresco et al. (2012)
	Norovirus	Flocculation	Santa Catarina Island-Brazil	Moresco et al. (2012)
	River water	Total coliforms	Membrane filtration (CFU/100 ml)	(Río Cúpira, Venezuela); (Danube river)
Most probable number (MPN)			(Río Nazas, Mexico); (Machángara and Monjas rivers, Ecuador)	Romero et al. (2009); Campaña et al. (2017)
Fecal coliforms		Membrane filtration (CFU/100 ml)	(Río Cúpira, Venezuela); (Luyanó River, Cuba)	Koschelov & Briedis (2013); Romeu-Álvarez et al. (2012)
		Most probable number (MPN)	(Río Nazas, Mexico); (Machángara and Monjas rivers, Ecuador)	Romero et al. (2009); Campaña et al. (2017)
Fecal enterococci		Membrane filtration (CFU/100 ml)	(Danube river)	Kirschner et al. (2009)
<i>Escherichia coli</i>		Membrane filtration (CFU/100 ml)	(Danube river); (Río Luyanó, Cuba); (Puerto Rico); (Umgeni River Catchment, South Africa); (Rivers of Poland)	Kirschner et al. (2009); Romeu-Álvarez et al. (2012); Wade et al. (2015); Baker et al. (2015); Lenart-Boroń et al. (2017)
		Bacteroides HF183	PCR	(Puerto Rico)
<i>A. bifidobacterium</i>		Fluorescence	(Pettaquamscutt River, USA)	Atoyan et al. (2011)
<i>Pseudomonas aeruginosa</i> and <i>Salmonella</i>				

Source: Developed by the authors.

enteroviruses are significantly associated with precipitation, sewage discharges, and temperature (Janelidze et al. 2011).

Microbiological indicators of drinking water quality

Purification systems are characterized by their efficiency in the elimination of bacteria but not in viruses. Thus, the above performance does not guarantee the removal of viruses and parasites since they can be resistant to chlorine disinfection systems (Campos-Pinilla et al. 2008). Viruses, parasites, and bacteria that indicate fecal contamination must be considered to evaluate the microbiological quality of water sources. Various studies have associated the presence of viruses with bacterial indicators of fecal contamination (Rusiñol et al. 2015).

There is low confidence in populations in developing countries to drink tap water for it not complying with sanitary conditions. This is due to contamination in distribution networks (WHO 2011). For bottled water, it is essential to highlight the following origins of the bacterial flora: (1) bacteria belonging to the point of emergence are found (autochthonous microflora); (2) the bacteria 'added' to the water during the packaging process (non-native microflora) (Payares et al. 2013). In the absence of fecal indicators in treated water, the heterotrophic plaque count (HPC) serves to assess its overall microbiological quality (Bartram et al. 2004).

Although it is not an indicator of adverse effects on human health, its increase may indicate problems with raw water quality, water treatment, or the distribution system (WHO 2012). Sometimes it is high, and operational decision-makers are faced with the problem of long lists of zero-count samples. Since water distribution systems are not sterile, this reflects that the conventional approach is a limited diagnostic tool (Gillespie et al. 2014). One of the most promising developments in microbiological surveillance of water quality is flow cytometry (FCM) (Gillespie et al. 2014). Switzerland was the first country to incorporate a standardized method for FCM in the water industry to determine the count of different total cells and populations (Harry et al. 2016).

The number of bacteria detected in drinking water by FCM is greater than the number of HPC (Hammes et al. 2008). Van Nevel et al. (2017) exposes multiple reasons (including cost) why FCM is a suitable alternative to replace

HPC. However, FCM has disadvantages such as its high investment cost due to the nature of the equipment, the procedures involved, the need for expensive dyes (used for sample staining), and the high initial capital cost of the equipment. Therefore, the use of FCM is inappropriate to be applied in field investigations (Bridgeman et al. 2015). Fluorescence analysis is a quick technique that requires low sample volumes. They are typically 10–1,000 times more sensitive than those of UV absorption spectroscopy with the detection of individual molecules (Henderson et al. 2009).

A fluorescence device with LED technology has been proposed to facilitate the monitoring of water quality (Bridgeman et al. 2015; Sorensen et al. 2018). Besides, it is essential to include such techniques to evaluate the microbiological quality of drinking water since the information obtained from the microorganisms is in real-time and it does not take long to generate the data (as in conventional techniques) (Sorensen et al. 2015; Bonadonna et al. 2019).

It must be ensured that the supplied drinking water does not pose any health risk to consumers (Bridgeman et al. 2015). Tables 4 and 5 show the recommended microorganisms for microbiologically evaluating the quality of drinking water.

CONCLUSIONS

The application of the most used microbiological indicators to assess the microbiological quality of drinking and recreational water has been described, highlighting reliable techniques for analyzing pathogenic microorganisms, considering technical and economic aspects. It is necessary to include other microbiological indicators to strengthen water

Table 4 | Microorganisms recommended as indicators of the quality of drinking water

Bacterias	Total coliforms Fecal coliforms <i>Escherichia coli</i> Heterotrophic microorganisms <i>Clostridium perfringens</i>
Viruses	Somatic coliphages Coliphages F specific Phages
Parasites	<i>Giardia lamblia</i> <i>Cryptosporidium parvum</i>

Source: Developed by the authors.

Table 5 | Criteria for evaluating the microbiological quality of drinking water

Fecal indicator	Allowable limit	Normative
Total coliforms	Not detectable	Resolution 2115/2007 (Colombia)
Fecal coliforms		U.S. EPA
Fecal coliforms		WHO
<i>Escherichia coli</i>		WHO
<i>Giardia lamblia</i>		Resolution 2115/2008 (Colombia)
Legionella		U.S. EPA
Enteric viruses		U.S. EPA
Legionella		U.S. EPA
Enteric viruses		U.S. EPA

Source: Developed by the authors.

quality research and guarantee the safety of its consumption and use in different recreational activities. These could be some biological agents that are not usually included in international regulations, such as viruses and parasites.

ACKNOWLEDGEMENTS

The authors thank COLCIENCIAS, the Soil, Environment and Society Research group of the Universidad del Magdalena, and the Magdalena Government. This study was funded by COLCIENCIAS and the Magdalena governorate with the code 2019-01. No potential conflicts are reported by the authors.

DATA AVAILABILITY STATEMENT

All relevant data are included in the paper or its Supplementary Information.

REFERENCES

- Ahmed, W., Harwood, V. J., Gyawali, P., Sidhu, J. P. S. & Toze, S. 2015 Comparison of concentration methods for quantitative detection of sewage-associated viral markers in environmental waters. *Applied and Environmental Microbiology* **81** (6), 2042–2049. doi:10.1128/AEM.03851-14.
- Atoyan, J. A., Herron, E. M. & Amador, J. A. 2011 Evaluation of microbiological water quality in the Pettaquamscutt River (Rhode Island, USA) using chemical, molecular and culture-dependent methods. *Marine Pollution Bulletin* **62** (7), 1577–1583. doi:10.1016/j.marpolbul.2011.04.032.
- Badilla-Aguilar, A. & Mora-Alvarado, D. A. 2019 Análisis de la calidad bacteriológica de dos playas tropicales: relación de indicadores de contaminación fecal entre el agua de mar y las arenas (Analysis of the bacteriological quality of two tropical beaches: relationship of indicators of fecal contamination between seawater and sands). *Revista Tecnología en Marcha* **37** (2), 38–45. doi:10.18845/tm.v32i10.4879.
- Baird, C. & Cann, M. 2014 *Química Ambiental*. Routledge, Madrid, Spain.
- Baker, A., Cumberland, S. A., Bradley, C., Buckley, C. & Bridgeman, J. 2015 To what extent can portable fluorescence spectroscopy be used in the real-time assessment of microbial water quality? *Science of the Total Environment* **532**, 14–19. doi:10.1016/j.scitotenv.2015.05.114.
- Bartram, J., Cotruvo, J., Exner, M., Fricker, C. & Glasmacher, A. 2004 Heterotrophic plate count measurement in drinking water safety management: report of an Expert Meeting Geneva, 24–25 April 2002. *International Journal of Food Microbiology* **92** (3), 241–247. doi:10.1016/j.ijfoodmicro.2003.08.005.
- Bonadonna, L., Briancesco, R. & La Rosa, G. 2019 Innovative analytical methods for monitoring microbiological and virological water quality. *Microchemical Journal* **150**, 104160. doi:10.1016/j.microc.2019.104160.
- Bonamano, S., Madonia, A., Borsellino, C., Stefanì, C., Caruso, G., De Pasquale, F., Peirmattei, V., Zappala, G. & Marcelli, M. 2015 Modeling the dispersion of viable and total *Escherichia coli* cells in the artificial semi-enclosed bathing area of Santa Marinella (Latium, Italy). *Marine Pollution Bulletin* **95** (1), 141–154. doi:10.1016/j.marpolbul.2015.04.030.
- Bridgeman, J., Baker, A., Brown, D. & Boxall, J. B. 2015 Portable LED fluorescence instrumentation for the rapid assessment of potable water quality. *Science of the Total Environment* **524**, 338–346. doi:10.1016/j.scitotenv.2015.04.050.
- Campaña, A., Gualoto, E. & Chiluisa-Utreras, V. 2017 Physico-chemical and microbiological evaluation of the water quality of the Machángara and Monjas rivers in the water network of the Quito metropolitan district. *Bionatura* **2** (2), 305–310. doi:10.21931/RB/2017.02.02.6.
- Campos-Pinilla, C., Cárdenas-Guzmán, M. & Guerrero-Cañizares, A. 2008 Behavior of the indicators of faecal contamination in different types of waters of the Bogotá savanna (Colombia). *Universitas Scientiarum* **13** (2), 103–108. Available from: <https://www.redalyc.org/pdf/499/49913201.pdf>
- Carrasquero Ferrer, S., Muñoz Colina, C., Tuvifíez Morales, P., Vargas Torres, R., Vargas Castellano, C. & Marín Leal, J. 2020 Calidad fisicoquímica y microbiológica del agua de piscinas de dos complejos recreacionales del Estado Zulia. Ve.scielo.org (Physicochemical and microbiological quality of the swimming pool water of two recreational complexes in Zulia State). Retrieved 20 October 2020, Available from:

- http://ve.scielo.org/scielo.php?script=sci_arttext&pid=S1690-46482016000200010
- Dudziak, M., Wyczarska-Kokot, J., Łaskawiec, E. & Stolarczyk, A. 2019 Application of ultrafiltration in a swimming pool water treatment system. *Membranes* **9** (3), 44. doi:10.3390/membranes9030044.
- Fernández, A., Molina, M., Alvarez, A., Alcántara, M. & Espigares, A. 2001 Phacohydric transmission and hepatitis A virus. *Higiene y Sanidad Ambiental*. Scielo.org.co. Retrieved 18 October 2020, Available from: http://www.scielo.org.co/scielo.php?script=sci_nlinks&ref=000100&pid=S1794-2470201300020000600023&lng=en
- Gillespie, S., Lipphaus, P., Green, J., Parsons, S., Weir, P., Juskowiak, K., Jefferson, B., Jarvis, P. & Nocker, A. 2014 Assessing microbiological water quality in drinking water distribution systems with disinfectant residual using flow cytometry. *Water Research* **65**, 224–234. doi. 10.1016/j.watres.2014.07.029.
- Hammes, F., Berney, M., Wang, Y., Vital, M., Köster, O. & Egli, T. 2008 Flow-cytometric total bacterial cell counts as a descriptive microbiological parameter for drinking water treatment processes. *Water Research* **42** (1–2), 269–277. doi:10.1016/j.watres.2007.07.009.
- Harry, I. S., Ameh, E., Coulon, F. & Nocker, A. 2016 Impact of treated sewage effluent on the microbiology of a small brook using flow cytometry as a diagnostic tool. *Water, Air, & Soil Pollution* **227** (2), 57. doi:10.1007/s11270-015-2723-9.
- Henderson, R. K., Baker, A., Murphy, K. R., Hambly, A., Stuetz, R. M. & Khan, S. J. 2009 Fluorescence as a potential monitoring tool for recycled water systems: a review. *Water Research* **43** (4), 863–881. doi:10.1016/j.watres.2008.11.027.
- Janelidze, N., Jaiani, E., Lashkhi, N., Tskhvediani, A., Kokashvili, T., Gvarishvili, T., Jgenti, D., Mikashvidze, E., Diasamidze, R., Narodny, S., Obiso, R., Whitehouse, C. A., Huq, A. & Tediashvili, M. 2011 Microbial water quality of the Georgian coastal zone of the Black Sea. *Marine Pollution Bulletin* **62** (3), 573–580. doi:10.1016/j.marpolbul.2010.11.027.
- Kirschner, A. K., Kavka, G. G., Velimirov, B., Mach, R. L., Sommer, R. & Farnleitner, A. H. 2009 Microbiological water quality along the Danube River: integrating data from two whole-river surveys and a transnational monitoring network. *Water Research* **43** (15), 3673–3684. doi:10.1016/j.watres.2009.05.034.
- Koschelov, V. S. & Briedis, G. S. 2013 Evaluation of the water quality of the Cúpira river (La Cumaca, Carabobo state, Venezuela) using microbiological bioindicators and physicochemical parameters. *Interciencia*. Redalyc.org. Retrieved 17 October 2020, Available from: <https://www.redalyc.org/pdf/339/33928556002.pdf>
- Lamparelli, C. C., Pogreba-Brown, K., Verhougstraete, M., Sato, M. I. Z., de Castro Bruni, A., Wade, T. J. & Eisenberg, J. N. 2015 Are fecal indicator bacteria appropriate measures of recreational water risks in the tropics: a cohort study of beach goers in Brazil? *Water Research* **87**, 59–68. doi:10.1016/j.watres.2015.09.001.
- Lenart-Boroń, A., Wolanin, A., Jelonkiewicz, E. & Żelazny, M. 2017 The effect of anthropogenic pressure shown by microbiological and chemical water quality indicators on the main rivers of Podhale, southern Poland. *Environmental Science and Pollution Research* **24** (14), 12938–12948. doi:10.1007/s11356-017-8826-7.
- Liberatore, L., Murmura, F. & Scarano, A. 2015 Bathing water profile in the coastal belt of the province of Pescara (Italy, Central Adriatic Sea). *Marine Pollution Bulletin* **95** (1), 100–106. doi:10.1016/j.marpolbul.2015.04.035.
- Lugo, J. L. & Lugo, E. R. 2018 Socioenvironmental benefits from water purification in the palafitic towns of the Ciénaga Grande de Santa Marta-Colombia. *Revista UDCA Actualidad & Divulgación Científica* **21** (1), 259–264. doi:10.31910/rudca.v21.n1.2018.685.
- Lugo-Arias, J., Lugo-Arias, E., Ovallos-Gazabon, D., Arango, J., de la Puente, M. & Silva, J. 2020 Effectiveness of the mixture of nopal and cassava starch as clarifying substances in water purification: a case study in Colombia. *Heliyon* **6** (6), e04296. doi:10.1016/j.heliyon.2020.e04296.
- Lyons, B. P., Devlin, M. J., Hamid, S. A., Al-Otiabi, A. F., Al-Enezi, M., Massoud, M. S., Al-Zaidan, A. S., Smith, A. J., Morris, S., Bersuder, P., Barber, J. L., Papachlimitzou, A. & Al-Sarawi, H. A. 2015 Microbial water quality and sedimentary faecal sterols as markers of sewage contamination in Kuwait. *Marine Pollution Bulletin* **100** (2), 689–698. doi:10.1016/j.marpolbul.2015.07.043.
- Miravet Sánchez, B. L., García Rivero, A. E., López Del Castillo, P., Alayón García, G. & Salinas Chávez, E. 2016 Water quality of the Ariguanabo river according to physical-chemical indices and bioindicators. *Ingeniería Hidráulica y Ambiental*. Scielo.sdl.cu. Retrieved 12 March 2020, Available from: http://scielo.sld.cu/scielo.php?pid=S1680-03382016000200009&script=sci_arttext&lng=pt
- Moreno, H. S., Bolívar-Anillo, H. J., Soto-Varela, Z. E., Aranguren, Y., González, C. P., Daza, D. A. V. & Anfusio, G. 2019 Microbiological water quality and sources of contamination along the coast of the Department of Atlántico (Caribbean Sea of Colombia). Preliminary results. *Marine Pollution Bulletin* **142**, 303–308. doi. 10.1016/j.marpolbul.2019.03.054.
- Moresco, V., Viancelli, A., Nascimento, M. A., Souza, D. S. M., Ramos, A. P. D., Garcia, L. A. T., Simoes, C. M. O. & Barardi, C. R. M. 2012 Microbiological and physicochemical analysis of the coastal waters of southern Brazil. *Marine Pollution Bulletin* **64** (1), 40–48. doi:10.1016/j.marpolbul.2011.10.026.
- Noble, R. T., Moore, D. F., Leecaster, M. K., McGee, C. D. & Weisberg, S. B. 2003 Comparison of total coliform, fecal coliform, and enterococcus bacterial indicator response for ocean recreational water quality testing. *Water Research* **37** (7), 1637–1643. doi:10.1016/S0043-1354(02)00496-7.
- Payares, B. M. B., Villasmil, K. J. F., Matos, L. C. R., Larreal, A. G. Á., Barboza, Y. & Levy, A. 2013 Calidad microbiológica del agua potable envasada en bolsas y botellas que se venden en la ciudad de Maracaibo, estado Zulia-Venezuela. (Physicochemical and microbiological quality of the

- swimming pool water of two recreational complexes in Zulia State) *Multiciencias*. Redalyc.org. Retrieved 24 February 2020, Available from: <https://www.redalyc.org/pdf/904/90428348002.pdf>
- Prasad, V. R., Srinivas, T. N. R. & Sarma, V. V. S. S. 2015 Influence of river discharge on abundance and dissemination of heterotrophic, indicator and pathogenic bacteria along the east coast of India. *Marine Pollution Bulletin* **95** (1), 115–125. doi:10.1016/j.marpolbul.2015.04.032.
- Praveena, S. M., Chen, K. S. & Ismail, S. N. S. 2013 Indicators of microbial beach water quality: preliminary findings from Teluk Kemang beach, Port Dickson (Malaysia). *Marine Pollution Bulletin* **76** (1–2), 417–419. doi:10.1016/j.marpolbul.2013.08.028.
- Pulido, M. D. P. A., de Navia, S. L. Á., Torres, S. M. E. & Prieto, A. C. G. 2005 Indicadores microbiológicos de contaminación de las fuentes de agua. (Microbiological indicators of contamination of water sources). *Nova* **3** (4), 69–79. doi:10.22490/24629448.338.
- Romero, A. M., Gómez, K. F., Sánchez, J. O. & García-Luján, C. 2009 Monitoring of the microbiological quality of water in the Río Nazas watershed, Mexico. *Química Viva*. Redalyc.org. Retrieved 26 March 2020, Available from: <https://www.redalyc.org/pdf/863/86311258005.pdf>
- Romeu-Álvarez, B., Larrea-Murrell, J., Lugo-Moya, D., Rojas-Hernández, N. & Heydrich-Pérez, M. 2012 Calidad microbiológica de las aguas del río Luyanó, La Habana, Cuba. (Microbiological qualities of the waters of the Luyanó River, Havana, Cuba) *Revista CENIC. Ciencias Biológicas*. Redalyc.org. Retrieved 2 March 2020, Available from: <https://www.redalyc.org/pdf/1812/181226874006.pdf>
- Rusiñol, M., Fernandez-Cassi, X., Timoneda, N., Carratalà, A., Abril, J. F., Silvera, C., Figueras, M. J., Gelati, E., Rodo, X., Kay, D., Wyn-Jones, P., Bofill-Mas, S. & Girones, R. 2015 Evidence of viral dissemination and seasonality in a Mediterranean river catchment: implications for water pollution management. *Journal of Environmental Management* **159**, 58–67. doi:10.1016/j.jenvman.2015.05.019.
- Samboni Ruiz, N. E., Carvajal Escobar, Y. & Escobar, J. C. 2007 A review of physical-chemical parameters as water quality and contamination indicators. *Ingeniería e Investigación*. Scielo.org.co. Retrieved 19 February 2020, Available from: http://www.scielo.org.co/scielo.php?script=sci_arttext&pid=s0120-56092007000300019
- Silva-Iñiguez, N., Corona, C. G. G., Miramontes, L. G. & Mendoza, A. L. 2007 The impact of tourist activity on the bacteriological quality of seawater. *Gaceta Ecológica* **82**, 69–76. Available from: <https://www.redalyc.org/pdf/539/53908206.pdf>
- Skibinski, B., Uhlig, S., Müller, P., Slavik, I. & Uhl, W. 2019 Impact of different combinations of water treatment processes on the concentration of disinfection byproducts and their precursors in swimming pool water. *Environmental Science & Technology* **53** (14), 8115–8126. doi:10.1021/acs.est.9b00491.
- Sorensen, J. P. R., Lapworth, D. J., Marchant, B. P., Nkhuwa, D. C. W., Pedley, S., Stuart, M. E., Bell, R. A., Chirwa, M., Kabika, J., Liemisa, M. & Chibesa, M. 2015 In-situ tryptophan-like fluorescence: a real-time indicator of faecal contamination in drinking water supplies. *Water Research* **81**, 38–46. doi. 10.1016/j.watres.2015.05.035.
- Sorensen, J. P. R., Vivanco, A., Ascott, M. J., Goody, D. C., Lapworth, D. J., Read, D. S., Rushworth, C. M., Bucknall, J., Herbert, K., Karapanos, I., Gumm, L. P. & Taylor, R. G. 2018 Online fluorescence spectroscopy for the real-time evaluation of the microbial quality of drinking water. *Water Research* **137**, 301–309. doi. 10.1016/j.watres.2018.03.001.
- Van Nevel, S., Koetzsch, S., Proctor, C. R., Besmer, M. D., Prest, E. I., Vrouwenvelder, J. S., Knezev, A., Boon, N. & Hammes, F. 2017 Flow cytometric bacterial cell counts challenge conventional heterotrophic plate counts for routine microbiological drinking water monitoring. *Water Research* **113**, 191–206. doi:10.1016/j.watres.2017.01.065.
- Vergaray, G., Méndez, C. R., Morante, H. Y., Heredia, V. I. & Béjar, V. R. 2007 *Enterococcus* y *Escherichia coli* como indicadores de contaminación fecal en playas costeras de Lima (Enterococcus and Escherichia coli as indicators of fecal contamination in coastal beaches of Lima). *Revista del Instituto de Investigación de la Facultad de Ingeniería Geológica, Minera, Metalúrgica Y Geográfica* **10** (20), 82–86. doi:10.15381/iigeo.v10i20.498.
- Wade, C., Otero, E., Poon-Kwong, B., Rozier, R. & Bachoon, D. 2015 Detection of human-derived fecal contamination in Puerto Rico using carbamazepine, HF183 Bacteroides, and fecal indicator bacteria. *Marine Pollution Bulletin* **101** (2), 872–877. doi:10.1016/j.marpolbul.2015.11.016.
- Wong, M., Kumar, L., Jenkins, T. M., Xagorarakis, I., Phanikumar, M. S. & Rose, J. B. 2009 Evaluation of public health risks at recreational beaches in Lake Michigan via detection of enteric viruses and a human-specific bacteriological marker. *Water Research* **43** (4), 1137–1149. doi. 10.1016/j.watres.2008.11.051.
- World Health Organization 2000 *Guidelines for Safe Recreational Water Environments*. WHO, Geneva, Switzerland. Available from: http://www.who.int/water_sanitation_health/bathing/recreaII-ch1.pdf?ua=1 (accessed 15 July 2019).
- World Health Organization 2003 *Hazard Characterization of Pathogens in Food and Water*. WHO, Geneva, Switzerland. Available from: <http://www.fao.org/3/a-at660s.pdf> (accessed 20 May 2019).
- World Health Organization 2011 *Guidelines for Drinking Water Quality*. WHO, Geneva, Switzerland. Available from: http://www.who.int/water_sanitation_health/dwq/gdwq3_es_full_lowres.pdf?ua=1 (accessed 15 July 2019).
- World Health Organization 2012 *Manual for the Inspection of Ships and Issuance of Health Certificates on Board*. WHO, Geneva, Switzerland. Available from: https://apps.who.int/iris/bitstream/handle/10665/44835/9789243548197_spa.pdf (accessed 15 July 2019).