

CONTEMPORARY REVIEW

Role of Macrophages in Acute Lung Injury and Chronic Fibrosis Induced by Pulmonary Toxicants

Debra L. Laskin,^{*,1} Rama Malaviya,^{*} and Jeffrey D. Laskin[†]^{*}Department of Pharmacology and Toxicology, Ernest Mario School of Pharmacy; and [†]Department of Environmental and Occupational Health, School of Public Health, Rutgers University, Piscataway, New Jersey¹To whom correspondence should be addressed at Department of Pharmacology and Toxicology, Ernest Mario School of Pharmacy, Rutgers University, 160 Frelinghuysen Road, Piscataway, NJ 08854. Fax: 1-732-445-0119. E-mail: laskin@eohsi.rutgers.edu.**ABSTRACT**

A diverse group of toxicants has been identified that cause injury to the lung including gases (eg, ozone, chlorine), particulates/aerosols (eg, diesel exhaust, fly ash, other combustion products, mustards, nanomaterials, silica, asbestos), chemotherapeutics (eg, bleomycin), and radiation. The pathologic response to these toxicants depends on the dose and duration of exposure and their physical/chemical properties. A common response to pulmonary toxicant exposure is an accumulation of proinflammatory/cytotoxic M1 macrophages at sites of tissue injury, followed by the appearance of anti-inflammatory/wound repair M2 macrophages. It is thought that the outcome of the pathogenic responses to toxicants depends on the balance in the activity of these macrophage subpopulations. Overactivation of either M1 or M2 macrophages leads to injury and disease pathogenesis. Thus, the very same macrophage-derived mediators, released in controlled amounts to destroy injurious materials and pathogens (eg, reactive oxygen species, reactive nitrogen species, proteases, tumor necrosis factor α) and initiate wound repair (eg, transforming growth factor β , connective tissue growth factor, vascular endothelial growth factor), can exacerbate acute lung injury and/or induce chronic disease such as fibrosis, chronic obstructive pulmonary disease, and asthma, when released in excess. This review focuses on the role of macrophage subsets in acute lung injury and chronic fibrosis. Understanding how these pathologies develop following exposure to toxicants, and the contribution of resident and inflammatory macrophages to disease pathogenesis may lead to the development of novel approaches for treating lung diseases.

Key words: Macrophages; lung injury; fibrosis; inflammatory mediators; cytokines; oxidants.

The idea that macrophages accumulating at sites of injury play a role in the pathogenic response to xenobiotics was proposed more than 100 years ago by Eli Metchnikoff. Describing the inflammatory response as a “salutary reaction against some injurious influence,” he hypothesized that “ferments” released by cells at inflammatory sites might contribute to tissue damage (Metchnikoff, 1968). Since that time, there have been numerous publications supporting this concept in all tissues of the body. In this review, we focus on the role of macrophages and inflammatory mediators they release in acute and chronic lung injury and disease pathogenesis induced by pulmonary toxicants. For

brevity, we have not included discussion of the impact of exposure dose, duration or frequency, animal species, or mechanism of action. Although we recognize that these factors influence the inflammatory response, herein we present a general overview that may be more broadly applicable to multiple toxicants with distinct mechanisms of action.

INFLAMMATORY MACROPHAGES

Inflammatory macrophages are mononuclear phagocytes that play an essential role in host defense and in innate immune




Figure 1. Proinflammatory/cytotoxic (M1) and anti-inflammatory/wound repair (M2) macrophages in acute injury and chronic fibrosis. Immature inflammatory mouse macrophages originating from blood and precursors express the integrin CD11b, high or low levels of the surface antigen Ly6C, and chemokine receptors, CCR2 or CX3CR1, respectively. In response to environmental cues (eg, cytokines, growth factors, TLR agonists, and lipids) and intracellular/nuclear regulatory pathways (eg, kinases, transcription factors, epigenetic regulators, metabolic factors) that are activated, these CD11b⁺ cells mature into F4/80⁺Ly6C^{hi}CCR2⁺ proinflammatory/cytotoxic M1 macrophages, which express markers such as inducible nitric oxide synthase (iNOS) and ARL11, or F4/80⁺Ly6C^{lo}CX3CR1⁺ anti-inflammatory/wound repair M2 macrophages which express CD68, CD163, CD206, Arginase (Arg-1), Ym1, and/or Fizz1. The extent to which macrophages develop into these subpopulations depends on the activities of M1 and M2 inducers/regulatory factors. As these vary over the course of the inflammatory response, macrophages exist along a continuum with subpopulations expressing varying levels of M1 and M2 markers and activities. The process of M1 and M2 macrophage activation is also highly dynamic; thus, as environmental cues, signaling molecules, transcription factors, and cellular metabolism change, macrophages readily modify their phenotype and function. The outcome of inflammatory responses to tissue injury depends on the relative activities of M1 and M2 macrophage subpopulations. In this context, overactivation of M1 macrophages and/or aberrant M2 anti-inflammatory/wound repair activity can exacerbate and perpetuate acute injury, while excessive activity of M2 macrophages can lead to chronic diseases such as fibrosis.

responses to noxious stimuli. In contrast to resident tissue macrophages (eg, alveolar macrophages), which originate mainly from embryonic precursors during development and self-renew during adulthood, inflammatory macrophages are largely derived from blood and bone marrow precursors. They express specific chemokine receptors such as CCR2 or CX3CR1 and accumulate in tissues in response to chemokines released from injured cells and tissues (Tsou et al., 2007; Vannella and Wynn, 2017). Inflammatory macrophages also express the integrin CD11b, a marker of migratory cells. Once localized at sites of tissue injury, these newly recruited macrophages become activated by mediators they encounter in the tissue microenvironment developing into subpopulations exhibiting varying levels of pro-inflammatory/cytotoxic (M1) or anti-inflammatory/wound repair (M2) activity (Arora et al., 2018; Hussell and Bell, 2014; Laskin et al., 2011) (Figure 1). The process of macrophage activation into M1 and M2 subsets is tightly controlled and involves specific signaling pathways and transcription factors, and posttranslational regulatory networks (Lawrence and Natoli, 2011; Wang et al., 2014a) (Table 1).

Proinflammatory/cytotoxic M1 macrophages, also referred to as classically activated macrophages, develop in response to interferon (IFN) γ alone, or in concert with toll-like receptor (TLR) 4 agonists (eg, lipopolysaccharide [LPS]; high-mobility group protein [HMGB1]) or other cytokines (eg, tumor necrosis factor [TNF] α and granulocyte macrophage-colony stimulating factor [GM-CSF]). Activated M1 macrophages release proinflammatory cytokines (eg, TNF α , interleukin [IL]-1 β , IL-6, IL-12, IL-15, IL-23), and generate cytotoxic reactive oxygen species (ROS) and reactive nitrogen species (RNS), proteolytic enzymes, and bioactive lipids. The activity of M1 macrophages is balanced by M2 macrophages, which downregulate inflammation and initiate

wound repair. These actions are mediated by anti-inflammatory cytokines (eg, IL-4, IL-10, IL-13), pro-resolving eicosanoids (eg, lipoxins, resolvins), and growth factors (eg, transforming growth factor [TGF] β , vascular endothelial growth factor [VEGF], epidermal growth factor [EGF], connective tissue growth factor [CTGF], fibroblast growth factor [FGF], platelet-derived growth factor [PDGF]). M2 macrophages have been subdivided into M2a macrophages, activated by IL-4 and IL-13, M2b macrophages, activated by immune complexes and LPS, M2c macrophages, activated by IL-10, TGF β , or glucocorticoids, and M2d macrophages, activated by IL-6 and adenosines (Roszer, 2015). Although acute lung injury and persistent inflammation involves a prolonged or exaggerated response of M1 macrophages and defective M2 macrophage-mediated tissue repair, the development of chronic diseases such as fibrosis and cancer is thought to be a consequence of hyper-responsive subpopulations of M2 macrophages.

Although originally considered two phenotypically and functionally distinct subpopulations at opposite ends of the activation spectrum, it is now recognized that macrophages exist on a continuum with subpopulations in between expressing varying levels of M1 and M2 markers and activities. Moreover, the extent of macrophage activation toward an M1 or M2 phenotype is regulated not only by environmental cues including cytokines, growth factors, TLR agonists, and lipids, but also by various signaling pathways (ie, kinases), downstream transcription factors (eg, nuclear factor-kappa B [NF- κ B], activator protein [AP]-1, interferon regulatory factor [IRF]s, signal transducer and activator of transcription [STAT]s), peroxisome proliferator-activated receptor [PPAR] γ , and epigenetic regulators (eg, microRNAs, chromatin modifiers) that are activated within the cells, as well as by posttranslational regulatory networks, and changes in

Table 1. Regulators of Macrophage Activation and Polarization

Regulator	M1	M2	Supplementary References
Extracellular			
Cytokines	IL-1 β , IL-6, IL-12, IL-23, IFN γ , TNF α , MIP-1 α	IL-4, IL-13, IFN α , IL-1 RA	Martinez and Gordon (2014), Amici et al. (2017), Murray (2017), Parisi et al. (2018)
Growth factors	GM-CSF	M-CSF, TGF β	Martinez and Gordon (2014), Wynn and Vannella (2016)
Eicosanoids/bioactive lipids	LTB $_4$; 12-HETE, 5-HETE	Lipoxins, resolvins, thromboxane, PGI $_2$	Masoodi et al. (2015), Robb et al. (2016), Amici et al. (2017)
TLR-4 agonists	DAMPs, PAMPs, LPS		Martinez and Gordon (2014)
NOD agonists	Peptidoglycans		Zhou et al. (2015)
Intracellular			
Oxidative stress	ROS, RNS		Martinez and Gordon (2014), Amici et al. (2017), Murray (2017), Parisi et al. (2018)
Metabolism	Anaerobic glycolysis, glucose uptake, fatty acid synthesis	Oxidative glucose metabolism, fatty acid oxidation, and uptake	Jha et al. (2015), Murray (2017); Van den Bossche et al. (2017); Van den Bossche and Saraber (2018)
Signaling pathways	AKT2, NOTCH1/2	AKT1, RTKs	Amici et al. (2017)
Nuclear			
Transcription factors	NF κ B, AP-1, STAT1, IRF1, IRF5, IRF8, HIF-1 α	IRF4, KLF-4, c-myc, PPAR γ , RXRs, LXRs, STAT3, STAT6, IRF3, IRF4, HIF2 α	Escribese et al. (2012), Pello et al. (2012), Martinez and Gordon (2014); Roszer (2015), Chistiakov et al. (2018), Parisi et al. (2018), Saradna et al. (2018)
Epigenetic			
MicroRNA	miRNA-155, miRNA-125b, miRNA-27b, miRNA-127, miRNA-223, miRNA-106a	miRNA-146a/b, miRNA-21, miRNA-511-3p, miRNA-124, miRNA-125a/b, miRNA-24, miRNA-34a, let-7c	Ponomarev et al. (2011), Amici et al. (2017), Parisi et al. (2018), Saradna et al. (2018), Shapouri-Moghaddam et al. (2018)
Histone modifications	HDAC3		
DNA Methylation	DNMT3b, DNMT1	H3K27 demethylase, DNMT3a, DNMT3al	

Abbreviations: AKT, protein kinase B; AP, activator protein; DAMP, damage-associated molecular pattern; DNMT, DNA methyltransferase; GM-CSF, granulocyte-macrophage colony stimulating factor; HDAC, histone deacetylase; HETE, hydroxyeicosatetraenoic acid; HIF, hypoxia-inducible factor; IFN, interferon; IL, interleukin; IRF, interferon regulatory factor; KLF, Kruppel-like factors; LPS, lipopolysaccharide; LT, leukotriene; LXR, liver X receptor; M-CSF, macrophage colony stimulating factor; MIP, macrophage inflammatory protein; miRNA, micro-RNA; NF- κ B, nuclear factor-kappa B; NOTCH, neurogenic locus notch homolog protein; PAMP, pathogen-associated molecular pattern; PG, prostaglandin; PPAR, peroxisome proliferator-activated receptor; RNS, reactive nitrogen species; ROS, reactive oxygen species; RTK, receptor tyrosine kinase; RXR, retinoid X receptor; STAT, signal transducer and activator of transcription; TGF, transforming growth factor; TNF, tumor necrosis factor.

intracellular metabolism (Murray, 2017) (Table 1). It should be noted, however, that M1 and M2 macrophage activation is a highly dynamic process; thus, as environmental cues, signaling molecules, transcription factors, epigenetic regulators, and cellular metabolism change in response to pathophysiological conditions, macrophages readily modify their phenotype and function. Accordingly, cells that initially promote proinflammatory and cytotoxic reactions, can undergo phenotypic switching, subsequently participating in the resolution of inflammation and injury (Porcheray et al., 2005; Parisi et al., 2018). In this context, the transition of macrophages from a proinflammatory M1 to an anti-inflammatory/wound repair M2 phenotype appears to be critical for the progression of normal wound healing and tissue regeneration. This is exemplified by findings that noxious pulmonary stimuli such as hypoxia impair M1 to M2 macrophage phenotypic switching, a response associated with prolonged tissue injury (Faulknor et al., 2017). Also of importance, is the observation that while there are some unique characteristics of the activation profile of M1 and M2 macrophages, there are also many overlapping activities. These findings underscore the dynamic nature of the macrophage activation process and the notion that within any inflamed tissue, mixed phenotype macrophages co-exist with M1 and M2 macrophages, their specific function depending on the

balance of activating and inhibiting activities and the tissue microenvironment (Martinez and Gordon, 2014).

INFLAMMATORY MACROPHAGES IN ACUTE LUNG INJURY AND REPAIR

Acute injury to the lung is associated with disruption of endothelial and epithelial barriers (Muller-Redetzky et al., 2014; Tam et al., 2011). It is characterized by an accumulation of protein-rich edema fluid, sloughing of the bronchial epithelium, the appearance of necrotic or apoptotic type I cells, denuded basement membrane, enlarged edematous interstitium, injured endothelial cells, and an accumulation of cellular debris in the tissue. Cellular characteristics include loss of alveolar-capillary membrane integrity, transepithelial migration of neutrophils and macrophages, and increases in proinflammatory/cytotoxic proteins. The outcome of acute lung injury depends on the nature of the toxicant, the dose and duration of exposure and the specific tissue location. Although after some exposures, lung structure and function return to normal, in other instances, there is persistence and/or progression of injury, leading to pulmonary vascular destruction, fibrosing alveolitis, multiple organ failure and death.

Initial evidence suggesting a role of macrophages in acute lung injury was based on findings that numbers of these cells increased in the tissue following exposure of animals to diverse pulmonary toxicants (eg, ozone, particulate matter, bleomycin, endotoxin, nanomaterials, mustard vesicants, silica, asbestos, and radiation) (Bekki *et al.*, 2016; Hiraiwa and van Eeden, 2013; Laskin *et al.*, 2011; Weinberger *et al.*, 2011). In addition, macrophages accumulating in the lung early after injury induced by pulmonary toxicants were found to be activated toward a proinflammatory M1 phenotype, as evidenced by morphologic changes including increased size and vacuolization, and expression of inducible nitric oxide synthase (iNOS) and TNF α , prototypical marks of proinflammatory/cytotoxic macrophages (Aggarwal *et al.*, 2014; Laskin *et al.*, 2011; Malaviya *et al.*, 2017; Sugiura and Ichinose, 2011). Macrophages responding to toxicant-induced injury also generated increased quantities of cytotoxic and proinflammatory mediators, which induce and/or amplify the pathogenic response, including ROS, RNS, IL-1, IL-6, IL-18, TNF α , chemokines, eicosanoids, proteases, and bioactive lipids.

Direct evidence for a role of proinflammatory/cytotoxic M1 macrophages in acute lung injury comes from findings that tissue damage is directly correlated with macrophage functional status. Thus, in a number of experimental models using a variety of approaches (eg, pharmacologic, genetic, or macrophage-deficient mice), pulmonary damage is ameliorated or prevented by suppressing or depleting macrophages (Table 2). For example, when M1 macrophage cytotoxic/proinflammatory activity is blocked with anti-inflammatory steroids, lung damage induced by ozone, silica, residual oil fly ash, bleomycin, sulfur mustard, endotoxin, oleic acid, or hydrogen sulfide is reduced (Al-Harbi *et al.*, 2016; Chen *et al.*, 2006; Geng *et al.*, 2018; Huang *et al.*, 2014; Laskin *et al.*, 2011; Samet *et al.*, 2000; Wigenstam *et al.*, 2009). Similarly, the accumulation of macrophages in the lung and subsequent toxicity of ozone, cigarette smoke, carbon nanotubes, particulate matter, radiation, and silica are mitigated by pretreatment of animals with gadolinium chloride, which blocks M1 macrophage activation or clodronate liposomes, administered intravenously, which depletes newly recruited inflammatory macrophages (Frank *et al.*, 2015; Laskin *et al.*, 2011; Marchini *et al.*, 2016; Nemmar *et al.*, 2005; Perez-Rial *et al.*, 2013; Poole *et al.*, 2012). Bleomycin-induced acute lung injury and expression of iNOS are also abrogated in CCR4 knockout mice, which cannot generate M1 macrophages (Trujillo *et al.*, 2008). Similar protective effects against dust have been described in IL-18R knockout mice, and against ozone in TNFR1 and galectin-3 knockout mice, which also display defective M1 macrophage development (Bauer *et al.*, 2013; Laskin *et al.*, 2011; Sunil *et al.*, 2015). Production of inflammatory mediators and lung injury induced by endotoxin are also reduced in mice lacking CD40, a cell surface receptor required for M1 macrophage activation and iNOS expression (Hashimoto *et al.*, 2004). Mice deficient in TLR4, a key receptor/signaling pathway regulating M1 inflammatory macrophage activation, are also protected from toxicity induced by ozone, endotoxin, and particulate matter (Connor *et al.*, 2012; Hollingsworth *et al.*, 2004). Likewise, loss of downstream TLR4 signaling molecules including MyD88 or NF- κ B, protects animals from lung injury induced by ozone, bleomycin, silica, and particulate matter (He *et al.*, 2016; Laskin *et al.*, 2011; Re *et al.*, 2014).

Protection against damage induced by ozone has also been described in animals treated with cyclophosphamide, which depletes bone marrow-derived monocytic precursors (Bhalla *et al.*, 1992), consistent with the origin of proinflammatory

macrophages. Analogous protection against LPS-induced lung toxicity, has been observed in CD11b-diphtheria toxin (DT) receptor mice, depleted of blood monocyte precursors by administration of DT (Dhaliwal *et al.*, 2012; Lu *et al.*, 2018). CCR2 knockout mice, which are defective in M1 macrophage migratory activity to sites of injury, are also protected from oxidative stress and tissue injury induced by ozone, radiation, bleomycin, and endotoxin (Francis *et al.*, 2017; Gharaee-Kermani *et al.*, 2003; Laskin *et al.*, 2011; Wiesemann *et al.*, 2018; Yang *et al.*, 2010).

Additional support for proinflammatory M1 macrophage involvement in the pathogenesis of acute lung injury comes from findings that activation of these cells exacerbates tissue damage induced by pulmonary toxicants. Hence, pretreatment of rodents with macrophage activators such as LPS or *Bacillus Calmette-Guerin* augments acute lung injury induced by endotoxin, radiation, bleomycin, and ozone (Laskin *et al.*, 2011). Similar increases in toxicity in response to radiation, ozone, and bleomycin have been observed in mice lacking the pulmonary collectin, surfactant protein D, which under homeostatic conditions, functions to suppress lung macrophage proinflammatory activity (Casey *et al.*, 2005; Groves *et al.*, 2012; 2013; Malaviya *et al.*, 2015a).

The resolution of acute injury and the return to normal lung structure and function is an active coordinated process. Evidence suggests that it is largely mediated by M2 macrophages, which stimulate counter-regulatory mechanisms that suppress the release of proinflammatory mediators and activate tissue repair processes. Anti-inflammatory/wound repair M2 macrophages have been reported to increase in the lung following exposure of animals to ozone, mustards, bleomycin, particulate matter, radiation, endotoxin, cigarette smoke, silica and asbestos (Arora *et al.*, 2018; Duru *et al.*, 2016; Francis *et al.*, 2017; Kambara *et al.*, 2015; Laskin *et al.*, 2011; Malaviya *et al.*, 2015b; Murthy *et al.*, 2015; Venosa *et al.*, 2016; Xiang *et al.*, 2016). However, their appearance is delayed relative to M1 macrophages, consistent with a role for these cells in tissue repair. These macrophages express prototypical M2 markers including CD68, CD163, CD206, Ym-1, Arg-1, and Fizz-1.

Increases in M2 macrophages in the lung following exposure to ozone, endotoxin, mustard, diesel exhaust, asbestos, silica, or hyperoxia are correlated with upregulation of IL-4, IL-10, and IL-13 (Hussell and Bell, 2014; Laskin *et al.*, 2011; Venosa *et al.*, 2016). These cytokines dampen macrophage production of proinflammatory/cytotoxic mediators and stimulate the generation of extracellular matrix proteins and growth factors important in wound healing (Wynn and Vannella, 2016). Notably, administration of IL-13, which is key for M2 macrophage development, protects mice from lethal endotoxemia, whereas anti-IL-13 antibodies significantly reduce survival of these mice (Matsukawa *et al.*, 2000). Additionally, cytotoxicity and mortality are increased in response to hyperoxia in IL-13 null mice (Bhandari *et al.*, 2007). Increases in susceptibility to ozone have also been described in mice deficient in IL-10, a potent anti-inflammatory cytokine known to activate M2 macrophages (Backus *et al.*, 2010). The importance of M2 macrophages in suppressing inflammation and initiating wound repair is also evidenced by findings that depletion of these cells by administration of DT to CD206-DT receptor transgenic mice is associated with increased expression of proinflammatory cytokines including TNF α , IL-1 β , IL-6, and MCP-1/CCL2 following endotoxin administration, a response correlated with exacerbation of lung inflammation and injury (Kambara *et al.*, 2015).

The origin of M2 macrophages has not been clearly established. The fact that their appearance in injured tissues is, like

M1 macrophages, dependent in large part on CCR2/CCL2, suggests that they are derived from bone marrow and monocyte precursors. However, whether they originate from M1 macrophages via phenotypic switching or from a separate monocytic precursor pool is unclear. In this regard, recent studies have demonstrated an accumulation of bone marrow-derived macrophages in the lung following ozone inhalation which co-express the M2 macrophage chemokine receptor, CX3CR1, and CD206 (Francis *et al.*, 2017). It has also been suggested that at least some populations of M2 repair macrophages are derived from proliferating resident lung macrophages (Alber *et al.*, 2012; Tighe *et al.*, 2011). In this regard, while both monocyte and tissue-derived M2 macrophages express Arg1, Ym-1, and Fizz-1, only monocyte-derived M2 macrophages express CX3CR1, CD206, Raldh2, and PD-L2 (Gundra *et al.*, 2014).

INFLAMMATORY MACROPHAGES AND PULMONARY FIBROSIS

Chronic pulmonary fibrosis refers to a range of disorders characterized by irreversible destruction and remodeling of lung architecture that occurs as a consequence of excess deposition of collagen and other extracellular matrix components. This leads to scarring of the airways and difficulty in breathing. Evidence suggests that dysregulated wound repair is a key factor contributing to pulmonary fibrosis, and that this is due, at least in part, to an imbalance in the actions of M1 and M2 macrophages after prolonged inflammation or pneumonitis (Alber *et al.*, 2012; Byrne *et al.*, 2016; Herold *et al.*, 2011) (Figure 1). This imbalance is associated with reduced production of antifibrotic cytokines (eg, CXCL10) and MMPs by M1 macrophages, which promote the resolution of scarring and matrix degradation, and excessive release of profibrotic mediators (eg, TGF β , CTGF) by M2 macrophages, which induce fibroproliferative tissue remodeling.

The importance of overactive macrophages in the development of pulmonary fibrosis has been recognized since the mid-1980s, and macrophage phenotype has more recently emerged as an important contributor to the process (Laskin *et al.*, 2011). This is largely based on findings that macrophages are persistently increased in the lung in close proximity to collagen producing fibroblasts, during the aberrant wound healing phase of fibrogenesis in animals treated with bleomycin, asbestos, silica, mustards, carbon nanotubes, or radiation (Duke and Bonner, 2018; Huang *et al.*, 2017; Laskin *et al.*, 2011; Malaviya *et al.*, 2015b; Murthy *et al.*, 2015), and that these macrophages express markers of an anti-inflammatory/profibrotic M2 phenotype including Arg-1, Fizz-1, Ym-1, CD68, CD163, and/or mannose receptor (CD206). In a number of these experimental models, macrophages are also enlarged and foamy in appearance, a characteristic feature of a profibrotic phenotype (Romero *et al.*, 2015; Venosa *et al.*, 2016). Increases in M2 macrophages in areas of fibrotic tissue have similarly been noted in lungs of patients with idiopathic pulmonary fibrosis, chronic obstructive pulmonary disease and cystic fibrosis; moreover, number of these cells in the tissue correlate directly with a worsening prognosis.

M2 macrophages accumulating in the lung during fibrogenesis in both humans and rodents have been identified as the major source of a number of key profibrotic mediators (eg, TGF β , CTGF, and CCL18), which stimulate fibroblast proliferation and collagen synthesis (Table 3) (Bickelhaupt *et al.*, 2017; Prasse *et al.*, 2006; Pulichino *et al.*, 2008; Shvedova *et al.*, 2005). M2 macrophages are also thought to promote fibrosis through the release

of TNF α , IL-1, IL-10, IL-13, IL-33, PDGF, FGF, and fibronectin, which are increased in patients and animals with fibrotic lung disease (Wynn and Vannella, 2016). Arginase, which favors polyamine and proline biosynthesis and promotes cell growth and collagen formation, is also upregulated in M2 macrophages localized in fibrotic lesions (Murthy *et al.*, 2015).

The strongest evidence supporting a role of hyperactive M2 macrophages in the development of pulmonary fibrosis comes from findings that the response to fibrogenic toxicants (eg, bleomycin, radiation, silica, carbon nanotubes) is exacerbated in animals overexpressing IL-10 or IL-13, or by exogenous administration of IL-33, which promote M2 macrophage activation (Table 3) (Fulkerson *et al.*, 2006; Laskin *et al.*, 2011; Li *et al.*, 2014; Lumsden *et al.*, 2015; Luzina *et al.*, 2013; Wang *et al.*, 2014b). Conversely, collagen production and fibrosis are reduced in animals lacking M-CSF or those treated with chlodronate liposomes or anti-CSF1R antibody, which deplete M2 macrophages (Baran *et al.*, 2007; Gibbons *et al.*, 2011; Meziani *et al.*, 2018; Murray *et al.*, 2011), or serum amyloid P, which inhibits the development of M2 macrophages in the lung (Laskin *et al.*, 2011; Murray *et al.*, 2011; Pilling *et al.*, 2007), and in mice deficient in the M2 macrophage-inducing cytokines IL-4, IL-10, IL-13, IL-18, IL-33, or their receptors (Hoshino *et al.*, 2009; Huaux *et al.*, 1998, 2003; Laskin *et al.*, 2011; Li *et al.*, 2014; Liu *et al.*, 2004; Lumsden *et al.*, 2015; Wang *et al.*, 2014b; Zhao *et al.*, 2018), chemokine receptor CCR4 (Trujillo *et al.*, 2008), Fizz-1, which induces M2 macrophage recruitment and myofibroblast differentiation (Liu *et al.*, 2014), or C/EBP homologous protein, which is required for M2 macrophage activation and TGF β production (Buck and Chojkier, 2011; Yao *et al.*, 2016). A role for IL-1R1 and MyD88 signaling has also been described in bleomycin-induced lung fibrosis (Gasse *et al.*, 2007).

There is also some experimental evidence that proinflammatory monocyte-derived M1 macrophages drive pulmonary fibrosis (Byrne *et al.*, 2016; Gibbons *et al.*, 2011). Thus, prednisone is effective in reducing TGF β production and collagen deposition in bleomycin-induced fibrosis, but only if it is administered beginning one day after bleomycin administration, a time associated with a prominent proinflammatory/cytotoxic M1 macrophage response (Chaudhary *et al.*, 2006). Additionally, selective depletion of circulating proinflammatory monocytes, defined in mice by expression of high levels of the surface marker Ly6C (Ly6C^{hi}), leads to decreases in M2 macrophages in the lung, and abrogation of bleomycin-induced pulmonary fibrosis; moreover, adoptive transfer of these cells during fibrogenesis exacerbates fibrosis (Gibbons *et al.*, 2011; Ji *et al.*, 2016). Studies with mice lacking CCR2 or CCL2, or wild-type mice treated with MCP-1 (CCL2) neutralizing antibodies, which have blunted M1-monocyte-derived macrophage emigration to sites of injury, have confirmed the importance of these cells in the development of fibrosis. Thus, in these experimental models, significantly fewer M2 macrophages are observed in the lung following administration of bleomycin or radiation, a response correlated with reduced TGF β , CTGF, and fibrosis (Baran *et al.*, 2007; Gharaee-Kermani *et al.*, 2003; Gibbons *et al.*, 2011; Groves *et al.*, 2018). In contrast, when proinflammatory monocytes are adoptively transferred into mice during the acute reversible inflammatory response to bleomycin, they have a resolution promoting role (Gibbons *et al.*, 2011). These data demonstrate that there is considerable heterogeneity in lung macrophage functioning during different phases of the fibrogenic process (Misharin *et al.*, 2013).

A clear causal link between inflammatory monocyte-derived lung macrophages and the pathobiology of fibrosis has recently

Table 2. Approaches for Assessing the Role of Macrophages in Acute Lung Injury and Fibrosis

Proinflammatory/cytotoxic M1 macrophages in acute injury			
Approach	M1 Macrophage Suppression	Outcome: Decreased Injury	Supplementary References
Pharmacologic	Glucocorticoids	Ozone, silica, PM, bleomycin, mustard, endotoxin, oleic acid, hydrogen sulfide	Haddad et al. (1995), DiMatteo and Reasor (1997), Samet et al. (2000), Chen et al. (2003b), Chen et al. (2006), Wigenstam et al. (2009), Huang et al. (2014), Wang et al. (2014a), Al-Harbi et al. (2016); Geng et al. (2018)
	Gadolinium chloride	Ozone, hyperoxia, PM, endotoxin	Jankov et al. (2003), Arimoto et al. (2005), Laskin et al. (2011), Duke-Novakovski et al. (2013)
	Chlodronate liposomes	Cigarette smoke, carbon nanotubes, endotoxin, PM, silica	Elder et al. (2004), Nemmar et al. (2005), Laskin et al. (2011), Dhaliwal et al. (2012), Poole et al. (2012), Perez-Rial et al. (2013), Frank et al. (2015), Marchini et al. (2016)
	Cyclophosphamide	Ozone, silica	Bhalla et al. (1992), Bassett et al. (2001), Nemmar et al. (2005)
	Anti-CCR2 antibody CCl ₂ -specific inhibitor	Endotoxin Radiation	Dhaliwal et al. (2012) Wiesemann et al. (2018)
Genetic	CCR4 ^{-/-} mice	Bleomycin	Trujillo et al. (2008)
	IL-18R ^{-/-} mice	PM	Bauer et al. (2013)
	TNF α ^{-/-} or TNFR ^{-/-} mice	Ozone	Cho et al. (2007), Fakhrzadeh et al. (2008b)
	Galectin-3 ^{-/-} mice	Ozone	Sunil et al. (2015)
	CD40 ^{-/-} mice	Endotoxin	Hashimoto et al. (2004)
	TLR2 ^{-/-} and TLR4 ^{-/-} mice	Ozone, endotoxin, PM	Kleeberger et al. (2000, 2001), Gilmour et al. (2004), Hollingsworth et al. (2004), Cho et al. (2005), Williams et al. (2007), Garantziotis et al. (2010), Bauer et al. (2011), Li et al. (2011), Connor et al. (2012); Connor et al. (2013)
	MyD88 ^{-/-} mice	Ozone, PM, bleomycin, silica	Gasse et al. (2007), Williams et al. (2007), Li et al. (2011), Bauer et al. (2013), Re et al. (2014), He et al. (2016a)
	NF- κ Bp50 ^{-/-} mice	Ozone	Fakhrzadeh et al. (2004b)
	CD11b-DT transgenic mice	Endotoxin	Dhaliwal et al. (2012), Lu et al. (2018)
	CCR2 ^{-/-} mice	Ozone, radiation, bleomycin, endotoxin	Moore et al. (2001), Gharaee-Kermani et al. (2003), Maus et al. (2003), Okuma et al. (2004), Yang et al. (2010), Francis et al. (2017), Wiesemann et al. (2018)
Macrophage-deficient mice	IL1R ^{-/-} mice	Bleomycin	Gasse et al. (2007)
	IL13 ^{-/-} mice	Hyperoxia	Bhandari et al. (2007)
Macrophage-deficient mice	MAFIA	Ricin	Lindauer et al. (2009)
Anti-inflammatory/profibrotic M2 macrophages in fibrosis			
Approach	M2 Macrophage Activation	Outcome: Increased Injury	Supplementary References
Pharmacologic	Endotoxin	Endotoxin, radiation, ozone	Wollert et al. (1994), Johnston et al. (2004), Haque et al. (2009)
	<i>Bacillus Calmette-Guerin</i>	Endotoxin, bleomycin	Chyczewska et al. (1993), Tasaka et al. (1995)
	Anti-IL-13 antibody	Endotoxin	Matsukawa et al. (2000)
Genetic	SP-D ^{-/-} mice	Ozone, radiation, bleomycin	Casey et al. (2005), Kierstein et al. (2006), Groves et al. (2012), Groves et al. (2013), Malaviya et al. (2015a)
	IL-10 ^{-/-} mice ^a CD206-DT transgenic mice	Ozone Endotoxin	Backus et al. (2010) Kambara et al. (2015)
Approach	M2 Macrophage Suppression	Outcome: Decreased Fibrosis	Supplementary References
Pharmacologic	Chlodronate liposomes	Bleomycin	Gibbons et al. (2011)
	Anti-CSF-1R antibody	Radiation	Meziani et al. (2018)
	Serum amyloid P	Bleomycin	Pilling et al. (2007), Laskin et al. (2011)

Table 2. (continued)

Anti-inflammatory/profibrotic M2 macrophages in fibrosis

Approach	M2 Macrophage Suppression	Outcome: Decreased Fibrosis	Supplementary References
Genetic	Glucocorticoid	Bleomycin	Chaudhary et al. (2006)
	Liposome-encapsulated spironolactone	Bleomycin	Ji et al. (2016)
	IL-13-recombinant immunotoxin	Silica	Ferreira et al. (2013)
	Anti-IL-33 antibody	Bleomycin	Li et al. (2014)
	IL-4 ^{-/-} or IL-4R ^{-/-} mice	Silica	Huaux et al. (2003), Laskin et al. (2011)
	IL-10 ^{-/-} mice	Silica	Huaux et al. (1998), Barbarin et al. (2004)
	IL-13R α 2 overexpression	Bleomycin	Lumsden et al. (2015)
	IL-33 ^{-/-} or IL-33R (ST2) ^{-/-} mice	Carbon nanotubes, bleomycin	Li et al. (2014), Wang et al. (2014b)
	Fizz-1 ^{-/-} mice	Bleomycin	Liu et al. (2014)
	C/EBP ^{-/-} mice	Bleomycin	Buck and Chojkier (2011), Yao et al. (2016)
	CCR2 or CCL2 ^{-/-} mice ^b	Bleomycin, radiation	Moore et al. (2001), Gharaee-Kermani et al. (2003), Okuma et al. (2004), Baran et al. (2007), Groves et al. (2018)
	CX3CR1 ^{-/-} mice	Bleomycin	Tighe et al. (2011), Ishida et al. (2017)
	CCR4 ^{-/-} mice	Bleomycin	Trujillo et al. (2008)
	IL-18 ^{-/-} or IL18R α ^{-/-} mice	Bleomycin	Hoshino et al. (2009)
MyD88 ^{-/-} mice	Bleomycin	Gasse et al. (2007)	
IL1R ^{-/-} mice	Bleomycin	Gasse et al. (2007)	
M-CSF ^{-/-} mice	Bleomycin	Baran et al. (2007)	
Macrophage-deficient mice	MAFIA	Bleomycin	Redente et al. (2014)
Approach	M2 Macrophage Activation	Outcome: Increased Fibrosis	Supplementary References
Pharmacologic	IL-33	Bleomycin	Luzina et al. (2013), Li et al. (2014)
Genetic	IL-10 over-expressing mice	Silica	Barbarin et al. (2005)
	MyD88 ^{-/-} mice	Silica	Re et al. (2014)
	SP-D ^{-/-} mice	Bleomycin	Casey et al. (2005), Kierstein et al. (2006), Groves et al. (2012, 2013), Malaviya et al. (2015a)

PM, particulate matter (ash, dust, diesel exhaust).

^aM1 macrophage activation secondary to loss of M2 macrophages.

^bIndirect suppression.

Abbreviations: CCL, chemokine ligand; CCR, chemokine receptor; C/EBP, CCAAT/enhancer-binding protein; CSF-1R, colony-stimulating factor-1 receptor; DT, diphtheria toxin; Fizz, found in inflammatory zone; IL, interleukin; LPS, lipopolysaccharide; MAFIA, macrophage Fas-induced apoptosis; M-CSF, macrophage-colony stimulating factor; SP-D, surfactant protein-D; TLR, Toll-like receptor; TNF, tumor necrosis factor; TNFR, tumor necrosis factor receptor.

been established by Misharin et al. (2017). Using novel lineage tracking systems, these investigators showed that lung fibrosis is ameliorated when inflammatory monocytes are driven to necroptosis during their differentiation into macrophages. A similar attenuation of pulmonary fibrosis was observed using an inducible transgenic system, which specifically depletes proinflammatory M1 macrophages (McCubrey et al., 2018). These data provide support for the notion that profibrotic M2 macrophages are largely derived from proinflammatory M1 macrophages. This is in accord with reports that proinflammatory monocytes are also precursors of circulating fibrocytes, which have been shown to be important in pulmonary fibrosis (Gibbons et al., 2011). It should be noted, however, that at present, direct evidence for a precursor relationship between proinflammatory M1-like (Ly6C^{hi}CCR2⁺) monocytes and M2 macrophages in the lung during fibrosis is limited. In this context, it cannot be ruled out that M2 macrophages are derived from anti-inflammatory monocytes, which express low levels of Ly6C (Ly6C^{lo}) and chemokine receptor CX3CR1, and migrate into the lung from the blood (Gibbons et al., 2011). The fact that

reduced number of M2 macrophages are observed in the lungs of bleomycin-treated mice lacking CX3CR1 and that fibrosis is blunted, are consistent with this idea (Ishida et al., 2017). It is also possible that, at least some M2 macrophages originate from proliferating resident alveolar macrophages. This idea is in accord with reports that M2 macrophage accumulation in the lung is reduced in bleomycin-treated CSF-1 knockout mice, which have defective resident macrophage proliferation relative to wild-type mice, a response characterized by reduced CTGF expression and collagen deposition (Baran et al., 2007).

Of note are findings that the M1 macrophage marker, iNOS, as well as the M1-associated chemokine, CXCL10/IP10, are increased in M2-suppressed animals, consistent with the idea that M2 macrophages are important in downregulating M1 macrophage activity, and that it is a balance in the activity between these cell populations that dictates the outcome of the response to fibrogenic stimuli. This is supported by findings that the development of fibrosis in a granulomatous lung disease model is associated with upregulation of arginase-positive M2 macrophages, and that administration of the proinflammatory

Table 3. Macrophage Mediators Implicated in Lung Injury and Fibrosis

Mediator	Toxicant	Supplementary References
Proinflammatory/cytotoxic M1 macrophage mediators		
ROS (eg, superoxide anion, hydrogen peroxide, hydroxide radicals)	Ozone	Fakhrzadeh et al. (2004a), Connor et al. (2012), Sunil et al. (2012), Kumarathasan et al. (2015), Zhu et al. (2016), Francis et al. (2017)
	Radiation	Rabbani et al. (2005), Lee et al. (2015)
	Endotoxin	Dang et al. (2018)
	Bleomycin	Yamazaki et al. (1998), Gu et al. (2015), Zhou et al. (2018)
	PM	Morio et al. (2001), Laskin et al. (2003), Li et al. (2003), Naota et al. (2010), Shvedova et al. (2013), Bekki et al. (2016), He et al. (2017)
	Asbestos	Simeonova and Luster (1995), Fattman et al. (2006), Blake et al. (2007), Nagatomo et al. (2007), He et al. (2011)
	Silica (Talc)	Rimal et al. (2005), Hu et al. (2006), Kaewamatawong et al. (2006), Nagatomo et al. (2006), Langley et al. (2011), Shim et al. (2015)
	Nanoparticles	Shvedova et al. (2005), Chou et al. (2008), Coccini et al. (2012)
	Chemical warfare agents	Malaviya et al. (2010), McGovern et al. (2011), Malaviya et al. (2015b), Lam et al. (2016), Elfsmark et al. (2018)
RNS (eg, NO, peroxydinitrite)	Ozone	Pendino et al. (1995), Kleeberger et al. (2001), Laskin et al. (2001), Fakhrzadeh et al. (2002), Fakhrzadeh et al. (2004b), Malaviya et al. (2010), Sunil et al. (2012)
	Radiation	Nozaki et al. (1997), Malaviya et al. (2015a)
	Endotoxin	Pendino et al. (1993b), Wizemann et al. (1994), Kristof et al. (1998), Moncao-Ribeiro et al. (2011), Tu et al. (2017), Dang et al. (2018)
	Bleomycin	Yamazaki et al. (1998), Gurujeyalakshmi et al. (2000), Chen et al. (2003a), Genovese et al. (2005)
	PM	Morio et al. (2001), Becher et al. (2007), Kumarathasan et al. (2015)
	Asbestos	Tanaka et al. (1998), Dorger et al. (2002), Zeidler and Castranova (2004)
	Silica	Srivastava et al. (2002), Zeidler et al. (2004), Rimal et al. (2005), Cruz et al. (2016)
	Nanoparticles	Coccini et al. (2012), Lee et al. (2012)
	Chemical warfare agents	Martin et al. (2003), Malaviya et al. (2010), Venosa et al. (2016), Weinberger et al. (2016)
	Proinflammatory cytokines and chemokines (eg, TNF α , IL-1, IL-2, IL-6, IL-12, IL-18, CCL2, CCL3, CCL4, CCL5, CCL17, CXCL1, CXCL2, CXCL5, CXCL10)	Ozone
Radiation		Zhang et al. (2008), Lee et al. (2015), Sohn et al. (2015), Li et al. (2017)
Endotoxin		Jordan et al. (2001), Moncao-Ribeiro et al. (2011), Takashima et al. (2014), Tu et al. (2017), Dang et al. (2018)
Bleomycin		Failla et al. (2006), Baran et al. (2007), Gasse et al. (2007), Trujillo et al. (2008), Hoshino et al. (2009), Yu et al. (2015), Wei et al. (2016), Zhou et al. (2018)
PM		Morio et al. (2001), Becher et al. (2007), Gowdy et al. (2008), Churg et al. (2009), Naota et al. (2010), Hiraiwa and van Eeden (2013), Shvedova et al. (2013), Snow et al. (2014), Bekki et al. (2016), He et al. (2016b), He et al. (2017)
Asbestos		Simeonova and Luster (1995), Fisher et al. (2000)
Silica		Srivastava et al. (2002), Zeidler et al. (2004), Barbarin et al. (2005), Rimal et al. (2005), Hu et al. (2006), Langley et al. (2011), Ferreira et al. (2013), Kawasaki (2015), Cruz et al. (2016)
Nanoparticles		Shvedova et al. (2005), Chou et al. (2008), Hsieh et al. (2012), Frank et al. (2015), Sager et al. (2016)
Chemical warfare agents		Sunil et al. (2011), Malaviya et al. (2015b), Venosa et al. (2016), Elfsmark et al. (2018)
Proteases (eg, MMPs, TIMPs, cathepsin)		Ozone
	Radiation	Flehsig et al. (2010), Lee et al. (2015), Li et al. (2017)
	Endotoxin	Franco et al. (2002), Moncao-Ribeiro et al. (2011), Takashima et al. (2014)
	Bleomycin	Yu et al. (2015), Liu et al. (2017)
	PM	Adamson et al. (2003), Zhang et al. (2010), Shimada et al. (2015), Bekki et al. (2016)
	Asbestos	Tan et al. (2006)
	Silica	Langley et al. (2011), Cruz et al. (2016)
	Nanoparticles	Hsieh et al. (2012)
	Chemical warfare agents	Guignabert et al. (2005), Anderson et al. (2009), Malaviya et al. (2010), Sunil et al. (2014), Venosa et al. (2016)
	Hydrogen sulfide	Wang et al. (2014a)
Oleic acid	Yeh et al. (2009)	

Table 3. (continued)

Mediator	Toxicant	Supplementary References
Bioactive lipids (eg, lipid peroxides, PAF, prostaglandins and leukotrienes)	Ozone	Giri <i>et al.</i> (1975), Tan and Bethel (1992), Pendino <i>et al.</i> (1993a), Kaneko <i>et al.</i> (1995), Stevens <i>et al.</i> (1995), Devlin <i>et al.</i> (1996), Kinney <i>et al.</i> (1996), Nakano <i>et al.</i> (2000), Fakhrzadeh <i>et al.</i> (2002), Connor <i>et al.</i> (2012), Sunil <i>et al.</i> (2012), Francis <i>et al.</i> (2017)
	Radiation	Lee <i>et al.</i> (2015)
	Endotoxin	Ermert <i>et al.</i> (2000), Inoue <i>et al.</i> (2004a)
	Bleomycin	Chen <i>et al.</i> (2003a), Chen <i>et al.</i> (2006), Failla <i>et al.</i> (2006)
	PM	Kuhn <i>et al.</i> (1993), Samet <i>et al.</i> (2000), Shvedova <i>et al.</i> (2013), Yanamala <i>et al.</i> (2013), Bekki <i>et al.</i> (2016), Salama <i>et al.</i> (2016)
	Silica	Mohr <i>et al.</i> (1992)
	Nanoparticles	Shvedova <i>et al.</i> (2005), Coccini <i>et al.</i> (2012), Lee <i>et al.</i> (2012)
Chemical warfare agents	Chemical warfare agents	Wigenstam <i>et al.</i> (2009), Malaviya <i>et al.</i> (2010), McGovern <i>et al.</i> (2011), Sunil <i>et al.</i> (2014), Venosa <i>et al.</i> (2016)
Anti-inflammatory/profibrotic M2 macrophage mediators		
Profibrotic cytokines and Chemokines (eg, IL-4, IL-10, IL-13, IL-17, IL-33 CXCL-9, fractalkine, CCL-18)	Ozone	Backus <i>et al.</i> (2010), Zhu <i>et al.</i> (2016)
	Radiation	Inoue <i>et al.</i> (2004b), Chung <i>et al.</i> (2016), Groves <i>et al.</i> (2016), Zhang <i>et al.</i> (2018)
	Hyperoxia	Bhandari <i>et al.</i> (2007)
	Endotoxin	Matsukawa <i>et al.</i> (2000), Hocke <i>et al.</i> (2006), D'Alessio <i>et al.</i> (2016), Tu <i>et al.</i> (2017)
	Bleomycin	Pochetuhnen <i>et al.</i> (2007), Li <i>et al.</i> (2014), Wei <i>et al.</i> (2016)
	PM	Gowdy <i>et al.</i> (2008)
	Silica	Barbarin <i>et al.</i> (2004), Barbarin <i>et al.</i> (2005), Ferreira <i>et al.</i> (2013)
Nanoparticles	Nanoparticles	Shvedova <i>et al.</i> (2005)
	Chemical warfare agents	Venosa <i>et al.</i> (2016)
Growth factors (eg, CTGF, TGF β , TGF α , PDGF α , M-CSF)	Radiation	Rube <i>et al.</i> (2000), Chung <i>et al.</i> (2014), Lee <i>et al.</i> (2015), Sohn <i>et al.</i> (2015), Bickelhaupt <i>et al.</i> (2017), Li <i>et al.</i> (2017), Zhang <i>et al.</i> (2018)
	Endotoxin	Tu <i>et al.</i> (2017)
	Bleomycin	Baran <i>et al.</i> (2007), Yu <i>et al.</i> (2015), Wei <i>et al.</i> (2016), Yao <i>et al.</i> (2016), Liu <i>et al.</i> (2017), Xie <i>et al.</i> (2017)
	PM	Yanamala <i>et al.</i> (2013)
	Asbestos	Nishimura <i>et al.</i> (2013), Murthy <i>et al.</i> (2015)
	Silica	Barbarin <i>et al.</i> (2004), Barbarin <i>et al.</i> (2005), Ferreira <i>et al.</i> (2013), Cruz <i>et al.</i> (2016)
	Nanoparticles	Shvedova <i>et al.</i> (2005)
Chemical warfare agents	Sunil <i>et al.</i> (2011), Venosa <i>et al.</i> (2016)	

Abbreviations: CCL, chemokine ligand; CTGF, connective tissue growth factor; IL, interleukin; M-CSF, macrophage-colony stimulating factor; MMPs, matrix metalloproteinases; NO, nitric oxide; PAF, platelet-activating factor; PDGF, platelet-derived growth factor; PM, particulate matter; RNS, reactive nitrogen species; ROS, reactive oxygen species; TGF, transforming growth factor; TIMP, tissue inhibitor of metalloproteinases; TNF, tumor necrosis factor.

cytokine, IL-12, results in increases in M1 macrophages, overexpression of iNOS, and reduced fibrosis (Hesse *et al.*, 2000).

Evidence suggests that proinflammatory cytokines produced by M1 macrophages including IL-1 and TNF α play a role in the development of pulmonary fibrosis. Thus, following administration of silica, bleomycin, or nitrogen mustard, IL-1 and TNF α are rapidly upregulated in the lung (Byrne *et al.*, 2015). Moreover, administration of IL-1 β or transient overexpression of IL-1 β in the lung recapitulates many of the features of bleomycin-induced fibrosis in mice, while blockade or loss of IL-1R1 reduces fibrosis (Byrne *et al.*, 2016; Gasse *et al.*, 2007). Silica-induced fibrosis in rodents is also blunted by neutralization of IL-1 β (Guo *et al.*, 2013). Similarly, although infusion of recombinant TNF α produces alterations characteristic of pulmonary fibrosis including fibroblast proliferation, collagen deposition, and cell necrosis, blockade or loss of TNF α activity early in the pathogenic process mitigates fibrosis induced by bleomycin, silica, asbestos, and mustard vesicants, a response associated with decreased expression of TGF β (Laskin *et al.*, 2011; Liu and Brody, 2001; Malaviya *et al.*, 2015b; Ortiz *et al.*, 1998). Conversely, pulmonary delivery of TNF α to mice with established fibrosis reduces

their fibrotic burden and improves lung function and architecture, a response linked to decreases in profibrotic M2 macrophages (Redente *et al.*, 2014). These data demonstrate that the contribution of M1 macrophages to fibrosis is distinct during different phases of the fibrogenic process.

MACROPHAGE-DERIVED MEDIATORS IMPLICATED IN ACUTE LUNG INJURY AND CHRONIC FIBROSIS

Mediators released by inflammatory macrophages are generally not target specific. Thus, their actions depend on the amount and location of release and their persistence in the tissue. In this regard, they may exacerbate acute tissue injury and/or promote chronic fibrosis. Among the more well-established proinflammatory/cytotoxic mediators implicated in tissue injury are ROS (eg, superoxide anion, hydrogen peroxide, hydroxyl radicals) and RNS (eg, nitric oxide, peroxytrite), proteases (eg, MMPs, TIMPs), lipid mediators (eg, lipid peroxides, PGE $_2$, PAF), and cytokines (eg, TNF α , IL-1, IL-6, IL-12, IL-18)/chemokines (eg, CCL2, CCL3, CCL4, CCL5, CXCL1). Macrophage mediators

implicated in chronic fibrosis include cytokines (eg, IL-4, IL-13, IL-33) and growth factors (eg, TGF β , CTGF, VEGF). As illustrated in Table 3, similar macrophage-derived mediators contribute to the pathogenic effects of diverse toxicants with distinct mechanisms of action. Complicating the understanding of the role of macrophage mediators in the response to toxicants is the fact that they likely act in concert to promote tissue damage and fibrosis. It should also be noted that macrophages are not the only source of proinflammatory/cytotoxic and profibrotic mediators in the lung. Thus, following toxicant exposure, airway and alveolar epithelial cells and fibroblasts also have the capacity to generate cytotoxic/proinflammatory and profibrotic mediators, a response due in part to macrophage-derived products (Hiraiwa et al., 2013). The contribution of mediators released by these cells to pulmonary toxicity, relative to macrophages, remains to be established.

RESIDENT ALVEOLAR MACROPHAGES IN ACUTE LUNG INJURY AND FIBROSIS

Phenotypically and functionally distinct subpopulations of macrophages are localized throughout the lung of healthy individuals and experimental animals; these resident tissue macrophages have been identified as alveolar macrophages, interstitial macrophages, pleural macrophages, intravascular macrophages, and airway macrophages (reviewed in Laskin et al., 2015). The largest and most well characterized are alveolar macrophages, which play a key role in recycling of surfactant. Resident alveolar macrophages are also central to pulmonary immune defense, serving as sentinels, strategically located to respond to invading pathogens and inhaled toxicants. Like other tissue-resident macrophages, they possess an M2-like phenotype, which is key to their role in maintaining homeostasis and protecting against injury and infection (Alber et al., 2012; Morales-Nebreda et al., 2015). In healthy lung, alveolar macrophages are sustained in a restrained state by their interaction with the alveolar epithelium and molecules such as surfactant protein D, CD200, IL-10, TGF β and a transmembrane glycoprotein, MUC-1, via macrophage receptors like TLR4, CD200R, and SIRP α (Herold et al., 2011; Morales-Nebreda et al., 2015; Snelgrove et al., 2011). Disturbances in the alveolar macrophage-epithelium contact are considered critical events in early inflammatory signaling (Alber et al., 2012). Evidence suggests that resident alveolar macrophages play a role in triggering acute inflammatory responses to noxious stimuli. They have the capacity to recognize danger signals (eg, DAMPs) released from injured and/or necrotic epithelial cells and initiate the inflammatory phase of wound healing. It has been suggested that following injury, epithelial cells release calcium which upregulates macrophage NADPH oxidase leading to hydrogen peroxide production; this triggers the release of HMGB1 and other DAMPs (eg, ATP, hyaluronan, heat shock proteins, heparan sulfate) from necrotic cells (Minutti et al., 2017). DAMPs are recognized by macrophage TLR4 and purigenic receptors; binding to these receptors initiates a signaling process resulting in the release of inflammatory mediators and chemokines and the recruitment of inflammatory cells to sites of injury.

Resident alveolar macrophage activation following pulmonary exposure to noxious stimuli can also occur via inflammasomes (eg, NLRP3). These are cytosolic multiprotein complexes that oligomerize and activate caspase-1 resulting in the generation and release of IL-1 and IL-18, which promote inflammation. Full activation of the NLRP3 inflammasome requires both

priming and activating stimuli. Although priming is initiated by DAMPs binding to pattern recognition receptors (PRR) such as TLR4, which causes upregulation of pro-IL-1 and pro-IL-18 and other components of the inflammasome (adaptor protein ASC, procaspase-1), activation involves the assembly of these components into the inflammasome structure followed by production of proinflammatory interleukins. In sterile inflammation, NLRP3 activation occurs following phagocytosis of crystals (eg, cholesterol, urate), particles (eg, silica, titanium), or nanomaterials (eg, carbon nanotubes) (Hosseini et al., 2015; Nakayama, 2018). Studies suggest that the NLRP3 inflammasome contributes to acute lung injury induced by ozone, titanium nanoparticles, asbestos, silica, and particulate matter (Che et al., 2016; Kim et al., 2017; Michaudel et al., 2016; Sayan and Mossman, 2016).

There are also data showing that the inflammasome and inflammasome-linked cytokines released from resident alveolar macrophages are involved in the development of chronic lung diseases including asthma, COPD, and fibrosis (Grailer et al., 2014). In this regard, acute pulmonary fibrotic changes are associated with increased levels of IL-1 β and IL-18 in alveolar macrophages from patients with idiopathic pulmonary fibrosis, as well as asbestos-induced fibrosis (Sayan and Mossman, 2016). Fibrogenic toxicants including cigarette smoke, asbestos, silica, carbon nanotubes, engineered nanomaterials, and bleomycin have all been shown to directly activate the NLRP3 inflammasome in alveolar-resident macrophages leading to IL-1 β secretion (Sayan and Mossman, 2016). IL-1 β stimulates the release of TGF β , which triggers activation, proliferation, and differentiation of epithelial cells and fibroblasts into collagen-producing myofibroblasts (dos Santos et al., 2012). Recent studies have demonstrated that depletion of resident macrophages just prior to the onset of bleomycin-induced fibrosis did not alter the severity of the pathology (Misharin et al., 2017). These data suggest that the contribution of resident macrophages to fibrogenesis is more prominent in early phases of the disease process.

Consistent with a role of resident alveolar macrophages in early responses to toxicants and pathogens are recent findings demonstrating that these cells are prone to undergo regulated cell death (ie, pyroptosis, necroptosis, METosis) following toxic exposures or infections (Doster et al., 2018; Ginhoux et al., 2017). These tightly regulated cell death pathways are highly inflammatory and immunogenic. They can be induced by a variety of factors including TLR4 ligands (eg, LPS, HMGB1), RAGE activation, inflammasomes and cytokines like IL-1 β and TNF α (Li et al., 2016; Xu et al., 2014), and involve caspase-1 receptor-interacting serine/threonine-protein kinases (RIPK1, RIPK3), mixed lineage kinase domain-like protein, and gasdermin D (Doster et al., 2018; Ginhoux et al., 2017). Resident alveolar macrophage death is thought to be key in triggering the recruitment of circulating monocytes and neutrophils to sites of injury and infection and initiating inflammatory responses (Fan and Fan, 2018). There is increasing recognition that macrophage death and inflammation reciprocally affect one another, forming an auto-amplification loop which exaggerates inflammation (Linkermann et al., 2014).

CONCLUSIONS AND FUTURE DIRECTIONS

Because the respiratory tract is in direct contact with the external environment, it is particularly vulnerable to the adverse effects of inhaled pathogens and toxicants. Macrophages represent an essential host immune defense mechanism against these harmful xenobiotics. However, effective host defense,

wound healing, and restoration of homeostasis requires that the activity of macrophages be carefully controlled. In the absence of operative control mechanisms, macrophages become overactivated, resulting in exacerbation of acute injury and/or the development of chronic lung disease. This is complicated by the fact that macrophages do not consist of one homogeneous cell population; rather subpopulations with unique phenotypic and functional properties. Moreover, as macrophages are highly plastic cells, they have the capacity to rapidly change their phenotype. The use of approaches that allow identification and characterization of individual macrophages within a mixed population (eg, single cell RNAseq or western blotting), as well as a conditional knockdown of macrophages (eg, Cre-lox) at different times during the pathogenic process will be particularly valuable for assessing the role of these cells in the development of lung disease. Also important will be a focus on the interactions between lung macrophages and epithelial cells in the response to toxicants. In this regard, recent studies demonstrating that epithelial cell-derived microvesicles generated following hyperoxia or toxicant-induced lung injury are key in proinflammatory activation of macrophages, represent an exciting new avenue of research (Lee et al., 2016, 2017). Another area of interest for future investigations is the role of extracellular traps (fibers composed of DNA and proteins) released from inflammatory macrophages (METs) in response to ROS, proteases, and TNF α in the pathogenic response to pulmonary toxicants (Boe et al., 2015; Doster et al., 2018). It has been suggested that a failure to remove METs may prolong inflammatory responses contributing to chronic tissue injury and disease (Boe et al., 2015; King et al., 2017). Understanding pathways regulating macrophage activation and the mediators they release may lead to more efficacious approaches for treating lung diseases and disorders caused by inhaled toxicants.

SUPPLEMENTARY DATA

Supplementary data are available at Toxicological Sciences online.

FUNDING

National Institutes of Health (U54AR055073, R01ES004738, and P30ES005022).

REFERENCES

- Aggarwal, N. R., King, L. S., and D'Alessio, F. R. (2014). Diverse macrophage populations mediate acute lung inflammation and resolution. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **306**, L709–L725.
- Al-Harbi, N. O., Imam, F., Al-Harbi, M. M., Ansari, M. A., Zoheir, K. M. A., Korashy, H. M., Sayed-Ahmed, M. M., Attia, S. M., Shabanah, O. A., and Ahmad, S. F. (2016). Dexamethasone attenuates LPS-induced acute lung injury through inhibition of NF- κ B, COX-2, and pro-inflammatory mediators. *Immunol. Invest.* **45**, 349–369.
- Alber, A., Howie, S. E., Wallace, W. A., and Hirani, N. (2012). The role of macrophages in healing the wounded lung. *Int. J. Exp. Pathol.* **93**, 243–251.
- Arora, S., Dev, K., Agarwal, B., Das, P., and Syed, M. A. (2018). Macrophages: Their role, activation and polarization in pulmonary diseases. *Immunobiology* **223**, 383–396.
- Backus, G. S., Howden, R., Fostel, J., Bauer, A. K., Cho, H.-Y., Marzec, J., Peden, D. B., and Kleeburger, S. R. (2010). Protective role of interleukin-10 in ozone-induced pulmonary inflammation. *Environ. Health Perspect.* **118**, 1721–1727.
- Baran, C. P., Opalek, J. M., McMaken, S., Newland, C. A., O'Brien, J. M., Hunter, M. G., Bringardner, B. D., Monick, M. M., Brigstock, D. R., Stromberg, P. C., et al. (2007). Important roles for macrophage colony-stimulating factor, CC chemokine ligand 2, and mononuclear phagocytes in the pathogenesis of pulmonary fibrosis. *Am. J. Respir. Crit. Care Med.* **176**, 78–89.
- Bauer, C., Kielian, T., Wyatt, T. A., Romberger, D. J., West, W. W., Gleason, A. M., and Poole, J. A. (2013). Myeloid differentiation factor 88-dependent signaling is critical for acute organic dust-induced airway inflammation in mice. *Am. J. Respir. Cell Mol. Biol.* **48**, 781–789.
- Bekki, K., Ito, T., Yoshida, Y., He, C., Arashidani, K., He, M., Sun, G., Zeng, Y., Sone, H., Kunugita, N., et al. (2016). PM2.5 collected in China causes inflammatory and oxidative stress responses in macrophages through the multiple pathways. *Environ. Toxicol. Pharmacol.* **45**, 362–369.
- Bhalla, D. K., Daniels, D. S., and Luu, N. T. (1992). Attenuation of ozone-induced airway permeability in rats by pretreatment with cyclophosphamide, FPL 55712, and indomethacin. *Am. J. Respir. Cell Mol. Biol.* **7**, 73–80.
- Bhandari, V., Choo-Wing, R., Homer, R. J., and Elias, J. A. (2007). Increased hyperoxia-induced mortality and acute lung injury in IL-13 null mice. *J. Immunol.* **178**, 4993–5000.
- Bickelhaupt, S., Erbel, C., Timke, C., Wirkner, U., Dadrich, M., Flechsig, P., Tietz, A., Pfohler, J., Gross, W., Psechke, P., et al. (2017). Effects of CTGF blockade on attenuation and reversal of radiation-induced pulmonary fibrosis. *J. Natl. Cancer Inst.* **109**, djw3339.
- Boe, D. M., Curtis, B. J., Chen, M. M., Ippolito, J. A., and Kovacs, E. J. (2015). Extracellular traps and macrophages: new roles for the versatile phagocyte. *J. Leukoc. Biol.* **97**, 1023–1035.
- Buck, M., and Chojkier, M. (2011). C/EBP β -Thr217 phosphorylation signaling contributes to the development of lung injury and fibrosis in mice. *PLoS One* **6**, e25497.
- Byrne, A. J., Maher, T. M., and Lloyd, C. M. (2016). Pulmonary macrophages: A new therapeutic pathway in fibrosing lung disease? *Trends Mol. Med.* **22**, 303–316.
- Byrne, A. J., Mathie, S. A., Gregory, L. G., and Lloyd, C. M. (2015). Pulmonary macrophages: key players in the innate defence of the airways. *Thorax* **70**, 1189–1196.
- Casey, J., Kaplan, J., Atochina-Vasserman, E. N., Gow, A. J., Kadire, H., Tomer, Y., Fisher, J. H., Hawgood, S., Savani, R. C., and Beers, M. F. (2005). Alveolar surfactant protein D content modulates bleomycin-induced lung injury. *Am. J. Respir. Crit. Care Med.* **172**, 869–877.
- Chaudhary, N. I., Schnapp, A., and Park, J. E. (2006). Pharmacologic differentiation of inflammation and fibrosis in the rat bleomycin model. *Am. J. Respir. Crit. Care Med.* **173**, 769–776.
- Che, L., Jin, Y., Zhang, C., Lai, T., Zhou, H., Xia, L., Tian, B., Zhao, Y., Liu, J., Wu, Y., et al. (2016). Ozone-induced IL-17A and neutrophilic airway inflammation is orchestrated by the caspase-1-IL-1 cascade. *Sci. Rep.* **6**, 18680.
- Chen, F., Gong, L., Zhang, L., Wang, H., Qi, X., Wu, X., Xiao, Y., Cai, Y., Liu, L., Li, X., et al. (2006). Short courses of low dose dexamethasone delay bleomycin-induced lung fibrosis in rats. *Eur. J. Pharmacol.* **536**, 287–295.
- Connor, A. J., Laskin, J. D., and Laskin, D. L. (2012). Ozone-induced lung injury and sterile inflammation. Role of toll-like receptor 4. *Exp. Mol. Pathol.* **92**, 229–235.

- Dhaliwal, K., Scholefield, E., Ferenbach, D., Gibbons, M., Duffin, R., Dorward, D. A., Morris, A. C., Humphries, D., MacKinnon, A., Wilkinson, T. S., et al. (2012). Monocytes control second-phase neutrophil emigration in established lipopolysaccharide-induced murine lung injury. *Am. J. Respir. Crit. Care Med.* **186**, 514–524.
- dos Santos, G., Kutuzov, M. A., and Ridge, K. M. (2012). The inflammasome in lung diseases. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **303**, L627–L633.
- Doster, R. S., Rogers, L. M., Gaddy, J. A., and Aronoff, D. M. (2018). Macrophage extracellular traps: A scoping review. *J. Innate Immun.* **10**, 3–13.
- Duke, K. S., and Bonner, J. C. (2018). Mechanisms of carbon nanotube-induced pulmonary fibrosis: a physicochemical characteristic perspective. *Wiley Interdiscip. Rev. Nanomed. Nanobiotechnol.* **10**, e1498.
- Duru, N., Wolfson, B., and Zhou, Q. (2016). Mechanisms of the alternative activation of macrophages and non-coding RNAs in the development of radiation-induced lung fibrosis. *World J. Biol. Chem.* **7**, 231–239.
- Fan, E. K. Y., and Fan, J. (2018). Regulation of alveolar macrophage death in acute lung inflammation. *Respir. Res.* **19**, 50.
- Faulknor, R. A., Olekson, M. A., Ekwueme, E. C., Krzyszczyk, P., Freeman, J. W., and Berthiaume, F. (2017). Hypoxia impairs mesenchymal stromal cell-induced macrophage M1 to M2 transition. *Technology* **5**, 81–86.
- Francis, M., Groves, A. M., Sun, R., Cervelli, J. A., Choi, H., Laskin, J. D., and Laskin, D. L. (2017). CCR2 regulates inflammatory cell accumulation in the lung and tissue injury following ozone exposure. *Toxicol. Sci.* **155**, 474–484.
- Frank, E. A., Birch, M. E., and Yadav, J. S. (2015). MyD88 mediates in vivo effector functions of alveolar macrophages in acute lung inflammatory responses to carbon nanotube exposure. *Toxicol. Appl. Pharmacol.* **288**, 322–329.
- Fulkerson, P. C., Fischetti, C. A., Hassman, L. M., Nikolaidis, N. M., and Rothenberg, M. E. (2006). Persistent effects induced by IL-13 in the lung. *Am. J. Respir. Cell Mol. Biol.* **35**, 337–346.
- Gasse, P., Mary, C., Guenon, I., Noulin, N., Charron, S., Schnyder-Candrian, S., Schnyder, B., Akira, S., Quesniaux, V. F. J., Lagente, V., et al. (2007). IL-1R1/MyD88 signaling and the inflammasome are essential in pulmonary inflammation and fibrosis in mice. *J. Clin. Invest.* **117**, 3786–3799.
- Geng, P., Ma, T., Xing, J., Jiang, L., Sun, H., Zhu, B., Zhang, H., Xiao, H., Wang, J., and Zhang, J. (2018). Dexamethasone ameliorates H₂S-induced acute lung injury by increasing claudin-5 expression via the PI3K pathway. *Hum. Exp. Toxicol.* **37**, 626–635.
- Gharaee-Kermani, M., McCullumsmith, R. E., Charo, I. F., Kunkel, S. L., and Phan, S. H. (2003). CC-chemokine receptor 2 required for bleomycin-induced pulmonary fibrosis. *Cytokine* **24**, 266–276.
- Gibbons, M. A., MacKinnon, A. C., Ramachandran, P., Dhaliwal, K., Duffin, R., Phythian-Adams, A. T., van Rooijen, N., Haslett, C., Howie, S. E., Simpson, A. J., et al. (2011). Ly6C^{hi} monocytes direct alternatively activated profibrotic macrophage regulation of lung fibrosis. *Am. J. Respir. Crit. Care Med.* **184**, 569–581.
- Ginhoux, F., Bleriot, C., and Lecuit, M. (2017). Dying for a cause: Regulated necrosis of tissue-resident macrophages upon infection. *Trends Immunol.* **38**, 693–695.
- Grailer, J. J., Canning, B. A., Kalbitz, M., Haggadone, M. D., Dhond, R. M., Andjelkovic, A. V., Zetoune, F. S., and Ward, P. A. (2014). Critical role for the NLRP3 inflammasome during acute lung injury. *J. Immunol.* **192**, 5974–5983.
- Groves, A. M., Gow, A. J., Massa, C. B., Hall, L., Laskin, J. D., and Laskin, D. L. (2013). Age-related increases in ozone-induced injury and altered pulmonary mechanics in mice with progressive lung inflammation. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **305**, L555–L568.
- Groves, A. M., Gow, A. J., Massa, C. B., Laskin, J. D., and Laskin, D. L. (2012). Prolonged injury and altered lung function after ozone inhalation in mice with chronic lung inflammation. *Am. J. Respir. Cell Mol. Biol.* **47**, 776–783.
- Groves, A. M., Johnston, C. J., Williams, J. P., and Finkelstein, J. N. (2018). Role of infiltrating monocytes in the development of radiation-induced pulmonary fibrosis. *Radiat. Res.* **189**, 300–311.
- Gundra, U. M., Girgis, N. M., Ruckerl, D., Jenkins, S., Ward, L. N., Kurtz, Z. D., Wiens, K. E., Tang, M. S., Basu-Roy, U., Mansukhani, A., et al. (2014). Alternatively activated macrophages derived from monocytes and tissue macrophages are phenotypically and functionally distinct. *Blood* **123**, e110–e122.
- Guo, J., Gu, N., Chen, J., Shi, T., Zhou, Y., Rong, Y., Zhou, T., Yang, W., Cui, X., and Chen, W. (2013). Neutralization of interleukin-1 β attenuates silica-induced lung inflammation and fibrosis in C57BL/6 mice. *Arch. Toxicol.* **87**, 1963–1973.
- Hashimoto, N., Kawabe, T., Imaizumi, K., Hara, T., Okamoto, M., Kojima, K., Shimokata, K., and Hasegawa, Y. (2004). CD40 plays a crucial role in lipopolysaccharide-induced acute lung injury. *Am. J. Respir. Cell Mol. Biol.* **30**, 808–815.
- He, M., Ichinose, T., Song, Y., Yoshida, Y., Bekki, K., Arashidani, K., Yoshida, S., Nishikawa, M., Takano, H., Shibamoto, T., et al. (2016). Desert dust induces TLR signaling to trigger Th2-dominant lung allergic inflammation via a MyD88-dependent signaling pathway. *Toxicol. Appl. Pharmacol.* **296**, 61–72.
- Herold, S., Mayer, K., and Lohmeyer, J. (2011). Acute lung injury: how macrophages orchestrate resolution of inflammation and tissue repair. *Front. Immunol.* **2**, 65.
- Hesse, M., Cheever, A. W., Jankovic, D., and Wynn, T. A. (2000). NOS-2 mediates the protective anti-inflammatory and antifibrotic effects of the Th1-inducing adjuvant, IL-12, in a Th2 model of granulomatous disease. *Am. J. Pathol.* **157**, 945–955.
- Hiraiwa, K., and van Eeden, S. F. (2013). Contribution of lung macrophages to the inflammatory responses induced by exposure to air pollutants. *Mediators Inflamm.* **2013**, 619523.
- Hollingsworth, J. W., Cook, D. N., Brass, D. M., Walker, J. K. L., Morgan, D. L., Foster, W. M., and Schwartz, D. A. (2004). The role of Toll-like receptor 4 in environmental airway injury in mice. *Am. J. Respir. Crit. Care Med.* **170**, 126–132.
- Hoshino, T., Okamoto, M., Sakazaki, Y., Kato, S., Young, H. A., and Aizawa, H. (2009). Role of proinflammatory cytokines IL-18 and IL-1 β in bleomycin-induced lung injury in humans and mice. *Am. J. Respir. Cell Mol. Biol.* **41**, 661–670.
- Hosseini, N., Cho, Y., Lockey, R. F., and Kolliputi, N. (2015). The role of the NLRP3 inflammasome in pulmonary diseases. *Ther. Adv. Respir. Dis.* **9**, 188–197.
- Huang, B., Wang, D. X., and Deng, W. (2014). Protective effects of dexamethasone on early acute lung injury induced by oleic acid in rats. *Int. J. Clin. Exp. Med.* **7**, 4698–4709.
- Huang, Y., Zhang, W., Yu, F., and Gao, F. (2017). The cellular and molecular mechanism of radiation-induced lung injury. *Med. Sci. Monit.* **23**, 3446–3450.
- Huax, F., Liu, T., McGarry, B., Ullenbruch, M., and Phan, S. H. (2003). Dual roles of IL-4 in lung injury and fibrosis. *J. Immunol.* **170**, 2083–2092.

- Huax, F., Louahed, J., Hudspeth, B., Meredith, C., Delos, M., Renaud, J. C., and Lison, D. (1998). Role of interleukin-10 in the lung response to silica in mice. *Am. J. Respir. Cell Mol. Biol.* **18**, 51–59.
- Hussell, T., and Bell, T. J. (2014). Alveolar macrophages: Plasticity in a tissue-specific context. *Nat. Rev. Immunol.* **14**, 81–93.
- Ishida, Y., Kimura, A., Nosaka, M., Kuninaka, Y., Hemmi, H., Sasaki, I., Kaisho, T., Mukaida, N., and Kondo, T. (2017). Essential involvement of the CX3CL1-CX3CR1 axis in bleomycin-induced pulmonary fibrosis via regulation of fibrocyte and M2 macrophage migration. *Sci. Rep.* **7**, 16833.
- Ji, W.-J., Ma, Y.-Q., Zhang, X., Zhang, L., Zhang, Y.-D., Su, C.-C., Xiang, G.-A., Zhang, M.-P., Lin, Z.-C., Wei, L.-Q., et al. (2016). Inflammatory monocyte/macrophage modulation by liposome-entrapped spironolactone ameliorates acute lung injury in mice. *Nanomedicine* **11**, 1393–1406.
- Kambara, K., Ohashi, W., Tomita, K., Takashina, M., Fujisaka, S., Hayashi, R., Mori, H., Tobe, K., and Hattori, Y. (2015). *In vivo* depletion of CD206⁺ M2 macrophages exaggerates lung injury in endotoxemic mice. *Am. J. Pathol.* **185**, 162–171.
- Kim, B. G., Lee, P. H., Lee, S. H., Park, M. K., and Jang, A. S. (2017). Effect of TiO₂ nanoparticles on inflammasome-mediated airway inflammation and responsiveness. *Allergy Asthma Immunol. Res.* **9**, 257–264.
- King, P. T., Sharma, R., O'Sullivan, K. M., Callaghan, J., Dousha, L., Thomas, B., Ruwanpura, S., Lim, S., Farmer, M. W., Jennings, B. R., et al. (2017). Deoxyribonuclease 1 reduces pathogenic effects of cigarette smoke exposure in the lung. *Sci. Rep.* **7**, 12128.
- Laskin, D. L., Malaviya, R., and Laskin, J. D. (2015). Pulmonary macrophages. In *Comparative Biology of the Normal Lung* (R. A. Parent, Ed.), pp. 629–649. Academic Press, San Diego, CA.
- Laskin, D. L., Sunil, V. R., Gardner, C. R., and Laskin, J. D. (2011). Macrophages and tissue injury: agents of defense or destruction? *Ann. Rev. Pharmacol. Toxicol.* **51**, 267–288.
- Lawrence, T., and Natoli, G. (2011). Transcriptional regulation of macrophage polarization: enabling diversity with identity. *Nat. Rev. Immunol.* **11**, 750–761.
- Lee, H., Zhang, D., Wu, J., Otterbein, L. E., and Jin, Y. (2017). Lung epithelial cell-derived microvesicles regulate macrophage migration via microRNA-17/221-induced integrin β 1 recycling. *J. Immunol.* **199**, 1453–1464.
- Lee, H., Zhang, D., Zhu, Z., Dela Cruz, C. S., and Jin, Y. (2016). Epithelial cell-derived microvesicles activate macrophages and promote inflammation via microvesicle-containing microRNAs. *Sci. Rep.* **6**, 35250.
- Li, D., Guabiraba, R., Besnard, A.-G., Komai-Koma, M., Jabir, M. S., Zhang, L., Graham, G. J., Kurowska-Stolarska, M., Liew, F. Y., McSharry, C., et al. (2014). IL-33 promotes ST2-dependent lung fibrosis by the induction of alternatively activated macrophages and innate lymphoid cells in mice. *J. Allergy Clin. Immunol.* **134**, 1422–1432.e11.
- Li, Z., Scott, M. J., Fan, E. K., Li, Y., Liu, J., Xiao, G., Li, S., Billiar, T. R., Wilson, M. A., Jiang, Y., et al. (2016). Tissue damage negatively regulates LPS-induced macrophage necroptosis. *Cell Death Differ.* **23**, 1428–1447.
- Linkermann, A., Stockwell, B. R., Krautwald, S., and Anders, H. J. (2014). Regulated cell death and inflammation: an auto-amplification loop causes organ failure. *Nat. Rev. Immunol.* **14**, 759–767.
- Liu, J. Y., and Brody, A. R. (2001). Increased TGF- β 1 in the lungs of asbestos-exposed rats and mice: reduced expression in TNF- α receptor knockout mice. *J. Environ. Pathol. Toxicol. Oncol.* **20**, 97–108.
- Liu, T., Jin, H., Ullenbruch, M., Hu, B., Hashimoto, N., Moore, B., McKenzie, A., Lukacs, N. W., and Phan, S. H. (2004). Regulation of found in inflammatory zone 1 expression in bleomycin-induced lung fibrosis: Role of IL-4/IL-13 and mediation via STAT-6. *J. Immunol.* **173**, 3425–3431.
- Liu, T., Yu, H., Ullenbruch, M., Jin, H., Ito, T., Wu, Z., Liu, J., and Phan, S. H. (2014). The *in vivo* fibrotic role of FIZZ1 in pulmonary fibrosis. *PLoS One* **9**, e88362.
- Lu, H. L., Huang, X. Y., Luo, Y. F., Tan, W. P., Chen, P. F., and Guo, Y. B. (2018). Activation of M1 macrophages plays a critical role in the initiation of acute lung injury. *Biosci. Rep.* **38**, 1–13.
- Lumsden, R. V., Worrell, J. C., Boylan, D., Walsh, S. M., Cramton, J., Counihan, I., O'Beirne, S., Medina, M. F., Gauldie, J., Fabre, A., et al. (2015). Modulation of pulmonary fibrosis by IL-13 α 2. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **308**, L710–L718.
- Luzina, I. G., Kopach, P., Lockett, V., Kang, P. H., Nagarsekar, A., Burke, A. P., Hasday, J. D., Todd, N. W., and Atamas, S. P. (2013). Interleukin-33 potentiates bleomycin-induced lung injury. *Am. J. Respir. Cell Mol. Biol.* **49**, 999–1008.
- Malaviya, R., Gow, A. J., Francis, M., Abramova, E. V., Laskin, J. D., and Laskin, D. L. (2015a). Radiation-induced lung injury and inflammation in mice: Role of inducible nitric oxide synthase and surfactant protein D. *Toxicol. Sci.* **144**, 27–38.
- Malaviya, R., Sunil, V. R., Venosa, A., Verissimo, V. L., Cervelli, J. A., Vayas, K. N., Hall, L., Laskin, J. D., and Laskin, D. L. (2015b). Attenuation of nitrogen mustard-induced pulmonary injury and fibrosis by anti-tumor necrosis factor- α antibody. *Toxicol. Sci.* **148**, 71–88.
- Malaviya, R., Laskin, J. D., and Laskin, D. L. (2017). Anti-TNF α therapy in inflammatory lung diseases. *Pharmacol. Ther.* **180**, 90–98.
- Marchini, T., Wolf, D., Michel, N. A., Mauler, M., Dufner, B., Hoppe, N., Beckert, J., Jackel, M., Magnani, N., Duerschmied, D., et al. (2016). Acute exposure to air pollution particulate matter aggravates experimental myocardial infarction in mice by potentiating cytokine secretion from lung macrophages. *Basic Res. Cardiol.* **111**.
- Martinez, F. O., and Gordon, S. (2014). The M1 and M2 paradigm of macrophage activation: time for reassessment. *F1000Prime Rep.* **6**, 13.
- Matsukawa, A., Hogaboam, C. M., Lukacs, N. W., Lincoln, P. M., Evanoff, H. L., Strieter, R. M., and Kunkel, S. L. (2000). Expression and contribution of endogenous IL-13 in an experimental model of sepsis. *J. Immunol.* **164**, 2738–2744.
- McCubbrey, A. L., Barthel, L., Mohnning, M. P., Redente, E. F., Mould, K. J., Thomas, S. M., Leach, S. M., Danhorn, T., Gibbings, S. L., Jakubzick, C. V., et al. (2018). Deletion of c-FLIP from CD11b^{hi} macrophages prevents development of bleomycin-induced lung fibrosis. *Am. J. Respir. Cell Mol. Biol.* **58**, 66–78.
- Metchnikoff, E. 1968. *Lectures on the Comparative Pathology of Inflammation Delivered at the Pasteur Institute in 1891*. Dover, New York.
- Meziani, L., Mondini, M., Petit, B., Boissonnas, A., Thomas de Montpreville, V., Mercier, O., Vozenin, M. C., and Deutsch, E. (2018). CSF1R inhibition prevents radiation pulmonary fibrosis by depletion of interstitial macrophages. *Eur. Respir. J.* **51**, doi: 10.1183/13993003.13902120-13992017.
- Michaudel, C., Couturier-Maillard, A., Chenuet, P., Maillet, I., Mura, C., Couillin, I., Gombault, A., Quesniaux, V. F., Huaux, F., and Ryffel, B. (2016). Inflammasome, IL-1 and

- inflammation in ozone-induced lung injury. *Am. J. Clin. Exp. Immunol.* **5**, 33–40.
- Minutti, C. M., Knipper, J. A., Allen, J. E., and Zaiss, D. M. (2017). Tissue-specific contribution of macrophages to wound healing. *Semin. Cell Dev. Biol.* **61**, 3–11.
- Misharin, A. V., Morales-Nebreda, L., Mutlu, G. M., Budinger, G. R., and Perlman, H. (2013). Flow cytometric analysis of macrophages and dendritic cell subsets in the mouse lung. *Am. J. Respir. Cell Mol. Biol.* **49**, 503–510.
- Misharin, A. V., Morales-Nebreda, L., Reyfman, P. A., Cuda, C. M., Walter, J. M., McQuattie-Pimentel, A. C., Chen, C.-I., Anekalla, K. R., Joshi, N., Williams, K. J. N., et al. (2017). Monocyte-derived alveolar macrophages drive lung fibrosis and persist in the lung over the life span. *J. Exp. Med.* **214**, 2387–2404.
- Morales-Nebreda, L., Misharin, A. V., Perlman, H., and Budinger, G. R. (2015). The heterogeneity of lung macrophages in the susceptibility to disease. *Eur. Respir. Rev.* **24**, 505–509.
- Muller-Redetzky, H. C., Suttorp, N., and Witzentrath, M. (2014). Dynamics of pulmonary endothelial barrier function in acute inflammation: Mechanisms and therapeutic perspectives. *Cell Tissue Res.* **355**, 657–673.
- Murray, L. A., Chen, Q., Kramer, M. S., Hesson, D. P., Argentieri, R. L., Peng, X., Gulati, M., Homer, R. J., Russell, T., van Rooijen, N., et al. (2011). TGF-beta driven lung fibrosis is macrophage dependent and blocked by Serum amyloid P. *Int. J. Biochem. Cell Biol.* **43**, 154–162.
- Murray, P. J. (2017). Macrophage polarization. *Annu. Rev. Physiol.* **79**, 541–566.
- Murthy, S., Larson-Casey, J. L., Ryan, A. J., He, C., Kobzik, L., and Carter, A. B. (2015). Alternative activation of macrophages and pulmonary fibrosis are modulated by scavenger receptor, macrophage receptor with collagenous structure. *Faseb J.* **29**, 3527–3536.
- Nakayama, M. (2018). Macrophage recognition of crystals and nanoparticles. *Front. Immunol.* **9**, 1–8.
- Nemmar, A., Nemery, B., Hoet, P. H., Van Rooijen, N., and Hoylaerts, M. F. (2005). Silica particles enhance peripheral thrombosis: Key role of lung macrophage-neutrophil cross-talk. *Am. J. Respir. Crit. Care Med.* **171**, 872–879.
- Ortiz, L. A., Lasky, J., Hamilton, R. F., Holian, A., Hoyle, G. W., Banks, W., Peschon, J. J., Brody, A. R., Lungarella, G., and Friedman, M. (1998). Expression of TNF and the necessity of TNF receptors in bleomycin-induced lung injury in mice. *Exp. Lung Res.* **24**, 721–743.
- Parisi, L., Gini, E., Baci, D., Tremolati, M., Fanuli, M., Bassani, B., Farronato, G., Bruno, A., and Mortara, L. (2018). Macrophage polarization in chronic inflammatory diseases: Killers or builders? *J. Immunol. Res.* **2018**, 8917804.
- Perez-Rial, S., del Puerto-Nevado, L., Terron-Exposito, R., Giron-Martinez, A., Gonzalez-Mangado, N., and Peces-Barba, G. (2013). Role of recently migrated monocytes in cigarette smoke-induced lung inflammation in different strain of mice. *PLoS One* **8**, e72975.
- Pilling, D., Roife, D., Wang, M., Ronkainen, S. D., Crawford, J. R., Travis, E. L., and Gomer, R. H. (2007). Reduction of bleomycin-induced pulmonary fibrosis by serum amyloid P. *J. Immunol.* **179**, 4035–4044.
- Poole, J. A., Gleason, A. M., Bauer, C., West, W. W., Alexis, N., van Rooijen, N., Reynolds, S. J., Romberger, D. J., and Kielian, T. L. (2012). CD11c⁺/CD11b⁺ cells are critical for organic dust-elicited murine lung inflammation. *Am. J. Respir. Cell Mol. Biol.* **47**, 652–659.
- Porcheray, F., Viaud, S., Rimaniol, A. C., Leone, C., Samah, B., Dereuddre-Bosquet, N., Dormont, D., and Gras, G. (2005). Macrophage activation switching: an asset for the resolution of inflammation. *Clin. Exp. Immunol.* **142**, 481–489.
- Prasse, A., Pechkovsky, D. V., Toews, G. B., Jungraithmayr, W., Kollert, F., Goldmann, T., Vollmer, E., Müller-Quernheim, J., and Zissel, G. (2006). A vicious circle of alveolar macrophages and fibroblasts perpetuates pulmonary fibrosis via CCL18. *Am. J. Respir. Crit. Care Med.* **173**, 781–792.
- Pulichino, A.-M., Wang, I.-M., Caron, A., Mortimer, J., Auger, A., Boie, Y., Elias, J. A., Kartono, A., Xu, L., Menetski, J., et al. (2008). Identification of transforming growth factor β 1-driven genetic programs of acute lung fibrosis. *Am. J. Respir. Cell Mol. Biol.* **39**, 324–336.
- Re, S. L., Giordano, G., Yakoub, Y., Devosse, R., Uwambayinema, F., Coullin, I., Ryffel, B., Marbaix, E., Lison, D., and Huaux, F. (2014). Uncoupling between inflammatory and fibrotic responses to silica: Evidence from MyD88 knockout mice. *PLoS One* **9**, e99383.
- Redente, E. F., Keith, R. C., Janssen, W., Henson, P. M., Ortiz, L. A., Downey, G. P., Bratton, D. L., and Riches, D. W. H. (2014). Tumor necrosis factor- α accelerates the resolution of established pulmonary fibrosis in mice by targeting profibrotic lung macrophages. *Am. J. Respir. Cell Mol. Biol.* **50**, 825–837.
- Romero, F., Shah, D., Duong, M., Penn, R. B., Fessler, M. B., Madenspacher, J., Stafstrom, W., Kavuru, M., Lu, B., Kallen, C. B., et al. (2015). A pneumocyte-macrophage paracrine lipid axis drives the lung toward fibrosis. *Am. J. Respir. Cell Mol. Biol.* **53**, 74–86.
- Roszer, T. (2015). Understanding the mysterious M2 macrophage through activation markers and effector mechanisms. *Mediators Inflamm.* **2015**, 1–16.
- Samet, J. M., Ghio, A. J., Costa, D. L., and Madden, M. C. (2000). Increased expression of cyclooxygenase 2 mediates oil fly ash-induced lung injury. *Exp. Lung Res.* **26**, 57–69.
- Sayan, M., and Mossman, B. T. (2016). The NLRP3 inflammasome in pathogenic particle and fibre-associated lung inflammation and diseases. *Part. Fibre Toxicol.* **13**, 51.
- Shvedova, A. A., Kisin, E. R., Mercer, R., Murray, A. R., Johnson, V. J., Potapovich, A. I., Tyurina, Y. Y., Gorelik, O., Arepalli, S., Schwegler-Berry, D., et al. (2005). Unusual inflammatory and fibrogenic pulmonary responses to single-walled carbon nanotubes in mice. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **289**, L698–L708.
- Snelgrove, R. J., Godlee, A., and Hussell, T. (2011). Airway immune homeostasis and implications for influenza-induced inflammation. *Trends Immunol.* **32**, 328–334.
- Sugiura, H., and Ichinose, M. (2011). Nitrate stress in inflammatory lung diseases. *Nitric Oxide* **25**, 138–144.
- Sunil, V. R., Francis, M., Vayas, K. N., Cervelli, J. A., Choi, H., Laskin, J. D., and Laskin, D. L. (2015). Regulation of ozone-induced lung inflammation and injury by the β -galactoside-binding lectin galectin-3. *Toxicol. Appl. Pharmacol.* **284**, 236–245.
- Tam, A., Wadsworth, S., Dorscheid, D., Man, S. F., and Sin, D. D. (2011). The airway epithelium: more than just a structural barrier. *Ther. Adv. Respir. Dis.* **5**, 255–273.
- Tighe, R. M., Li, Z., Potts, E. N., Frush, S., Liu, N., Gunn, M. D., Foster, W. M., Noble, P. W., and Hollingsworth, J. W. (2011). Ozone inhalation promotes CX3CR1-dependent maturation of resident lung macrophages that limit oxidative stress and inflammation. *J. Immunol.* **187**, 4800–4808.
- Trujillo, G., O'Connor, E. C., Kunkel, S. L., and Hogaboam, C. M. (2008). A novel mechanism for CCR4 in the regulation of macrophage activation in bleomycin-induced pulmonary fibrosis. *Am. J. Pathol.* **172**, 1209–1221.

- Tsou, C.-L., Peters, W., Si, Y., Slaymaker, S., Aslanian, A. M., Weisberg, S. P., Mack, M., and Charo, I. F. (2007). Critical roles for CCR2 and MCP-3 in monocyte mobilization from bone marrow and recruitment to inflammatory sites. *J. Clin. Invest.* **117**, 902–909.
- Vannella, K. M., and Wynn, T. A. (2017). Mechanisms of organ injury and repair by macrophages. *Annu. Rev. Physiol.* **79**, 593–617.
- Venosa, A., Malaviya, R., Choi, H., Gow, A. J., Laskin, J. D., and Laskin, D. L. (2016). Characterization of distinct macrophage subpopulations during nitrogen mustard-induced lung injury and fibrosis. *Am. J. Respir. Cell Mol. Biol.* **54**, 436–446.
- Wang, N., Liang, H., and Zen, K. (2014a). Molecular mechanisms that influence the macrophage m1-m2 polarization balance. *Front. Immunol.* **5**, 614.
- Wang, X., Shannahan, J. H., and Brown, J. M. (2014b). IL-33 modulates chronic airway resistance changes induced by multi-walled carbon nanotubes. *Inhal. Toxicol.* **26**, 240–249.
- Weinberger, B., Laskin, J. D., Sunil, V. R., Sinko, P. J., Heck, D. E., and Laskin, D. L. (2011). Sulfur mustard-induced pulmonary injury: Therapeutic approaches to mitigating toxicity. *Pulm. Pharmacol. Ther.* **24**, 92–99.
- Wiesemann, A., Ketteler, J., Slama, A., Wirsdorfer, F., Hager, T., Rock, K., Engel, D. R., Fischer, J. W., Aigner, C., Jendrossek, V., et al. (2018). Inhibition of radiation-induced Ccl_2 signaling protects lungs from vascular dysfunction and endothelial cell loss. *Antioxid. Redox Signal.*, doi: 10.1089/ars.2017.7458.
- Wigenstam, E., Rocksén, D., Ekstrand-Hammarström, B., and Bucht, A. (2009). Treatment with dexamethasone or liposome-encapsulated vitamin E provides beneficial effects after chemical-induced lung injury. *Inhal. Toxicol.* **21**, 958–964.
- Wynn, T. A., and Vannella, K. M. (2016). Macrophages in tissue repair, regeneration, and fibrosis. *Immunity* **44**, 450–462.
- Xiang, G.-A., Zhang, Y.-D., Su, C.-C., Ma, Y.-Q., Li, Y.-M., Zhou, X., Wei, L.-Q., and Ji, W.-J. (2016). Dynamic changes of mononuclear phagocytes in circulating, pulmonary alveolar and interstitial compartments in a mouse model of experimental silicosis. *Inhal. Toxicol.* **28**, 393–402.
- Xu, J., Jiang, Y., Wang, J., Shi, X., Liu, Q., Liu, Z., Li, Y., Scott, M. J., Xiao, G., Li, S., et al. (2014). Macrophage endocytosis of high-mobility group box 1 triggers pyroptosis. *Cell Death Differ.* **21**, 1229–1239.
- Yang, D., Tong, L., Wang, D., Wang, Y., Wang, X., and Bai, C. (2010). Roles of CC chemokine receptors (CCRs) on lipopolysaccharide-induced acute lung injury. *Respir. Physiol. Neurobiol.* **170**, 253–259.
- Yao, Y., Wang, Y., Zhang, Z., He, L., Zhu, J., Zhang, M., He, X., Cheng, Z., Ao, Q., Cao, Y., et al. (2016). Chop deficiency protects mice against bleomycin-induced pulmonary fibrosis by attenuating M2 macrophage production. *Mol. Ther.* **24**, 915–925.
- Zhao, J., Okamoto, Y., Asano, Y., Ishimaru, K., Aki, S., Yoshioka, K., Takuwa, N., Wada, T., Inagaki, Y., Takahashi, C., et al. (2018). Sphingosine-1-phosphate receptor-2 facilitates pulmonary fibrosis through potentiating IL-13 pathway in macrophages. *PLoS One* **13**, e0197604.