

Treatment-resistant depression

Amy Werremeyer, PharmD, BCPP¹

¹Vice Chair and Associate Professor, Pharmacy Practice, College of Pharmacy, Nursing and Allied Sciences, North Dakota State University, Fargo, ND

KEYWORDS

treatment-resistant, depression

The term “resistant” does not usually bode well in the healthcare setting. “Antimicrobial resistant,” “resistant to care,” “chemotherapy resistant cancer”—all of these may prompt specific focused interventions or programs, often involving multi-disciplinary team approaches, and creative, out-of-the box solutions, to attempt achievement of positive outcomes for the patients involved.

Treatment-resistant depression is no different. It is a diagnosis that is often dreaded by patients, caregivers, and providers alike. It is responsible for increased hospitalizations, increased outpatient provider visits, greater use of psychotropic medications, and an average six times greater total healthcare cost incurrence compared to non-treatment-resistant depressed patients.¹ Yet, the definition of treatment-resistant depression (TRD) remains somewhat controversial. In its simplest form, TRD can be defined as lack of significant improvement after two adequate trials of two different antidepressants from two different pharmacologic classes.² However, the definition can quickly become quite complicated. According to Berlim, one can find more than 10 different definitions of TRD throughout the published literature.³ In addition, various authors have proposed staging systems for further description and characterization of TRD.^{4,5,6,7,8} Still others have argued that perhaps many instances of TRD actually represent undiagnosed bipolar disorder,⁹ nonadherence with therapy, and/or inadequate dose or duration of antidepressant therapy.¹⁰

The published literature focusing on treatment approaches for the management of TRD remains somewhat muddled due to the varying methods for defining TRD and how these definitions are employed in intervention studies. Despite the confusion and difficulty to extract conclusions on the data, some progress is being made in the treatment of TRD. This issue highlights some of that progress. Articles highlighting the use of lithium and second-generation antipsychotics for treatment of TRD provide a solid review of these important treatment

options. Additional pieces discussing emerging therapies in TRD and the pharmacogenomic foundations upon which future TRD treatment decisions may be made highlight the directions in which we are heading. A toolbox summarizing pharmacologic treatment options and their advantages and disadvantages in TRD is also included and may serve as an excellent reference when working with this patient population.

REFERENCES

1. Crown WH, Finkelstein S, Berndt ER, Ling D, Poret AW, Rush AJ, et al. The impact of treatment-resistant depression on healthcare utilization and cost. *J Clin Psychiatry*. 2002;63 (11):963-71.
2. Berlim MT, Turecki G. Definition, assessment, and staging of treatment-resistant refractory major depression: a review of current concepts and methods. *Can J Psychiatry*. 2007;52(1):46-54. PubMed PMID: [17444078](#).
3. Berlim MT, Turecki G. What is the meaning of treatment resistant/refractory major depression (TRD)? A systematic review of current randomized trials. *Eur Neuropsychopharmacol*. 2007;17(11):696-707. DOI: [10.1016/j.euroneuro.2007.03.009](#). PubMed PMID: [17521891](#).
4. Sackeim HA, Prudic J, Devanand DP, Decina P, Kerr B, Malitz S. The impact of medication resistance and continuation pharmacotherapy on relapse following response to electroconvulsive therapy in major depression. *J Clin Psychopharmacol*. 1990;10(2):96-104. PubMed PMID: [2341598](#).
5. Thase ME, Rush AJ. When at first you don't succeed: sequential strategies for antidepressant nonresponders. *J Clin Psychiatry*. 1997;58 Suppl 13:23-9. PubMed PMID: [9402916](#).
6. Souery D, Amsterdam J, de Montigny C, Lecrubier Y, Montgomery S, Lipp O, et al. Treatment resistant depression: methodological overview and operational criteria. *Eur Neuropsychopharmacol*. 1999;9(1-2):83-91. PubMed PMID: [10082232](#).
7. Fava M. Diagnosis and definition of treatment-resistant depression. *Biol Psychiatry*. 2003;53(8):649-59. PubMed PMID: [12706951](#).
8. Fekadu A, Wooderson S, Donaldson C, Markopoulou K, Masterson B, Poon L, et al. A multidimensional tool to quantify treatment resistance in depression: the Maudsley staging method. *J Clin Psychiatry*. 2009;70(2):177-84. PubMed PMID: [19192471](#).
9. Sackeim HA. The definition and meaning of treatment-resistant depression. *J Clin Psychiatry*. 2001;62 Suppl 16:10-7. PubMed PMID: [11480879](#).
10. Hirschfeld RM, Keller MB, Panico S, Arons BS, Barlow D, Davidoff F, et al. The National Depressive and Manic-Depressive Association consensus statement on the undertreatment of depression. *JAMA*. 1997;277(4):333-40. PubMed PMID: [9002497](#).

How to cite this editor-reviewed article

Werremeyer A. Treatment-resistant depression. *Ment Health Clin* [Internet]. 2014;4(5):211. Available from: <http://dx.doi.org/10.9740/mhc.n207177>