

# Use of very high resolution climate model data for hydrological modelling: estimation of potential evaporation

Alison C. Rudd and Alison L. Kay

## ABSTRACT

Climate model data are increasingly used to drive hydrological models, to assess the possible impacts of climate change on river flows. Hydrological models often require potential evaporation (PE) from vegetation, alongside precipitation, but PE is not usually output by climate models so has to be estimated from other meteorological variables. Here, the Penman–Monteith formula is applied to estimate PE using data from a 12 km Regional Climate Model (RCM) and a nested very high resolution (1.5 km) RCM covering southern Britain. PE estimates from RCM runs driven by reanalysis boundary conditions are compared to observation-based PE data, to assess performance. The comparison shows that both the 1.5 and 12 km RCMs reproduce observation-based PE well, on daily and monthly time-steps, and enables choices to be made about application of the formula using the available data. Data from Current and Future RCM runs driven by boundary conditions from a Global Climate Model are then used to investigate potential future changes in PE, and how certain factors affect those changes. In particular, the importance of including changes in canopy resistance is demonstrated. PE projections are also shown to vary to some extent according to how aerosols are modelled in the RCMs.

**Key words** | canopy conductance, climate change, high resolution, potential evaporation, Regional Climate Model, stomata

**Alison C. Rudd** (corresponding author)  
**Alison L. Kay**  
Centre for Ecology & Hydrology,  
Maclean Building, Crowmarsh Gifford,  
Wallingford,  
Oxfordshire OX10 8BB,  
UK  
E-mail: [alirud@ceh.ac.uk](mailto:alirud@ceh.ac.uk)

## INTRODUCTION

There is increasing concern about the potential impacts of climate change on the hydrological cycle (Stocker *et al.* 2013). Modelling the possible hydrological impacts is particularly important as changes in the water cycle can affect people and ecosystems both directly (e.g. via changes in water availability and flood frequency) and indirectly (e.g. via changes in food and energy production) (Jiménez Cisneros *et al.* 2014).

Using Regional Climate Model (RCM) data as an input to hydrological models allows investigation of how climate change may affect river flows (e.g. Ott *et al.* 2013; Kay & Jones 2012). Many hydrological models require inputs of potential evaporation (PE) from vegetated surfaces, alongside precipitation (Bartholomeus *et al.* 2015). Unlike

precipitation, PE is usually not available directly from RCMs, so has to be estimated from other variables. The meteorological variables that influence PE are temperature, radiation, humidity and wind speed. In addition, vegetation factors such as leaf area and roughness affect the transpiration component (Kay *et al.* 2013). Many formulae exist for estimating PE, ranging from the physically based Penman–Monteith formula (Monteith 1965) to much simpler empirical formulae like that of Oudin *et al.* (2005), and the choice can affect the results of subsequent hydrological modelling (e.g. Kay & Davies 2008; Seiller & Ancil 2014). There is much disagreement on the best approach when deriving PE from climate model data for future periods, with concerns about empirical formulae not explicitly

including changes in all the influencing variables, but also concerns about data quality when using more complex formulae (see discussion by [Kay \*et al.\* \(2013\)](#)).

As part of a recent Natural Environment Research Council (NERC) Changing Water Cycle project, CONVEX, the Met Office ran a very high resolution (1.5 km) RCM for southern Britain, nested in a 12 km RCM driven by global atmospheric reanalysis (ERA-Interim) boundary conditions (1989–2008). They also ran Current (1996–2009) and Future (~2100 s) climate simulations, with both aerosol climatology and full aerosol modelling setups, nesting the RCMs in a Global Climate Model (GCM). [Kendon \*et al.\* \(2012\)](#) found that rainfall in the 1.5 km RCM is more realistic than in the 12 km RCM. In the 12 km RCM, there is a tendency for heavy rain events to be too persistent and widespread, and not heavy enough. Conversely, the 1.5 km RCM has a tendency for heavy rain to be too intense, but it still gives a much better representation of duration and spatial extent.

This paper uses the RCM data from the CONVEX project to estimate PE for short grass, using the Penman–Monteith formula. The following questions are considered: how do RCM estimates of PE compare with observation-based PE; how do the 12 and 1.5 km estimates of PE compare; and how might PE change in the future due to climate change, and what factors influence this PE change?

PE estimates from the ERA-driven RCM runs are compared against observation-based PE from the Met Office Rainfall and Evaporation Calculation System (MORECS) ([Hough \*et al.\* 1996](#)), as MORECS is the closest to an observational estimate of PE and is widely used by the hydrological community in Britain. Also, the RCM PE estimates are required for an investigation of the use of very high resolution data for hydrological modelling, in which the hydrological models to be used are tuned using MORECS PE along with observed precipitation data ([Bell \*et al.\* 2012](#); [Crooks \*et al.\* 2014](#)). However, MORECS PE is only produced on a 40 × 40 km grid of squares across the UK. The RCM data thus provides the opportunity to estimate PE using a much finer resolution. The comparison includes an assessment of several choices available within the PE estimation method.

Future changes in PE are investigated using the GCM-driven Current and Future RCM simulations. Relatively few studies have looked at potential future changes in PE

in Britain, and even fewer have looked at historical changes, either in Britain or globally, but the studies that do exist generally suggest increases ([Kay \*et al.\* 2013](#)). However, most of these studies have calculated PE changes only from changes in (some of) the meteorological variables; PE can also be affected by increases in the atmospheric concentration of CO<sub>2</sub> via changes in stomatal resistance ([Bell \*et al.\* 2011](#); [Pan \*et al.\* 2015](#)). The effect of changes in stomatal resistance is considered here, as is the influence of the method of including aerosols in the RCMs.

Although the focus of this paper is use of RCM data to produce PE estimates that will subsequently be used to drive hydrological models, the issues highlighted will be of wider interest. For example, PE can be an important component in crop modelling ([Lovelli \*et al.\* 2010](#)) and ecological modelling ([Fisher \*et al.\* 2011](#)).

## METHODOLOGY

### The RCM

The 1.5 km RCM is a climate version of the UK Met Office 1.5 km weather forecast model (UKV) ([Kendon \*et al.\* 2012](#)) but with a smaller domain, spanning southern England and Wales ([Figure 1](#)). The 1.5 km RCM lateral boundary conditions are supplied by the 12 km RCM, which is a limited-area (European domain) atmosphere-only version of the Met Office Hadley Centre Global Environmental Model. For the ERA-driven simulations, the 12 km RCM is driven at its lateral boundaries by the latest European Centre for Medium-Range Weather Forecasts Re-Analysis (ERA-Interim) ([Dee \*et al.\* 2011](#)) for the period 1989 to 2008. For the GCM-driven simulations the 12 km RCM uses boundary conditions from the HadGEM3 GCM, for Current and Future periods. [Table 1](#) summarises the various RCM simulations available, including two versions of the GCM-driven runs with different aerosol formulations (see below). [Kendon \*et al.\* \(2012\)](#) found that it takes a few months for soil moisture to spin up in the 1.5 km RCM, therefore only data from January of the first full year of each run are included.

For the Current and Future GCM-driven runs, there are both aerosol climatology and fully coupled aerosol



**Figure 1** | Map showing the 1.5 km RCM domain (dotted line) and the locations of three MORECS sites used in the analysis (points).

modelling runs. The aerosol modelling runs differ from the aerosol climatology runs in that aerosol concentrations are calculated interactively, with advection and deposition of aerosols described. In the case of the 12 km RCM, the aerosols are fully coupled to the microphysics, so that aerosol concentrations determine the number of cloud condensation nuclei. A comparison is made of PE from both pairs of model runs for the 12 km RCM, as data are available for the full period of ~13 years. The equivalent 1.5 km RCM aerosol modelling runs are only about 5 years in

length (Table 1) and are therefore not included in this analysis.

### Estimating PE from atmospheric data

PE is generally considered as the amount of water that would be lost to the atmosphere if there were no limits to soil-moisture supply (Kay *et al.* 2013). The Penman–Monteith method is recommended by the United Nations Food and Agriculture Organization (FAO) for deriving grass reference PE (Pereira *et al.* 1999) and is used by the UK Climate Projection 09 weather generator (Jones *et al.* 2009). Penman–Monteith PE (mm/s) is given by

$$PE = \frac{1}{\lambda} \frac{\Delta R_n + \rho_a c_a (e_s - e_d) / r_a}{\Delta + \gamma (1 + r_s / r_a)}, \quad (1)$$

where  $\lambda$  is the latent heat of vaporisation ( $\text{J kg}^{-1}$ ),  $\Delta$  is the rate of change of saturated vapour pressure with temperature ( $\text{kPa } ^\circ\text{C}^{-1}$ ),  $R_n$  is the net radiation ( $\text{J m}^{-2} \text{s}^{-1}$ ),  $\rho_a$  is the near surface air density ( $\text{kg m}^{-3}$ ),  $c_a$  is the specific heat of air ( $\text{J kg}^{-1} ^\circ\text{C}^{-1}$ ),  $e_s$  is the saturation vapour pressure at screen temperature (kPa),  $e_d$  is the screen vapour pressure (kPa),  $\gamma$  is the psychrometric constant ( $\text{kPa } ^\circ\text{C}^{-1}$ ),  $r_a$  is the aerodynamic resistance to vapour transfer in the atmosphere ( $\text{sm}^{-1}$ ) and  $r_s$  is the bulk surface (canopy or bare soil) resistance ( $\text{sm}^{-1}$ ). The saturation vapour pressure  $e_s$  at temperature  $T$  ( $^\circ\text{C}$ ) is given by

$$e_s(T) = 0.611 \exp\left(\frac{17.27T}{T + 237.3}\right), \quad (2)$$

$$\text{so } \Delta = de_s/dT = 17.27 \times 237.3 e_s(T) / (T + 237.3)^2.$$

**Table 1** | Summary of 12 and 1.5 km RCM runs, with run IDs and time periods

RCM run	Run ID		Time period	Other details
	12 km	1.5 km		
ERA-driven baseline	ajtyr	akigd	April 1989 – November 2008	
GCM-driven Current (clim)	alqtj	alxmc	May 1996 – November 2009	aerosol climatology
GCM-driven Future (clim)	alqtk	alxme	as above but for ~2100s	aerosol climatology, RCP 8.5 emissions
GCM-driven Current (mod)	alqtl	alxmk	May 1996 – November 2009 (12 km). June 1996 – March 2002 (1.5 km)	aerosol modelling
GCM-driven Future (mod)	alqtm	alxml	as above but for ~2100 s	aerosol modelling, RCP 8.5 emissions

Two different formulations are considered for calculating vapour pressure  $e_d$  from relative humidity RH (%) and temperature ( $^{\circ}\text{C}$ ); one uses mean temperature  $T$

$$e_d = \frac{\text{RHe}_s(T)}{100}, \quad (3)$$

and the other uses the min and max temperature ( $T_{\min}$ ,  $T_{\max}$ ) (Allen *et al.* 1994)

$$e_d = \frac{\text{RH}}{\left(\frac{50}{e_s(T_{\min})} + \frac{50}{e_s(T_{\max})}\right)}. \quad (4)$$

The aerodynamic resistance  $r_a$  is calculated from the 10 m wind speed  $U_{10}$  ( $\text{ms}^{-1}$ ) using

$$r_a = \frac{243.489}{U_{10}}, \quad (5)$$

which includes a logarithmic correction for wind height (Hough *et al.* 1996), and surface resistance  $r_s$  is calculated using

$$r_s = \frac{1}{((1-A)/r_{sc}) + (A/r_{ss})}, \quad (6)$$

where  $A = 0.7^L$ ,  $L$  is leaf area index (LAI),  $r_{sc}$  is crop resistance and  $r_{ss}$  is bare soil resistance ( $100 \text{ sm}^{-1}$ ) (Hough *et al.* 1996). MORECS monthly values of  $r_{sc}$  and LAI for short grass are used (Hough *et al.* 1996) (Table 2).

The climate model variables used for the calculation of PE are thus 1.5 m temperature, 1.5 m relative humidity, 10 m wind speed and net surface downward longwave and shortwave radiation (which sum to  $R_n$ ).

### Estimating PE for future RCM runs

PE can be affected not just by changes in meteorological inputs, but also by changes in the behaviour of vegetation. In particular, higher  $\text{CO}_2$  concentrations can lead to plant stomata opening less widely, resulting in higher stomatal resistance, but can also enhance plant growth, leading to a greater leaf area and more stomata (e.g. Bunce 2004; Keenan *et al.* 2013). Here, following Kruijt *et al.* (2008),

**Table 2** | Monthly leaf area index (LAI) and crop resistance for Current ( $r_{sc,M}$ ) and Future ( $r_{sc,F}$ ) periods, for short grass. LAI and  $r_{sc,M}$  are from MORECS, and  $r_{sc,F}$  is calculated from  $r_{sc,M}$  using Equation (7)

Month	LAI	Current crop resistance $r_{sc,M}$	Future crop resistance $r_{sc,F}$
January	2.0	80	168
February	2.0	80	168
March	3.0	60	126
April	4.0	50	105
May	5.0	40	84
June	5.0	60	126
July	5.0	60	126
August	5.0	70	147
September	4.0	70	147
October	3.0	70	147
November	2.5	80	168
December	2.0	80	168

changes in crop resistance are considered but changes in leaf area are not, as it was found by Bunce (2004) that there is a lack of response in LAI due to elevated  $\text{CO}_2$  concentrations in all functional types, except trees; only grass PE is considered here.

For the Future RCM runs, the equation of Kruijt *et al.* (2008) is used to estimate appropriate values of grass  $r_{sc}$  and then  $r_s$ . Kruijt *et al.* (2008) found that the average change in grass and herbal crop conductance ( $g_{sc} = 1/r_{sc}$ ) per 1 ppm increase in atmospheric  $\text{CO}_2$  concentration is  $-9.3 \times 10^{-2}\% \pm 1.5 \times 10^{-2}\%$  (under well-watered conditions). The increase in atmospheric  $\text{CO}_2$  for RCP 8.5 emissions up to year 2100 is 562 ppm, therefore the increase in  $r_{sc}$  is 109.5%, or

$$r_{sc,F} = 2.095r_{sc,M}, \quad (7)$$

where  $r_{sc,M}$  are the monthly MORECS grass  $r_{sc}$  values and  $r_{sc,F}$  are the values adjusted for the future climate (Table 2). Surface resistance  $r_s$  is then calculated from  $r_{sc,F}$  using Equation (6). The subsequent  $r_s$  values compare well with other approaches to estimating future  $r_s$ , such as taking the average change in  $r_s$  from members of the UKCP09 RCM ensemble (Murphy *et al.* 2009) and applying pattern scaling (Mitchell 2003) to allow for differences in emissions scenarios (not shown), and  $r_s$  changes used by Moratier *et al.* (2011).

## RESULTS

### ERA-driven RCM runs

To choose a calculation of vapour pressure (Equations (3) and (4)), the 12 and 1.5 km estimates of PE are compared with MORECS PE at different timescales, daily and

monthly, for three sites in Britain (Figure 1) for the year 1990. The sites were chosen to give spatial coverage across the UK; a southern site (Lyneham), a Midlands site (Nottingham) and a northern site (Galashiels) (not covered by the 1.5 km domain). Figure 2 shows the PE using Equation (3) for vapour pressure; equivalent figures for PE calculated using Equation (4) are similar (not shown). Both estimates




Figure 2 | Comparison of daily and monthly MORECS PE and ERA-driven RCM PE for 1990, for three MORECS sites.



of PE are comparable to MORECS, however Equation (3) was chosen as it compares slightly better to MORECS PE.

To estimate monthly PE, two different averaging methods were compared (for 1990 data): (a) calculating daily PE from daily meteorological variables and then averaging, and (b) calculating monthly averages of the meteorological variables and then calculating monthly PE from them. The two methods yield almost identical results (not shown) so daily PE are made and then the monthly average is calculated (the monthly plots in Figure 2 use this method). This test was done as, although daily data were available here, it can be the case that only monthly data are available, or that it is impractical to obtain all the required data at a daily rather than monthly time-step. In such situations, use of monthly data to calculate monthly PE is unlikely to cause problems (as also suggested by Allen *et al.* (1998)).

Figure 3 presents maps of the seasonal mean PE for MORECS and the 12 and 1.5 km ERA-driven RCMs for 1990–2007. It shows that the RCMs compare well with MORECS and with each other. The comparison looks to be best for spring (MAM) and winter (DJF). The 12 km RCM might be slightly overdoing summer (JJA) PE in the south-east compared to MORECS and the 1.5 km RCM.

### Current and Future RCM runs


Figure 4 shows maps of seasonal mean PE from the Current and Future aerosol climatology RCM runs, and the percentage change between them. The Current PE for the 12 km and 1.5 km RCMs are similar to each other in magnitude and spatial pattern, with the highest PE in the summer in the southeast and the lowest PE in winter, as for MORECS and the ERA-driven runs (Figure 3). The Future PE from the 12 and 1.5 km RCMs is also similar, but consistently higher than Current PE; PE increases across the country and throughout the year. The percentage change in PE for the 1.5 km run looks larger in winter compared to the 12 km run, however the other seasons look more similar between the two resolutions.

Figure 5 shows the seasonal mean percentage change in PE for three different setups. The left hand column is for the aerosol climatology 12 km RCM with fixed MORECS values of crop resistance ( $r_{sc}$ ), the middle column is for the aerosol



**Figure 3** | Comparison of seasonal mean PE (mm/day) from MORECS and the 12 and 1.5 km ERA-driven RCMs, for 1990–2007.

climatology 12 km RCM with adjusted values of future  $r_{sc}$ , and the right hand column shows the equivalent plot for the aerosol modelling 12 km RCM with adjusted future  $r_{sc}$ . This shows that the percentage changes in seasonal mean PE are much more similar between the aerosol climatology and full aerosol modelling runs (middle and right) than between the adjusted future  $r_{sc}$  and fixed MORECS  $r_{sc}$  runs (middle and left). For the adjusted future  $r_{sc}$  runs, the summer and autumn percentage changes in PE are larger in the aerosol climatology run than the aerosol modelling run, although the winter and spring changes are similar. Not accounting for the change in stomatal resistance gives a much larger increase in PE from the Current to the Future (left column of Figure 5).



**Figure 4** | Seasonal mean PE (mm/day) under climate change, including the effect of changes in stomatal resistance. The averages are for 1997–2008 for the Current RCM and for 12 years in the 2100s for the Future RCM. The percentage change between the Current and Future PE is also shown.


To take a closer look at the difference in PE changes between the RCMs, **Figure 6** shows how the monthly mean percentage change in PE varies for the three MORECS sites (**Figure 1**). As well as re-emphasising that the percentage change in PE using the fixed MORECS  $r_{sc}$  values is consistently higher than the other two runs, these plots also show that the percentage change in PE for the full aerosol modelling run is lower than that from the aerosol climatology run, for June to September. For the other months the values are more comparable. Fully modelling the aerosols appears to have the effect of lowering the summer PE.

**Figure 7** shows the seasonal mean percentage change in shortwave (solar) and longwave radiation for the aerosol climatology and aerosol modelling runs. It shows a larger percentage change in summer radiation in the aerosol climatology run (left column) compared to the aerosol modelling run, especially for shortwave radiation. The seasonal mean percentage change in temperature, relative

humidity and wind are more similar for the aerosol climatology and aerosol modelling runs (not shown), so the differences seen in the estimated PE are likely to be due to the radiation differences.


## DISCUSSION AND CONCLUSIONS

Estimates of PE from high-resolution (12 and 1.5 km) RCM data have been produced, using the physically based Penman–Monteith formula, with the aim of making PE suitable to use as input for hydrological modelling. Using ERA-driven RCM data, and a comparison with the observation-based MORECS PE, choices were made about the method to use. The comparison shows that both the 1.5 and 12 km RCM PE estimates are very similar to each other and comparable to MORECS, spatially and at selected locations, at daily and monthly time-steps.



**Figure 5** | Seasonal mean percentage change in PE for the 12 km RCM. Left is fixed MORECS  $r_{sc}$  for both Current and Future, with aerosol climatology. Middle is MORECS  $r_{sc}$  for Current and adjusted  $r_{sc}$  for Future, with aerosol climatology. Right is MORECS  $r_{sc}$  for Current and adjusted  $r_{sc}$  for Future, with full aerosol modelling.


PE estimates from Current and Future RCM runs driven by GCM boundary conditions have been used to investigate potential future changes in PE, and how certain factors affect those changes. It is found that the seasonal mean PE and its change are in close agreement between the 1.5 and 12 km models; this is perhaps not surprising because the large scale changes in humidity, temperature and circulation are common to both RCMs (inherited from the driving GCM). It is also found that future PE is likely to be larger than at present, with the largest increases in summer, autumn and (to a lesser extent) winter. This PE change is influenced by the changes in temperature, relative humidity, wind and radiation as well as stomatal influences through increased  $\text{CO}_2$ . Not accounting for the



**Figure 6** | Comparison of percentage change in mean monthly 12 km RCM PE for aerosol climatology and full aerosol modelling runs for three MORECS sites.

stomatal influences can inflate the future PE estimates, consistent with the findings of Bell *et al.* (2011) for UK annual and seasonal mean PE, and such differences can affect subsequent flow projections (e.g. Bell *et al.* 2012; Prudhomme *et al.* 2014).





**Figure 7** | Seasonal mean percentage change in net downward surface radiation (left: shortwave, right: longwave) for the aerosol climatology and full aerosol modelling runs of the 12 km RCM.

PE projections are also shown to vary to some extent according to how aerosols are modelled in the RCMs. The PE differences from aerosol differences are in the summer/early autumn and this could potentially be important for hydrological modelling as autumn PE that is too high could delay the rewetting of soils in the lead-up to the main flood season in Britain (Bayliss & Jones 1993). The smaller percentage changes

in PE for the aerosol modelling run appear to be related to smaller changes in radiation. The potential hydrological importance of aerosol concentrations, via their influence on radiation and so evaporation, is also shown by Gedney *et al.* (2014), who identify a link between solar dimming, due to rising atmospheric concentrations of aerosols from 1980, and increases in runoff.

Use of the Penman–Monteith PE formulation enabled specific application of changes to crop resistance alongside future changes in meteorological variables. This is not possible (at least in such a straightforward way) with most of the simpler PE formulae, where empirically derived coefficients replace many of the factors present in the more physically based formulae (see for example the summary of 17 PE variations given by Oudin *et al.* (2005)). There are already concerns about the application of empirical formulae under changing climates (e.g. Donohue *et al.* 2010; Bartholomeus *et al.* 2015), and the need to also consider changes in canopy resistance is an added complication (Kay *et al.* 2013). The use of fixed crop coefficients to estimate crop PE from reference PE is an additional factor that requires consideration under climate change (Bartholomeus *et al.* 2015), both for hydrological modelling and crop modelling, as different crops may react differently to the same change in CO<sub>2</sub> concentrations (Kruijt *et al.* 2008). Even for hydrological models that do not specifically require PE inputs, similar considerations are likely to apply in the model's internal calculation of evaporation when applied under changing climatic conditions.

Future work will involve running CLASSIC-GB (Crooks *et al.* 2014) with the high-resolution RCM data to investigate the effect of model resolution, and future climate changes, on peak river flows in southern Britain. As the 1.5 km full aerosol modelling run is not full length, the aerosol climatology run will be used to get sufficient length to look at flood frequencies.

## ACKNOWLEDGEMENTS

The authors would like to thank Elizabeth Kendon and Richard Jones (Met Office Hadley Centre) for providing the data and help with analysis of results. This work was funded by the Natural Hazards science area of the NERC-CEH Water and Pollution Science programme.

## REFERENCES

- Allen, R., Smith, M., Perrier, A. & Pereira, L. 1994 An update for the definition of reference evapotranspiration. *ICID Bulletin of the International Commission on Irrigation and Drainage* **43**, 1–35.
- Allen, R. G., Pereira, L. S., Raes, D. & Smith, M. 1998 *Crop evapotranspiration (guidelines for computing crop water requirements)*. FAO Irrigation and Drainage Paper No. 56, Food and Agriculture Organization (FAO) of the United Nations, Rome, Italy.
- Bartholomeus, R. P., Stagge, J. H., Tallaksen, L. M. & Witte, J. P. M. 2015 Sensitivity of potential evaporation estimates to 100 years of climate variability. *Hydrol. Earth Syst. Sci.* **19**, 997–1014.
- Bayliss, A. & Jones, R. 1993 *Peaks-over-threshold flood database: Summary statistics and seasonality*. 121. IH Report.
- Bell, V., Gedney, N., Kay, A., Smith, R., Jones, R. & Moore, R. 2011 Estimating potential evaporation from vegetated surfaces for water management impact assessments using climate model output. *Journal of Hydrometeorology* **12**, 1127–1136.
- Bell, V., Kay, A., Cole, S., Jones, R., Moore, R. & Reynard, N. 2012 How might climate change affect river flows across the Thames Basin? An area-wide analysis using the UKCP09 Regional Climate Model ensemble. *Journal of Hydrology* **442**, 89–104.
- Bunce, J. 2004 Carbon dioxide effects on stomatal responses to the environment and water use by crops under field conditions. *Oecologia* **140**, 1–10.
- Crooks, S., Kay, A., Davies, H. & Bell, V. 2014 From catchment to national scale rainfall-runoff modelling: demonstration of a hydrological modelling framework. *Hydrology* **1**, 63–88.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N. & Vitart, F. 2011 The ERA-Interim reanalysis: configuration and performance of the data assimilation system. *Quarterly Journal of the Royal Meteorological Society* **137**, 553–597.
- Donohue, R. J., McVicar, T. R. & Roderick, M. L. 2010 Assessing the ability of potential evaporation formulations to capture the dynamics in evaporative demand within a changing climate. *Journal of Hydrology* **386**, 186–197.
- Fisher, J. B., Whittaker, R. J. & Malhi, Y. 2011 ET Come home: potential evapotranspiration in geographical ecology. *Global Ecology and Biogeography* **20**, 1–18.
- Gedney, N., Huntingford, C., Weedon, G., Bellouin, N., Boucher, O. & Cox, P. 2014 Detection of solar dimming and brightening effects on Northern Hemisphere river flow. *Nature Geoscience* **7**, 796–800.
- Hough, M., Palmer, S., Weir, A., Lee, M. & Barrie, I. 1996 *The Meteorological Office Rainfall and Evaporation Calculation System: MORECS version 2.0 (1995)*. An update to Hydrological Memorandum 45, Met Office Hadley Centre, Exeter, UK.
- Jiménez Cisneros, B. E., Oki, T., Arnell, N. W., Benito, G., Cogley, J. G., Döll, P., Jiang, T. & Mwakalila, S. S. 2014 Freshwater

- resources. In: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (C. B. Field *et al.* eds). Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 229–269.
- Jones, P., Kilsby, C., Harpham, C., Glenis, V. & Burton, A. 2009 *UK Climate Projections Science Report: Projections of Future Daily Climate for the UK from the Weather Generator*.
- Kay, A. L. & Davies, H. N. 2008 [Calculating potential evaporation from climate model data: a source of uncertainty for hydrological climate change impacts](#). *Journal of Hydrology* **358**, 221–239.
- Kay, A. L. & Jones, R. G. 2012 [Comparison of the use of alternative UKCP09 products for modelling the impacts of climate change on flood frequency](#). *Climatic Change* **114**, 211–230.
- Kay, A., Bell, V., Blyth, E., Crooks, S., Davies, H. & Reynard, N. 2013 [A hydrological perspective on evaporation: historical trends and future projections in Britain](#). *Journal of Water and Climate Change* **4**, 193–208.
- Keenan, T. F., Hollinger, D. Y., Bohrer, G., Dragoni, D., Munger, J. W., Schmid, H. P. & Richardson, A. D. 2013 [Increase in forest water-use efficiency as atmospheric carbon dioxide concentrations rise](#). *Nature* **499**, 324–327.
- Kendon, E., Roberts, N., Senior, C. & Roberts, M. 2012 [Realism of rainfall in a very high-resolution regional climate model](#). *Journal of Climate* **25**, 5791–5806.
- Kruijt, B., Witte, J.-P., Jacobs, C. & Kroon, T. 2008 [Effects of rising atmospheric CO<sub>2</sub> on evapotranspiration and soil moisture: a practical approach for the Netherlands](#). *Journal of Hydrology* **349**, 257–267.
- Lovelli, S., Perniola, M., Di Tommaso, T., Ventrella, D., Moriondo, M. & Amato, M. 2010 [Effects of rising atmospheric CO<sub>2</sub> on crop evapotranspiration in a Mediterranean area](#). *Agricultural Water Management* **97**, 1287–1292.
- Mitchell, T. 2003 [Pattern scaling: an examination of the accuracy of the technique for describing future climates](#). *Climatic Change* **60**, 217–242.
- Monteith, J. 1965 [Evaporation and environment](#). *Symp. Soc. Exp. Biol.* **19**, 205–234.
- Moratiel, R., Snyder, R., Durán, J. & Tarquis, A. 2011 [Trends in climatic variables and future reference evapotranspiration in Duero Valley \(Spain\)](#). *Natural Hazards and Earth System Science* **11**, 1795–1805.
- Murphy, J., Sexton, D., Jenkins, G., Boorman, P., Booth, B., Brown, K., Clark, R., Collins, M., Harris, G. & Kendon, L. 2009 *UK Climate Projections Science Report: Climate change projections*. Met Office Hadley Centre, Exeter, UK.
- Ott, I., Duethman, D., Liebert, J., Berg, P., Feldmann, H., Ihringer, J., Kunstmann, H., Merz, B., Schaedler, G. & Wagner, S. 2013 [High-resolution climate change impact analysis on medium-sized river catchments in Germany: an ensemble assessment](#). *Journal of Hydrometeorology* **14**, 1175–1193.
- Oudin, L., Hervieu, F., Michel, C., Perrin, C., Andreassian, V., Anctil, F. & Loumagne, C. 2005 [Which potential evapotranspiration input for a lumped rainfall-runoff model? Part 2 – Towards a simple and efficient potential evapotranspiration model for rainfall-runoff modelling](#). *Journal of Hydrology* **303**, 290–306.
- Pan, S., Tian, H., Dangal, S. R. S., Yang, Q., Yang, J., Lu, C., Tao, B., Ren, W. & Ouyang, Z. 2015 [Responses of global terrestrial evapotranspiration to climate change and increasing atmospheric CO<sub>2</sub> in the 21st century](#). *Earth's Future* **3**, 15–35.
- Pereira, L., Perrier, A., Allen, R. & Alves, I. 1999 [Evapotranspiration: concepts and future trends](#). *Journal of Irrigation and Drainage Engineering* **125**, 45–51.
- Prudhomme, C., Giuntoli, I., Robinson, E. L., Clark, D. B., Arnell, N. W., Dankers, R., Fekete, B. M., Franzen, W., Gerten, D., Gosling, N. H., Hagemann, S., Hannah, D. M., Kim, H., Masaki, Y., Satoh, Y., Stacke, T., Wada, Y. & Wisser, D. 2014 [Hydrological droughts in the 21st century, hotspots and uncertainties from a global multimodel ensemble experiment](#). *PNAS* **111**, 3262–3267.
- Seiller, G. & Anctil, F. 2014 [Climate change impacts on the hydrologic regime of a Canadian river: comparing uncertainties arising from climate natural variability and lumped hydrological model structures](#). *Hydrol. Earth Syst. Sci.* **18**, 2033–2047.
- Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V. & Midgley, P. M. (eds) 2013 *IPCC, 2013: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK, New York, NY, USA, p. 1535.

First received 16 February 2015; accepted in revised form 5 May 2015. Available online 23 June 2015