

# Comparative evaluation of 24 reference evapotranspiration equations applied on an evergreen-broadleaved forest

Athanassios Bourletsikas, Ioannis Argyrokastritis and Nikolaos Proutsos

## ABSTRACT

Reference evapotranspiration ( $ET_0$ ) is a major component of the hydrological cycle. Its use is essential both for the hydrological rainfall–runoff assessment models and determination of water requirements in agricultural and forest ecosystems. This study investigates the performance of 24 different methods, which produce  $ET_0$  or potential evapotranspiration estimates above a grass-covered ground in a Mediterranean forest environment in Greece and compares the derived results with those of the presumed most accurate and scientifically acceptable Penman–Monteith method ( $ET_{P-M}$ ). Their performance was evaluated on a daily basis for a period of 17 years, using 17 different statistical parameters of goodness of fit. The results showed that some empirical methods could serve as suitable alternatives. More specifically, Copais ( $ET_{COP}$ ), Hargreaves original ( $ET_{HAR}$ ), and Valiantzas2 ( $ET_{VA2}$ ) methods, exhibited very good values of the model efficiency index, EF (0.934, 0.932, and 0.917, respectively) and the index of agreement,  $d$  (0.984, 0.982, and 0.977, respectively). Additionally, the differences of the estimated mean daily value against the respective  $ET_{P-M}$  value (rt index) for all methods had a range of  $-27.8\%$  (Penman –  $ET_{PEN}$ ) to  $+59.5\%$  (Romanenko –  $ET_{ROM}$ ), while Copais ( $ET_{COP}$ ), Hargreaves–Samani modified1 ( $ET_{HS1}$ ), and STU ( $ET_{STU}$ ) yielded the best values ( $-0.06\%$ ,  $+0.06\%$ , and  $0.22\%$ , respectively).

**Key words** | comparative analysis, evergreen-broadleaved forest, FAO56 Penman–Monteith, Mediterranean climate, reference evapotranspiration, western Greece

**Athanassios Bourletsikas** (corresponding author)  
**Nikolaos Proutsos**  
 Department of Forest Ecology and Hydrology,  
 Institute of Mediterranean Forest Ecosystems,  
 Terma Alkmanos, Ilissia,  
 Athens,  
 Greece  
 E-mail: [mpat@fria.gr](mailto:mpat@fria.gr)

**Ioannis Argyrokastritis**  
 Department of Natural Resources Development  
 and Agricultural Engineering,  
 Agricultural University of Athens,  
 Athens,  
 Greece

## INTRODUCTION

Evapotranspiration (ET) is an important component of both the water and energy cycles. ET is used in agricultural and forest hydrometeorology and in urban planning. It is also used in many rainfall–runoff and ecosystem models (Vörösmarty *et al.* 1998; Hay & McCabe 2002; Oudin *et al.* 2005a, 2005b), even for the estimation of ecosystem productivity (Currie 1991). ET also affects significantly regional water availability and use (Zhang *et al.* 2001; Sun *et al.* 2006). In order to carry out a long-term study of hydrological, environmental, and ecological processes, reliable estimates and/or accurate measurements of ET are required (Rosenberry *et al.* 2007; Tabari *et al.* 2011).

The concept of ‘reference evapotranspiration’ ( $ET_0$ ) was defined by Doorenbos & Pruitt (1977) as ‘the rate of evapotranspiration from an extensive surface of 8 to 15 cm tall, green

grass cover of uniform height, actively growing, completely shading the ground and not short of water’. Allen *et al.* (1998) evolved an  $ET_0$  definition based on a reference surface (a hypothetical grass and/or alfalfa reference crop) to define unique evaporation parameters for each crop and growth stage. Hence, the FAO Expert Consultation on Revision of FAO Methodologies for Crop Water Requirements accepted the following unambiguous definition for the reference surface: ‘A hypothetical crop with an assumed height of 0.12 m having a surface resistance of  $70 \text{ s m}^{-1}$  and an albedo of 0.23, closely resembling the evaporation of an extension surface of green grass of uniform height, actively growing and adequately watered’ (Allen *et al.* 1998). The method proposed by Allen *et al.* (1998) was the FAO56 Penman–Monteith ( $ET_{P-M}$ ) equation, which was accepted worldwide as the best

doi: 10.2166/nh.2017.232

estimator of  $ET_0$  in many regions and in different weather conditions (Droogers & Allen 2002; Xu & Singh 2002; Oudin et al. 2005a, 2005b; Alexandris et al. 2006; Gavilán et al. 2006; Tabari et al. 2011; Rahimikhoob et al. 2012; Khoshravesh et al. 2015; Valipour 2015a, 2015b).

As  $ET_0$  is considered to be the most difficult component to estimate, the wealth of  $ET_0$  methods and empirical equations proposed by many researchers have certain strong points and limitations depending on the methods' applications and assumptions (Rana & Katerji 2000; Grismer et al. 2002; Valipour 2014a, 2014b). Additionally, past studies at various scales proved that different  $ET_0$  methods gave widely different values at particular locations (Federer et al. 1996; Vörösmarty et al. 1998). This means that all these methods cannot be used globally as they need calibration for regional application (Kolka & Wolf 1998; Grismer et al. 2002; Xu & Singh 2002; Rosenberry et al. 2004; Lu et al. 2005; Tabari et al. 2011; Rahimikhoob et al. 2012; Xu et al. 2013; Bogawski & Bednorz 2014; Samaras et al. 2014; Valipour & Eslamian 2014; Valipour 2015c). The selection of the appropriate method based on the availability of data, its cost, estimation accuracy, operational time and space scales, is challenging.

There are many methods in the literature used for the estimation of  $ET_0$ . These methods can be grouped into categories depending on the variables needed for input. The main categories reported in the literature are: mass-transfer, temperature-based, radiation-based, pan-evaporation, and combination. Many researchers reported overviews by using these methods and categories (Jensen et al. 1990; Xu & Singh 2002; Rosenberry et al. 2004, 2007; Oudin et al. 2005a; Alexandris et al. 2008; Trajkovic & Kolakovic 2009; Tabari et al. 2011; Xystrakis & Matzarakis 2011; Xu et al. 2013; Valipour 2014c) in different areas and environments.

It would be really interesting to investigate the performance of the  $ET_0$  methods in a forested area because: (1) forests affect the climatic status variability since they influence the hydrological and carbon cycles at regional and global scale (Houghton 1991; Musselman & Fox 1991; Nepsstad et al. 1994) and (2) it is well established that forested sites and catchments have higher ET rates than grassed catchments (Zhang et al. 2001). In the literature there is a lack of studies dealing with  $ET_0$  in forests. This is due to the costly equipment requirements (lysimeters, eddy covariance towers, etc.) and to the fact that the main interest of the

scientific community focuses on agriculture (Fisher et al. 2005; Alexandris et al. 2008).

Such studies have been performed mainly in coniferous species. Indicatively, McNaughton & Black (1973) measured the ET in a Douglas fir forest for 18 days and came up with a proposed estimation method. Scholl (1976) determined the ET in a Chaparral stand. Spittlehouse & Black (1980) used the Bowen ratio/energy balance method to measure the ET of a thinned Douglas fir forest. Riekerk (1985) measured the ET of a young splash pine stand (*Pinus elliottii*) with lysimeters for 2 years. Stannard (1993) evaluated three ET models in a sparsely vegetated, semi-arid rangeland. Federer et al. (1996) used specific coefficients to estimate the potential evapotranspiration (PET) in different forested areas. Farahani & Ahuja (1996) worked in partial canopy/residue-covered fields. Kolka & Wolf (1998) modified the Thornthwaite model in order to estimate the actual ET in 29 forested sites. Fisher et al. (2005) compared five models of PET in a mixed (dominant species *Pinus ponderosa*) coniferous forest. Ha et al. (2015) worked in semi-arid high-elevation disturbed ponderosa pine forests and compared ET between eddy covariance measurements and meteorological and remote sensing-based models.

To our knowledge, except for Gebhart et al. (2012) who studied some temperature-based and radiation-based methods in northern Greece, there have been no reports for comparative evaluation of the behavior of  $ET_0$  methods in forested areas of Greece. Other studies conducted in Greece used meteorological data from the Greek National Meteorological Service (Xystrakis & Matzarakis 2011; Samaras et al. 2014) exclusively for urban and agricultural areas. The meteorological stations that provided the data did not always follow the protocols imposed by the FAO (Alexandris et al. 2013). Moreover, there has not been such a study in the Mediterranean forests containing evergreen sclerophyllous broadleaved species.

For the above reasons, the main objective of this study was to test and evaluate the accuracy of different  $ET_0$  estimation equations, taking into account the data requirements for each model and making the assumptions that: (1) the  $ET_{P-M}$  model is the best estimator for the  $ET_0$  and (2) the environmental conditions of the site approximate the conditions for the application of the  $ET_{P-M}$  model. The 24 selected equations are very common, extensively used in other studies, and

represent the four main categories (mass-transfer, temperature-based, radiation-based, and combination).

The models tested in this work produce  $ET_0$  or PET estimates above ground covered with grass in a Mediterranean forest environment in Greece. The results will be useful to other researchers for incorporating them as input into hydrological, environmental, and soil models applied on similar Mediterranean vegetative and climatic conditions.

## MATERIALS AND METHODS

### Site description

The study was carried out in a small experimental forest watershed (1.23 km<sup>2</sup>) covered by evergreen sclerophyllous broadleaved vegetation (*maquis* vegetation) in Western Greece close to Varetada village (Figure 1). This is a multi-layer, dense coppice forest with canopy closure 1.2–1.3 (tree canopies overlap). The understorey is dominated by *Phillyrea latifolia*, *Arbutus unedo*, and *Erica arborea*. Sporadic stems of *Cercis siliquastrum* and *Erica verticilata* are also present. In the upper storey there is a number of *Quercus ilex* stems distributed almost uniformly (Baloutsos et al. 2009). The height of all species varies from 4 to 15 m. The terrain is hilly and the soil is a Haplic Luvisol one (FAO 1988) and its parent material is flysch.

The site receives a mean annual amount of precipitation of 1,174 mm in the form of rain which ranges from 696 to

2,230 mm, as calculated from climatic data of the period 1996–2012. The wettest months are October (161 mm) and November (130.8 mm) and the driest ones July (46 mm) and June (46.5 mm). The mean annual air temperature is 15.6 °C. The coldest month is January with a mean monthly value of 7.1 °C with August being the hottest (25.5 °C) one. The mean annual relative humidity is 67.5% with an average of 80% during December and 59% during June. The average annual wind speed is 1.9 m/s with a monthly average value of 2.4 m/s during July and 1.5 m/s during December.

The wider area is classified in the Csa climatic type according to the Köppen–Geiger updated world map (Kottek et al. 2006), which shows seasonal variability (warm temperate rainy climates with mild winters and very hot dry summers).

The particular site was selected for forest research purposes for two reasons: (1) the *maquis* vegetation is one of the most representative vegetations in Mediterranean forest ecosystems and (2) the watershed is a long way from urban and industrial areas, so it is not likely that it receives any kind of pollution. Additionally, the entire existing forest has not been managed by the local Forest Service for over 40 years; therefore, there has not been any land use change for the same period of time.

### Meteorological data (evaluation and processing)

The meteorological data were collected from an automatic meteorological station (latitude 38° 50' 35", longitude 21° 18' 25", elevation 332 m a.s.l.) installed in a natural forest opening inside the watershed. The main advantage




Figure 1 | The location of the forest meteorological station in Greece.

of the location and the ground vegetation of the meteorological station is that they meet the specifications imposed by the FAO to avoid (as much as possible) significant and systematic cumulative errors in determining  $ET_0$  (from 27% up to 47% during the warm season – [Alexandris \*et al.\* 2013](#)). Meteorological variables, such as air temperature ( $T_{aver}$ ,  $T_{max}$ ,  $T_{min}$ ), relative humidity ( $RH_{aver}$ ,  $RH_{max}$ ,  $RH_{min}$ ), solar radiation ( $R_s$ ), wind speed ( $u_2$ ), and precipitation (PR) were continuously recorded for a time period of 17 years (1996–2012). All sensors were set at 2 m above the ground level except for the rain gauge which was at a height of 1.3 m. The sampling period for all the monitored variables was set up to 15 min and the collected data were stored in a digital datalogger connected to the sensors. The data were periodically downloaded to be summarized and provide hourly, daily, monthly, and annual averages. These values constituted the input data used for the estimation of the daily values of  $ET_0$  in all of the equations.

For the present study, daily data covering the 87.5% of the total length of the 17-year period (5,433 days) were used. The existing gaps (12.5% or 777 daily values: 356 in winters, 114 in springs, 100 in summers, and 207 in autumns) were randomly distributed and were excluded from the statistical analysis, since any gap filling could possibly affect the reliability of the results. The gaps were due to the lack of measurements of some parameters that made the application of the  $ET_{P-M}$  equation prohibitory. In some cases, some of the methods included in this study estimated a negative  $ET_0$  daily value. These values were also excluded from the statistical analysis. A summary of the notations, definitions, and the units of the symbols used are shown in [Table 1](#).

### ET<sub>0</sub> estimation equations

The 24 different equations used in this study were categorized in the following groups: five mass-transfer (Albrecht,

**Table 1** | Notations, definitions, and units used in all equations for the estimation of the  $ET_0$

| Notation        | Definition   | Unit  |
|-----------------|--|---|
| ET              | Evapotranspiration   | mm/day  |
| $\Delta$        | Slope of vapor pressure curve                                  | kPa/°C  |
| $\lambda$       | Latent heat of vaporization                                    | MJ/kg   |
| $\rho$          | Water density  | =1.0 kg/l   |
| $\gamma$        | Psychrometric constant   | kPa/°C  |
| $e_s$           | Saturation vapor pressure                                      | kPa   |
| $e_a$           | Actual vapor pressure  | kPa   |
| $u_2$           | Wind speed at 2 m above ground surface                         | m/s   |
| $T_{aver}$      | Mean daily air temperature                                     | °C  |
| $R_n$           | Net solar radiation  | MJ/m <sup>2</sup> /day  |
| $C_1$ and $C_2$ | Functions of the attributes $R_s$ , $T_{aver}$ and $RH_{aver}$ | mm/day  |
| G               | Soil heat flux density   | =0 MJ/m <sup>2</sup> /day for daily computations ( <a href="#">ASCE-EWRI 2005</a> ) |
| $R_s$           | Incident shortwave solar radiation flux                        | MJ/m <sup>2</sup> /day  |
| $R_a$           | Extraterrestrial solar radiation                               | MJ/m <sup>2</sup> /day  |
| $T_{max}$       | Maximum daily air temperature                                  | °C  |
| $T_{min}$       | Minimum daily air temperature                                  | °C  |
| PR              | Precipitation  | mm  |
| N               | Maximum possible duration                                      | hrs   |
| $RH_{aver}$     | Mean daily relative humidity                                   | %   |
| $\varphi$       | Latitude   | Rad   |
| $\alpha$        | Albedo   | =0.23   |

Note: For details needed for the computation of the parameters which were not measured directly, refer to [Allen \*et al.\* \(1998\)](#).

Mahringer, Penman, Romanenko, and WMO), four combination (Copais, Solar Thermal Unit, Valiantzas (1) and (2)), ten radiation-based (Abtew, Caprio, De Bruin–Keijman, FAO24 Radiation, Jensen–Haise, Hansen, Makkink, McGuinness–Bordne, Priestley–Taylor and Turc), and five temperature-based (Hargreaves original, Hargreaves–Samani, two modified Hargreaves–Samani, and modified Thornthwaite) methods. The formulas of the equations are presented in Table 2 along with their references. The conversion of the units is in agreement with the units shown in Table 1.

### Statistical analysis

There are many widely used statistical indices and coefficients to evaluate the systematic quantification of the accuracy of compared models (Willmott 1982; Berengena & Gavilán 2005; Alexandris *et al.* 2008; Valiantzas 2013). A great number of them was selected in this study aiming to facilitate further comparison of the results with those of other studies.

The computational formulas for all indices and coefficients except of mean value (MV), standard deviation (SD), coefficient of determination ( $R^2$ ), and the coefficients of the linear trend line  $a$  (slope) and  $b$  (intercept) are the following:

Correlation

$$\text{Correl} = \frac{\sum_{i=1}^n (E_i - E_{\text{aver}})(O_i - O_{\text{aver}})}{\sqrt{\sum_{i=1}^n (E_i - E_{\text{aver}})^2 \sum_{i=1}^n (O_i - O_{\text{aver}})^2}}$$

$$\text{Mean bias error} = \text{MBE} = \frac{\sum_{i=1}^n (E_i - O_i)}{n}$$

$$\text{Mean absolute error} = \text{MAE} = \frac{\sum_{i=1}^n |E_i - O_i|}{n}$$

$$\text{Root mean square error} = \text{rMSE} = \sqrt{\frac{1}{n} \sum_{i=1}^n (E_i - O_i)^2}$$

$$\text{Average absolute error} = \text{AAE} = \sqrt{\frac{1}{n} \sum_{i=1}^n |E_i - O_i|}$$

$$\text{Relative mean square error} = \text{RMSE} = \frac{\sum_{i=1}^n ((E_i - O_i)/E_i)^2}{n}$$

$$\text{Relative mean absolute error} = \text{RMAE} = \frac{\sum_{i=1}^n |(E_i - O_i)/E_i|}{n}$$

Index of agreement =

$$d = 1 - \frac{\sum_{i=1}^n (E_i - O_i)^2}{\sum_{i=1}^n (|E_i - O_{\text{aver}}| + |O_i - O_{\text{aver}}|)^2}$$

$$\text{Weighted determination coefficient} = \text{wR}^2 = \frac{R^2}{a},$$

where  $a > 1$ ,  $\text{wR}^2 = R^2 \times a$ , where  $a < 1$

Variance of distribution of differences =

$$S_d^2 = \frac{\{\sum_{i=1}^n (E_i - O_i - \text{MBE})\}^2}{n - 1}$$

$$\text{Model efficiency} = \text{EF} = 1 - \frac{\sum_{i=1}^n (E_i - O_i)^2}{\sum_{i=1}^n (O_i - O_{\text{aver}})^2}$$

$$\text{Long-term average ratio} = \text{rt} = \frac{E_{\text{aver}}}{\text{Long term Mean of ET}_{\text{P-M}}}$$

where  $E_i$  and  $E_{\text{aver}}$  are the predicted daily and the average of the  $\text{ET}_0$  method values, respectively,  $O_i$  and  $O_{\text{aver}}$  are the calculated daily and the average of the  $\text{ET}_{\text{P-M}}$  values, respectively, and  $n$  is the total number of data.

Specifically, Correl,  $R^2$ ,  $a$  and  $b$  indices, of the least squared regression analysis, are commonly used correlation measures. The MV and SD indices provide a general view of the models' performance. For more efficient model assessment, Krause *et al.* (2005) suggest the use of the combined index  $\text{wR}^2$ . For the mean error evaluation, MBE,  $S_d^2$ , and rMSE indices were used (Fox 1981; Berengena & Gavilán 2005). For the absolute and/or relative errors' estimation, MAE, AAE, RMSE, and RMAE indices were also calculated, so as to facilitate the discussion in this work, since they are widely reported in the literature (Xystrakis & Matzarakis 2011; Gebhart *et al.* 2012; Kisi 2014; Samaras *et al.* 2014). The descriptive  $d$  index was used for the

**Table 2** | Formulas, symbols, and references of the ET<sub>0</sub> equations

| Method                           | Symbol                       | Equation   | References   |
|----------------------------------|------------------------------|--|--|
| <b>Benchmark equation</b>        |                              |  |  |
| 1                                | FAO56<br>Penman-<br>Monteith | $ET_{P-M} = \frac{0.408 \Delta(R_n - G) + \gamma \frac{900}{T_{aver} + 273} u_2 (e_s - e_a)}{\Delta + \gamma(1 + 0.34u_2)}$                    | Allen et al. (1998)  |
| <b>Mass-transfer equations</b>   |                              |  |  |
| 2                                | Albrecht                     | $ET_{ALB} = F(e_s - e_a)$ , where $F = 0.4$ if $u_2 \geq 1$ m/s and $F = 0.1005 + 0.297 u_2$ if $u_2 < 1$ m/s                                  | Albrecht (1950) and Friesland et al. (1998)                      |
| 3                                | Mahringer                    | $ET_{MAH} = 0.15072 \sqrt{3.6 u_2} (e_s - e_a)$  | Mahringer (1970) and Tabari et al. (2011)                        |
| 4                                | Penman                       | $ET_{PEN} = 0.35 \left( 1 + \frac{0.98}{100 u_2} \right) (e_s - e_a)$  | Penman (1948) and Tabari et al. (2011) <sup>a</sup>              |
| 5                                | Romanenko                    | $ET_{ROM} = 4.5 \left[ 1 + \left( \frac{T_{aver}}{25} \right) \right]^2 \left( 1 - \frac{e_a}{e_s} \right)$                                    | Oudin et al. (2005a, 2005b) and Xystrakis & Matzarakis (2011)    |
| 6                                | WMO                          | $ET_{WMO} = (0.1298 + 0.0934 u_2)(e_s - e_a)$  | WMO (1966) and Tabari et al. (2011) <sup>b</sup>                 |
| <b>Combinations equations</b>    |                              |  |  |
| 7                                | Copais                       | $ET_{COP} = m_1 + m_2 C_2 + m_3 C_1 + m_4 C_1 C_2$<br>where $m_1 = 0.057$ , $m_2 = 0.277$ , $m_3 = 0.643$ , $m_4 = 0.0124$                     | Alexandris et al. (2006, 2008) <sup>c,d</sup>                    |
| 8                                | Solar Thermal Unit           | $ET_{STU} = 6.1 \times 10^{-9} R_s (1.8 T_{aver} + 1)$   | Caprio (1974)  |
| 9                                | Valiantzas (1)               | $ET_{VA1} = 0.0393 R_s \sqrt{T_{aver} + 9.5} - 0.19 R_s^{0.6} \varphi^{0.15} + 0.078 (T_{aver} + 20) \left( 1 - \frac{RH_{aver}}{100} \right)$ | Valiantzas (2013)  |
| 10                               | Valiantzas (2)               | $ET_{VA2} = 0.0393 R_s \sqrt{T_{aver} + 9.5} - 0.19 R_s^{0.6} \varphi^{0.15} + 0.0061 (T_{aver} + 20) (1.12 T_{aver} - T_{min} - 2)^{0.7}$     | Valiantzas (2013)  |
| <b>Radiation-based equations</b> |                              |  |  |
| 11                               | Abtew                        | $ET_{ABT} = \frac{1}{56} \frac{R_s T_{max}}{\lambda}$  | Abtew (1996) and Samaras et al. (2014)                           |
| 12                               | Caprio                       | $ET_{CAP} = 1.65 \frac{\Delta}{\Delta + \gamma} \left( \frac{R_n - G}{\lambda} \right)$  | Caprio (1974) and Samaras et al. (2014)                          |
| 13                               | De Bruin-Keijman             | $ET_{DBK} = \left( \frac{\Delta}{0.85\Delta + 0.63\gamma} \right) \left( \frac{R_n - G}{\lambda} \right)$                                      | DeBruin & Keijman (1979) and Rosenberry et al. (2007)            |
| 14                               | FAO24 Radiation              | $ET_{F24} = b \left( \frac{\Delta}{\Delta + \gamma} R_s \right) - 0.3$   | Doorenbos & Pruitt (1977) and Frevert et al. (1983) <sup>e</sup> |

(continued)

Table 2 | continued

| Method                            | Symbol     | Equation  | References   |
|-----------------------------------|------------|---|--|
| 15 Hansen                         | $ET_{HAN}$ | $= 0.7 \left( \frac{\Delta}{\Delta + \gamma} \right) \left( \frac{R_s}{\lambda} \right)$  | Hansen (1984) and Xu & Singh (2002)                        |
| 16 Jensen-Haise                   | $ET_{J-H}$ | $= \left( \frac{R_s}{\lambda} \right) (0.025 T_{aver} + 0.08)$  | Rosenberg et al. (1983) and Xystrakis & Matzarakis (2011)  |
| 17 Makkink                        | $ET_{MAK}$ | $= 0.61 \left( \frac{\Delta}{\Delta + \gamma} \right) \left( \frac{R_s}{\lambda} \right) - 0.12$  | Rosenberry et al. (2004) and Alexandris et al. (2008)      |
| 18 McGuinness-Bordne              | $ET_{MGB}$ | $= \left( \frac{R_a}{\lambda \rho} \right) \left( \frac{T_{aver} + 5}{68} \right)$  | McGuinness & Bordne (1972) and Oudin et al. (2005a, 2005b) |
| 19 Priestley-Taylor               | $ET_{P-T}$ | $= a \left( \frac{\Delta}{\Delta + \gamma} \right) \left( \frac{R_n}{\lambda} \right)$  | Priestley & Taylor (1972) <sup>f</sup>                     |
| 20 Turc                           | $ET_{TUR}$ | $= 0.013 \left( \frac{T_{aver}}{T_{aver} + 15} \right) (R_s + 50)$ , for $RH_{aver} > 50\%$<br>$= 0.013 \left( \frac{T_{aver}}{T_{aver} + 15} \right) (R_s + 50) \left( 1 + \frac{50 - RH_{aver}}{70} \right)$ , for $RH_{aver} < 50\%$ | Turc (1961) and Lu et al. (2005)                           |
| 21 Hargreaves (original)          | $ET_{HAR}$ | $= 0.0135 R_s (T_{aver} + 17.8)$  | Hargreaves (1975)  |
| 22 Hargreaves-Samani              | $ET_{H-S}$ | $= 0.0023 \sqrt{(T_{max} - T_{min})} (T_{aver} + 17.8) 0.408 R_a$   | Hargreaves & Samani (1985)                                 |
| 23 Hargreaves-Samani (modified 1) | $ET_{HS1}$ | $= 0.0030 (T_{max} - T_{min})^{0.4} (T_{aver} + 20) 0.408 R_a$  | Droogers & Allen (2002)                                    |
| 24 Hargreaves-Samani (modified 2) | $ET_{HS2}$ | $= 0.0025 \sqrt{(T_{max} - T_{min})} (T_{aver} + 16.8) 0.408 R_a$   | Droogers & Allen (2002)                                    |
| 25 Thornthwaite (modified)        | $ET_{THO}$ | $= 0.533 \frac{N}{12} \left( \frac{10 T_{aver}}{WI} \right)^A$ , where $WI = 33.617$ and $A = 1.033$  | Siegert & Schrodter (1975)                                 |

<sup>a</sup> $e_s$  and  $e_a$  are in mmHg and  $u_2$  is in miles/day.

<sup>b</sup> $e_s$  and  $e_a$  are in hPa.

<sup>c</sup>In the original paper (Alexandris et al. 2006) the coefficient  $m_2$  is 0.227 due to a misprint and should be replaced with the correct value 0.277 (Alexandris et al. 2008).

<sup>d</sup> $C_1 = 0.6416 - 0.00784 RH_{aver} + 0.372 R_s - 0.00264 RH_{aver}$ ;  $C_2 = -0.0033 + 0.00812 T_{aver} + 0.101 R_s + 0.00584 R_s T_{aver}$ .

<sup>e</sup> $b = 1.066 - 0.13 \times 10^{-2} (RH_{aver}) + 0.045 (u_2) - 0.20 \times 10^{-3} (RH_{aver} \times u_2) - 0.135 \times 10^{-4} (RH_{aver})^2 - 0.11 \times 10^{-2} (u_2)^2$ .

<sup>f</sup> $a = 1.26 =$  Priestley-Taylor's empirically constant, dimensionless.

cross-comparison between the models, expressing the degree to which a model's predictions are error free (Willmott 1982). Finally, the EF index specifies the relationship between calculated and predicted mean deviations (Greenwood et al. 1985), while  $rt$  returns a long-term value of the predicted against the calculated MVs.

## RESULTS AND DISCUSSION

On an annual basis, the average observed  $ET_0$  rate calculated from the  $ET_{P-M}$  for the period 1996–2012, was found to be 1,190 mm. This value is a little different from the mean annual precipitation (1,174 mm) and specifically

indicates that the water requirements of a reference crop are totally sufficed. Additionally, the mean seasonal values of  $ET_0$  showed seasonal variation, as expected for the Mediterranean climate (Csa), varying from 99 mm (winter) to 558 mm (summer) and moderate values during the transitional periods of spring and autumn (307 mm and 226 mm, respectively).

From the analysis of the annual values, the best approaches seem to give the  $ET_{HS2}$ ,  $ET_{COP}$ , and  $ET_{STU}$  methods, in which the percentages of  $ET_0$  average annual values diverge by +0.47%, -0.69%, and -2.27%, respectively. In contrast, the largest annual deviations appeared in the  $ET_{ROM}$  (+60.85%),  $ET_{MAK}$  (-26.47%), and  $ET_{PEN}$  (-26.23%) methods. These results were in line with Federer *et al.* (1996), who reported that different methods gave widely differing estimates of annual  $ET_0$  at particular locations which sometimes were up to several hundreds of millimeters.

From the analysis of 5,433 daily values with 18 different statistical parameters and indices, the 24 tested methods were evaluated comparatively. Their performance against the  $ET_{P-M}$  method is shown in Figure 2 and Table 3 and the obtained results are the following.

### Mass-transfer equations

Among the five examined methods, the  $ET_{PEN}$  showed average daily  $ET_0$  equal to 2.44 mm, a value significantly lower (-27.8%,  $rt = 0.722$ ) compared to the corresponding  $ET_{P-M}$ . Simultaneously, it displayed the smallest correlation coefficient ( $R^2 = 0.826$ ). The approaches of the average daily  $ET_0$  of the  $ET_{ALB}$ ,  $ET_{WMO}$ , and  $ET_{MAH}$  were very similar, with deviations of +9.1% ( $rt = 1.091$ ), -9.9% ( $rt = 0.901$ ), and +13.0% ( $rt = 1.131$ ), respectively. However, their  $R^2$ s are not considered satisfactory (0.830, 0.855, and 0.860 respectively). Finally, the  $ET_{ROM}$  method displayed the worst statistics for almost all of the evaluation indices.

It is worth noting that this category of methods resulted in the smallest EF index ranging from -0.437 ( $ET_{ROM}$ ) to 0.759 ( $ET_{WMO}$ ). Overall, the statistical indices of the mass-transfer equations were not satisfactory (Table 3). Hence, these methods cannot be recommended for use without calibration, in models that need the input of  $ET_0$  daily values. Similar results were reported by Valipour (2014b, 2015c), who examined  $R^2$  and MBE and suggested new calibrated

mass-transfer equations for the provinces of Iran which relatively improved the performance of the original models.

In our study (forest environment with warm humid Mediterranean climate), except for  $ET_{PEN}$ , the high summer daily values (>5 mm) are overestimated with the mass-transfer equations. Although Tabari *et al.* (2011) reported that the majority of the mass-transfer empirical equations they tested, had also the worst performances but at the same time underestimated  $ET_{P-M}$  in humid environments. This could be attributed to the fact that VPD ( $e_s - e_a$ ) presents significant variations among locations, as stated by Irmak *et al.* (2006), who performed a sensitivity analysis of the Penman-Monteith method for several regions with different climate types, in the USA.

### Combination equations

Combination methods, in general, showed much better statistical indices compared to the methods of all other categories (Table 3). The four methods examined here presented sufficiently strong correlations ( $R^2$  is ranging from 0.937 in  $ET_{COP}$  to 0.958 in  $ET_{VA1}$ ), compared to  $ET_{P-M}$ . However, the  $rt$  was significantly lower in  $ET_{COP}$  deviating only by -0.6% ( $rt = 0.994$ ), compared to the other three methods, while  $ET_{VA1}$  presented the greatest divergence (+11.3%,  $rt = 1.113$ ). Impressive was the finding that despite the relatively small  $R^2$  of  $ET_{COP}$ , most of the other statistical indices outweighed the respective indices which derived from the analysis of all 24 tested methods (Table 3). Similar results for the  $ET_{COP}$ ,  $ET_{VA1}$ , and  $ET_{VA2}$  equations were presented by Kisi (2014) in a Mediterranean environment in Turkey. The findings of Valipour (2015b) in Iran for the Valiantzas' equations are also in line with the results presented here.


The  $ET_{STU}$  method also gave satisfactory results ( $rt = 0.978$ ,  $R^2 = 0.955$ ) despite the small requirements in input data. These methods are strongly recommended for use in models which need the input of  $ET_0$  daily values because they have a very satisfactory EF index ranging from 0.894 ( $ET_{STU}$ ) to 0.934 ( $ET_{COP}$ ).

### Radiation-based equations


In this category ten methods were tested. The radiation-based equations overall performed better than the mass-transfer




Mass transfer equations



Combination equations



Radiation-based equations



Temperature-based equations




Figure 2 | R<sup>2</sup> coefficient, slope, and intercept of the daily values of ET<sub>0</sub> (in mm) estimated by various methods (x axis) vs. the standardized ET<sub>p-m</sub> (y axis). The continuous line in all figures indicates the trendline of the scatter and the dashed line indicates the 1:1 slope trend. The point shapes for each category differ.

**Table 3** | Statistical analysis of the 24 methods versus the FAO56 PM model for estimating daily  $ET_0$  during the study period (1996–2012)

| A/A | Categories                | Symbol            | N     | Mean (mm)           | SD (mm) | rt                  | Correl              | MBE (mm)             | MAE (mm)            | rMSE (mm)           | rMAE (mm)           | RMSE (mm)           | RMAE (mm)           | d                   | wR <sup>2</sup>     | S <sub>d</sub> <sup>2</sup> (mm) | EF                  |
|-----|---------------------------|-------------------|-------|---------------------|---------|---------------------|---------------------|----------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|---------------------|----------------------------------|---------------------|
| 1   |                           | ET <sub>P-M</sub> | 5,433 | 3.380               | 2.196   |                     |                     |                      |                     |                     |                     |                     |                     |                     |                     |                                  |                     |
| 2   | Mass-transfer methods     | ET <sub>ALB</sub> | 5,414 | 3.687               | 2.900   | 1.091               | 0.911               | 0.379                | 0.901               | 1.348               | 0.949               | 1.381               | 0.863               | 0.928               | 0.567               | 1.673                            | 0.625               |
| 3   |                           | ET <sub>MAH</sub> | 5,414 | 3.821               | 3.265   | 1.131               | 0.927               | 0.523                | 0.984               | 1.593               | 0.992               | 1.066               | 1.078               | 0.914               | 0.531               | 2.265                            | 0.475               |
| 4   |                           | ET <sub>PEN</sub> | 5,414 | 2.440               | 1.895   | 0.722               | 0.909               | -0.897               | 0.990               | 1.283               | 0.995               | 0.757               | 0.737               | 0.905               | 0.654               | 0.843                            | 0.660               |
| 5   |                           | ET <sub>ROM</sub> | 5,414 | 5.392               | 3.611   | 1.595               | 0.919               | 2.046                | 2.099               | 2.642               | 1.449               | 0.293               | 0.604               | 0.811               | 0.478               | 3.117                            | -0.437              |
| 6   |                           | ET <sub>WMO</sub> | 5,414 | 3.046               | 2.635   | 0.901               | 0.925               | -0.270               | 0.773               | 1.080               | 0.879               | 2.026               | 1.278               | 0.949               | 0.652               | 1.094                            | 0.759               |
| 7   | Combination methods       | ET <sub>COP</sub> | 5,428 | <b><u>3.360</u></b> | 2.241   | <b><u>0.994</u></b> | 0.968               | <b><u>-0.022</u></b> | <b><u>0.442</u></b> | <b><u>0.564</u></b> | <b><u>0.665</u></b> | 0.438               | 0.325               | <b><u>0.984</u></b> | <b><u>0.921</u></b> | 0.318                            | <b><u>0.934</u></b> |
| 8   |                           | ET <sub>STU</sub> | 5,426 | 3.306               | 2.681   | 0.978               | 0.977               | -0.079               | 0.569               | 0.715               | 0.755               | 1.940               | 0.704               | 0.978               | 0.764               | 0.513                            | 0.894               |
| 9   |                           | ET <sub>VA1</sub> | 5,425 | 3.763               | 2.423   | 1.113               | <b><u>0.978</u></b> | 0.375                | 0.523               | 0.649               | 0.723               | 1.214               | 0.294               | 0.980               | 0.849               | 0.281                            | 0.913               |
| 10  |                           | ET <sub>VA2</sub> | 5,367 | 3.051               | 2.023   | 0.903               | 0.972               | -0.355               | 0.468               | 0.635               | 0.684               | 2.024               | 0.364               | 0.977               | 0.847               | 0.278                            | 0.917               |
| 11  | Radiation-based methods   | ET <sub>ABT</sub> | 5,430 | 2.783               | 2.186   | 0.824               | 0.976               | -0.600               | 0.623               | 0.766               | 0.789               | 1.857               | 0.827               | 0.970               | <b><u>0.926</u></b> | <b><u>0.227</u></b>              | 0.878               |
| 12  |                           | ET <sub>CAP</sub> | 5,433 | 3.772               | 2.579   | 1.116               | 0.953               | 0.391                | 0.697               | 0.913               | 0.835               | 0.183               | 0.259               | 0.963               | 0.737               | 0.681                            | 0.827               |
| 13  |                           | ET <sub>DBK</sub> | 5,433 | 2.930               | 1.959   | 0.867               | 0.949               | -0.452               | 0.597               | 0.836               | 0.773               | 0.337               | 0.318               | 0.959               | 0.762               | 0.495                            | 0.855               |
| 14  |                           | ET <sub>F24</sub> | 5,260 | 3.763               | 2.668   | 1.114               | 0.976               | 0.296                | 0.622               | 0.783               | 0.788               | 2.676               | 0.975               | 0.974               | 0.753               | 0.526                            | 0.877               |
| 15  |                           | ET <sub>HAN</sub> | 5,433 | 2.974               | 1.884   | 0.880               | 0.961               | -0.408               | 0.556               | 0.765               | 0.746               | 0.714               | 0.339               | 0.965               | 0.762               | 0.418                            | 0.879               |
| 16  |                           | ET <sub>J-H</sub> | 5,431 | 3.488               | 2.719   | 1.032               | <b><u>0.978</u></b> | 0.105                | 0.575               | 0.740               | 0.758               | 1.697               | 0.475               | 0.977               | 0.755               | 0.537                            | 0.886               |
| 17  |                           | ET <sub>MAK</sub> | 5,399 | 2.487               | 1.635   | 0.736               | 0.901               | -0.912               | 0.926               | 1.192               | 0.962               | 2.603               | 0.731               | 0.907               | 0.662               | 0.590                            | 0.707               |
| 18  |                           | ET <sub>MGB</sub> | 5,432 | 3.876               | 2.361   | 1.147               | 0.930               | 0.618                | 0.815               | 1.063               | 0.903               | <b><u>0.133</u></b> | 0.264               | 0.945               | 0.863               | 0.749                            | 0.766               |
| 19  |                           | ET <sub>P-T</sub> | 5,433 | 2.881               | 1.969   | 0.852               | 0.953               | -0.501               | 0.608               | 0.843               | 0.780               | 0.407               | 0.345               | 0.959               | 0.775               | 0.460                            | 0.853               |
| 20  |                           | ET <sub>TUR</sub> | 5,419 | 2.970               | 1.956   | 0.879               | 0.974               | -0.414               | 0.513               | 0.674               | 0.716               | 0.316               | 0.397               | 0.974               | 0.819               | 0.283                            | 0.906               |
| 21  | Temperature-based methods | ET <sub>HAR</sub> | 5,433 | 3.139               | 2.116   | 0.929               | 0.972               | -0.244               | <b><u>0.427</u></b> | <b><u>0.572</u></b> | <b><u>0.653</u></b> | 0.780               | 0.321               | <b><u>0.982</u></b> | 0.884               | <b><u>0.268</u></b>              | <b><u>0.932</u></b> |
| 22  |                           | ET <sub>H-S</sub> | 5,433 | 3.123               | 1.910   | 0.924               | 0.931               | -0.167               | 0.603               | 0.832               | 0.777               | 0.171               | 0.248               | 0.958               | 0.701               | 0.664                            | 0.856               |
| 23  |                           | ET <sub>HS1</sub> | 5,433 | <b><u>3.400</u></b> | 1.969   | <b><u>1.006</u></b> | 0.932               | 0.116                | 0.615               | 0.808               | 0.784               | <b><u>0.111</u></b> | <b><u>0.231</u></b> | 0.961               | 0.726               | 0.640                            | 0.865               |
| 24  |                           | ET <sub>HS2</sub> | 5,433 | 3.300               | 2.034   | 0.976               | 0.931               | <b><u>0.016</u></b>  | 0.590               | 0.802               | 0.768               | 0.154               | <b><u>0.237</u></b> | 0.963               | 0.748               | 0.643                            | 0.867               |
| 25  |                           | ET <sub>THO</sub> | 5,423 | 2.777               | 1.608   | 0.822               | 0.915               | -0.532               | 0.895               | 1.111               | 0.946               | 0.942               | 0.593               | 0.913               | 0.558               | 0.951                            | 0.744               |

The two best fitted methods for each index or coefficient are in bold and underlined.

**N**: Sample days; **Mean**: Time series average; **SD**: Standard deviation; **rt**: long-term average ratio; **Correl**: Pearson's correlation; **MBE**: Mean bias error; **MAE**: Mean absolute error; **rMSE**: Root mean square error; **rMAE**: Root mean absolute error; **RMSE**: Relative mean square error; **RMAE**: Relative mean absolute error; **d**: Index of agreement; **wR<sup>2</sup>**: Weighted determination; **S<sub>d</sub><sup>2</sup>**: Coefficient variance of distribution of differences; **EF**: Model efficiency.

equations, since a more important role is expected for  $R_s$  when estimating  $ET_0$  in humid climates (Irmak et al. 2006) and in forest environments (Gebhart et al. 2012).  $ET_{J-H}$  and  $ET_{ABT}$  ( $R^2 = 0.956$  and  $0.953$ , respectively) methods presented satisfactory correlations. Additionally, the  $ET_{J-H}$  method showed the lowest deviation from the daily MV (+3.2%,  $rt = 1.032$ ), while  $ET_{ABT}$  had the best indices,  $wR^2$  and  $S_d^2$ , over all of the 24 tested methods. These findings for  $ET_{J-H}$  are in contrast to Tabari et al. (2011), who worked in humid environments.

$ET_{TUR}$  can be considered as the best performing equation in this category, in terms of its EF (0.906), rMAE (0.716), RMSE (0.674), and MAE (0.513). The best performance of the equation was also found by Lu et al. (2005) in forest watersheds with warm and humid climates in the southeastern USA, and by Trajkovic & Kolakovic (2009) who recommended  $ET_{TUR}$  for use under humid conditions, and by Gebhart et al. (2012) who suggested the use of  $ET_{TUR}$  for the southern regions of Central Macedonia in Greece.

In contrast, the  $ET_{MGB}$  and  $ET_{DBK}$  methods gave the worst correlations among all the radiation-based equations ( $R^2 = 0.865$  and  $R^2 = 0.901$ , respectively). Under humid conditions, similar results were reported for  $ET_{MGB}$  by Tabari et al. (2011). The largest deviations from the daily MV were displayed by  $ET_{MAK}$  (-26.4%,  $rt = 0.736$ ) and  $ET_{ABT}$  (-17.6%,  $rt = 0.824$ ). The EF index ranged from 0.707 ( $ET_{MAK}$ ) to 0.886 ( $ET_{J-H}$ ). The poor performance of  $ET_{MAK}$  was also reported by Lu et al. (2005).

From the above, it can be concluded that some methods ( $ET_{TUR}$ ,  $ET_{J-H}$ , and  $ET_{ABT}$ ) of this category can be satisfactorily accepted for use in models needing the input of  $ET_0$  daily values.

### Temperature-based equations

In this category five methods were tested. The best correlation was exhibited by  $ET_{HAR}$  with  $R^2 = 0.945$  and the worst by  $ET_{THO}$  ( $R^2 = 0.837$ ). Concerning the deviation from the daily MV, the  $ET_{HS1}$  gave the best of all 24 tested methods (along with  $ET_{COP}$ ), with  $rt = 1.006$ . These findings were similar to those of Valipour & Eslamian (2014) and Valipour (2015a) who gave specific ranges of the meteorological parameters used in 11 temperature-based equations for Iran's provinces and found a better performance of  $ET_{H-S}$ ,  $ET_{HS1}$ , and  $ET_{HS2}$  against  $ET_{THO}$ . The results in our study

indicated a slight underestimation in  $ET_0$  daily values for all of the tested temperature-based equations (except of  $ET_{THO}$  which underestimated by 17.8%). Valipour (2015a) also found a slight underestimation in  $ET_0$  daily values with  $ET_{H-S}$  and  $ET_{HS1}$  equations, while the  $ET_{HS2}$  equation showed overestimation but not a significant one.

Contrary to these results, Trajkovic & Kolakovic (2009) and Tabari et al. (2011) found overestimation when using  $ET_{HS1}$ ,  $ET_{HS2}$ , and  $ET_{H-S}$  equations, on a monthly time-step analysis and under humid conditions. They also found very poor performance of the  $ET_{THO}$  equation. Additionally, Lu et al. (2005) suggested careful calibration and verification when applying the  $ET_{THO}$  equation.

The  $ET_{HAR}$  method, despite the small data requirements, exhibited in general impressive statistical indices (e.g., best MAE, rMSE, rMAE,  $d$ ,  $S_d^2$ , and EF values) compared to all tested methods of all categories (Table 3). Gebhart et al. (2012) also proposed the  $ET_{HAR}$  equation as a good alternative for  $ET_0$  estimations in northern regions of Central Macedonia in Greece. From the above mentioned, it can be concluded that this category of equations seems to have similar performance (with the exception of  $ET_{THO}$ ) and they are recommended for use in forest environments.

## CONCLUSIONS

This study attempted to investigate and evaluate the best-fit methods for the estimation of daily  $ET_0$  in a humid Mediterranean evergreen broadleaved forest environment. Twenty-four different equations classified in four categories were tested and seventeen different statistical indices were used for the evaluation.

At the category level, the combination equations seem to have the best performance followed by temperature-based and radiation-based methods. The mass-transfer methods have the worst coefficients and overestimate  $ET_0$ , especially for the high summer daily values (>5 mm).

At the method level, the most accurate and consistent estimates of  $ET_0$  derived from  $ET_{COP}$  and  $ET_{HAR}$ , followed by  $ET_{VA1}$  and  $ET_{VA2}$ . The methods  $ET_{TUR}$ ,  $ET_{HS2}$ ,  $ET_{STU}$ , and  $ET_{J-H}$  are also proposed for use because their  $ET_0$  estimations compared quite well with those of the  $ET_{P-M}$  method. The latter ones can also be used for the annual estimation of the

ET<sub>0</sub>. Concerning the rest of the tested methods, it is suggested that calibration should be made for local conditions, mainly at a seasonal time-step to obtain more reliable daily estimates. Especially, calibration is necessary for all the mass-transfer equations and the ET<sub>MAK</sub> and ET<sub>THO</sub> methods. There is an ongoing work by the authors, in which new coefficients will be proposed for some of these methods. The results of this study will be useful to a multidisciplinary community working on similar climates. More specifically, the best performing equations in Figure 2 could be tested further in order to optimize the ET<sub>0</sub> obtaining values of the empirical models in the Mediterranean. Further research is needed in order to evaluate the performance of the proposed modified equations in other areas with different climates. Finally, evaluation is needed for the performance of the models on a different time scale (monthly and seasonal).

## ACKNOWLEDGEMENTS

The authors would like to express their sincere thanks to the Greek 'Ministry of Agriculture and Rural Development' and the European Union that funded various research projects from which the data used in this work were derived. The contribution of the anonymous reviewers is also greatly acknowledged.

## REFERENCES

- Abtew, W. 1996 *Evapotranspiration measurements and modeling for three wetland systems in South Florida*. *Journal of American Water Resources Association* **32** (3), 465–473.
- Albrecht, F. 1950 *Die Methoden zur Bestimmung der Verdunstung der natürlichen Erdoberfläche (The methods for the determination of the natural surface evaporation)*. *Archiv für Meteorologie, Geophysik und Bioklimatologie, Serie B* **2** (1), 1–38.
- Alexandris, S., Kerkides, P. & Liakatas, A. 2006 *Daily reference evapotranspiration estimates by the Copais approach*. *Agricultural Water Management* **82** (3), 371–386.
- Alexandris, S., Stricevic, R. & Petkovic, S. 2008 *Comparative analysis of reference evapotranspiration from the surface of rain-fed grass in central Serbia, calculated by six empirical methods against the Penman-Monteith formula*. *European Water* **21/22**, 17–28.
- Alexandris, S., Proutsos, N., Caravitis, Ch., Tsiros, I. & Stamatakos, D. 2013 *Reasons for non-rational estimates of reference evapotranspiration in Greece*. In: *Proceedings of the 8th National Agricultural Conference*, 25–26 September, Volos, Greece, pp. 110–116 (in Greek).
- Allen, R., Pereira, L., Raes, D. & Smith, M. 1998 *Crop Evapotranspiration. Guidelines for Computing Crop Water Requirements*. FAO Irrigation and Drainage Paper No. 56, FAO, Rome, p. 300.
- ASCE-EWRI 2005 *The ASCE Standardized Reference Evapotranspiration Equation. Technical Committee Report to the Environmental and Water Resources Institute of the American Society of Civil Engineers from the Task Committee on Standardization of Reference Evapotranspiration*. ASCE-EWRI, Reston, VA, p. 173.
- Baloutsos, G., Bourletsikas, A. & Baltas, E. 2009 *Development of a simplified model for the estimation of hydrological components in areas of maquis vegetation in Greece*. *WSEAS Transactions on Environment and Development* **5** (3), 310–320.
- Berengena, J. & Gavilán, P. 2005 *Reference ET estimation in a highly advective semi-arid environment*. *Journal of Irrigation and Drainage Engineering* **131** (2), 147–163.
- Bogawski, P. & Bednorz, E. 2014 *Comparison and validation of selected evapotranspiration models for conditions in Poland (Central Europe)*. *Water Resources Management* **28** (14), 5021–5038.
- Caprio, J. 1974 *The solar thermal unit concept in problems related to plant development and potential evapotranspiration*. *Phenology and Seasonality Modeling* **8**, 353–364.
- Currie, D. J. 1991 *Energy and large-scale patterns of animal and plant species richness*. *The American Naturalist* **137** (1), 27–49.
- DeBruin, H. A. R. & Keijman, J. Q. 1979 *The Priestley-Taylor evaporation model applied to a large, shallow lake in the Netherlands*. *Journal of Applied Meteorology* **18**, 898–903.
- Doorenbos, J. & Pruitt, W. O. 1977 *Guidelines for Predicting Crop Water Requirements*. FAO Irrigation and Drainage Paper No. 24, FAO, Rome, pp. 144.
- Droogers, P. & Allen, R. G. 2002 *Estimating reference evapotranspiration under inaccurate data conditions*. *Irrigation and Drainage Systems* **16** (1), 33–45.
- FAO 1988 *FAO/Unesco Soil Map of the World, Legend, World Soil Resources Report 60*. FAO, Rome.
- Farahani, H. J. & Ahuja, L. R. 1996 *Evapotranspiration modeling of partial canopy/residue-covered fields*. *Transactions of the ASAE* **39** (6), 2051–2064.
- Federer, C. A., Vörösmarty, C. & Fekete, B. 1996 *Intercomparison of methods for calculating potential evaporation in regional and global water balance models*. *Water Resources Research* **32**, 2315–2321.
- Fisher, J. B., DeBiase, T. A., Ye, O., Ming, X. & Goldstein, A. H. 2005 *Evapotranspiration models compared on a Sierra Nevada forest ecosystem*. *Environmental Modelling & Software* **20**, 783–796.
- Fox, D. G. 1981 *Judging air quality model performance: a summary of the AMS workshop on dispersion model performance*. *Bulletin of American Meteorological Society* **62**, 599–609.

- Frevert, D. K., Hill, B. W. & Braaten, B. C. 1983 Estimation of FAO evapotranspiration coefficients. *Journal of Irrigation and Drainage Engineering ASCE* **109**, 265–270.
- Friesland, H., Kersebaum, K.-C. & Löpmeier, F.-J. 1998 Operational use of Irrigation Models Using Medium Range Weather Forecast. Report of COST 711: Operational Applications of Meteorology to Agriculture, Including Horticulture, p 79.
- Gavilán, P., Lorite, I. J., Tornero, S. & Berengena, J. 2006 Regional calibration of Hargreaves equation for estimating reference ET in a semiarid environment. *Agricultural Water Management* **81** (3), 257–281.
- Gebhart, S., Radoglou, K., Chalivopoulos, G. & Matzarakis, A. 2012 Evaluation of potential evapotranspiration in Central Macedonia by EmPEst. In: *Advances in Meteorology, Climatology and Atmospheric Physics* (C. G. Helmis & P. T. Nastos, eds). Springer Atmospheric Sciences, Springer-Verlag, Berlin Heidelberg, pp. 451–456.
- Greenwood, D. J., Neteson, J. J. & Draycott, A. 1985 Response of potatoes N fertilizer: dynamic model. *Plant and Soil* **85**, 185–205.
- Grismer, M. E., Orang, M., Snyder, R. & Matyac, R. 2002 Pan evaporation to reference evapotranspiration conversion methods. *Journal of Irrigation and Drainage Engineering* **128** (3), 180–184.
- Ha, W., Kolb, T. E., Springer, A. E., Dore, S., O'Donnell, F. C., Morales, R. M., Masek Lopez, S. & Koch, G. W. 2015 Evapotranspiration comparisons between eddy covariance measurements and meteorological and remote-sensing-based models in disturbed ponderosa pine forests. *Ecohydrology* **8**, 1335–1350.
- Hansen, S. 1984 Estimation of potential and actual evapotranspiration. *Nordic Hydrology* **15**, 205–212.
- Hargreaves, G. H. 1975 Moisture availability and crop production. *Transactions of the ASAE* **18** (5), 980–984.
- Hargreaves, G. L. & Samani, Z. A. 1985 Reference crop evapotranspiration from temperature. *Applied Engineering in Agriculture* **1** (2), 96–99.
- Hay, L. E. & McCabe, G. J. 2002 Spatial variability in water balance performance in the conterminous United States. *Journal of the American Water Resources Association* **38**, 847–860.
- Houghton, R. A. 1991 Tropical deforestation and atmospheric carbon dioxide. *Climatic Change* **19**, 99–118.
- Irmak, S., Payero, J. O., Martin, D. L., Irmak, A. & Howell, T. A. 2006 Sensitivity analyses and sensitivity coefficients of standardized daily ASCE Penman-Monteith equation. *Journal of Irrigation and Drainage Engineering* **132** (6), 564–578.
- Jensen, M. E., Burman, R. D. & Allen, R. G. 1990 *Evapotranspiration and Water Requirements, ASCE Manual 70*, American Society of Civil Engineers, New York, p. 332.
- Khoshravesh, M., Sefidkouhi, M. A. G. & Valipour, M. 2015 Estimation of reference evapotranspiration using multivariate fractional polynomial, Bayesian regression, and robust regression models in three arid environments. *Applied Water Science*, doi:10.1007/s13201-015-0368-x.
- Kisi, O. 2014 Comparison of different empirical methods for estimating daily reference evapotranspiration in Mediterranean climate. *Journal of Irrigation and Drainage Engineering* **40** (1). 10.1061/(ASCE)IR.1943-4774.0000664, 04013002.
- Kolka, R. K. & Wolf, A. T. 1998 *Estimating Actual Evapotranspiration for Forested Sites: Modifications to the Thornthwaite Model*. Res. Note SRS-6, USDA Forest Service, Southern Research Station, p. 7.
- Kottek, M., Grieser, J., Beck, C., Rudolf, B. & Rubel, F. 2006 World map of the Köppen-Geiger climate classification updated. *Meteorologische Zeitschrift* **15** (3), 259–263.
- Krause, P., Boyle, D. P. & Base, F. 2005 Comparison of different efficiency criteria for hydrological model assessment. *Advances in Geosciences* **5**, 89–97.
- Lu, J. B., Sun, G., McNulty, S. G. & Amatya, D. M. 2005 A comparison of six potential evapotranspiration methods for regional use in the southeastern United States. *Journal of American Water Resources Association* **41** (3), 621–633.
- Mahringer, W. 1970 Verdunstungsstudien am Neusiedler See. *Archiv für Meteorologie, Geophysik und Bioklimatologie, Serie B* **18**, 1–20.
- McGuinness, J. L. & Bordne, E. F. 1972 *A Comparison of Lysimeter-Derived Potential Evapotranspiration with Computed Values*. Technical Bulletin 1452, Agricultural Research Service, USDA, Washington, DC, p. 71.
- McNaughton, K. G. & Black, T. A. 1975 A study of evapotranspiration from a Douglas fir forest using the energy balance approach. *Water Resources Research* **9** (6), 1579–1590.
- Musselman, R. C. & Fox, D. G. 1991 A review of the role of temperate forests in the global CO<sub>2</sub> balance. *Journal of the Air & Waste Management Association* **41** (6), 798–807.
- Nepstad, D. C., de Carvalho, C. R., Davidson, E. A., Jipp, P. H., Lefebvre, P. A., Negreiros, G. H., da Silva, E. D., Stone, T. A., Trumbore, S. E. & Vieira, S. 1994 The role of deep roots in the hydrological and carbon cycles of Amazonian forests and pastures. *Nature* **372**, 666–669.
- Oudin, L., Hervieu, F., Michel, C., Perrin, C., Andreassian, V., Anctil, F. & Loumagne, C. 2005a Which potential evapotranspiration input for a lumped rainfall-runoff model? Part 2—Towards a simple and efficient potential evapotranspiration model for rainfall-runoff modeling. *Journal of Hydrology* **303**, 290–306.
- Oudin, L., Michel, C. & Anctil, F. 2005b Which potential evapotranspiration input for a lumped rainfall-runoff model? Part 1—Can rainfall-runoff models effectively handle detailed potential evapotranspiration inputs? *Journal of Hydrology* **303**, 275–289.
- Penman, H. L. 1948 Natural evaporation from open water, bare soil and grass. *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences* **193** (1032), 120–145.
- Priestley, C. H. B. & Taylor, R. J. 1972 On the assessment of surface heat flux and evaporation using large-scale parameters. *Monthly Weather Review* **100** (2), 81–92.

- Rahimikhoob, A., Behbahani, M. R. & Fakheri, J. 2012 An evaluation of four reference evapotranspiration models in a subtropical climate. *Water Resources Management* **26** (10), 2867–2881.
- Rana, G. & Katerji, N. 2000 Measurement and estimation of actual evapotranspiration in the field under Mediterranean climate: a review. *European Journal of Agronomy* **13**, 125–153.
- Riekerk, H. 1985 Lysimetric evaluation of pine forest evapotranspiration for water balances. In: *The Forest – Atmosphere Interaction* (B. A. Hutchison & B. B. Hicks, eds). Reidel Publishing Co., Boston, MA, pp. 293–308.
- Rosenberg, N., Blad, B. & Verma, S. 1983 *Microclimate: The Biological Environment*, Wiley, New York, p. 495.
- Rosenberry, D. O., Stannard, D. I., Winter, T. C. & Martinez, M. L. 2004 Comparison of 13 equations for determining evapotranspiration from a prairie wetland, Cottonwood Lake area, North Dakota, USA. *Wetlands* **24** (3), 483–497.
- Rosenberry, D. O., Winter, T. C., Buso, D. C. & Likens, G. E. 2007 Comparison of 15 evaporation methods applied to a small mountain lake in the northeastern USA. *Journal of Hydrology* **340**, 149–166.
- Samaras, D. A., Reif, A. & Theodoropoulos, K. 2014 Evaluation of radiation-based reference evapotranspiration models under different Mediterranean climates in Central Greece. *Water Resources Management* **28**, 207–225.
- Scholl, D. G. 1976 Soil moisture flux and evaporation determined from soil hydraulic properties in a chaparral stand. *Soil Science Society of America Journal* **40** (1), 14–18.
- Siegert, E. & Schrodter, H. 1975 Erfahrungen mit dem Wasserbilanzschreiber nach Klausung (Experiences on the water balance recorder according to Klausung). *Deutsche Gewasserkundliche Mitteilungen* **19**, 167–171.
- Spittlehouse, D. L. & Black, T. A. 1980 Evaluation of the Bowen ratio/energy balance method for determining forest evapotranspiration. *Atmosphere-Ocean* **18** (2), 98–116.
- Stannard, D. I. 1993 Comparison of Penman-Monteith, Shuttleworth-Wallace, and modified Priestley-Taylor evapotranspiration models for wildland vegetation in semiarid rangeland. *Water Resources Research* **29** (5), 1379–1392.
- Sun, G., Zhou, G., Zhang, Z., Wei, X., McNulty, S. G. & Vose, J. M. 2006 Potential water yield reduction due to reforestation across China. *Journal of Hydrology* **328**, 548–558.
- Tabari, H., Grismer, M. E. & Trajkovic, S. 2011 Comparative analysis of 31 reference evapotranspiration methods under humid conditions. *Irrigation Science* **31** (2), 107–117.
- Trajkovic, S. & Kolakovic, S. 2009 Evaluation of reference evapotranspiration equations under humid conditions. *Water Resources Management* **23**, 3057–3067.
- Turc, L. 1961 Estimation of irrigation water requirements, potential evapotranspiration: a simple climatic formula evolved up to date. *Annals of Agronomy* **12**, 13–49.
- Valiantzas, J. D. 2013 Simple ET<sub>0</sub> forms of Penman's equation without wind and/or humidity data. II: comparisons with reduced set-FAO and other methodologies. *Journal of Irrigation and Drainage Engineering* **139** (1), 9–19.
- Valipour, M. 2014a Analysis of potential evapotranspiration using limited weather data. *Applied Water Science*, doi:10.1007/s13201-014-0234-2.
- Valipour, M. 2014b Application of new mass transfer formulae for computation of evapotranspiration. *Journal of Applied Water Engineering and Research* **2** (1), 33–46.
- Valipour, M. 2014c Use of average data of 181 synoptic stations for estimation of reference crop evapotranspiration by temperature-based methods. *Water Resources Management* **28** (12), 4237–4255.
- Valipour, M. 2015a Temperature analysis of reference evapotranspiration models. *Meteorological Applications* **22**, 385–394.
- Valipour, M. 2015b Investigation of Valiantzas' evapotranspiration equation in Iran. *Theoretical and Applied Climatology* **121** (1), 267–278.
- Valipour, M. 2015c Calibration of mass transfer-based models to predict reference crop evapotranspiration. *Applied Water Science*, doi:10.1007/s13201-015-0274-2.
- Valipour, M. & Eslamian, S. 2014 Analysis of potential evapotranspiration using 11 modified temperature-based models. *International Journal of Hydrology Science and Technology (IJHST)* **4** (3), 192–207.
- Vörösmarty, C. J., Federer, C. A. & Schloss, A. L. 1998 Potential evaporation functions compared on US watersheds: possible implications for global-scale water balance and terrestrial ecosystem modeling. *Journal of Hydrology* **207**, 147–169.
- Willmott, C. J. 1982 Some comments on the evaluation of model performance. *Bulletin American Meteorological Society* **63** (11), 1309–1313.
- WMO 1966 Measurement and estimation of evaporation and evapotranspiration. Technical Note No. 83, Geneva (WMO), *Quarterly Journal of the Royal Meteorological Society* **95**, 444.
- Xu, C. Y. & Singh, V. P. 2002 Cross comparison of empirical equations for calculating potential evapotranspiration with data from Switzerland. *Water Resources Management* **16** (3), 197–219.
- Xu, J., Peng, S., Ding, J., Wei, Q. & Yu, Y. 2013 Evaluation and calibration of simple methods for daily reference evapotranspiration estimation in humid East China. *Archives of Agronomy and Soil Science* **59** (6), 845–858.
- Xystrakis, F. & Matzarakis, A. 2011 Evaluation of 13 empirical reference potential evapotranspiration equations on the island of Crete in Southern Greece. *Journal of Irrigation and Drainage Engineering* **137** (4), 211–222.
- Zhang, L., Dawes, W. R. & Walker, G. R. 2001 Response of mean annual evapotranspiration to vegetation changes at catchment scale. *Water Resources Research* **37** (3), 701–708.

First received 14 September 2016; accepted in revised form 7 March 2017. Available online 22 April 2017