

Lessons From Pediatric HIV: A Case for Curative Intent in Pediatric Cancer in LMICs

Henry Miller, BA,^a Jeremy S. Slone, MD, MPH,^{b,c} Eric Raabe, MD, PhD,^d Nader Kim El-Mallawany, MD,^{b,c} Parth Mehta, MD,^{b,c} B. Ryan Phelps, MD, MPH^a

Antiretroviral medications (ARVs) are now being provided to nearly 50% of children living with HIV (CLHIV) in low- and middle-income countries (LMICs), an increase of 30% in the past 5 years.¹ The provision of ARVs in LMICs demonstrates that effective therapy can be delivered to children with complex medical issues. An estimated 96% of CLHIV and 80% of children <15 years of age living with cancer reside in LMICs, where 94 900 pediatric AIDS deaths and 120 000 pediatric cancer deaths occur annually.^{2,3} The health system infrastructure gains in the fight against HIV and other communicable diseases represent a promising platform for long-overdue, transformational cancer care in LMICs.

Similar to combination antiretroviral therapy in the context of pediatric HIV, the advent of combination chemotherapy in the mid-1960s revolutionized pediatric cancer treatment. Before chemotherapy, pediatric cancer was uniformly fatal, and care was focused on palliation. By the 1970s, over half of children diagnosed with cancer in high-income countries (HICs) were cured. Today ~4 out of 5 survive.⁴

In LMICs, however, the story is different. An estimated 215 000 children are diagnosed with cancer

globally each year, and fewer than half of these children currently survive.⁵ Many countries do not capture pediatric cancer diagnoses or related mortality, meaning that even the current numbers may reflect an underestimate of the total burden of pediatric cancer in LMICs.⁶ The reasons behind the lack of progress in treating pediatric cancers in LMICs are due in part to beliefs that chemotherapy is too expensive, that the disease burden is too low to warrant investment, and that pediatric cancer care is too complex.⁶

Although the costs of treating childhood cancer vary by cancer type, treating certain common pediatric malignancies, such as acute lymphoblastic leukemia (ALL), is relatively affordable. A 2016 systematic review of costs for ALL treatment revealed differences among LMICs, showing a range of \$4400 to \$6100 spent. The variation was largely due to the costs of treating complications rather than to drug expenses. Significant differences in regimens and drug costs exist among LMICs, making direct comparisons difficult.⁷ Successful Burkitt lymphoma (BL) therapy in Malawi, for instance, costs <\$50.⁶ These examples suggest that curative therapy for ALL and BL is similar in cost to suppressive lifetime



NIH

^aOffice of HIV/AIDS, United States Agency for International Development, Washington, DC; ^bDepartment of Pediatrics, Baylor College of Medicine, Houston, Texas; ^cTexas Children's Cancer and Hematology Centers, Houston, Texas; and ^dDivision of Pediatric Oncology, Johns Hopkins University School of Medicine, Baltimore, Maryland

The views and opinions expressed in this article are solely those of the authors and do not necessarily reflect those of the US Agency for International Development, Baylor College of Medicine, Texas Children's Hospital, or Johns Hopkins University School of Medicine.

Drs El-Mallawany, Mehta, Phelps, Raabe, and Slone conceptualized the commentary, provided initial data analysis, and drafted the initial manuscript; Mr Miller collected data, assisted with analysis, and reviewed and critically revised the manuscript; and all authors approved the final manuscript as submitted.

DOI: <https://doi.org/10.1542/peds.2017-0525>

Accepted for publication Apr 20, 2017

Address correspondence to Henry Miller, Office of HIV/AIDS, United States Agency for International Development, 1300 Pennsylvania Ave, NW, Washington, DC 20523. E-mail: hmliller@usaid.gov

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2017 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

To cite: Miller H, Slone JS, Raabe E, et al. Lessons From Pediatric HIV: A Case for Curative Intent in Pediatric Cancer in LMICs. *Pediatrics*. 2017;140(4):e20170525

antiretroviral therapy for CLHIV and can be accomplished in LMICs.^{6,8}

We propose that health programs adopt established, successful, cost-effective treatment strategies to deliver high-quality, curative therapy to all children with cancer in LMICs. Expansion of financially reasonable, LMIC-appropriate regimens now provides a backbone on which improvements can be made and allows for extension of chemotherapy to other cancer types.

Government and nongovernment stakeholders should work together to ensure adequate access to essential pediatric chemotherapy agents. As was true with pediatric ARVs in the early 2000s,¹ expanding access to chemotherapy agents is crucial. Furthermore, to ensure stable access to these medicines, appropriate forecasting, procurement, and delivery systems must be in place.^{6,9}

To properly institute a treatment program, a scale-up of professional training is necessary. The African Pediatric Fellowship Program based at the University of Cape Town and the Red Cross War Memorial Children's Hospital trains physicians from across Africa in pediatric oncology. In Uganda and Rwanda, new cancer facilities have been established, with more space for patients and research. These facilities have also allowed for new training programs, which have begun

educating physicians and nurses.⁹ Twinning programs between HIC hospitals and several countries (Uganda, Bangladesh, Costa Rica) have furthered improvements in pediatric oncology service delivery. These collaborations have not only developed training for oncologists but have also provided an avenue for knowledge-sharing, including internet-based learning.⁹

Pediatric cancer programs in LMICs should build on the strengths of existing health systems, such as the pediatric HIV treatment platform. Baylor College of Medicine and Texas Children's Hospital, through the Baylor International Pediatric AIDS Initiative, have established comprehensive pediatric HIV treatment centers throughout southern Africa. In Botswana, pediatric hematology-oncology services are now an extension of comprehensive pediatric HIV programs based at tertiary referral hospitals. This program provides direct patient care while training local health care workers and conducting clinical research.¹⁰ In Rwanda, investment in the health system to combat HIV/AIDS has strengthened cancer management, including the treatment of childhood malignancies in rural areas.¹¹ In each positive example, a shared commitment among governments and other stakeholders, domestic and international, to improve both the

access to and the quality of cancer care has led to progress despite many challenges.

The provision of ARVs to pediatric patients worldwide demonstrates that the treatment of children with complex disease processes can occur in challenging settings. Widespread adoption of established, low-cost, curative treatment regimens for high-burden malignancies is a reasonable first step for many LMICs. Even a relatively conservative 60% ALL and 50% BL cure rate would mean 48 000 child lives saved annually.⁶ If an 80% survival rate were achieved across all diagnoses, as occurs in HIC settings,³ 172 000 children could achieve life-long cures each year. The ultimate goal is to provide comprehensive curative therapy, equivalent to standards in HICs, to all children with cancer throughout the world, following the successful model of pediatric HIV treatment now in place in many LMICs.

ABBREVIATIONS

ALL: acute lymphoblastic leukemia

ARV: antiretroviral medication

BL: Burkitt lymphoma

CLHIV: children living with HIV

HIC: high-income country

LMIC: low- and middle-income country

FUNDING: This article was supported by a National Cancer Institute core grant to the Johns Hopkins Sidney Kimmel Comprehensive Cancer Center (P30 CA006973). Funded by the National Institutes of Health (NIH).

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

REFERENCES

- UNAIDS. Number of people receiving ART, children (0-14). AIDSinfo. Available at: <http://aidsinfo.unaids.org/>. Accessed March 29, 2016
- UNAIDS. Number of AIDS-related deaths, children (0-14). AIDSinfo. Available at: <http://aidsinfo.unaids.org/>. Accessed March 29, 2016
- Kellie SJ, Howard SC. Global child health priorities: what role for paediatric oncologists? *Eur J Cancer*. 2008;44(16):2388–2396
- Robison LL, Armstrong GT, Boice JD, et al. The Childhood Cancer Survivor Study: a National Cancer Institute-supported resource for outcome and intervention research. *J Clin Oncol*. 2009;27(14):2308–2318
- International Agency for Research on Cancer. International Childhood Cancer Day: much remains to be done to fight childhood cancer [press release]. Geneva, Switzerland: World Health Organization; February 15, 2016

6. Gupta S, Howard SC, Hunger SP, et al. Treating childhood cancer in low- and middle-income countries. In: Gelband H, Jha P, Sankaranarayanan R, Gauvreau CL, Horton S, eds. *Cancer: Disease Control Priorities*. Vol 3. 3rd ed. Washington, DC: The International Bank for Reconstruction and Development/The World Bank; 2015
7. Hayati H, Kebriaeezadeh A, Ehsani M, et al. Treatment costs for pediatrics acute lymphoblastic leukemia; comparing clinical expenditures in developed and developing countries: a review article. *Int J Pediatr*. 2016;4(12):4033–4041
8. Wandeler G, Johnson LF, Egger M. Trends in life expectancy of HIV-positive adults on antiretroviral therapy across the globe: comparisons with general population. *Curr Opin HIV AIDS*. 2016;11(5):492–500
9. Binağwaho A, Wagner CM, Farmer PE. A vision for global cancer medicine: pursuing the equity of chance. *J Clin Oncol*. 2016;34(1):3–5
10. Slone JS, Agrawal AK, Mehta PS, et al. Reply to “childhood cancer in Africa”. *Pediatr Blood Cancer*. 2014;61(8):1524
11. Stulac S, Mark Munyaneza RB, Chai J, et al. Initiating childhood cancer treatment in rural Rwanda: a partnership-based approach. *Pediatr Blood Cancer*. 2016;63(5):813–817