

# Molecular Regulation of Fas Expression in $\beta$ -Cells

Martine I. Darville, Dongbo Liu, Meng-Chi Chen, and Décio L. Eizirik

**F**as-mediated cell death may play an important role in the autoimmune destruction of  $\beta$ -cells in type 1 diabetes.  $\beta$ -Cells usually do not express Fas, but Fas expression is detected in  $\beta$ -cells from patients with recent-onset type 1 diabetes (1) and in cytokine-exposed mouse and human islets. Fas expression renders the  $\beta$ -cells susceptible to Fas ligand-induced apoptosis (2–4). It has been proposed that cytokine-induced Fas expression in human islets is mediated by nitric oxide (NO) production (4).

The aim of the present study was to investigate Fas regulation by cytokines in human, mouse, and rat  $\beta$ -cells. For this purpose, we analyzed Fas mRNA expression by reverse transcriptase–polymerase chain reaction in isolated human islets and fluorescence-activated cell sorter (FACS)-purified rat  $\beta$ -cells exposed to cytokines with or without  $N^G$ -monomethyl-L-arginine (L-MA), an inhibitor of inducible NO synthase (iNOS). Fas mRNA expression was also determined in islets isolated from wild-type and iNOS knockout mice. (A complete version of the present study will be published elsewhere [5].) In the second part of the study, regulation of Fas promoter activity was determined in FACS-purified rat  $\beta$ -cells.

In human islets, Fas mRNA expression was induced by interferon- $\gamma$  (IFN- $\gamma$ ), a cytokine that did not induce iNOS expression. A combination of IFN- $\gamma$  and interleukin-1 $\beta$  (IL-1 $\beta$ ) induced both iNOS and Fas expression, but L-MA did not prevent Fas mRNA expression. In purified rat  $\beta$ -cells, Fas

expression was induced by IL-1 $\beta$  alone and was not prevented by L-MA. Islets isolated from wild-type or iNOS $^{-/-}$  mice similarly expressed Fas mRNA after exposure to a mixture of three cytokines (IL-1 $\beta$  + IFN- $\gamma$  + tumor necrosis factor- $\alpha$ ). These results indicate that cytokine induction of Fas expression is NO-independent in human, mouse, and rat pancreatic cells.

Transfection of rat Fas promoter–luciferase reporter constructs into purified rat  $\beta$ -cells showed that IL-1 $\beta$  induction of promoter activity is mediated by a region between nucleotides –223 and –54 that contains putative binding sites for the transcription factors NF- $\kappa$ B, CCAAT/enhancer binding protein, and Ets. The individual contributions of these transcription factors for Fas expression is presently under study.

## REFERENCES

1. Moriwaki M, Itoh N, Miyagawa J, Yamamoto K, Imagawa A, Yamagata K, Iwahashi H, Nakajima H, Namba M, Nagata S, Hanafusa T, Matsuzawa Y: Fas and Fas ligand expression in inflamed islets in pancreas sections of patients with recent-onset type I diabetes mellitus. *Diabetologia* 42:1332–1340, 1999
2. Yamada K, Takane-Gyotoku N, Yuan X, Ichikawa F, Inada C, Nonaka K: Mouse islet cell lysis mediated by IL-1-induced Fas. *Diabetologia* 39:1306–1312, 1996
3. Zumsteg U, Frigerio S, Holländer GA: Nitric oxide production and Fas surface expression mediate two independent pathways of cytokine-induced murine  $\beta$ -cell damage. *Diabetes* 49:39–47, 2000
4. Stassi G, De Maria R, Trucco G, Rudert W, Testi R, Galluzzo A, Giordano C, Trucco M: Nitric oxide primes pancreatic beta cells for Fas-mediated destruction in insulin-dependent diabetes mellitus. *J Exp Med* 186:1193–1200, 1997
5. Liu D, Pavlovic D, Chen M-C, Flodström M, Sandler S, Eizirik DL: Cytokines induce apoptosis in  $\beta$ -cells isolated from mice lacking the inducible isoform of nitric oxide synthase (iNOS $^{-/-}$ ). *Diabetes* 49:1116–1123, 2000

From the Diabetes Research Center, Vrije Universiteit, Brussels, Belgium.

Address correspondence and reprint requests to Martine I. Darville, Diabetes Research Center, Vrije Universiteit Brussel, Laarbeeklaan 103, B-1090-Brussels, Belgium. E-mail: mdarv@mebo.vub.ac.be.

Received for publication 27 May 2000 and accepted 19 June 2000.

This article is based on a presentation at a symposium. The symposium and the publication of this article were made possible by an unrestricted educational grant from Les Laboratoires Servier.

FACS, fluorescence-activated cell sorter; IL-1 $\beta$ , interleukin-1 $\beta$ ; INF- $\gamma$ , interferon- $\gamma$ ; iNOS, inducible NO synthase; L-MA,  $N^G$ -monomethyl-L-arginine; NO, nitric oxide.