

Index

- Argillaceous Saluda rocks, 1, 6–8, 11, 14,
19, 20, 22, 23, 24
- Australia, 13, 15
- Bahamas, 17, 18
- Bardstown, Kentucky, 7, 8
- Barriers, 1, 13, 14, 17, 18, 20, 21, 24, 25
coralline, 1, 13, 14, 17, 18, 20, 21, 24, 25
environmental isolation by, 13, 14, 17,
18, 20, 24, 25
- Bedding, 1, 7, 9–14, 16, 18–20, 23
- Bonaire, Netherlands Antilles, 14, 17
- Brachiopods, 9, 11, 23
- Brassfield Limestone, 6
- Brine, magnesium-rich, 17
- Bryozoans, 9, 23
- Burrows, 7–11, 18–20, 22, 23
- Calcite/dolomite ratio, 1, 6–9, 12, 15, 16,
19, 24
- Calcitic dolomite, 1, 6–8, 12, 24
- Carbonate rocks, classification of, 6
- Chattanooga Shale, 6
- Cincinnatian Series, 1, 3
- Clay shale, 1, 7, 11, 23, 24
- Conodonts, 23
- Coorong lagoon, 15–17, 22
- Coralline zone, 1, 10–14, 16–21, 24, 25
consolidation of, 14, 18, 20
profusion of corals in, 10, 20, 25
thickness of, 10, 13, 20
- Corals, 1, 3, 10, 11, 13, 14, 17, 18, 20, 21,
24, 25
growth form, 10, 20
orientation, 1, 10, 18, 20
paleoecology, 20, 21
- Depth of water, 1, 12, 13, 16, 20
- Desiccation cracks, 1, 8, 12, 13, 16, 18
- Diagenesis, 1, 14, 15, 17–19, 24, 25
- Dillsboro Formation, 4
- Dolomite, 1, 5–9, 11–19, 24, 25
penecontemporaneous, 17–19, 24, 25
primary, 1, 14–19, 24
rhombohedra, 7, 8, 11, 17
- Dolomitic micritic limestone, 6–10, 12, 13,
24
- Elkhorn Formation, 4
- Environment of deposition, 1, 3, 4, 12–25
- Evaporation, 1, 13, 14, 16, 17, 19, 24
- Exposures, 3, 4, 6, 8–10
distribution, 4
typical, 3, 4
- Favistella*, 3, 10, 20, 21
- Florida Keys, 18
- Fossils, paucity of, 1, 6–9, 12, 14, 15, 18, 19,
21–24
- Franklin County, Indiana, 4, 6
- Great Inagua, Bahamas, 17
- Hypersaline lakes, 15–17
- Jefferson County, Indiana, 3, 8, 10
- Jefferson County, Kentucky, 6, 10
- Laminae, 1, 7, 10, 12–14, 16, 18–20, 23
- Lebanon Group, 3
- Leperditella*, 9, 21, 22
- Leperditia*, 9, 21, 22
- Liberty Formation, 3
- Madison Beds, 3
- Madison, Indiana, 6, 10
- Manlius Formation, 18
- Marion County, Kentucky, 4, 6
- Maysville Group, 4
- Micrite, 1, 6–13, 19, 20, 23, 24
- Micritic limestone, 9
- Micritic skeletal limestone, 9, 11, 12, 19,
20, 23
- Ostracodes, 8, 9, 11, 21, 22
- Paleoecology, 20–23
- Penesaline lagoons, modern, 13–17, 22
- Penesalinity, 1, 13–19, 22–25
- Persian Gulf, 18
- Polychaetia, 22, 23
- Primitia*, 9, 21, 22
- Reefs, modern, 13, 21
- Restriction, environmental, 1, 13, 14, 17,
18, 20, 21, 24, 25
- Richmond Group, 3, 4
- Ripple marks, 1, 8, 12, 13
modern, 12, 13
Saluda, 1, 8, 12, 13
- Salinity, 1, 13–25
- Saluda-Whitewater contact, 6, 7, 10, 21,
24
- Scolecodonts, 22, 23
- Sediment composition, 7–11
- Shale break, 1, 7–9, 19, 22, 24
- Silurian, 4, 6
- Skeletal micritic limestone, 9, 11, 12, 19,
23
- South Australia, 15
- Stratigraphic classification, 3, 4, 24
- Stromatoporoids, 1, 10
- Subaerial exposure, 1, 12, 16–19, 24
- Subsurface Saluda, 6, 8
- Tanners Creek Formation, 3, 4, 6, 10–12,
19, 24
- Tanners Creek–Saluda contact, 6, 10, 12,
24
- Terrigenous detritus, dearth of, 12, 14, 18,
20, 24
- Tetradium*, 10, 20, 21

Thickness and distribution, 6
Time transgression, 1, 7, 24
Turbulence, 1, 12, 13, 18–24
Versailles, Indiana, 8

Washington County, Indiana, 6
Waynesville Formation, 3
Whitewater Formation, 4, 6, 7, 10–12, 21,
24

